



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.


### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

582

C219

The Hopkins Library  
presented to the  
Leland Stanford Junior University  
by Timothy Hopkins.







**DE CANDOLLE**  

---

**PRODROMUS**  
**SYSTEMATIS NATURALIS**  
**REGNI VEGETABILIS**  
**PARS XVI**  
**SECTIO PRIOR**

---

PARISIIS. — TYPIS C. LAHURE  
via dicta de Fleurus, 9

---

**DE CANDOLLE**

---

**PRODROMUS**  
*811 J 977*  
**SYSTEMATIS NATURALIS  
REGNI VEGETABILIS**

**SIVE**

**ENUMERATIO CONTRACTA  
ORDINUM, GENERUM, SPECIERUMQUE PLANTARUM**

**HUC USQUE COGNITARUM, JUXTA METHODI NATURALIS NORMAS DIGESTA**

**EDITORE ET PRO PARTE**

**AUCTORE**

**ALPHONSO DE CANDOLLE**

**PARS DECIMA SEXTA**

**SECTIO PRIOR**

**Sistens Urticaceas, Piperaceas, etc.**

---

**PARISIIS**

**SUMPTIBUS VICTORIS MASSON ET FILII**

**FORO DICTO DE L'ÉCOLE DE MÉDECINE, N° 17**

**VENITQUE APUD EOSDEM LIPSIE, PROCURANTE F. WAGNER**

**OCT. M DCCC LXIX**



HB 618

## SIGNORUM PRODROMI EXPLICATIO.

- ⊙ Planta monocarpica duratione ignota.
- ④ Planta monocarpica annua.
- ② Planta monocarpica biennis.
- ✕ Planta rhizocarpica, seu perennis herbacea.
- ♂ Planta lignosa, cujus magnitudo est ignota.
- ♂ Suffrutex, nempe planta lignosa cujus altitudo ad summum bipedalis.
- ♂ Frutex, nempe planta lignosa, a basi ramosa, plus quam bipedalis.
- ♂ Arbuscula, nempe planta lignosa, trunco donata et infra 25 ped. altitudinis.
- ♂ Arbor, nempe lanta lignosa, trunco donata et 25 ped. superans.
- ⊂ Scandens.
- ✕ Forma hybrida.
- ♂ Planta seu flos masculus.
- ♀ Planta seu flos femineus.
- ♂♀ Planta seu flos hermaphroditus.
- ∞ Numerus indefinitus, sic ovula ∞, id est plurima non numero definito.
- ? Signum dubitationis.
- ! In synonymia aut loco natali indicat specimen ab auctore ipso aut viatore inscriptum.
- v. s. c. Visa sicca culta.
- v. s. sp. aut v. s. Visa sicca spontanea.
- v. v. sp. aut v. v. Visa viva spontanea.
- v. v. c. Visa viva culta.
- v. in h. Visa in herbario.

---

---

# PRODROMUS

## SYSTEMATIS UNIVERSALIS

### REGNI VEGETABILIS.

---

#### Ordo CLXXIX. DAPHNIPHYLLACEÆ (1).

Rhamnacearum genus dubium *Blume Bijdr. p. 1153; Endl. Gen. pl. n. 5755.* — Euphorbiacearum genus *Baill. Etud. gén. Euphorb. p. 564. t. 21. fig. 25-27.*

Flores unisexuales. Calycis laciniae æstivatione imbricativæ. Petala (in genere unico noto) nulla. Stamina in centro florum masculorum inserta. Ovarium superum, compositum, carpidiis 2 constructum (2-loculare). Loculi ovarii apice 2-ovulati. Ovula pendula, geminatim collateralia, anatropa; raphe interior st. angulum interiorem loculorum spectans; micropyle exterior et superior (omnino ut in Euphorbiaceis). Fructus syncarpeus, haud aperiens. Semina pendula, albuminosa. Albumen oleagineum. Embryo apicalis, i. e. in apice albuminis nidulans, albumine multoties brevior. Radicula superior. — Arbores orbis antiqui.

A proximis Euphorbiaceis et Buxaceis simul embryone exiguo in apice albuminis nidulante distinguuntur et a Buxaceis insuper directione contraria micropyles et raphes ovulorum diversæ sunt. — Ob situm staminum non perigynum, similiter ac Buxaceæ, in vicinitate Euphorbiacearum servandæ sunt et ad seriem ordinum Calyciflorarum, speciatim ad Illicineas, quibus characteribus prope accedunt, relegari nequeunt.

DAPHNIPHYLLUM *Bl. Bijdrag. p. 1153; Endl. Gen. plant. n. 5755; Baill. Etud. gén. Euphorb. p. 564. t. 21. fig. 25-27.* — *Goughia Wight Icon. plant. Ind. orient. t. 1877-1878; Benth. in Hook. Journ. of Bot. 1854. p. 8.* — *Gyandra Wall. Cat. (non Geisel.).*

Flores unisexuales. Laciniae calycinæ utriusque sexus æstivatione imbricativæ. Petala 0. Discus florum masc. 0, fœmineorum evolutus v. suppressus. Stamina in receptaculo convexo centralia; antheræ biri-

---

(1) Auctore J. Müller, Argoviensi.

mosæ; oculi longitrorsum adnati. Rudimentum ovarii 0. Ovarii liberi loculi biovulati. Ovula euphorbiacea, i. e. ex apice loculorum sub obturatore pendula, anatropa, raphe introrsa, micropyle exterior et superior. Fructus haud capsulari-aperiens. Semina ocarunculata, albuminosa. Embryo sub apice albuminis nidulans, exiguus, albumine multoties brevior. Radicula cotyledonibus semicylindricis multo longior easque latitudine æquans. — Arbores et frutices in Asia meridionali continentali, insulis sundaicis et in Japonia, una in Africa occidentali, crescentes. Rami teretes, densiuscule foliosi, glabri. Folia alterna, sæpissime stipulis destituta, petiolata, simplicia, penninervia, integra v. dentata, reticulato-venosa, tenuiter coriacea, plus minusve fusco-glaucrescentia et ea Lauracearum nonnullarum primo intuitu simulantia. Inflorescentiæ unisexuales, axillares. Flores dioici, racemosi. Pedicelli masc. mox articulado-decidui. Calyx utriusque sexus 3-8-partitus; laciniae florum masc. sæpius mox deciduæ, inciso-denticulatæ, femineorum marcescendo deciduæ aut persistentes. Glandulæ disci hypogyni 4-6, majusculæ, v. nullæ. Stamina 5-18; filamenta libera, e basi radiatim divergente adscendentia; antheræ ambitu variæ, semper tamen latiusculæ, nunc tetragonæ, nunc a dorso nunc a latere plus minusve compressæ. Ovarium 2-loculare. Stigmata 2, subsessilia, disciformi-reniformia. Fructus abortu monospermus, baccatus, drupam simulans; mesocarpium carnosum; endocarpium rigide chartaceum, vulgo rugoso-tuberculosum, intus album et plus minusve sericeo-fibrosus. Albumen olivaceo-nigrescens.

§ 1. *Antherarum rimæ introrsæ aut laterales. — Species asiaticæ.*

1° Calyx femineus deciduus (in nonnullis hucusque ignotus).

\* Antheræ a latere compressæ.

1. D. MAYUS (Müll. Arg. in Linnæa vol. 34. p. 76), foliis longo-petiolatis, racemis subcorymbiformibus, staminibus 8 a latere compressis intus acutis apice subincurvo breviter acuminatis. § In India orientali: martabanica prope Amherst (Wall. n. 7805) in hb. Wall. nunc in hb. Soc. Linn. londin., et in hb. DC.). Ricinus furfuraceus Wall. Cat. n. 7805 (ubi cum fructibus ex parte putredine corruptis, furfuraceis, Macarangæ hypoleucæ mixtum, unde nomen Wall. sched.). Ramuli glauci, cum omnibus reliquis partibus glabri. Petioli 10-15 cm. longi. Limbus foliorum 18-28 cm. longus, 8-14 cm. latus, oblongato-ellipticus, acutus v. breviter acuminatus, basi acutiusculus, rigidulus, utraque pagina tenuiter quidem sed prominenter laxè reticulato-venosus, supra viridis, subtus intense albido-cassius. Racemi masc. 2½-3½ cm. longi. Pedicelli masc. 12-16 mm. æquantas. Flores masc. aperti 3 mm. lati. — Flores fem. et fructus ignoti. (v. s.)
2. D. ZOLLINGERI, foliis mediocriter petiolatis coriaceo-membranaceis, racemis utriusque sexus subfasciculifloris gracilibus, rhachi tenella, calycibus utriusque sexus deciduis, staminibus circ. 10, antheris a latere compressis triente longioribus quam latis apiculatis, fructu ellipsoideo utrinque obtuso. § In insula Java (Zolling. ! absque n°). Rami subgraciles, obscure lurido-vides. Petioli 2-3 cm. longi, graciles, cum omnibus reliquis partibus glabri. Limbus foliorum 10-14 cm. longus, 2½-3 cm. latus, oblongo-lanceolatus, integer, prominenter dense reticulato-venosus, supra fusciscenti-lurido-viridis, subtus fuscidulo-glaucus. Racemi utriusque sexus circ. 5 cm. longi. Pedicelli fem. 7 mm. longi, fructigeri et masculi ½ cm. subæquantas, utriusque sexus filiformes. Antheræ 1½ mm. longæ. Stigmata ovario glauco-pruinoso bis longiora, erecto-patula. Fructus 9 mm. longus, 7 mm. latus, siccus undulato-subrugulosus. (v. s.)

**\*\* Antheræ tetragonæ, nec a dorso nec a latere distincte compressæ (in *D. Teysmanni* ignotæ).**

3. *D. BLUMEANUM*, foliis subcoriaceis subtus prominenter reticulato-venosis, antheris 7-9 tetragono-cylindricis duplo longioribus quam latis parvulis. § In insula Java (Zolling. n. 2140! in hb. Mus. paris.). *D. Blumeanum* Baill. Etud. gén. Euphorb. p. 565 (nomen). Totum glabrum. Petioli 1½ cm. longi. Limbus foliorum 9-11 cm. longus, sæpius 2½-3 cm. latus, anguste lanceolatus, longe acuminatus, basi acutus, margine undulatus, florendi tempore subtus prominenter reticulatus. Racemi masc. breves, congesti. Flores breviter pedicellati. Calyces masc. 6-fidi; laciniae breves, dentatae. — Primo intuitu quibusdam speciminibus *D. glaucescentis* simile, sed folia longius acuminata et antheræ non a dorso compressæ. (v. s.)
4. *D. BÄNGERI*, foliis tenuiter coriaceis, antheris 5-8 subduplo longioribus quam latis nec a dorso nec a latere compressis sed latere dorsali paulo latioribus basi leviter angustatis minutissime papilloso-tuberculatis. § In Japonia (Burger! in hb. lugdun. batav.). Foliorum limbus obovato-lanceolatus, breviter acuminatus. Racemi masc. foliis multoties breviores. Pedicelli antheras casu calycis masc. denudatas circiter bis æquantes. — Specimina masc. perfecte similia iis *D. glaucescentis*, sed antheris minoribus statim dignoscuntur. Antheræ magis opacæ, haud læves, non a dorso compressæ, sectione transversali rhomboideæ, extus paulo latiores, intus distincte angustiores, cæterum fere tetragonæ. A *D. Teysmanni* etiam in Japonia lecto, cuius specimina mascula hucusque ignota, comparatis speciminibus differt consistentia et colore foliorum, petiolis paulo brevioribus et crassioribus. (v. s.)
5. *D. TEYSMANNI* (Sulp. Kurz in Teysm. et Binnend. Plant. nov. v. minus cogn. hort. bogor. p. 37), foliis coriaceis, racemis fœm. brevibus paucifloris apice umbellatam florigeris. § In Java, ex Japonia allata (Zolling. n. 1029. 21). Ramuli validi, dense foliosi, fusco-nigricantes. Petioli 1-1½ v. rarius usque 2 cm. longi. Limbus foliorum 7-9 cm. longus, 2-3 cm. latus, lanceolatus, acuminatus, basi cuneato-angustatus, margine recurvus, utraque pagina prominenter laxe reticulato-venosus, supra nitidus, lurido-viridis, subtus cassio-glaucus. Racemi fœminæ in ramis biennibus siti, vix 2 cm. superantes. Pedicelli fœm. 5-7 mm. longi. Calycis fœm. laciniae ovatae, subirregulares. Disci hypogyni glandulæ conico-ovoideæ, calycem hinc inde excedentes. — A *D. glaucescente* differt foliis coriaceis, subtus cassio-glaucis, ovario et stylis subduplo brevioribus. (v. s.)

**\*\*\* Antheræ a dorso compressæ.**

6. *D. GLAUDESCENS* (Bl. Bijdr. p. 1153), foliis vix prominenter reticulato-venosis, calyce fœm. exiguo, antheris 7-10 a dorso compressis longitudine latitudinem haud superantibus caeva brevioribus. § v. s. In montosis insulæ Javæ (Bl. hb. lugd. bat.), in montibus Neilgherries Indiæ orientalis (Perrott. n. 29!), in Malabar (Storks et Law in Hook. et Thoms. hb. plant. Ind. orient.), in Loo-Choo (Wright n. 275!), Hongkong (hb. Hook.), et in Corea, ubi arbor magna (Wilford n. 644! in hb. Hook. et in hb. berol.). — Miq. Flor. Ind. bat. vol. 1. part. 2. p. 431 et Supplem. Sumatra p. 472. *D. Roxburghii* Baill. Etud. gén. Euphorb. p. 565, exclus. syn. Roxb. *Goughia Neilgherrensis* R. Wight Icon. plant. Ind. orient. t. 1878-1879 (antheræ quam in natura ambitu longiores). Arbor sæpius 20-30-pedalis. Truncus diametro sesquipedalis. Petioli in planta fœm. 1½-2 cm. longi, in planta mascula 2-4 cm. æquantes. Limbus foliorum sæpius 8-10 cm. longus, 3-4 cm. latus, oblongo-obovatus v. obovato-lanceolatus, apice nunc retundato-obtusius, nunc obtusiusculus, nunc acutus, basi subacutus, margine integro recurvus, subtus glaucus. Racemi laxiflori, petiolos subæquantes v. superantes. Pedicelli 4-5 mm. longi, validiusculi, antheras circiter ter æquantes. Laciniae calycis masc. lanceolatae, antheris paulo longiores. Antheræ 1½ mm. longæ, filamentis longiores. Fructus ellipsoidei, utrinque obtusi, 9-11 mm. longi, sicci 7-8 mm. lati. Endocarpium intus pallide sericeum. — *Phacourtia inermis* Roxb. Coromand. vol. 3. t. 222

a cl. Baill. l. c. hic relata non est hujus generis. — *Ludit concolor*, foliis subconcoloribus obscurius fuscis nitidulis subtus paulo pallidioribus nec gâucis utraque pagina distinctius reticulato-venosis. 5 In Ceylonia, altitudine 4000-6000 ped. (Thwait. n. 491!). D. Neilgherrense Thwait. Enum. pl. Zeyl. p. 290, exclus. syn. R. Wight. (v. s.)

7. D. HIMALAYENSE, foliis supra impresso-reticulato-venosis, floribus masc. 5-andris, antheris magnis oblongatis a dorso compressis, fructibus ellipsoideis, 5 In Indiæ orientalis regione temperata, in montibus Khasiæ altitud. 5000 ped. (Hook. f. et Thoms.!) et in Sikkim altit. 8-10 000 ped. (Hook. f. et Thoms.!). Goughia Himalensis Benth. in Hook. Journ. of Bot. 1854. p. 9. Daphniphyllum Benthami Baill. Etud. gén. Euphorb. p. 565 (nomen). Ramuli florigeri crassi, sæpius 5-7 mm. diametro æquantes. Petioli 2-2½ cm. longi. Limbus foliorum 10-15 cm. longus et 2½-3½ cm. latus, lanceolatus, v. lanceolato-spathulatus, basi apiceque acutus, integer, glaber, subtus fusco-glaucescens v. plus minusve concolor, nunc prominenter, nunc obsoletius reticulato-venosus. Racemi circ. 4½ cm. longi, angusti, multiflori. Pedicelli masc. antheras paulo superantes, mox decidui. Antheræ 4½ mm. longæ, fusco-nigricantes. Fructus siccati submaturi 11 mm. longi, utrinque rotundato-obtus. (v. s.)

2° Calyx foemineus persistens.

\* Baccæ tuberculato-asperæ.

8. D. LAURINUM (Baill. Etud. gén. Euphorb. p. 565. t. 21. fig. 25-27), foliis coriaceo-membranceis utraque pagina demum prominenter reticulato-venosis, calycis fœm. laciniiis denticulatis, floribus masc. decandris, antheris a latere compressis apice incurvis, fructibus ellipsoideis utrinque acutiusculis. 5 In Singapore (Wall. n. 8020!), in Malacca (Griffith! in hb. Hook.) et in Sumatra. Goughia laurina Benth. in Hook. Journ. of Bot. 1854. p. 9. Gyrandra laurina Wall. Cat. n. 8020 (nomen). Rami subtenués, florigeri diametro circ. 3-5 mm. æquantes, dense foliosi. Petioli 6-8 cm. longi, graciles. Limbus foliorum 15-20 cm. longus, 4-8 cm. latus, ovato-lanceolatus v. oblongo-ovatus v. oblongo-obovatus, basi apiceque acutus, glaber. Racemi laxiflori, petiolos paulo superantes. Pedicelli utriusque sexus 1½-2 cm. longi, masculi antheras pluries superantes. Calycis fœm. lacinia inæquales, persi-tentes, inciso-denticulatæ. Antheræ 1½ mm. longæ. Baccæ ut in congeneribus drupiformes, 1 cm. longæ, 6 mm. crassæ. — Folia occurrunt nunc fusciscentia, nunc glaucescentia. — A proximo D. calycino, cujus specimina masc. ignota, distinguitur foliis majoribus, paulo longius petiolatis et præsertim baccis lævioribus utrinque acutiusculis nec rotundato-obtus. (v. s.)
9. D. CALYCINUM (Benth. Flora hongkong. p. 316), foliis coriaceo-membranceis subtus vix prominenter venosis, calyce fœm. 4-partito, fructibus oblongo-ellipsoideis utrinque rotundato-obtusis sparse tuberculato-asperis. 5 In China circa Cantonem (Park! in hb. DC.), ibidem in Hongkong et prope Macao (Gaudich. n. 275! Champion n. 184! Calléry n. 77!). D. Gaudichaudianum Baill. Etud. gén. Euphorb. p. 565 (nomen). D. Calleryanum Baill. Etud. gén. Euphorb. p. 565 (nomen). Petioli 4-6 cm. longi. Limbus foliorum 9-12 cm. longus, 4-5 cm. latus, oblongo-obovatus v. oblongo-ellipticus, utrinque subobtus, subtus glaucescens. Racemi fructigeri 3-4 cm. longi, numerosi. Pedicelli fœm. fere 1 cm. æquantes. Lacinia calycis fructigeri late ovata, obtusæ, 2 mm. longæ. Baccæ maturæ 8 mm. longæ, 5½ mm. latæ, longiusculo tractu æquilatæ, plus minusve glauco-pruinosa. — Ad D. laurinum accedit sed fructus utrinque rotundato-obtus. A D. glaucescente, cui etiam simile, recedit calyce fœm. persistente. (v. s.)
10. D. BANCANUM (Sulp. Kurz in Teysm. et Binnend. Plant. nov. v. minus cogn. hort. bogor. p. 37), foliis coriaceo-chartaceis tenuiter venosis, calyce fœm. breviter lateque trilobo, antheris a latere compressis, fructibus oblongo-ellipsoideis utrinque obtusis. 5 In insula javanica Bangka (Teysm. qui nuper benevole misit, hb. DC.). Arbor 3-metralis. Stipulæ ad apicem axium lineari-lanceolatæ, deciduæ. Folia subverticillatim approximata. Petioli 2½-5½ cm. longi. Limbus foliorum 14-18 cm. longus, 3½-7 cm. latus



vulgo anguste lanceolatus, longiuscule acuminatus, basi rotundatus v. breviter acutatus, margine leviter revolutus, integer, subcor. color, viridis, nitidus. Racemi simplices, foeminei solitarii, masculi fasciculati. Pedicelli foem. 10-12 mm. longi, rigidi, masculi 1½-2 cm. æquantes, gracillimi. Calyx masc. vix 1 mm. longus, repando-lobatus v. subinde fere integer, foemineus masculo subbrevior, ovario arcte adpressus, sub fructu haud accrescens et distinctus late breviterque 3-lobus; lobi brevissime acutati. Antheræ similes iis D. laurini, sæpissime 10, circiter 1½ mm. longæ, subsessiles, erectæ, semilunatæ. Ovarium ellipsoideum, more congenerum 2-loculare, læve. Baccæ siccæ 8-9 mm. longæ, 5½-6 mm. latæ, similes iis D. calycini, atro-violaceæ. Embryo in albumine apicalis (nec basilaris ut a cl. auct. infauste dictum fuit), perexiguus, albidus; cotyledones visæ cœrulescenti-v. cœruleo-nervosæ. Radicula conica, acuta. (v. s.)

\*\* Fructus læves.

11. D. ACUTIFOLIUM, foliis coriaceo-membranaceis prominenter reticulato-venosis acutissime acuminatis, calycis foem. laciniis triangulari-lanceolatis subulato-acuminatis persistentibus, fructibus ellipsoideis utrinque obtusis lævibus. 5 In insula Java (hb. lugd. bat. a cl. Miq. benevole miss.). Ramuli validiusculi, dense v. apice densissime foliosi. Petioli 12-18 mm. longi, subtenues. Foliorum limbus 8-10 cm. longus, 2½-3 cm. latus, elliptico-lanceolatus, basi æquilaterali acutus, apice sensim acutissime acuminatus, margine leviuscule undulatus, fuscescens. Racemi foem. folia dimidia æquantes v. subæquantes, simplices, apice umbellatim florigeri, cæterum sparsiflori. Rhachis inferne longiuscule pedunculiformi-denudata, apice quasi in umbellam 4-6-floram desinens. Pedicelli foem. circ. 7-9 mm. longi, crassiusculi, patentes, cum omnibus reliquis partibus glabri. Fructus immaturi 8-10 mm. longi, circ. 6 mm. lati, siccando undulato-rugosi, haud tuberculato-asperi, plus minusve nigro-fusciscentes. Flores masc. ignoti. — Ab omnibus congeneribus differt apice foliorum acute subcuspidato-acuminato, cæterum simile D. calycino, sed calyce foemineo et fructibus diversum. (v. s.)

## § 2. *Antherarum rimæ extrorsæ. — Species africana.*

12. D. AFRICANUM (Müll. Arg. in Flora ratish. 1864. p. 536), foliis reticulato-venosis, antheris 9-18 a dorso leviter compressis subsessilibus late ellipsoideis rigidis. 5 In Africa occidentali æquatoriali ad River Kongin (G. Mann n. 1811 in hb. Hook.). Arbor 30-pedalis (Mann sched.). Ramuli teretes, cum tota planta glabri. Petioli 1½-2 cm. longi. Limbus foliorum 9-16 cm. longus, 4-7 cm. latus, lanceolato-ellipticus, breviter et obtuse cuspidato-acuminatus, basi acutiusculus, rigidulus. Racemi masc. 4-5 cm. longi, a basi æqualiter florigeri, circ. 12-15-flori, fusco-nigricantes. Bractæ late triangulari-ovatæ, acutæ, 1-floræ. Pedicelli masc. calycem æquantes. Calyx masc. apertus 5 mm. latus, basi planiusculus, 4-6-fidus, fuscus. Antheræ 1½ mm. longæ, rimis distincte extrorsis insignitæ, in receptaculo planiusculo sitæ. Flores foem. et fructus ignoti. (v. s.)

## *Species dubii loci.*

13. D. MACROPODUM (Miq. in Ann. Mus. bot. lugd. bat. 3. p. 129. 1867), foliis coriaceis, racemis fem. elongatis, fructu quam pedicellus brevior ellipsoideo. 5 In Japoniæ sylvis montanis ins. Kiusiu (Pierot, in hb. lugd. bat.). Ramuli validi, superne angulati et foliosi, cæterum teretiusculi, cortice atrofusco et cicatricibus petiolorum amplis obcordato-rotundatis pedunculorumque his arcte contiguis notati. Petioli vulgo 2-1½ poll. longi, antice clauso-canaliculati. Foliorum limbus 8-5 poll. longus, 2-3, vulgo 2½ latus, e basi acuta elliptico-oblongus v. sublanceolatus, apice abrupte brevissime et acute duro-apiculatus, glaber, subtus subglauclus v. fuscus, siccus supra atro-fuscus, margine extremo leviter recurvus; costulæ utrinque 12-15, erecto-patulæ, teneræ. Racemi fructigeri 3-pollicares, tenues; pedicelli distantes. Fructus succosi, sicci

nigricantes; endocarpium læve, firmum, monospermum. Stigmata 2, divaricato-adpressa, lanceolata, acuta. Semen ellipsoideum; testa tenuis; albumen durum, nunc nigrescens. — Ex cl. Miq. l. c.

*Species excludenda, nondum recognita, sub Goughia edita.*

GOUGHIA GRIFFITHIANA (H. Wight Icon. plant. Ind. orient. 1878 sub adnot. ad genus Goughiam). 5 In Malacca (Griff.).

---

## Ordo CLXXX. BUXACEÆ (1).

*Tithymalorum* genus *Adans. Fam. des plantes, vol. 2. p. 355.* — *Tricocarum* genus *L. Ordin. nat. plant. p. 474 et p. 480.* — *Euphorbiarum* genus *A. L. Juss. Gen. plant. p. 388.* — *Tithymaloidearum* genus *Vent. Tabl. vol. 3. p. 491.* — *Euphorbiacearum* genera *buxea Bartl. Ord. nat. p. 370. pr. p.* — *Euphorbiacearum* trib. 6. *Buxinæ Dumort. Analyse des fam. nat. p. 45. pr. p.* — *Euphorbiacearum* trib. *Buxæ Endl. Gen. p. 1122. pr. p.; Meissn. Gen. Comment. p. 255. pr. p.; Spach Hist. des végét. vol. 2. p. 486. pr. p.; Reichenb. Conspect. p. 196. pr. p.; Lindl. Veget. Kingd. p. 282. pr. p.* — *Buxaceæ Klotzsch, Linnæ's Pflanzenklasse Tricocceæ p. 12. pr. p.* — *Buxaceæ F. Pke Types des Familles; Baill. Monogr. des Buxac. et Stylocerac. p. 45; Agardh Theor. syst. plant. p. 292; Le Maout et Decaisne, Traité gén. de bot. p. 498.* — *Stylocereæ Baill. Monogr. des Buxac. et Stylocerac. p. 72.*

Flores unisexuales. Calycis laciniae utriusque sexus aestivatione imbricativæ, basi connatæ v. masc. nullæ. Petala et discus in generibus notis 0. Stamina in centro florum v. circa basin rudimenti ovarii inserta. Ovarium superum (carpidiis 2-3), compositum, pluri (2-3)-loculare. Loculi duo in placenta parietali, dein ex apice v. sub apice 1-2-ovulati. Ovula pendula, geminatim collateralia v. solitaria, anatropa; raphe exterior st. peripheriam loculorum spectans, micropyle interior et superior, obturatore quasi operculata. Fructus syncarpus, capsularis aut baccatus, capsularis loculicide aperiens, carpidia a columella centrali secedentia. Semina pendula, albuminosa. Albumen oleagineum. Embryo in albumine centralis, longitudine albumen subæquans. Radicula supera. — Arborea v. frutices, rarius herbæ perennes, in utroque hemisphærio crescentes, sæpius in montosis vigentes. Lignum valde compactum, durissimum. Succus aquosus, parvus, nunquam lacteus. Rami tetragoni v. teretes. Folia opposita v. alterna, exstipulata, penninervia sæpeque tripli-quintuplinervia, integra v. rarius superne grosse serrata, sæpissime coriacea, glabra, nitidula, vulgo sempervirentia. Inflorescentiæ axillares aut rarius terminales, racemiformes v. plus minusve capitato-spiciformes, bisexuales aut raro sexu distinctæ, fœminæ nunc ad florem unicum reductæ. Flores sæpius monoici, fœminei inflorescentiarum bisexualium nunc apicem occupantes et solitarii, nunc masculis inferiores, semper tamen masculis minus numerosi. Pedicelli utriusque sexus bracteati, v. fœminei terminales ebracteati, masculi ebracteolati v. bibrac-

---

(1) Auctore J. Müller, Argoviensi.

teolati, bracteolæ masculorum cum laciniis calycinis duabus inferioribus decussatim alternantes sæpeque ab iis magnitudine forma et consistentia discrepantes, foemineorum autem sæpius plures, inferiores vulgo decussatim oppositæ, superiores gradatim in calycem foemineum abeuntes. Calyx masc. profunde 4-partitus v. rarius nullus, foemineus profunde 4-6-partitus; laciniæ utriusque sexus imbricativæ, florum masc. decussatim oppositæ, foemineorum decussatim oppositæ v. ternatæ. Petala et discus utriusque sexus nulla, sed verticillus alter ex situ staminum tantum suppressus. Stamina, ubi 4, laciniis calycinis opposita, nunc iis magis numerosa, 6-30, semper in fundo calycis inserta; filamenta in alabastro erecta, libera, valida; antheræ liberæ, rimis duabus introrsis longitudinaliter dehiscentes, oblongatæ, dorso supra basin v. ima basi fixæ; loculi longitrorsum connectivo adnati. Connectivum haud peculiari-ter evolutum. Grana pollinis discreta, subglobosa, siccando globoso-tetraëdra, creberrime punctulato-aspera, nunc insuper stellatim subreticulata. Staminodia nulla. Ovarii rudimentum florum masculorum evolutum aut nullum. Ovarium 2-3-loculare, in genere unico dissepimentis spuriis mox 4-6-loculare. Styli 2-3, sæpissime omnino discreti et basi inter se distantes, persistentes, intus canaliculati, integri aut bilobi. Loculi ovarii biovulati, in Stylocerate autem dissepimento intercedente mox spurie 1-ovulati, in Simmondsia vere uniovulati. Fructus in aliis baccati, in aliis capsulares, 3-2-loculares v. abortu 1-loculares, 6-4-v. abortu 5-1-spermi, aperientes centro columella præditi, 3-2-cocci, coccorum endocarpium coriaceum, demum secedens. Semina nuda aut arillata; testa subcrustacea, lævis. Albumen copiosum, carnosum-farinaceum. Embryo magnus; radícula elongata; cotyledones complanatæ, carnosæ; plumula inconspicua.

Ordo parvus, diu cum Euphorbiaceis fere ab omnibus auctoribus unitus, sed capsulis loculicide aperientibus, situ primum parietali ovulorum et directione centripetali micropyles diversus est, cæterum Euphorbiaceas inter et Phytocrenaceas quasi medium tenet. Cum Illicineis, quibus a cl. Baillon approximatus fuit, ob situm hypogynum staminum, analogiam potius quam veram affinitatem præbet, et in vicinitate Euphorbiacearum retinendus est. A Celastrineis dein characteribus gravissimis pluribus distat. — A Forstiera, typo reducto unisexuali et apetalò Oleinearum, ubi ovula similia, cui characteribus proxime accedit (conf. ad Tol. in Annal. des Sc. nat. 1851. p. 264), distinguitur præsentia obturatoris, indole fructuum, et vulgo situ stylium subperipherico, defectu punctorum poriformium foliorum.

*Conspectus diagnosticus tribuum, subtribuum et generum.*

**Trib. I. Buxææ.** Loculi ovarii 2-ovulati.

Subtrib. I. *Stylocerææ*. Stamina centralia. — Loculi 3-2 ovarii mox dissepimento spurio in loculos 6-4 spurie uniovulatos mutati.

1. *Styloceras*. — Arbores alternifoliæ.

Subtrib. II. *Eubuxææ*. Stamina circa rudimentum ovarii inserta. — Loculi 2-3 ovarii absque dissepimento spurio.

2. *Sarcococca*. Fructus baccatus, haud aperiens. Grana pollinis stelatim subreticulata. — Frutices alternifolii. Inflorescentiæ inferne fœmineæ.

3. *Buxus*. Fructus capsulari-aperiens. Grana pollinis haud reticulata. — Frutices oppositifolii. Inflorescentiæ apice fœmineæ.

4. *Pachysandra*. Fructus capsulari-aperiens. Grana pollinis stelatim subreticulata. — Herbæ alternifoliæ. Inflorescentiæ inferne fœmineæ.

Trib. II. *Simmondsiææ*. Loculi ovarii 1-ovulati.

5. *Simmondsia*. Fructus capsularis. — Frutices oppositifolii.

Trib. I. *BUXEÆ*. — Buxacæ Baill. *Monogr. des Buxac. et Stylocer.* p. 45, et *Stylocerææ* Baill. l. c. p. 72. — Loculi ovarii biovulati.

Subtrib. I. *STYLOCEREÆ*. — Ordo *Stylocerææ* Baill. *Monogr. des Buxac. et Styloc.* p. 76.

Stamina centralia. — Flores masc. calyce destituti, 6-3-andri; antheræ ima basi fixæ; ovarii rudimentum nullum. Loculi ovarii primum 2-ovulati, mox dissepimento in loculos geminos uniovulatos divisi et tum stylis duplo numerosiores.

I. *STYLOCERAS* Adr. *Juss. Tent. Euphorb.* p. 53. t. 17. fig. 56; *Kunth in Humb. et Bonpl. Nov. gen. et spec. plant. vol. 7.* p. 172. t. 637-638; *Kunth Syn. pl. æquinoc. vol. 4.* p. 206; *Endl. Gen. n.* 5773; *Baill. Étud. gén. Euphorb.* p. 665. t. 20. fig. 25-37 et in *Monogr. Buxac. et Styloc.* p. 77. — *Tropis* sp. *Willd. Spec. plant. vol. 4.* p. 733.

Flores masc. calyce destituti, fœminei eo præditi, utriusque sexus petalis et disco carentes. Stamina subnumerosa, centralia; antheræ birimosæ, ima basi fixæ. Grana pollinis haud reticulata. Rudimentum ovarii 0. Ovarium compositum, 2-3-loculare; loculi 2-ovulati, mox dein dissepimento spurio divisi et quisque in loculos spurios 2 uniovulatos abiens. Fructus carnosio-sublignosi, haud aperiens. Semina albuminosa. — Arbores Americæ austro-occidentalis, glaberrimæ. Folia alterna, exstipulata, petiolata, plus minusve oblongo-obovata, integra, subtripplinervia, præter basin penninervia, coriacea. Inflorescentiæ axillares, spiciformes. Spicæ unisexuales aut bisexuales et tum flores masculii inferiores. Flores masc. basi unibracteati, fœminei bracteolis nonnullis præditi, bractææ et calyces fœm. parvi. Lacinia calycis fœm. 4, decussatim oppositæ, æstivatione imbricativæ. Stamina 6-30. Filamenta libera, brevissima. Antheræ longitrorsum et introrsum birimosæ, apice connec-

tivo obtase apiculatæ. Grana pollinis puncticulato-aspera. Ovarii loculi 2-3 mox dissepimento spurio in 4-6 conversi. Ovula ex apice loculorum pendula, micropyle oblique axin ovarii spectans. Styli 2 v. rarius 3, tot quot loculi veri ovarii, validi, liberi v. inferne connati. Fructus edules, demum sicci, 2-3-cornuti. Seminum testa lævis, fragilis. Embryo longitudine albuminis.

### § 1. Styli liberi, inter se discreti.

1. *S. KUNTHIANUM* (Adr. Juss. Tent. Euphorb. p. 117. t. 17. fig. 56), spicis bisexualibus, antheris oblongo-obovoideis molliusculis, ovario globoso-ehipsoideo. § In prov. Quito ad radicem montis Tunguraguæ altit. 1400 hexapod. mense Junio (Bonpl. n. 3208! in hb. Mus. paris.), prope Huayaquil (hb. Pav.). — Kunth in Humb. et Bonpl. Nov. gen. et spec. pl. vol. 7. p. 172. t. 637; Baill. Etud. gén. Euphorb. Atl. p. 37. t. 20. fig. 32-37 et Monogr. Buxac. et Styloc. p. 78. *Trophis laurifolia* Willd. Spec. pl. vol. 4. p. 733. Foliorum limbus oblongo-obovatus, basi acutus, apice breviuscule acute acuminatus. Spicæ bisexualis, quam masculæ speciei sequentis multo breviores, petiolos parum superantes eoque vix æquantes, sæpius 5-7-floræ. Flores præter differentias indicatas cum iis sequentis quadrant. Antheræ superne latiores quam in sequente, multo minus rigidæ, cæterum ut in illo scabræ. Folia utriusque magnitudine, consistentia, nervatione et colore satis quadrant, sed hujus sunt acutiora et minus oblongata. Ovarium mox 4-loculare, rarissime 6-loculare (v. s.).
2. *S. LAURIFOLIUM* (Kunth in Humb. et Bonpl. Nov. gen. et spec. pl. vol. 7. p. 173. t. 638), spicis unisexualibus, antheris cylindrico-obovoideis chartaceo-rigidis, ovario globoso. § In regno Novo-Granatensi, in regione Andina Bogotensi. altit. 2600-3800 metror. (Bonpl. Triana n. 3564! in hb. Mus. paris., Goudot n. 1! Holton n. 847!). — Baill. Etud. gén. Euphorb. Atl. p. 37. t. 20. fig. 25-31 et Monogr. des Buxac. et Styloc. p. 79. Arbor 6-8-metralis (Triana). Petioli  $1\frac{1}{2}$ -2 cm. longi. Foliorum limbus 12-15 cm. longus, 5-6 cm. latus, oblongo-v. lanceolato-obovatus, basi cuneato-angustatus, apice obtusiusculus. Costæ secundariæ inferiores elongatæ, prominulæ. Spicæ masc. circ. 3-6 cm. longæ, petiolos duplo triplove superantes, circiter 10-14-floræ. Pedunculi fem. petiolis breviores, apice incrassato uniflori. Bractæe latissime ovatæ, subcutæ, glabræ, margine ciliolatæ, circ. 2 mm. longæ, inferiores nonnullæ steriles. Calyx fœm. 4-partitus, subinde ab ovario subremotus, vix  $1\frac{1}{2}$  mm. longus; lacinie ambitu latissimæ, acutiusculæ. Antheræ 3-4 mm. longæ, magis cylindricæ quam in specie præcedente. Ovarium subpuberulum. Fructus haud omnino maturi diametro 2 cm. æquantes, longe bicornuti, globosi, subignosi. — Ludit: *Dombeyanum*, foliis pallidioribus supra fere eventis costisque sæpe minus prominentibus, pedunculis fœm. paulo brevioribus 6-8 mm. æquantibus. § In Peruvia (Dombey! in hb. Mus. paris.). *S. Dombeyanum* Baill. Etud. gén. Euphorb. p. 666, et Monogr. Buxac. et Styloc. p. 80. (v. s.)

### § 2. Styli inferne in columnam connati.

3. *S. COLUMNARE*, ovario globoso-pyriformi apice truncato-obtuso, stylis circ. quarta v. tertia parte longitudinalis in columnam connatis. § In Bolivia Andina prov. Larecaja, prope Sorata inter Cochipata et Milipaya, alt. 10 000 ped. (G. Mandon n. 1084!). specimina similia iis *S. laurifolii* et absque fructu aut flore fœm. æge ab iis recognoscenda. Fructus demum 2 cm. longi, superne totidem lati, basin versus brevissime acutati. Styli duo, vulgo circ. 18 mm. longi. Semina 10 mm. longa, 8 mm. lata, nitida. — A duabus præcedentibus forma et magnitudine subduplo majore fructuum et stylis inferne in columnam connatis diversa sed iis cæterum simillima. (v. s.)

### † Species ex ordine exclusa.

- S. MACROSTACHYUM* Baill. Monogr. Buxac. et Styloc. p. 81 (Peppigii n. 2545! in hb. Mus. paris.) ad Euphorbiaceas pertinet et eadem planta est ac *Alchornea cordata* Müll. Arg. in DC. Prodr. vol. 15. part. 2. p. 901.

Subtrib. II. EUBUXEÆ. — *Plée Types de chaque Famille*; Kirschleger *Flore d'Alsace*, vol. 2. p. 48; Baill. in *Bull. Soc. bot. de France*, 1856. p. 287, et *Monogr. Buxac. et Styloc.* p. 45; Agardh *Theor. syst. plant.* p. 292. pr. p. — Eubuxaceæ Baill. *Monogr. Buxac. et Styloc.* p. 76.

Stamina circa rudimentum ovarii inserta. — Flores masc. calyce præditi, 4-andri; antheræ dorso supra basin fixæ; ovarii rudimentum tetragonum. Loculi ovarii semper 2-ovulati, absque dissepimento spurio, tot quot styli.

II. SARCOCOCCA Lindl. in *Bot. Reg.* t. 1012; Endl. *Gen. pl.* n. 5875; Baill. *Monogr. Buxac. et Styloc.* p. 49. t. 3. fig. 15-30. — *Lepidopelma Klotzsch in Bot. Ergebn. d. Reise d. Prinz. Waldemar*, p. 118. t. 22. — Buxi sp. Don *Prodr. Flor. nepal.* p. 63. — *Pachysandra?* sp. Hook. *Flor. esot.* t. 148.

Flores utriusque sexus calyce præditi; laciniae æstivatione imbricativæ. Petala et discus 0. Stamina laciniiis calycis isomera iisque opposita; antheræ dorso supra basin fixæ. Grana pollinis stellatim subreticulata. Rudimentum ovarii evolutum. Ovarium 2-3-loculare. Loculi 2-ovulati. Fructus subbaccatus, haud aperiens. Semina albuminosa. — Frutices indici et sundaici, sempervirentes. Rami teretes. Folia alterna, exstipulata, triplinervia aut peminervia, integra. Inflorescentiæ axillares, racemiformes, plurifloræ, inferne fœminæ. Pedicelli 1-bracteati, bi-pluri-bracteolati. Bracteolæ pedicellorum masc. v. 2 v. 0, laciniiis calycinis similes, pedicellorum fœm. magis numerosæ et minores, superiores gradatim majores, decussatim oppositæ. Calyx masc. profunde 4-partitus; laciniae decussatim oppositæ, quarum exteriores in pedicello ebracteolato locum tenent bracteolarum, i. e. laterales evadunt; ubi autem in pedicellis masc. (ejusdem speciminis) bracteolæ duæ adsunt, tum laciniae exteriores calycis (cum bracteolis et cum duabus laciniiis interioribus calycis alternantes) evadunt altera postica altera antica, et interiores laterales, unde situs parium laciniarum calycis e præsentia aut e defectu bracteolarum pedicelli pendet. Calycis fœm. laciniae 4-6, decussatim oppositæ v. ternatæ. Filamenta libera, valida. Antheræ fere stantes; loculi fere tota longitudine connectivo latiusculo adnatæ, intus longitrorsum birimosæ, connectivo leviter producto subapiculatæ. Ovarii rudimentum tetragono-obovoideum, vertice suburceolatum ibidemque margine subintegrum. Styli tot quot loculi, integri v. apice subuncinato bilobi, elongati, validi, intus superne sulcati et papilloso. Obturator ovulorum quam in Buxo et Stylocerate longe tenuior, nec tumido-carnosulus. Baccæ 6-spermæ v. abortu sæpius 5-1-spermæ. Semina arillata. Radicula cotyledonibus late ovatis carnosis compressis paulo brevior. — Genus hoc characteribus valde accedit ad Buxum, sed ramuli teretes, folia alterna, inflorescentiæ inferne nec apice fœminæ, et, quod majoris momenti, fructus haud capsulares.

1° Folia triplinervia.

1. S. SALIGNA, pedunculis masc. 2-bracteolatis v. ebracteolatis, stylis patulis, mesocarpio carnoso, bacca siccando ambitu irregulari oblongata.

a. *coriacea*, foliis ovato-lanceolatis longe cuspidato-acuminatis, bracteis fl. masc. ovatis longe acuminatis majusculis lacinias calycinas semiattin-

gentibus superioribus elongatis, bracteolis fem. oblongo-triangulari-ovatis longe acuminatis, lacinis calycinis fem. 4 decussatim oppositis, ovario sæpius trimero.  $\frac{3}{4}$  In Nepalia (Wallich! in hb. DC.). Buxus coriacea Spreng. Syst. 4. p. 314 (1827). Pachysandra? coriacea Hook. Exot. Flora t. 148 (1825). Prunus Puddum hort. ex Hook. l. c. Folia 2-3½ cm. lata, absque petiolo 1-1½ cm. longo 5-10 cm. longa. Inflorescentiæ laxiores quam in reliquis var., petiolos subæquantes. Pedunculi fem. gemini v. terni, tenues, 6-8 mm. longi. Calycis masc. laciniae 4-5 mm. longæ, pallidæ, margine late albidæ, ciliolatæ, circiter triente majores quam in sequentibus varietatibus. (v. s.)

$\beta$ , *genuina*, foliis lineari-lanceolatis basi longe angustatis apice longissime cuspidato-acuminatis, bracteis pedunculorum masc. ovatis acutis abbreviatis, bracteolis oblongo-ovatis acuminatis, calyce fem. sæpius 6-partito, inflorescentiis petiolos breviusculos paulo superantibus, ovario sæpius trimero.  $\frac{3}{4}$  In Nepalia (Wallich n. 7979. Cl.) et in regione temperata Himalayæ bor. occid. (Th. Thomson!), ad Otypore (Griff.!). Buxus saligna Don Prodr. Flor. nepal. p. 63 (1825). Sarcococca salicifolia Baill. Monogr. Bux. p. 49. S. pruniformis  $\beta$  angustifolia Lindl. Bot. Reg. t. 1012 (1826); Baill. Monogr. Buxac. p. 50. Lepidopelma podocarpifolia Klotzsch in Bot. Ergebn. Reis. Prinz. Waldemar p. 118. t. 22. Differt a reliquis varietatibus foliis ejusdem longitudinis ac in var. præcedente sed tantum 7-18 mm. latis et basi longius angustatis. Petioli tantum 5-8 mm. longi. — Puncta nigro-fusca in utraque pagina hujus et sequentis varietatis occurrunt, forte eadem de quibus cl. Baillon mentionem fecit, sed non specifiçi momenti sunt et ad regnum animalium referenda sunt. (v. s.)

$\gamma$ , *latifolia*, foliis latius angustiusve ovato-lanceolatis basi acutis apice longius cuspidato-acuminatis, bracteis ovatis acutis, bracteolis oblongo-ovatis acuminatis subcoloratis, calyce fem. 4-6-partito, inflorescentia petiolos mediocres paulo superante, ovario sæpius dimero.  $\frac{3}{4}$  In Nepalia (Wallich n. 7979 A!), in Khasia (Griff.!), Sikkim altitud. 7-9000 ped. s. m. (J. D. Hook.!), in Bengalia orientali (Griff. n. 4896!). Tricera Nepalesis Wall. Cat. n. 7979 (nomen). Sarcococca Sumatrana Bl. in Mus. lugd. bat. 2. p. 191; Baill. Monogr. Bux. p. 51. S. pruniformis  $\alpha$  latifolia Lindl. Bot. Reg. t. 1012; Baill. Monogr. Buxac. p. 50. Folia 2½-3½ cm., imo usque ad 5 cm. lata. Petioli sæpius 7-12 mm. longi. — Hæc ludit : *petiolaris*, petiolis elongatis.  $\frac{3}{4}$  In Khasia Indiæ orientalis altitud. 4000-6000 ped. (J. D. Hooker et Th. Thomson!). Petioli 1½-2 cm. longi. Inflorescentiæ juniores graciles, petiolos paulo superantes. (v. s.)

$\delta$ , *densiflora*, foliis ovato-lanceolatis basi acutis apice breviter cuspidato-acuminatis, inflorescentiis quam petioli brevioribus quam in reliquis varietatibus densioribus subovoideis subpallidis, ovario sæpius dimero.  $\frac{3}{4}$  In montibus Neilgherries peninsulae Indiæ orientalis (Perrott. n. 1029! Hohenack. n. 1056! G. Thomson!), in prov. Canaræ (Hohenack. n. 484! et 484. al.). S. trinervia Wight Icon. pl. Ind. or. t. 1877, sed folia triplinervia sunt nec trinervia. A cl. Miq. in pl. Hohenack. exs. l. c. ad genus Myricam relata fuit. — A reliquis var. differt foliis sæpe firmioribus, minus longe cuspidato-acuminatis et inflorescentiis abbreviatis. Frutex est 10-15-pedalis (ex Perrott. sched.). (v. s.)

$\epsilon$ , *brevifolia*, foliis ovatis basi acutis apice obtuse acuminatis, inflorescentiis quam petioli brevioribus, pedunculis masculis ebracteolatis.  $\frac{3}{4}$  In insula Zeylonia (Thwait. n. 203 pr. p. ! Walker!). Petioli circ. 7-8 mm. longi. Limbus foliorum 4-5½ cm. longus, 2½-3½ cm. latus, tenuiter coriaceus. (v. s.)

2. S. ZEYLANICA (Baill. Monogr. Bux. p. 52), pedunculis masc. decussatim 4-bracteolatis, stylis arte uncinato-recurvis, mesocarpio coriaceo firmo, bacca siccando ambitu regulari breviter ellipsoidea.  $\frac{3}{4}$  In insula Zeylonia (Thwaites n. 203 pr. p. !). S. pruniformis Thwait. Enum. pl. Zeyl. p. 290 pr. p. Petioli 7-12 mm. longi. Limbus foliorum 4-8 cm. longus, 1-2 m. latus, ambitu valde variabilis, ovato-lineari-lanceolatus, basi acutus v. angustatus, apice longe cuspidato-acuminatus, firme membranaceus. Inflorescentiæ integræ petiolos



subsequentes lisse breviores, pars masc. demum secedens. Bracteolæ masc. summæ duæ inferioribus duabus duplo majores, orbiculari-ovatæ, acutæ, margine ut in congeneribus ciliatæ,  $2\frac{1}{2}$  mm. æquantes. Pelunculi masc. subnulli, feminei 2-3 mm. longi, hi decussatim pluribracteolati, bracteolis superioribus sensim majoribus. Ovarium dimerum. Fructus globoso-ellipsoidei, superficie regulares, tantum  $4-5\frac{1}{2}$  mm. longi. — A var.  $\beta$  genuina *Sarcococcæ* salignæ distinguitur floribus baccisque multo minoribus, stylis brevioribus et mesocarpio. (v. s.)

2° Folia penninervia.

3. *S. HOOKERIANA* (Baill. Monogr. Bux. p. 53), bracteis pedicellorum masc. longe acuminatis, pedunculis fem. pluribracteolatis, bracteolis ternatis, calycis fem. lacinii 6, ovario globoso, stylis suberectis, bacca globosa.  $\delta$  In Indiæ orientalis montibus Sikkim (J. D. Hook. et Th. Thomson! in hb. Mus. paris., DC. etc.), in Himalaya orientali (Griff. n. 4897!). Petioli 5-8 mm. longi. Limbus foliorum 5-7 v. 8 cm. longus, anguste lanceolatus, acuminatus, basi acutus, subcoriaceus, supra nitidulus, v. opacus, nervi secundarii subtus vix bene perspicui, par infimum reliquis haud dissimile. Nervi tertiarii haud distingui possunt. Inflorescentiæ breves, petiolum æquantes, laxifloræ. Pedunculi fem. 5-7 mm. longi, gemini v. ternati. Bractæe masc. ovato-lanceolatæ, lacinias calycinas bene semiatingentes, fem. ovato-lanceolatæ, longe acuminatæ, decussatim oppositæ. Calycis masc. lacinia 4, decussatæ, acuminatæ, utriusque sexus 4-5 mm. longæ, femineæ masculis angustiores pauloque longius acuminatæ. — Folia ut in *S. saligna*  $\beta$  genuina, sed non cuspidato-acuminata, nec triplinervia. (v. s.)
4. *S. ZOLLINGERI* (Baill. Monogr. Bux. p. 54), bracteis pedicellorum masc. acuminatis, pedicellis fem. 4-bracteolatis, lacinii calycis fem. 4 decussatim oppositis, ovario subgloboso, stylis suberectis apice hamato-recursis.  $\delta$  In insula Java (Zolling. n. 2316!). Petioli circ. 8 mm. longi. Limbus foliorum fere ut in *S. saligna*  $\epsilon$  brevifolia, sc. 4-7 cm. longus,  $2\frac{1}{2}-3\frac{1}{2}$  cm. latus, ovatus v. oblongo-ovatus, basi acutus, apice breviter acuteque acuminatus, firme membranaceus, opacus, subpenninervius. Nervi secundarii in utraque pagina minus perspicui, par infimi reliquis paulo longiores et juxta marginem foliorum arcuatim fere usque ad medium folii, præsertim in foliis junioribus, producti, sed iis haud validiores. Margo foliorum in speciminibus visis integer, sæpe quasi undulato-crenatus (remote obtuseque crenatus Baill. l. c.). Pedicelli ad basin inflorescentiarum 2-5-ni, ovario sæpius dimero paulo longiores. Bractæe fem. plus minusve lanceolatæ, decussatim oppositæ, acuminatæ, virides, inferior 4-5 mm. longa, reliquæ paulo breviores, masculæ ovato-lanceolatæ. Calycis masc. lacinia 4, decussatim oppositæ, oblongo-ovatæ, acuminatæ, femineæ  $3-3\frac{1}{2}$  mm. longæ, oblongo-ovatæ, acuminatæ. Rudimentum ovarii basi angustatum, vertice oblongo-tetragono urceolatum. Baccæ desunt. (v. s.)

III. *BUXUS* Tournef. (emend.); Spreng. Syst. veg. 3. p. 760. n. 3103 et pag. 847; Baill. Monogr. Buzac. p. 58. — *Buxus* et *Tricera* auct.

Flores utriusque sexus calyce præditi; lacinia æstivatione imbricativæ. Petala et discus 0. Stamina lacinii calycis isomera iisque opposita; antheræ dorso supra basin fixæ. Grana pollinis haud reticulata. Rudimentum ovarii evolutum. Ovarium 3-loculare. Loculi 2-ovulati. Fructus trilocci, capsulari-aperientes. Semina albuminosa. — Frutices v. rarius suffrutices sempervirentes, in utroque hemisphærio parce vigentes. Ramuli tetragoni. Folia opposita, exstipulata, penninervia, integra, lævia, nitida. Inflorescentiæ axillares, spiciformes, apice femineæ. Pedicelli masc. unibracteati, feminei sæpius 4-bracteolati; bracteolæ decussatim oppositæ, sæpe magis minusve numerosæ minusque regulariter sitæ. Flores monoici, feminei solitarii et inflorescentias terminantes. Calyx

mass. profunde 4-partitus, lacinii decussatim oppositis, femineus decussatim bitermarius, hujus laciniae tres interiores cum stigmatibus alternantes, demum 2-3-partito-ruptae. Stamina 4; filamenta libera, valida; antherae longitudinaliter introrsum birimosae; connectivum haud productum. Grana pollinis sicca humectaque globosa, sublaevia, saltem haud reticulata. Ovarii rudimentum tetragonum, subcompressum, superne paulo latius leviterque 4-lobulatum. Ovarium obovoideum, 3-gonum, vertice stylis ad angulos sitis brevibus intus sulcatis apice subbilobis tricornutum, inter cornua turgide tuberculatum. Capsula coriacea, loculicide trivalvis, 6- v. abortu 5-3-sperma. Semina trigona, laevia. Albumen copiosum. Embryo longitudine albuminis; radícula elongata, cotyledones oblongo-ellipticas crassas superans.

Sect. I. *TRICERA* Baill. *Monogr. Bux.* p. 66. — Genus *Tricera* Sw. *Flor. Ind. occid.* vol. 1. p. 333. t. 7; Schreb. *Gen. pl.* n. 1416; Endl. *Gen. pl.* n. 5868; A. Rich. *Flor. cub.* t. 71. — Genus *Crantzia* Sw. *Prodr.* p. 38; Vahl *Symb.* 2. p. 99 (non alior.)

Styli elongati, paginae foliorum cohaerentes. — Inflorescentiae partes fere semper haud dense imbricatae; characteres florum et fructuum caeterum admodum cum his sectionis Eubuxi conveniunt, sed flores mass. fere semper pedicellati, vulgo ebracteolati, ovarium semper longe 3-v. raro 4-rostratum, et bractae nec dense imbricatae nec scariosae. — Arillus apud Sw. et Vahl nihil est nisi coccorum endocarpium secedens.

### § I. Ovarium apice angustatum in stylos basi breviter connatos columnam spuriam formantes abiens.

1. *B. SUBCOLUMNARIS*, foliis demum subcoriaceis vix prominenter venosis, cymis subgrandifloris, floribus fem. sessilibus masculis elongato-pedicellatis, filamentis inferne dilatatis, ovario anguste ellipsoideo superne in columnam stylarem brevem abeunte, stylis elongatis longo tractu liberis. § In Martinica (Sieb. hb. fl. mart. n. 2081) et in Cuba (C. Wright n. 1920, a cl. Griseb. Cat. pl. cub. p. 14 ad *Triceram laevigatam* Sw. relata). Ramuli elongati, siccando angulosi, virides. Petioli 3-8 mm. longi, rigidi. Foliorum limbus 6-9 cm. longus, 1½-3 cm. latus, elliptico-lanceolatus, breviter acuminatus, basi acutus, pallide viridis, junior membranaceus, dein mox firmior, laevigatus, supra nitidulus. Cymae multiflorae, fere corymbiformes, varie pedunculatae v. subsessiles. Bractae oblongo-ovatae v. triangulari-lanceolatae, 2½-3 mm. longae. Calycis masculi laciniae lanceolato-ovatae v. ovato-lanceolatae, circiter 5 mm. longae, a filamentis validis superatae, mox reflexae; calycis feminei laciniae ovatae, ovario pluries breviores, vix 1½ mm. longae. Filamenta antheris latiora. Ovarium apice callosio-angustatum, in stylos centro se tangentes longo tractu erectos et quasi columnam spuriam ovarium longitudine aequantem eoque pluries angustiorum formantes abiens. Styli basi tantum connati, apice patuli saepeque ibidem revoluti, ovario longiores, in genere graciles. (v. s.)

### § 2. Ovarium apice late truncatum; styli basi inter se distantes.

α. Costae secundariae saepiusque et venae distinctae.

1° Flores masculi pedicellati, feminei sessiles.

\* Species grandiflorae, foliis venosis.

2. *B. CUBANA* (Baill. *Monogr. Bux.* p. 71), foliis basi subcordatis triplinerviis demum coriaceis supra lucidis, antheris cordato-ovoidis quam filamenta longioribus, ovario ovoidico globoso, stylis ovarium sequens sequentibus. § In insulae Cuba: saepe in Libano, in districtu Santiago de Cuba (Kinden n. 1081).

in hb. Benth. et in hb. Boiss.). *Tricera Cubana* A. Rich. in Ramon de la Sagra Flora Cub. edit. hisp. vol. 3. p. 217. t. 71. Tota glaberrima. Ramuli sicando angulosi. Petioli 2 mm. longi, crassi. Limbus foliorum  $3\frac{1}{2}$ -4½ cm. longus,  $2\frac{1}{2}$ -3½ cm. latus, orbiculari-ovatus, acutus, basi leviter cordatus. Inflorescentiæ in axillis ultimi paris foliorum sitæ sunt, bisexuales bifloræ, aut unisexuales masculæ 6-floræ; pedunculus bracteolis sterilibus præditus. Pedicelli masc. 3-8 mm. æquantes, calycem paulo superantes. Lacinia calycis late triangulari-ovata, acutiusculæ, femineæ masculis duplo et ultra minores. Styli integri, patuli, ovarium sesqui æquantes. — In flore masculino iconis antheræ errore 5 delineatæ sunt. (v. s.)

3. *B. PURDIANA* (Baill. Monogr. Bux. p. 70, 1859), foliis subsessilibus prominenter reticulato-venosis nitidis, filamentis calycem demum longe superantibus crassis, ovario glabro. ♂ In Jamaica ad St. Anna (Purdie in hb. Mus. Paris.). *Tricera fasciculata* Griseb. Flor. of Brit. West Ind. p. 31 (1859). Frutex humilis. Lignum durissimum, flavum. Cortex griseus, rugosus. Rami teretes. Ramuli tetragoni, virescentes, juniores tenues, glabri. Folia vix petiolata, 6-7 cm. longa,  $2\frac{1}{2}$ -3 cm. lata, e basi attenuata oblongo-ovata, v. breviter lanceolato-elliptica, utrinque acuta v. apice obtusiuscula, membranaceo-coriacea. Inflorescentiæ sæpius supraaxillares, abbreviatæ, subsessiles, quam petioli breves 2-3-plo longiores, 5-7-floræ. Pedicelli masc. sæpius bracteolati. Calycis glabri lacinia ovata, oblongata, subacutæ, 2 mm. longæ. Ovarium 4-6 mm. longum, calycem longe superans. (v. s.)

4. *B. CITRIFOLIA* (Spreng. Syst. veg. vol. 3. p. 847), foliis longius petiolatis firme membranaceis, filamentis antheras diametro bene æquantibus, ovario glabro, capsulis mediocribus.

α, *genuina*, foliis ovato-lanceolatis pallidis, inflorescentiis quam folia 4-5-plo brevioribus, pedicellis masc. calycem bis æquantibus, cornubus gracilibus capsula brevioribus. ♂ Ad Caracas (Bredemeyer in hb. Willd. fol. 17384 !). *Tricera citrifolia* Willd. Spec. pl. vol. 4. p. 338. Frutex 15-pedalis, facie Coffæ arabicæ (Bredem.). Ramuli glabri. Petioli circ. 5 mm. longi. Limbus foliorum 8-11 cm. longus, 4 cm. latus, basi acutus, apice acuta acuminatus, firme membranaceus, glaber. Pedunculi breves, basi subdecompressi bracteati, circiter 10-flori. Bractea inferioris steriles circiter 4-6, approximatae. Axis racemi  $1\frac{1}{2}$  cm. longus, fere tertia parte inferiore ananthus. Pedicelli masc. calycem bis et bractea quater-quinq. æquantes, 1-1½ cm. longi. Calycis masc. lacinia 4 mm. longæ v. longiores, oblongo-ellipticae, marginæ membranaceae, feminei lacinia autem fere triplo minores, oblongo-ovatae, acuminatae, puberulae. Filamenta calycem demum paulo superantia. (v. s.)

β, *fuscescens*, foliis lanceolato-ovatis subcoriaceis rufescenti-fuscis, racemis breviuscule pedunculatis, pedicellis masc. calycem paulo superantibus. ♂ In Portorico (Wydler n. 406!). *B. citrifolia* Baill. l. c. descript. pr. p. Folia 5-9 cm. longa,  $2\frac{1}{2}$ -4½ cm. lata, multo rigidiora quam in varietate α, et, si breviora essent et basi cordata, magis similia iis *B. Cubanae*. Calyces etiam sunt firmiores, masculi  $3\frac{1}{2}$ -4 mm. longi. Capsulae lignosae, absque stylis persistentibus et ea brevioribus 1 cm. longæ, scabrae. (v. s.)

γ, *brevipes*, foliis lanceolato-ellipticis utrinque subæqualiter acutatis viridibus v. dein nonnihil rufescentibus, cymis compactis subsessilibus dense imbricatis bracteatis. ♂ In Cuba (C. Wright n. 1919 pr. p. 1). Folia 6-8 cm. longa, 3-3½ cm. lata, juniora rubro-marginata; cætera junior rubra, dein expallens. Capsulae ignotæ. (v. s.)

5. *B. ACUMINATA*, foliis rigide membranaceis venosis, ovario globoso glabro, cornubus ovarium æquantibus distantibus subgracilibus dein erectis apice deflexis, capsulis parvis globoso-ellipsoideis, cornubus capsulae circiter triente brevioribus. ♂ In Cuba (C. Wright n. 1919, pr. p. 5). *Tricera acuminata* Griseb. in Nachricht. d. Königl. Gesellsch. d. Wissensch. d. Univ. Götting. 1865. p. 162. Limbus foliorum 8-9 cm. longus, 2-3 cm. latus, anguste ovato-lanceolatus, basi acutus, apicem versus longius acuminatus, in pagina inferiore argillaceo-fuscescens, viridi-maculatus. Cymæ breviter v. breviuscule pedunculatae, pævisformæ. Bractea triangularis, parva. Calycis lacinia lan-

ceolato-ovatæ, subacutæ, femineæ parvæ. Capsulæ absque cornubus 6 mm. longæ, fere 5 mm. latæ. — Medium quasi tenet inter *B. citrifoliam* et *B. lævigatam*, cum priore longitudine relativa cornuum, cum posteriore minutie fructuum quadrans. (v. s.).

6. *B. LÆVIGATA* (Spreng. Syst. veg. vol. 3. p. 847), foliis firme membranaceis venosis, filamentis antheras diametro superantibus, ovario puberulo, stylis ovarium his et ultra æquantibus, capsulis truncatis parvis, cornubus gracilibus quam capsula longioribus. ♂ In fruticetis montium Jamaicae occidentalis (Sw. in variis hb.). — Baill. Monogr. Bux. p. 66. *Tricera lævigata* Sw. Flor. Ind. occid. vol. 1. p. 333. t. 7; Adr. Juss. Tent. Euphorb. t. 1. fig. 3. B; Griseb. Flor. of Brit. West Ind. p. 31. *Crantzia lævigata* Sw. Prodr. p. 38. Frutex 2-3-pedalis (Sw.). Rami s. simpliciter, teretes, cinerei; ramuli tetragoni, glabri. Petioli 1-2 mm. longi. Foliorum limbus 4-7 cm. longus, 1-1½ cm. latus, elliptico-lanceolatus, utrinque acutus v. angustatus, fusciculus, fusco-maculatus, glaber, leviter nervosus, triplinervius; costæ secundariæ numerosæ, costas submarginales transgredientes. Inflorescentiæ paulo supra-axillares, quam folia subtriplo breviores, solitariae v. interdum geminae, totæ puberulo-scabridæ; pedunculi 1-1½ cm. longi, infra pedicellos masculos bracteis sterilibus 3-5 onusti, qui sub apice pedicellos masc. decussatis oppositos approximatos 4-8 quasi umbellam formantes gerunt. Bractæe ovato-lanceolatae, laciniiis calycinis similes, fertiles masculæ quam pedicelli fere duplo breviores. Calyces 3-4 mm. longi, sicci fusci, rigidi. Filamenta calycem paulo excedentia. Ovarium plus minusve puberulum. — A proxima *B. citrifolia* recognoscitur floribus minoribus, capsulis parvis, cornubus capsularum subduplo brevioribus. (v. s.)

\*\* Species parvifoliæ, foliis subobsolete venosis.

7. *B. VAHLII* (Baill. Monogr. Bux. p. 67, exclus. syn. Juss.), foliis subsessilibus coriaceis acute mucronulatis subtus subvenosis, filamentis demum quam antheræ multo angustioribus, ovario glabro stylis breviori, capsula cylindrico-ellipsoidea cornubus longiore. ♂ In insula Sancta Crucis (Richard in hb. Willd. fol. 17383! et West in hb. Puerari nunc in hb. DC.!), in Jamaica (hb. Mus. paris.). *Crantzia lævigata* Vahl Symb. 2. p. 99. *Tricera lævigata* Willd. Spec. vol. 4. p. 338 pr. p. *Crantzia Solandri* Vahl mss. ex Baill. l. c. et *Buxus cruciata* L. C. Rich. mss. ex Baill. l. c. *Buxus pulchella* Baill. Monogr. Bux. p. 68. Ramuli glabri. Folia brevissime petiolata 3-3½ cm. longæ, supra medium 1½ cm. lata, obovato-lanceolata, basi breviuscule angustata, apice obtusiusculo mucronulata. Inflorescentiæ axillares, quam folia 4-6-plo breviores, sessiles. Bractæe steriles infra flores decussatæ, sæpius 6, approximate, ovatae, puberulae, fertiles pedicellos masc. breves fere æquantes. Calycis vix 2 mm. longi laciniae oblongo-ovatae, acutæ. Capsulae 5-6 mm. longæ, minus longe tricornutæ. — A *B. lævigata* primo intuitu distinguitur rigiditate ramorum et ramulorum, foliis coriaceis sæpe rubro-maculatis v. striolatis et floribus subduplo minoribus. Pedicelli et præsertim filamenta præter alia omnino differunt. — Apud cl. Baill. Monogr. l. c. synonymon Jussieanum ad hanc, Willdenowianum autem (pro *Crantzia lævigata* Vahl Symb. 2. p. 99) ad *B. lævigatam* lapsu quodam translata sunt. (v. s.)
8. *B. GONOCLEADA*, foliis obsolete arcunerviis, capsulæ lobis lineari-spathulatis v. spathulatis obtusis septo sublongioribus. ♂ In parte occidentali insulae Cubæ, in savanis prope Guamacaro (C. Wright, ex cl. Griseb. l. c.). *Tricera gonocleada* Wright ex Griseb. Cat. pl. cub. p. 282. Frutex 2-6-pedalis. Foliorum limbus 4-5 cm. longus, 15-25 mm. latus, in petiolum latiusculum 2-4 mm. longum breviter attenuatus, lanceolato-oblongus, apice obtuso emarginatus, basi complicatus, supra lucidus, subtus virens. Pedunculi axillares, internodium æquantes, foliis triplo breviores, 3-5-flori. Pedicelli capsulis subæqui-longi. — Habitus et folia *B. retusæ*, ubi autem capsulæ lobi latiores, semiovati, obtusi, septo duplo breviores.
9. *B. RETUSA*, foliis coriaceis, bracteis triangularibus, capsula globoso-ellipsoidea divergenter tricornuta, cornubus quam capsula duplo brevioribus ipsis duplo latioribus quam longis rotundato-obtusis. ♂ In Cuba orientali, Mont-

verde (C. Wright n. 1921!). *Tricera retusa* Griseb. Nachricht. d. Königl. Gesellsch. d. Wissensch. d. Univ. Götting. 1865, p. 163. Ramuli circ. 15 cm. longi, paria circ. 6-7 foliorum gerentes, leviter tetragoni. Foliorum limbus 4-5 cm. longus, 15-22 mm. latus, anguste ellipticus, rotundato-obtusius v. obtusus, apice mutico plus minusve distincte emarginatus, basi acutus, nitidulus, lævigatus. Costæ secundariæ primum non prominentes, venæ juniorum foliorum obsolete, demum cum costis secundariis magis prominulæ. Cymæ sessiles. Pedicelli circ. 7 mm. longi. Bractæ parvæ, acutæ. Capsulæ absque cornubus horizontaliter patentibus 7 mm. longæ, 6 mm. latæ. — Similis *B. Vahlîi*, sed folia demum distinctius costata et venosa, haud abrupte mucronato-acuminata et capsularum cornua omnino alia. (v. s.)

2° Flores utriusque sexus sessiles.

10. *B. GLOMERATA*, foliis subsessilibus venosis, floribus sessilibus glomeratis, calycis masc. laciniis oblongato-ellipticis obtusis fem. ovatis acutis puberulis, antheris quam filamenta multo latioribus sesqui v. fere duplo longioribus quam latis papilloso-asperulis, stigmatibus ovario longioribus breviter stipitatis. ♂ In insula Cuba (C. Wright n. 1676! in hb. DC. et in hb. Hook.). *Tricera glomerata* Griseb. in Plant. Wright. p. 157. Rami flavido-cinerei, glabri, tenues quidem sed rigiduli, dense foliosi, breviter ramulosi; ramuli circiter 4-8-phylli. Limbus foliorum  $2\frac{1}{2}$ -3 cm. longus, 5-9 mm. latus, obovato-lanceolatus, acutus, basi cuneato-angustatus, integer, margine recurvus, haud dissimilis ei *B. sempervirentis* γ arborescentis, sed pallidior et vix distincte petiolatus. Calyces utriusque sexus circiter  $1\frac{1}{2}$  mm. tantum longi. Cornua ovarii 2 mm. longæ. — A reliquis speciebus hujus sectionis recedit et floribus omnibus sessilibus et foliis minoribus, quibus manifeste ad sectionem *Eubuxum* approximatur. (v. s.)

β. Costæ secundariæ et venæ omnino indistinctæ.

11. *B. WRIGHTII*, foliis subsessilibus coriaceis evenosis lævigatis, floribus masc. pedicellatis femineis sessilibus, laciniis calycis masc. accrescentibus demum sublineari-lanceolatis acuminatis fem. ter et ultra superantibus, antheris elongato-ovoideis, filamentis inferne latioribus sub apice quam antheræ multo angustioribus, stylis ovario duplo et ultra longioribus. ♂ In Cuba occidentali (C. Wright n. 1928!). *Tricera microphylla* Griseb. in Nachricht. d. Univ. Götting. 1865 p. 163 (non *B. microphylla* Sieb. et Zucc.). *Frutex humilis*, diffusus (Wright). Rami teretes, cinerascens; ramuli tetragoni, breviusculi, cum tota planta glabri. Limbus foliorum  $1\frac{1}{2}$ -2 $\frac{1}{2}$  cm. longus, 3 $\frac{1}{2}$ -5 mm. latus, elliptico-lanceolatus, breviter submucronato-acuminatus, basi acutus, uninerviis, supra nitidulus, intense viridis, utraque pagina evenosus, undique sed præsertim subtus minutissime bullato-puncticulatus. Cymæ circiter 2-7-floræ. Pedicelli breviusculi, limbo foliorum duplo, v. pluries breviores. Bractæ ovatæ, acuminatæ, calycibus multo breviores. Calyx florendi tempore circiter 2 $\frac{1}{2}$  mm. longus, femineus non accrescens, masculus demum 5-6 mm. longus, et ambitu dein multo angustior. Fructus ignoti. — Specimina parvifolia simulat *B. Vahlîi*, sed flores fem. sessiles præter alia differunt. (v. s.)

Sectio II. *EUBUXUS* Baill. Monogr. Buxac. p. 58. — Genus *Buxus* Tournef. Instit. p. 578. t. 345; Linn. Gen. n. 1053; Juss. Gen. p. 388; Schkuhr Handb. t. 287; Gaertn. Fruct. vol. 2. p. 125. t. 108; Lam. Illustr. vol. 4. t. 761. Adr. Juss. Tent. p. 13. t. 1. fig. 3. A; Nees Gen. t. 56; Endl. Gen. pl. n. 5869; Reichenb. Icon. Flor. germ. et helv. vol. 5. t. 153. fig. 4808-9; Plée Types de chaq. famille; Baill. in Bull. soc. bot. de France 1856. p. 285.

Styli ovario breviores; foliorum paginæ facile in duas facie se spectantes secedentes v. præter marginem segregatæ. — Inflorescentiæ partes dense imbricatæ.

## 1° Flores feminei pedicellati.

12. *B. MADAGASCARICA* (Baill. Monogr. Bux. p. 65), foliis obiter reticulato-venosis, floribus fem. pedicellatis, ovario vertice concavo, stylis subulatis bifidis. ♂ In Madagascaria (Dup. Thouars! in hb. Mus. paris.). Frutex ramosus; rami tetragoni; ramuli juniores tenuiter longitrorsum striati. Folia 3-5 cm. longa,  $1\frac{1}{2}$ - $2\frac{1}{2}$  cm. lata, oblongo-ovata v. obovata, obtusiuscula v. obtusa, basi cuneato-attenuata, tenuiter coriacea, lævia, glabra, nitida. Flores dioici?, masculi ignoti, feminei axillares, solitarii. Pedicelli fem. erecti, rigidi, apice incrassati, bracteolas nonnullas decussatas in axilla gemmiferas gerentes. Calyx 5-v. sæpius 4-partitus; laciniae ovatae, acutae, coriaceae, integrae. Styli e basi incrassata subulati, cornuti, apice bifidi, intus longitrorsum sulcati, papilloso. Capsula longe tricornuta. Semina albedo-arillata, nigra, lævia, nitida. — A reliquis hujus sectionis distat axillis ut videtur unisexualibus, turgescit inter bases styliorum sitis rudimentariis nec non floribus fem. pedunculatis; insuper reliquis rigidior est, tota lurido-viridis. (v. s.)

## 2° Flores feminei sessiles, masculi pedicellati.

13. *B. BALEARICA* (Willd. Spec. vol. 4. p. 337), inflorescentiis bracteis steriles suborbiculares scariousas densissimas gerentibus, floribus masc. longiuscule pedicellatis, calycibus dorso apice primum lanato-puberulis, stylis ovarium fere his æquantibus, antheris 4-5-plo longioribus quam latis. ♂ In insulis balearicis (Cambess! in hb. DC.). — Loudon Encycl. p. 704. t. 1378-9; Buch Dec. 6. t. 3. ex Baill.; Baill. Monogr. Bux. p. 62. B. sempervirens v. gigantea Duham. Traité vol. 1. p. 82. t. 23. Major et omnibus partibus multo validior quam B. sempervirens. Ramuli petiolique subglabri. Folia cum petiolo 3-5 mm. longo 3-4 cm. æquantia; sæpius  $1\frac{1}{2}$  cm. lata, sæpe in eodem specimine latiora et angustiora, oblongo-ovata v. oblongo-elliptica v. ovato-lanceolata ellipticave, basi in petiolum brevem angustata, apice rotundato-obtusum v. emarginata, rigida, lucida. Inflorescentiæ foliis 2-3-plo breviores, basi bracteis sterilibus densissimis auctæ. Pedicelli masc. calycem circ. 5 mm. longum subæquantes; filamenta eum fere bis æquantia. Calyx fem. terminalis sessilis a bracteis superatus et quasi obiectus, musculo brevior; laciniae angustae laciniate. Antheræ graciles, lineari-oblongæ, etiam in alabastro ambitu multo longiores quam in affini B. longifolia. Styli erecti, apice patuli. (v. s.)
14. *B. LONGIFOLIA* (Boiss. Diagn. pl. orient. 12. p. 107), inflorescentiis basi densissime bracteatis, floribus masculis breviter pedicellatis, calycis lacinii junioribus dorso lanato-puberulis fructigeris inferne indurato-scariosis, antheris subtriplo longioribus quam latis, stylis elongatis, capsulis longe trirostratis. ♂ In montibus ad meridiem Antiochiæ et inter hanc urbem et montem Cassium sitis (Boiss.!). — Baill. Monogr. Buxac. p. 63. Ramuli juniores leviter puberuli, mox glabrati. Folia  $2\frac{1}{4}$ -4 cm. longa, 7-12 mm. lata, lineari-elliptica v. lanceolata, apice obtusa v. acutiuscula, glabra, firma, nitida. Inflorescentiæ foliis 3-4-plo breviores, simplices. Calycis fem. laciniae sub fructu rigidae, ovatae, longius acuminatae, masculis paulo breviores. Capsulae 8-9 mm. longæ, sublæves, pallidae. — A proxima B. balearica differt præter formam foliorum: antheris fere duplo brevioribus, calyce fem. rigido, pedicellis masc. calyce subtriplo brevioribus. Ei cæterum inflorescentia et consistentia foliorum valde similis est. (v. s.)

## 3° Flores utriusque sexus sessiles.

15. *B. WALLICHIANA* (Baill. Monogr. Buxac. p. 63), floribus utriusque sexus sessilibus, calycis lacinii masculis glabris fructigeris membranaceis, antheris circ. triplo longioribus quam latis, rudimento ovarii brevi, stylis crassis erectis ovarium æquantibus, capsulis longe graciliterque tricornutis. ♂ In India orientali ad Kamaon (Wallich n. 7978!), ubi altitudine 8000 ped. viget, et in summa valle Jumnoh (Jacquem. hb. ind. n. 694), in Himalaya boreali-occidentali (Th. Thomson!). B. longifolia Jacquem Journ. ex Baill. l. c. Ramuli tenuiores quam in præcedente, obscure fulvo-pubescentes, demum gla-

brati. Folia 2-5½ cm. longa, 5-12 mm. lata, lineari-lanceolata, basi breviter angustata, apice truncata v. emarginata, submucronulata, firma quidem sed minus rigida quam in præcedente, basi margine cum petiolis primum pubescentia. Inflorescentiæ simplices, breves. Alabastra glabra, minus quam in præcedente pallida. Calycis masc. laciniae late ovatae, breviusculæ. (v. s.)

16. *B. SEMPERVIRENS* (L. Spec. pl. ed. 1. p. 983), inflorescentiis basi bracteis ovatis acutis præditis, floribus utriusque sexus sessilibus, antheris 2-3-plo longioribus quam latis, rudimento ovarii brevi, cornubus ovario circ. duplo brevioribus.

α, *grandifolia*, arborescens, ramis validis, ramulis dein glabris, foliis oblongo-v. lanceolato-ovatis basi longius angustatis apicem obtusum emarginatum versus paulo angustatis majusculis coriaceis, capsulis subdistincte longius quam alibi in specie 3-cornutis. ♂ In Lusitania (hb. Willd. n. 17381 pag. 41) et in Græcia (de Heldreich n. 2527! in hb. Boiss.), prope Constantinopolin (foliis minus coriaceis, Castagne! in hb. Boiss.) et circa mare nigrum ad Guriel (Szovics! in hb. DC., in aliis hb. pr. parte vix a varietate arborescente distinguenda). Folia 2-3½ cm. longa, 7-16 mm. lata, rigida. — Foliis majoribus subangustis nonnihil ad *B. Wallichianam* et *B. longifoliam* accedit. (v. s.)

β, *rotundifolia* (Baill. Monogr. Bux. p. 61, excl. syn. Linn.), caule arboreo, foliis rotundatis v. breviter ovatis utrinque rotundatis v. basi subattenuatis. ♂ Verisimiliter in hortis colebatur.

γ, *arborescens* (L. Spec. pl. ed. 1. p. 983), caule arborescente v. fruticoso altitudine vario, ramulis puberulis, foliis oblongo-ovatis basi angustatis apice vulgo obtusis v. retusis circiter duplo longioribus quam latis. ♂ In Europa media et præsertim australiore late distributa et præsertim colles calcareos siccis amans (Devos in Bull. Soc. roy. de Bot. de Belgique, v. 5. p. 69), nec non in Caucaso (C. Koch!). — Duham. Traité des arbres et arbustes 1. p. 82. t. 24 et *B. sempervirens* c. fruticosa ejusd. l. c. *B. sempervirens* Auct.; Gaertn. Fruct. vol. 2. p. 125. t. 108; Schkuhr Handb. t. 287; Engl. Bot. t. 1341; Reichenb. Icon. Flor. germ. et helv. vol. 5. t. 153. fig. 4808 et verisim. 4809. *R. arborescens* Lam. Encycl. vol. 1. p. 511. Folia glabra, ad petiolos utrinque ciliolata. Inflorescentiæ simplices, subglobosæ, basi bracteolis sterilibus quam bractæe florigeræ multo brevioribus munitæ, fertiles flores masculos stipantes lacinis calycinis similes iisque æquilongæ. Capsulæ subglobosæ v. breviter ellipsoideæ, more congenerum tricornutæ. — In hortis ludit: foliis argenteo-variegatis (subvar. *argentea* Loud. Encycl. p. 703; Baill. Monogr. Bux. p. 60), vel aureo-maculatis (subvar. *aurea* Loud. l. c., quæ subvar. *aurea maculata* Baill. l. c.), vel foliis in acumen luteum abeuntibus (subvar. *aurea acuminata* Baill. l. c.), vel aureo-puncticulatis (subvar. *aurea puncticulata* Baill. l. c.), vel aureo-marginatis (subvar. *marginata* Loud. l. c., quæ subvar. *aurea marginata* Baill. l. c. p. 61). (v. s.)

δ, *angustifolia* (Loud. Encycl. p. 703), caule arboreo, foliis lanceolatis circiter triplo longioribus quam latis obscurius viridibus. ♂ In hortis, et in montibus Ouled Madhi Algeriæ meridionalis (hb. Boiss.). — Baill. Monogr. Bux. p. 61 et *B. sempervirens* c. *tenuifolia* ejusd. l. c. excl. syn. Boerh. et L. hort. cliff. *B. tenuifolia* hortul. In iisdem speciminibus subinde folia occurrunt quæ ab iis var. arborescentis distingui haud possunt. Folia sæpius 2-2½ cm. longa, 5-8 mm. lata. — Similiter ac var. arborescens ludit: foliis concoloribus aut variegato-punctatis (subvar. *variegata punctata* Baill. Monogr. Bux. p. 61), vel variegato-maculatis (subvar. *variegata* Loud. Encycl. p. 703, quæ subvar. *variegata maculata* Baill. l. c.). (v. s.)

ε, *myrtifolia* (Loud. Encycl. p. 704), caule suffruticuloso humili, foliis parvis oblongis subangustis firmis. ♂ In hortis. — Baill. Monogr. Bux. p. 60. *B. myrtifolia* Lam. Encycl. 1. p. 511. *B. sempervirens* c. *leptophylla* Duham. Traité des Arb. p. 82. t. 23. fig. 3.

ζ, *rosmarinifolia* (Baill. Monogr. Bux. p. 61), caule suffruticuloso ramosis-

simo, foliis polymorphis lineari-lanceolato-spathulatis apice retusis basi angustatis subbullato-puncticulatis subtus albedo-viridibus firmis margine recurvis sæpius 4-6-plo longioribus quam latis.  $\S$  In hortis. Sponte crescentem in Galliæ prov. Saone leg. cl. Hénon (hb. DC.!). Folia 2-2½ cm. longa, 4-5 mm. lata (subvar. *major* Baill. l. c. B. rosmarinifolius Hort. fide Baill.) aut 1½-2 cm. longa, 1-4 mm. lata (subvar. *minor* Baill. l. c. p. 62). B. mucronata Hortul. fid. Baill., cui specimen spontaneum citatum referendum est). (v. s.)

- 7, *suffruticosa* (L. Spec. pl. ed. 1. p. 983), caule suffruticuloso humili subintricato-ramosissimo, ramulis subglabris, foliis ellipticis v. obovatis apice rotundato-obtusos emarginatis basi sæpius acutis submembranaceis longitudine diametrum circiter sesqui æquantibus quam in præcedentibus variet. minoribus.  $\S$  In hortis, vulgatissime pro cincturis arearum hortensium culta. — Loudon Encycl. p. 704; Baill. Monogr. Bux. p. 61. B. suffruticosa Lam. Encycl. vol. 1. p. 511. More præcedentium varietatum foliis ludit aureso-limbatis (subvar. aurea Baill. l. c.). (v. s.)
17. B. JAPONICA, floribus utriusque sexus sessilibus, antheris 2½-plo longioribus quam latis, rudimento ovarii longo calycem subæquante apice disciformi-dilatato et crenato longitrorsum sulcato, stylis crassis abbreviatis.  $\S$  In Japonia (Thunb. in hb. Upsal., Göring n. 254! Zolling. n. 517!). B. sempervirens (lapsu typographico virens) Thunb. Flor. Japon. p. 77 excl. syn. Linn.; Sieb. et Zucc. Flor. japon. fam. nat. p. 34. Ramuli glabri. Folia 10-15 mm. longa, 8-11 mm. lata, obovata, apice emarginata, basi acuta, ad petiolos puberula. Inflorescentiæ subglobosæ, basi bracteis sterilibus ovatis obtusis quam fertiles duplo et ultra brevioribus præditæ. — Tota simillima B. sempervirenti γ arborescenti, sed ovarii rudimento peculiariter evoluto facile recognoscitur. Folia cæterum colore ludunt ut in specie communi europæa. (v. s.)

β, *microphylla*, foliis spathulato-lanceolatis triplo et ultra longioribus quam latis.  $\S$  In Japonia (Göring n. 253! in hb. lugd. bat.). B. microphylla Sieb. et Zucc. Flor. Japon. fam. nat. p. 34; Abhandl. der Mathem. phys. Klass. d. Königl. Akad. d. Wissensch. vol. 4. part. 2. p. 142, ex Walp. Annal. bot. syst. vol. 1. p. 632; Baill. Monogr. Bux. p. 64. B. sempervirens Thunb. Flor. Jap. p. 77 pr. p. ex Sieb. et Zucc. l. c. excl. syn. Kæmpferi ad llicem spectante. — Foliorum longitudo ut in forma normali speciei, sed latitudo tantum 3-7 mm. æquans. Stigmata carnosa, emarginato-biloba. (v. s.)

### *Species non satis nota.*

18. STENOPHYLLA (Hance in Seem. Journ. of Bot. 1868. p. 331), suffrutescens, ramis erectis confertis cortice crasso albo-cinerascente suberoso fungoso fissis obtectis, foliis sessilibus oblongo-lanceolatis basi sensim angustatis apice acutiusculis marginibus incrassatis recurvis glaberrimis supra lucidis venis tenuissimis parum elevatis subtus opacis 6-8 lin. longis 1½-2 lin. latis, floribus dense glomeratis, staminibus calycem paulo superantibus.  $\S$  In Chinæ collibus theiferis An-Koe, versus fines occidentales prov. Fokien (Grijs, Hance exs. n. 6683). — Affinis præcedenti, sed species characteribus expositis haud specificis hucusque non satis stabiliata.

### *Species enucleandæ.*

- B. AUSTRALIS Cunningh.; Steud. Nomencl. p. 242; Baill. Monogr. Bux. p. 66 (sine charact.)
- B. ? CALIFORNICA Hort. ex Baill. Monogr. Bux. p. 66. An eadem ac Simmondsia Californica?

### † *Species a genere exclusæ.*

- B. CHINENSIS Link = Simmondsia Californica.
- B. CORIACEA Spreng. = Sarcococca saligna α coriacea.
- B. SALIGNA Don = Sarcococca saligna β genuina.



†† *Species ab ordine exclusæ.*

B. CORDIFOLIA Spreng. Syst. veg. vol. 3. p. 847; Baill. l. c; quæ eadem ac Tricera cordifolia Willd. Spec. vol. 4. p. 339 (fid. hb. Willd. fol. 17385!) est Myrtacea.

B. DIOICA Forsk. Descr. p. 159, quæ Nageia arabica Willd. Spec. vol. 4. p. 749, et Myrica arabica Vahl Symb. 2. p. 99 (fide specim. Forsk. ! in hb. holm.) est Myrsine Bottensis Alph. DC. Prodr. vol. 8. p. 93, ex Prodr. vol. 16. sect. 2. p. 154.

IV. PACHYSANDRA Mich. Flor. bor. amer. vol. 2. p. 177. t. 45; Adr. Juss. Tent. Euphorb. p. 13. t. 1. fig. 2; Endl. Gen. pl. n. 5870; Baill. Monogr. Bux. p. 155. t. 3. fig. 1-14.

Flores utriusque sexus calyce præditi; laciniaë æstivatione imbricativæ. Petala et discus utriusque sexus nulla. Stamina laciniaë calycis isomera iisque opposita; antheræ dorso supra basin fixæ. Grana pollinis stellatim subreticulata. Rudimentum ovarii evolutum. Ovarium 3-loculare; loculi biovulati. Fructus capsularis. — Herbæ perennes America septentrionalis et Japoniaë. Rami teretes. Folia alterna, exstipulata, ovata, superne grosse serrata. Inflorescentiaë aut in parte inferiore caulis ad axillas squamarum sitæ aut in parte superiore in axillis foliorum evolutorum sitæ, aut dein terminales, spiciformes. Pedicelli masc. bibracteolati v. ebracteolati, feminei pluribracteolati; bracteaë decussatim oppositaë. Flores monoici, calyculi in inflorescentia superioribus, numerosi, feminei inferiores. Calyx masc. decussatim profunde 4-partitus, femineus 4-6-partitus, hujus laciniaë 4 decussatim oppositaë v. 6 decussatim ternatæ. Filamenta valida, elongata, libera. Antheræ anguste oblongo-ovoideæ, fere tota longitudine adnatæ, introrsum longitudinaliter birimosæ, connectivo leviter producto obtuse apiculatæ. Ovarii rudimentum tetragono-obovoideum, vertice depressum. Ovarium parvum. Ovula dein placentam versus hemianatropa remanentia (omnino recurva melius evoluta haud vidi). Obturatores in ovario juvenili nondum evoluti erant, evolutos tamen, in floribus minus juvenilibus cl. Baillon observavit. Styli ovarium pluries superantes eoque crassiores, erecti, apice patuli, subbilobi. Fructus tricoccus, stylis persistentibus tricornutus, mihi haud obviuus, ex cl. Baillon cum iis Buxi conveniens. Semina oblonga, lævigata. — Etiam hic, ut in genere Sarcococca, situs laciniarum calycinarum masc. a præsentia v. defectu bracteolarum geminarum dependet. — A Buxo præter grana pollinis vix nisi characteribus vegetativis et inflorescentia differt.

## \* Inflorescentiaë terminales.

1. P. TERMINALIS (Sieb. et Zucc. Flor. Japon. fam. nat. p. 34), spicis terminalibus, laciniaë calycis masc. orbiculari-ovatis obtusis feminei triangulari-ovatis acutis. ½ In Japonia (Sieb. et Zucc. in hb. Lugd. bat. ! Zolling. n. 518! C. Wright!). — Sieb. et Zucc. in Abhandl. Math. phys. Kl. d. Königl. bai. Acad. d. Wissensch. vol. IV. 2. p. 142; Baill. Monogr. Bux. p. 57. Caules semipedales usque pedales, undique foliati, sed demum casu foliorum basi nudati, cum reliquis partibus glabri. Foliorum limbus 4-9 cm. longus, 3-6 cm. latus, in petiolum 1½-3 cm. longum longe decurrens, e basi cuneata obovatus, obtusus v. truncatus, v. etiam acutus, a medio sursum grosse et remote serratus, subcoriaceus; serraturæ deltoideæ, inæquales, acutæ, adpressæ. Spica terminalis, solitaria, minus multiflora quam in P. procumbente, erecta, subsimplex, i. e. pedicelli tantum uniflori et abbreviati, sed bracteolati. Flores feminei pauci; bracteaë ovataë, acutæ, coriaceæ, magnitudine

sensim in sepala abeuntes. Flores masc. numerosiores, vulgo bibracteolati; bracteolæ late ovatæ, cum sepalis margine tenuiter ciliolatæ, subcoriaceæ, utriusque sexus circiter 3½ mm. longæ. Stamina ut in altera hujus generis specie longe exserta, styli calycem longius superantes, reflexi. — Flores masc. ut in *Sarcococca saligna*, i. e. pedicelli breves, cum calyce decussatim 4-partito et duabus bracteis oppositis ex bracteis axilla evoluti sunt. (v. s.)

**\*\* Inflorescentiæ laterales.**

2. *P. PROCUMBENS* (Mich. Flor. Amer. bor. II. p. 178. t. 45), spicis lateralibus laciniiis calycis masc. late triangulari-ovatis acutis feminei oblongo-ovati cuspidato-acuminatis. ¼ In America septentrionali, in montibus Alleghanies (Mich.), in Carolina (Bosc! in hb. DC.), in Kentucky (hb. berol. ! *Rafinesque!* in hb. Deless. et in hb. Mus. paris.). Etiam in Guadeloupe (Beauportuis) indicatur. — Pursh Flor. of the North. Amer. vol. 1. p. 117; Adr. Juss. Tent. Euphorb. t. 1. fig. 2; Herb. Amat. fasc. 1. t. 3; Bot. Reg. t. 33; Bot. Mag. t. 1964; Loddig. Bot. cab. t. 910; A. Gray Manual of the Bot. of North. Unit. States p. 393. *P. erecta* Baill. Monogr. Bux. p. 56. Radices ad basin caulium subterraneorum valde densissimeque cæspitose-intricatæ. Caules humiles, sæpe longiuscule sub terra repentes ibique squamas pallidas tantum proferentes e quarum axillis progrediuntur spicæ inferiores, dein parte supraterraneæ procumbentes et superne folia alterna v. fere opposita gerentes. Petioli 2-4 cm. longi. Foliorum limbus 4-8 cm. longus, 3-7 cm. latus, late ovatus, basi cuneato-angustatus, apice subobtusius, superne distanter serratus, triplinervius, ad petiolos nervosque puberulus. Spicæ inferiores 8-12 cm. longæ, multifloræ, superiores si adsunt petiolis multo breviores, circiter 2 cm. longæ, ovoideæ. Bractæ laciniiis calycis masc. conformes, 4-5 mm. longæ, bractæ fem. paulo longiores, omnes margine ciliolatæ. Filamenta demum 9 mm. subæquantia, superne sæpe antheras diametro superantia. Ovarium stylis multoties brevius, glabrum. Styli calycem paulo superantes. — *P. erecta* Rafin. sched. ex Baill. l. c. ex ipsissimo specim. Rafin. (in hb. Mus. paris. servato!) nihil est nisi specimen junius nondum bene evolutum et hanc ob rem spicis duplo brevioribus densioribus præditum *P. procumbentis*. (v. s.)

**Tribus II. SIMMONDSIÆ.**

**Loculi ovarii uniovulati.**

- V. *SIMMONDSIA* Nutt. in Hook. Lond. Journ. of Bot. 1844. p. 400. — *Brocchia Mauri* in Tenore Cat. ort. Napol. 1845. p. 80. — *Buxi* sp. Link.

Flores utriusque sexus calyce præditi; lacinia æstivatione imbricativæ. Petala et discus utriusque sexus o. Stamina subnumerosa. Ovarium 3-4-loculare; loculi 1-ovulati. Fructus capsularis. Semina exalbuminosa. — Frutices californici. Folia decussatim opposita, penninervia, basi subquintuplinervia, integra, subcoriacea, viridula, exstipulata, semper-virentia. Indumentum simplex. Flores fem. in axillis foliorum solitarii. Calyces fem. et masc. quincunciales v. feminei biserialim 6-partiti, hi persistentes, foliacei, majusculi. Stamina 10-12, in basi elevata receptaculi inserta; filamenta abbreviata; antheræ longitrorsum birimosæ; connectivum haud productum; loculi longitrorsum adnati. Ovarium magnum, conico-acuminatum, lævigatum, sulcatum, consistentia rigide stramineo-coriaceum, glabrum. Styli 3, simplices, basi confluentes, carnosocoriacei, ovario multo molliores et obscuriores, hirsuto-papilloso. Loculi 1-ovulati vulgo 3, unus ante laciniam secundam calycis, secundus inter laciniam primam et quartam, tertius inter tertiam et quintam situs est. Ovula ex parte superiore, non ex apice loculorum pendula, rhaphe exterior, micropyle interior; obturator indistinctus. Capsulæ loculicide apertæ, sæpe abortu monospermæ, centra columelligeræ.

Cotyledones crassæ, carnosæ, complanatæ. Radicula superior, brevis. — Genus a cl. Nutt. dubitanter Garryaceis adscriptum, hinc autem Putranjivæ affine declaratum, a cl. Lindley ad Euphorbiaceas relatum, cum Buxaceis bene quadrat.

1. *S. CALIFORNICA* (Nutt. in Hook. Lond. Journ. of Bot. 1844. p. 400. t. 16). ♂ In California superiore, in collibus maritimis argillosis sterilibus prope St. Diego et hæud rara in parte meridionali Californiæ, et culta in hort. napolit. (Ten. in hb. DC.) et berol. (hb. reg. berol.) — Torr. Mex. bound. surv. p. 202. t. 49. *Brocchia dichotoma Mauri* in Tenore Cat. ort. napol. 1845. p. 80; Flor. ratisb. 1866. p. 236. *Buxus Chinensis* Link Enum. Hort. berol. 2. p. 386. (1822), quod nomen specificum prioritate gaudet sed falsum est et negligendum. Frutex humilis, diffuse ramosus. Rami divergentes dichotomi v. subdichotome ramosi. Ramuli teretes, juniores cum foliis et calyce fem. juvenili pilis brevibus aureis v. fulvis v. pallidis adpressis parviscule v. densius vestiti, mox autem cum reliquis partibus omnino v. fere omnino glabrati. Folia fere omnino sessilia v. petiolo  $\frac{1}{2}$ -1 $\frac{1}{2}$  mm. longo prædita, 2-4 cm. longa, 12-20 mm. lata, lanceolata-v. lineari-elliptica v.-obovata, subacuta, integra, rigida, utrinque pallide v. lutescenti-viridia, internodiis paulo longiora, margine plana et juniora ciliolata. Costæ secundariæ utroque latere circ. 4-6, infimæ approximate, reliquis longiores, unde limbus indistincte quintuplinervius. Flores masc. axillares, in pedunculis brevibus glomerato-thyrsoidei, feminei axillares, vulgo solitarii, breviter pedicellati, masculis multo majores. Pedicelli fem. calyce amplo pluries breviores, rigidi, pallide subsericeo-pubescentes, apice bracteolis 2-4 circ. 2 mm. longis lanceolato-ovatis onusti. Calyx fem. desloratus 12-18 mm. longus, coriaceo-membranaceus, pubescens v. demum glabratus, fructum juniorem superans. Fructus juvenilis ovoideo-conicus, acuminatus, nitidus, circ. 8 mm. longus, inferne circ. 6 $\frac{1}{2}$  mm. latus. Styli fructu juniore sic descripto triente breviores, fere a basi reflexi, graciles. Fructus maturi ignoti. (v. s.)

*Species non satis nota.*

2. *S. PABULOSA* (Kellog in Proceed. of the Californ. Acad. of nat. sc. vol. 2. p. 21. Oct. 1859). ♂ In fissuris et faucibus rupium maritarum Californiæ. — Frutex arborescens, 3-6-pedalis. Truncus valde flexuoso-curvus, cortice lævi albido tectus. Rami oppositi, horizontaliter patentes, valde intricati et flexuosi, subgrisei; ramuli nonnihil angulosi, griseo-pubescentes. Folia opposita, densa, oblongo-ovata, v. ovato-lanceolata, obtusa, submucronulata, hinc inde obovata, cum apice emarginato, basi nonnihil cuneata,  $\frac{1}{2}$ -1 $\frac{1}{2}$ -pollicaria, supra pallide glaucescentia, cum ramulis junioribus quodammodo succulenta, coriacea, viva subinconspuice trinervia, rigide rugulosa, sicca subtus sæpe ochracea, utraque pagina pubescentia, margine tenui acuto integro leviter scabrida. Petioli valde abbreviati, eglandulosi, hirsuti. Flores masc. numerosi, in cymas dichotomas axillares foliis breviores dispositi, rarius solitarii et sessiles v. etiam glomerulos parvos formantes. Pedunculi et pedicelli bracteati. Calyx fructiger accrescens,  $\frac{1}{2}$ - $\frac{3}{4}$ -pollicaris, basi bracteis 2-4 imbricatis auctus. Petala (sic) florum masculorum 5, imbricata, unguiculata, carinata, viridescencia, intus leviter villosa, eglandulosa, apice obtuso inflexa, ciliata, leviter eroso-dentata. Stamina vulgo 10, petalis breviora; filamenta antheris breviora, e lata basi subulata, interdum 3 v. plura basi cohærentia. Antheræ erectæ. Capsulæ similes nuci Coryli, obtuse triquetrae, stylis 3 subcapitatis terminatæ. Semina leviter amara, cæterum gustu Castaneæ, cum foliis a capris et corvis devorata, unde nomen specificum. — Si vera petala adsunt tum verisimiliter genus proprium.

## Ordo CLXXXI. EMPETRACEÆ (1).

Cisti sect. 3, *Adans. fam.* 2. p. 448. — *Ericis* affinia *Juss. gen.* p. 162. — *Empetridia* (1815) *Rafn. anal. nat.* p. 182 (*solum nomen et excl. Phytolacca*), *fl. tellur. cent.* p. 56. (*excl. plur. gen.*) — *Coniferarum* sectio *Empetreæ* (1818) *Nutt. gen. am.* 2. p. 233. — *Empetrideæ* *S. Fr. Gray* (1821) *arrang. brit. pl.* p. 732. — *Empetreæ* *D. Don, on affin. of Empetreæ in Edinb. new phil. mag.* 1826. p. 59; *Endl. gen.* p. 1105; *Schnizl. ic. fam. t.* 241. — *Rutacearum* subdiv. *Empetreæ* (1828) *Reichb. consp. fr.* p. 196. — *Empetrineæ* (1829) *Dumortier anal. fam.* p. 80. — *Empetraceæ* (1836) *Lindl. nat. syst.* p. 117; *A. Gray, Chlor. bor. am. in Mem. amer. Acad.* 1848. v. 3. p. 13. t. 1; *Nees gen. fl. germ. ic.* v. 1; *Lindl. veg. Kingd.* p. 285; *Ag. f. Theor.* p. 103. t. 10. (*tunica seminis omissa.*)

Flores polygami, in axillis foliorum superiorum vel bractearum sessiles, singuli bracteis cincti. Sepala 3-2, hypogyna, imbricata, bracteis subsimilia Petala 3-2, sepalis alterna, plerumque colorata, æstivatione imbricata. Discus 0. Stamina 3-2, petalis alterna, in fl. fem. plus minus abortientia; filamenta fl. masc. exserta, elongata; antheræ puncto insertæ, loculis 2 medio solum connatis, lateraliter dehiscentibus. Pollen e granis 4 coalitis constans. Ovarium liberum, ovoideum, 2-9-loculare, quando isomerum loculis cum staminibus alternis, in fl. masc. rudimentarium vel nullum. Stylus brevis. Stigmata numero loculorum vel dupla. Ovula in loculis solitaria, angulo interno versus basim aut medium appensa, anatropa, micropyle infera. Drupa pinguis vel fere sicca, mesocarpio copioso aquoso vel parco, endocarpio in pyrenas osseas vel cartilagineas mutato. Semen pyrenæ conforme, albumine carnosum. Embryo axilis, rectus vel paullo curvatus; radícula infera, elongata; cotyledones minimæ. — Fruticuli semperviventes, ericoidei, in frigidioribus et temperatis sparsi; foliis subverticillatis crebris integris, ex plicatura longitudinali subtus rima sulcatis et inde forma lineari, nervo medio sub pagina superiore vix perspicuo; petiolo brevissimo, apice articulato; stipulis nullis; floribus præcoccibus, dioicis vel monoicis, bracteis squamiformibus calyci plus minus simili superpositis, de qua inflorescentia confer ad Buchenau *Bot. Zeit.* 1862. p. 297. t. 10.

Affinitates diu controversæ, sed sine dubio inquirendæ cum ordinibus calyce et corolla donatis, sepalis petalisque distinctis et ovario libero. Juxta Euphorbiaceas et Buxaceas ideo fre-

quenter collocatæ, et equidem non nisi evolutione ovuli (radicula infera, nec supera) diversæ sunt. Eodem caractere ab Illicineis differunt, ubi præterea corollæ sæpius gamopetalæ. A Celastrineis non situ radiculæ, sed defectu disci, arilli et stipularum, cotyledonibusque non foliaceis magis distant. A Rutaceis situ radiculæ, ovulis solitariis atque absentia disci ut et habitu recedunt. Analogiæ quædam cum Ericaceis et Epacridaceis ex forma foliorum (nec eorum plicatura) et polline e granulis coalitis, sed corolla polypetalâ, ovula solitaria, etc., discrepant. Inde post Euphorbiaceas et Buxaceas lubenter adnumero.

I. EMPETRUM *Tourn. inst. p. 519. t. 421; L. gen. n. 775; Endl. gen. n. 5761; Ag. f. theor. syst. p. 103. t. 10. f. 1. 2. (de ovulis); Schumler, icon. fam. t. 241; Buchenau in Bot. Zeit. 1862, p. 297. t. 10. f. 1-10, et alii infra cit.*

Flores in axillis foliorum superiorum solitarii vel (rarius) gemini ternive, subsessiles, prophyllis 2 et bracteis 3-2 squamiformibus cincti. Sepala 3, late ovata, posticum in præfloratione externum. Petala 3, sepalis alterna, obovata, sepalis longiora. Discus nullus. Stamina 3, fertilia, filamento gracili cito accrescente exserta, anthera dorso introrsum filamento inserta, loculis ellipsoideis latere dehiscentibus. Ovarium subglobosum, 9-7-loculare; stylus brevis; stigma 9-6-partitum, lobis pinguibus forma variabili. Ovula in angulo interno cujusvis loculi versus medium inserta, solitaria, pendientia, anatropa. Drupa globosa; mesocarpio aquoso colorato; endocarpio in pyrenas distinctas mutato. Pyrenæ 6-9, extus punctulatæ, cuneiformes, demum distinctæ, dorso convexæ, vertice obtusæ, inferne acutiusculæ, prope basim angulo interno hilo falso notatæ. Semen putamini conforme, latere infra medium putaminis insertum, albumine carnosio copioso. Embryo axilis, gracilis, cylindricus, rectus vel paullo curvatus; radicula infera (ex plerisque auctoribus et obs. mea, invito aliis); cotyledones brevissimæ, lineares, erectæ. — Rami floriferi prophyllis bracteisque planis, interioribus majoribus, donati. Flores sæpe dioici, nunc monoici, feminei cum vestigio staminum, masculi cum rudimento ovarii. Petala purpureo-purpurea. Organa omnia persistentia. De inflorescentia et situ organorum confer ad egregias observationes cl. Buchenau, l. c.

E. NIGRUM (L. sp. p. 1450). ♂ In rupestribus et meridiem versus in uliginosis et paludosis, circa polum arcticum, nempe : in Europa a Spitzberg (Ch. Martins!) et Laponia (L., Wahlenb. etc.) usque ad Hiberniam merid. (Moore et More Cyb. trib. p. 258) et Verviers (Lejeune!), deinde in turfosis, montium Europæ mediæ et merid. usque ad Pyrenæos elatiores (Lap., DC. olim, Bordère! Zetterst. pl. Pyr. p. 231), Alpium merid. (Risso in Mut. fl. fr. 2 p. 271) et Apenninum Lucensem (Giannino in Bertol. fl. it. 10. p. 338); in Caucaso (Bieb. Szowicz! e Gouriél), Sibiria tota (Ledeb. fl. ross. 3. p. 556), Japonia (Sieb. et Zucc. fam. nat.), in insulis Kurils (Ledeb.) et Unalaschka (h. DC!); in America arctica usque ad Sitka (Bongard. Veget. Sitka p. 34) et Nootka (Lag. in h. DC!), montes Novæ Angliæ (Bigel. Boot!), Terram Novam (La Pyl.), et Islandiam (Linds. in Trans. bot. soc. Edinb. p. 142). — Engl. bot. t. 526, Fl. Dan. 6. t. 975; Lam. ill. t. 803, f. 1; Schk. Handb. t. 818; Sv. bot. t. 269; Nees gen. pl. fl. germ. ic. vol. 1 (ubi fig. 22. 23. 25 erroneæ); Reichb. ic. fl. germ. t. 158; Gaud. fl. helv. 6. p. 276, Wahlenb. fl. suec. 2. p. 649; Kirschl. fl. d'Als. 1. p. 157. E. procumbens L. hort. cliff. p. 470. E. crassifolium Rafin. new sylv. p. 51? Fruticulus decumbens vel

paulo adscendens, ramulis parce pubescentibus. Folia approximata, linearia vel oblonga, obtusa, 2-5 mill. longa, 1- $\frac{1}{2}$  mill. lata, rigida, ad lentem scabro-puberula, demum glabra, supra planiuscula, lateraliter rotundata, subtus ex marginibus reflexis conniventibus ciliatis sutura pallida nunc depressa nervum mentiente fere a basi et fere usque ad apicem percursa, unde folium sectum vacuum videtur, nervo tamen vero in medio præditum! Bacca matura magnit. Juniperi, nigra, ut dicunt botanici, potius intense rubro-violacea, nam color niger vix in plantis adest. (v. v.)

$\beta$ , *purpureum*, bacca rubra, pube foliorum.... In America artica, cum vero E. nigro mixtum, ad fretum Smith (Durand, enum. arctic. plant. coll. by Hayes, in Proceed. Acad. Nat. sc. Philad. 1863. p. 95), Labrador (Rafn., Ferland, ibid.) et Terra Nova (La Pyl. et Tuckerm. ibid. cit.). E. purpureum Rafn. New Sylva. p. 50. E. rubrum Willd., an var. E. nigri? (Durand l. c.). Identitatem cum forma australi affirmat cl. Durand; num vero folia magis pubescentia quam in vulgari fructu nigro non dixit.

$\gamma$ , *Andinum*, glabriusculum, bacca rubra? In montibus Chili, ad Chillan (Germain!) et alibi (Philippi!). E. Andinum Philippi! in h. meo. E. rubrum Hohen! in Philippi pl. chil. n. 90. Ramuli et folia glabriuscula ut in plerisque speciminibus hemisphærii borealis E. nigri, sed ex nomine ab Hohen. imposito et patria bacca verisimiliter rubra. Cl. Hook. f. in Fl. Antarct. rubrum in Chili merid. adesse ait, quod forsitan ad seq. var. referendum. (v. s.)

$\delta$ , *rubrum*, bacca rubra, ramulis puberulis, foliis plerumque ad margines præsertim lanuginoso-pubescentibus proce glabriusculis. — In Americæ antarcticæ, ins. Falkland (Darw.!) et Terra ignea (Hook. f. ! Lechler! n. 1057); in ins. Tristan d'Acunha (Pet. Th. fid. descript.). E. rubrum Willd. sp. 4. p. 713 (præeuntibus Vahl et Soland. in mss.); Lindl. bot. reg. t. 1783; Hook. f. fl. antarct. 2. p. 345. E. rubrum Pet. Th. fl. Tristan d'Acunha p. 39 (ubi descriptio fusa quæ cum speciminibus Americæ australis convenit). E. medium Carmich. descr. of Tristan d'Acunha p. 26. Pubes ejusdem naturæ quam in  $\alpha$ , sed in foliis plerumque longior et copiosa, in specimine Terræ igneæ a cl. Hook. f. comm. deficiens. Ut clar. bot. kewensis nullum characterem vere specificum detexi inter hoc et E. nigrum vulgare. (v. s.)

### *Species exclusæ.*

E. ACETOSUM Rumph. = Begonia tuberosa Lam.

E. ACICULARE Bertol. = Ceratiola ericoides.

E. CONRADII Tuckerm. = Corema Conradii.

E. PINNATUM Lam. = Margyricarpus setosus Ruiz et Pav.

II. COREMA D. Don in Edinb. n. phil. journ. oct. 1826. p. 63; A. Gray Chloris Americ. bor. in Mem. Amer. Acad. n. ser. 3. p. 14. t. 1. — Empetrum sp. Linn.; Gært. fr. 2. p. 167. t. 106. — Euleucum Rafn. fl. tellur. cens. 7. n. 634 (anno 1836) — Colema (errore pro Corema) Rafn. new sylva p. 49 (ubi sectiones 2 Euleuca et Endamnia) — Oakesia Tuckerm. in Hook. Lond. Journ. bot. 1842. p. 443. — Tuckermannia Klotzsch in Wieg. Archiv. 1842.

Flores glomerati, bracteis squamiformibus cincti. Sepala bractei-formia, 3 vel rarius 2, irregulariter fimbriata. Petala 3 vel rarius 2, sepalis tenuiora magisque fimbriata, colorata. Stamina sæpius 3, in fl. fem. plus minus abortientia. Ovarium sæpius 3-loculare. Stylus cylindricus. Stigmata linearia 3, nunc 2, 4, 5. Ovula loculorum angulo interno versus basim inserta. Drupa exsucca, globosa, pyrenis plus minus induratis. — Eruticuli europeo-americi, capitulis terminalibus; bracteis sepalis-

que caducis; petalis eleganter fimbriatis roseis; partibus omnibus in autopsia caducis, sæpeque numero variabili, sed flores regulariter trimeros frequenter vidi.

1. C. ALBUM (D. Don l. c.), ramis erectis, foliis elongatis, pyrenis osseis.  $\frac{3}{4}$  In locis præsertim arenosis littoralibus Hispaniæ austro-occid. (Willk. Strand und Stepp. Iber. p. 151), Lusitaniæ (Bourgeau! 1946, Welw. l. 111), Galiciæ (Colm. in Willk. cit.), ins. Azorensium Pico (Wats. in Hook. journ. 1844. p. 605) et forsan insul. Canariensium (Brouss. l. in h. DC. et h. Bouchet monsp. ubi schedulæ aiunt Teneriffæ, sed sæpe cl. viator specimina e Mogador, vel hispanica, cum canariensibus miscuit, et planta in Webb. fl. Canar. deest). *Empetrum album* L. sp. p. 1450; Gærtn. fr. t. 106 (ubi fructus); Lam. ill. t. 803 (e Gærtn. iter.); Torr. in Loud. gard. mag. t. 17, a Gray l. c. p. 56. *Eulucum album* Rafin. fl. tell. n. 634. *Colema* sect. *Euleuca* Rafin. new sylv. p. 49. Rami tenuiter pubescentes. Folia 5-10 mill. longa, 1 mill. lata, rigida, ad lentem passim aspero-puberula, per 2 annos perennantia. Bracteæ exteriores lanceolatae, 1-3 mill. longæ. Sepala obovata. Petala angustius obovata, sublongiora. Antheræ millim. longæ. Drupa 3-4 mill. longæ, sæpius alba, nunc (ex Brot. fl. lusit.) purpurea. (v. s.).
2. C. CONRADII (Torrey in Loudon gard. mag. v. 17, ex Gray l. c. p. 56), ramis diffusis basi prostratis inde adscendentibus, pyrenis cartilagineis.  $\frac{5}{8}$  In arenosis et aridis Novæ Angliæ (Oakes! Tuckerm. l. A. Gray!) et Terræ Novæ (Cormack, in h. olim Lamb. fide Tuckerm. in A. Gray chloris p. 4). *Empetrum Conradii* Torr. in Ann. Lyceum N. York 4. p. 83. *Oakesia Conradii* Tuckerm. in Hook. Lond. journ. of bot. 1. p. 446, Hook. ic. t. 501, A. Gray l. c. p. 4. t. 1. *Colema* (sect. *Endammia*) *arenaria* Rafin. new sylv. p. 49. *Tuckermannia Conradii* Klotzsch l. c. Fruticulus prostratus, ramosissimus, parce puberulus. Folia 4-5 mill. longa,  $\frac{1}{4}$ -1 mill. lata. Capitula similiter C. albæ minor, bracteis latioribus. Flores non solum polygami, sed quoad numerum partium variabiles, ut ex fusa descript. et ic. cl. A. Gray patet. Sepala 3-2, nec nisi ex analogia cum affini a bracteis distinguenda, uno externo majore. Loculi sæpius 3, nunc 4. (v. s.).

III. CERATIOLA Mich. fl. bor. am. 2. p. 221; Pursh fl. Am. sept. 1. p. 21 t. 13. (mala); Hook. in Bot. mag. t. 2758; A. Gray Chloris bor. am. in Mem. amer. Acad. n. ser. 3. p. 14.

Flores in axillis folii vel bracteæ 1-2-3, sessiles, singuli bracteolis 2 ovatis stipati. Sepala 2, bracteolis similia, iis alterna et longiora, imbricata. Petala 2, arcte ovario adpressa, imbricata, sepalis alterna iisque longiora et tenuiora. Stamina 2, petalis alterna, in fl. fem. abortientia. Ovarium ovoideum, biloculare. Stylus cylindricus. Stigmata 3-4, lineari-lanceolata, irregulariter passim dentato-lobata. Ovula in singulo loculo lateraliter inferne inserta. Drupa fere exsucca, ovoidea, organis floralibus basi cincta, pyrenis 2. Semen latere appensum. — Fruticulus Americæ borealis; foliis sulco inferiore foliorum ex reduplicacione limbi magis hiantes quam in aliis generibus.

C. ERICOIDES (Mich. l. c.). — In aridis sabulosis Georgiæ et Carolinæ (Micht. Bosc! etc.) et Alabama (Gates in Bertol.). *Empetrum aciculare* Bertol. misc. 12. p. 20. t. 1. f. 2 (pessima). Rami griseo-subtomentosi, ad origines foliorum torulosi. Folia centim. longa,  $\frac{1}{4}$  mill. lata, glabriuscula, margine obtusa, subtus interne pubescentia. Flores sæpius gemini, vix 2 mill. longi. Bracteolæ ovato-acutæ,  $\frac{1}{4}$  mill. longæ. Sepala ovata, longiora, petalæque obtusa et margine pubescenti-imbricata. Stamina exserta. Drupa 2 mill. longa (v. s.).

## Ordo CLXXXII. CANNABINÆ (1).

Castaneorum sectio III (cum Urtica, Moro, Ulmo, etc.) *Adans. fam. nat.* 2. p. 376 (1763). — Urticarum pars *Juss. gen.* p. 304 (1789). — Urticæ Cannabineæ *Blume bijdr.* p. 515 (1825); *Gaudich. voy. Freyc.* p. 507; *Kunth. fl. ber.* 2. p. 188. — Urticæ trib. Cannabineæ et Humulineæ *Dumort. anal. fam.* p. 17 (1829). — Lupulaceæ *Raspail phys. veg.* 2. p. 495. (1837). — Cannabineæ *E. Mey. Preuss. Gattung.* p. 80 (1839); *Endl. gen.* p. 286 et 1376; *Meissn. gen.* p. 348, *comm.* p. 259; *Brongn. enum.* p. 99; *Payer organ. t.* 61; *Schnizl. ic. fam.* n. 95; *Lemaout et Decsne Traité de bot.* p. 507. — Cannabineæ *Lindl. veg. Kingd.* (1846). — Cannabisaceæ *Ser. descr. Muriers* p. 188 (1855). — Cannabisaceæ *Al. Braun in Asch. Fl. Brandenb. Einleit.* p. 58 et *Asch. ibid.* p. 611 (1864).

Flores dioici. MASCULI. Perigonium e tepalis 5, liberis, imbricatione quincunciali constitutum, uno externo axim spectante (*Schnizl. Payer ll. cc.*). Stamina 5, tepalis opposita, libera, non inflexa; filamentis brevibus; antheris apice filamentis insertis, oblongis, bilocularibus, longitudinaliter dehiscentibus; polline lævi, (in aqua) globoso. FEMINEI. Perigonium membranaceum, tenerrimum, monophyllum, ovarium, amplexans, persistens. Ovarium liberum, uniloculare. Stylus brevissimus. Stigmata 2, elongata, linearia, sæpius inæqualia, undique papillosa, erga bracteam antice et postice sita (*Payer, Schnizl.*), cum stylo caduca. Ovulum unicum, pendens, campylotropum. Caryopsis in perigonio sacciformi tenuissimo inclusa. Semen exalbuminosum. Embryo curvatus. Radicula supera, dorso cotyledonis incumbens. Cotyledones lineares, radiculam longitudine subæquantes. — Herbæ in hemisphærio boreali crescentes, graveolentes, scabro-pilosæ, succo aqueo; foliis palminerviis, fasciculis continuis (monomeris, sensu a *Cas. DC. théor. feuille proposito*), sæpius lobatis, oppositis superioribusque sæpe alternis; stipulis persistentibus; inflorescentia e cymulis axillaribus et terminalibus; mascula laxa; feminea condensata, bracteis et bracteolis persistentibus.

De evolutione florum et ovulorum confer ad *Schleiden in Wiegmann. Archiv.* vol. 5. et *Beitr.* p. 40. t. 2. f. 19. 20; *Payer organ. t.* 61; *Gasparr. ricerche sull. embriol. del Canape, in Mem. bot.* p. 1. t. 1-3. — De inflorescentia confer ad *Wydł. in Flora* 1844 p. 735, 1851. p. 435, 1865. p. 318, et *Irmisch in Bot. Zeit.* 1848. p. 793.

---

(1) Auctore Alph. de Candolle.



- I. HUMULUS L. gen. ed. 1. p. 304; Juss. gen. p. 404; Lam. dict. 3. p. 138, ill. t. 815; Nees jun. ic. gen. fl. germ. fasc. 3 (perigonio fem. omisso); Kunth fl. ber. ed. 1838. v. 2. p. 189; Schleid. in Wiegmann Archiv. 5. p. 229. t. 7, Beitr. t. 2. f. 20; Endl. gen. p. 286, 1376; Asch. Fl. Brandenb. p. 611. — Lupulus Bauh.; Tourn. inst. p. 535. t. 309; Gärtn. fruct. t. 75. — Cannabis sp. Scop. Carn.

Stamina erecta. Embryo circinato-involutus. — Herbæ perennes, volubiles; stipulis connatis (supremis sæpe bifidis); inflorescentiæ masc. laxæ, bracteis lanceolatis; femineæ condensatæ, bracteis et stipulis amplis, distinctis, limbo medio fere arbortiente aut iis multo minore formatis, bifloris; bracteolis (stipulæ connatæ) quemque florem cingentibus, ovatis vel lanceolatis; bracteis bracteolisque plus minus in strobilo fructifero auctis. — De materie crystallina in textu foliorum confer ad Schleid. Grundz. p. 341. f. 103.

§ 1. *Strobili maturi bractæ et bracteolæ valde accretæ, scariosæ, glandulis luteis reniformibus lupulinam præbentibus basi externa scatentes. Caryopsis perigonio membranaceo glandulifero brevior.*

1. H. LUPULUS (L. sp. p. 1457), ramis petiolis foliisque supra retrorsum aculeato-vel scabro-pilosis, foliis cordatis ovatis vel rotundatis plerisque 3-5-fidis, superioribus indivisis, omnibus crenato-dentatis, strobilis ovoideis, bracteis junioribus laxis ovatis, demum amplis scariosis extus basi glanduligeris. ♀ ☉ In sepibus et dumetis Europæ, ab Angliæ sept. (Wats. Cyb. 3. p. 372), Scania (Hartm. fl. p. 195), Fennia et Rossia media (Ledeb. fl. 3. p. 635), ad Hispaniam centr. (Willk. et Lange Prodr. p. 254), montes demissos Sardinia (Moris fl. 3. p. 507), Sicilia (Guss. syn. 2. p. 608), Peloponesi (Orphan. 3626 in h. Boiss.), in Sibiria altaica, baikalensi et orientali (Ledeb. l. c.), prov. Caucasica (id.), Lazistan (h. Boiss.), circa Damas (Gail. 2235. in h. Boiss.), Karabagh (Szovicz!). In America boreali orient. sed ibi introd. (A. Gray bot. north. U. St. ed. 5. p. 446) et in Novo Mexico (Wright l. n. 1860 in h. Boiss.), ubi veris. culta vel spontanea facta. Colitur propter strobilorum vires, ad cerevisiam parandam. — Sm. Engl. bot. t. 427; Schkuhr Handb. t. 326; Fl. Dan. t. 1239; Svensk bot. t. 332; Hayne Arzn. Gew. 8. t. 36; Nees ic. pl. offic.; Reichb. ic. fl. germ. 12. t. 656; Nees jun. ic. gen. fl. germ. fasc. 3. Cannabis Lupulus Scop. Carn. 2. p. 263. Lupulus communis Gärtn. fr. t. 75. H. Americanus Nutt. in Journ. Acad. sc. Philad. n. ser. v. 1. p. 181 (fide num. Wrightii ibi cit. et Torreyi in Emory-Mexic. bound. p. 203). Turiones edules. Caulis (e centro spiræ obs.) dextrorsum involutus. Petioli limbo breviores. Lobi fol. ovato-acuminati vel acuti. Pagina inferior glabriuscula. Crenæ apiculo donatæ. Pedunculi petiolum subæquantes. De glandulis et lupulina confer ad Raspail nouv. syst. chim. org. ed. 2. p. 183; Personne in Journ. de pharm. ser. 3. v. 26; Mehu, étude sur le houblon, thèse, Montpellier. 1867. Culta varietates ex colore surculorum, forma et colore strobili præbet. (v. v.)

§ 2. *Strobili maturi bractæ et bracteolæ paulo accrescentes, non scariosæ, nec glanduligeræ. Caryopsis perigonium non superans.*

2. H. JAPONICUS (Sieb. et Zucc. fl. jap. fam. nat. 2. p. 89), ramis petiolis foliisque supra retrorsum aculeolato-vel scabro-pilosis, foliis ovatis rotundatisve profunde 5-7-fidis crenato-dentatis, petiolis elongatis, strobilis paucifloris bracteis et bracteolis deltoideo-acuminatis margine hispidis parum ampliatis foliaceis acuminatis eglandulosis. ♀ ☉ In Japonia (Thunb. Sieb. Maxim.)

in h. Boiss.), archipelago Coreensi (h. Kew., fide Benth. in lit.), circa Pekin, (id. ibid.), in China merid. (Seem., Hance in h. Kew. ex Benth. lit.) et Formosa (Oldham! 515 in h. Boiss.). — Seem. bot. Herald p. 512. t. 98. H. *Lupulus* Thunb. fl. jap. p. 113, non L. Planta ins. Chusan quam Cantor (Ann. and. mag. of nat. hist. 9, p. 271). H. *Lupulum* vocavit, propter patriam et H. *Lupulo* in China ex herb. Kew. deficiente, hic referenda videtur, ut et H. *aculeatus* Nutt. in Journ. of Acad. sc. Philad. ser. 2. v. 1. p. 182, e Canton Sinensium. Folia etiam suprema profunde lobata. Petioli 5-9 centim. aut in foliis infer. 2 decim. longi, limbo longiores; pagina inferior limbi puberula. Racemi fl. masc. elongati, laxiflori, pilosi. Perigonii masc. tepala lanceolata, extus hispida. Perigonium fem. (ex ic. Seem. et fructu jun. in h. Boiss.) ovarium omnino tegens. Strobilus maturus 1-1½ poll. longus, globosus vel ovoideus; bracteis 5-8 mill. longis, margine et interdum dorso pilis rigidis hispidis. Nuculæ ovato-lentiformes, 2-3 mill. longæ, læves, flavæ, passim obscure maculatæ. In specimine ins. Formosæ plures vidi bracteas ubi limbus lanceolatus, 2-4 millim. longus, dentatus, inter 2 lobos ampliores stipulaceos adest. (v. s.)

- 3 ? H. *CORDIFOLIUS* (Miq. in Ann. Mus. Lugd. bat. 2. p. 133), caule petioliis pedunculisque retrorsum minutissime aculeolatis, foliis ovatis cordatis breviter mucronato-acuminatis indivisis dentato-serratis subtus in nervis parce scabriusculis in parenchymate glandulis lepidiformibus puncticulatis, petiolis brevibus, strobilibus fem. ovoideis bracteis laxis ovato-triangularibus acuminatis. ♀ In Japoniæ montibus prov. Kiso. Petioli limbo duplo fere breviores. Limbi 2-2½ poll. longi, supra puncticulati (ex pilis deciduis?), subtus glandulis orbiculatis lepidoti. Flores masc. ignoti. Capitula fem. juniora globosa, magnit. nuclei cerasi, bracteis ovato-triangularibus extus puberulis, exterioribus basi extus albidopubescentibus. Pedunculus petiolo æqualis vel brevior. Strobili maturi 1½ poll. longi, bracteis maturis glabris et obtusioribus. — Nucula non descripta.

II. *CANNABIS* Tourn. inst. p. 535. t. 309; Linn. gen. ed. 1. p. 304; Gærtn. fruct. t. 75; Schkuhr. Handb. t. 325; Nees jun. ic. gen. fl. germ. fasc. 4 (omisso perigonio); Schleid. in Wieg. Arch. vol. 5, Beitr. p. 40. t. 2. f. 19 (de ovulo); Endl. gen. p. 286 et 1376; Payer organ. t. 61; Schnizl. icon. fam. t. 95; Asch. Fl. Brandeb. p. 611; Gasparr. sall. Canape in Mem. bot. p. 1. t. 1-3.

Stamina pendientia. Embryo genuflexus. — Herba annua, erecta, cujus femina elatior; fibris flexilibus, tenacissimis; stipulis liberis; foliis 5-7-9-sectis; inflorescentiæ masculæ laxæ bracteis linearibus, supremis sæpe deficientibus, femineæ bracteis etiam in cymula foliaceis, stipuligeris, bifloris; bracteolis quemque florem cingentibus ovato-lanceolatis, extus glanduloso-velutinis, e stipulis connatis formati, deinde fructu incluso paulo longioribus; materie glutinosa inebriante in inflorescentia præsertim scatente. — Perigonium, ex Schnizl., sæpe abortiens.

C. *SATIVA* L. sp. p. 1457; Lam. dict. 1. p. 695; Hayne Arzn. Gew. 8. t. 35; Spach ic. veg. phan. t. 133; Reichb. ic. fl. germ. 12. p. 11. t. 655; Dietr. fl. bor. t. 720, et aliæ ic. supra cit. C. *Indica* Lam. dict. l. c. C. *erratica* Siev. in Pall. ex Ledeb. fl. ross. C. *Chinensis* Del. ind. sem. h. Monsp. 1849; C. Koch in ind. sem. h. ber. 1853 (et Ann. sc. nat. ser. 4. v. 1. p. 352). Polygonum viridiflorum Poir. in Dict. enc. 6. p. 140 (fide Meissn. in Prodr. 14. p. 143). Kalengi Rheede Mal. 10. t. 60. ① In Asia temperata certe spontanea, ex gr. in pratensibus ad Irtysch et in deserto Soongoro-Kirghisico (Kar. et Kir. 1043, Bull. soc. Mosc. 1854. v. 1. p. 367), ad radices montium transbaicalensium (Turcz.!), olim ab indo-europæis culta et sub nominibus quorum radix an servatur per Asiam et Europam diffusa (Alph. DC. geogr. bot. p. 833), nunc ubique propter fila textilia aut vires inebriantes culta et passim fere spontanea facta. Qui formas australiores sedulo observaverunt, at Christison (On C. indica, in Ann. and Mag. of nat. hist. 1850. v. 5. p. 483 et

Trans. bot. soc. Edinb. 4. p. 59) et Dukerley (Note sur le kif, in Bull. soc. bot. Fr. 13. p. 401), diversitates specificas negant, sed formæ adsunt plus minus hereditariæ, nec non indoles, pro cultura et temperie mutata, quæ sub-varietates, aut variationes videntur potius quam varietates, nempe :

- α, *Kif*, caule 1-3-pedali, foliis oppositis approximatis, inflorescentia sub-condensata, semine minore quam in vulgari fulvescente venis reticulatis pallidioribus maculisque præterea nigricantibus consperso. — Culta in Algeria sub nom. *Kif* vel *Tekrouri*, propter inflorescentias admodum inebriantes (Dukerley l. c. ubi descriptio fusa). Specimen Tuneto cultum possideo, quod humile et habitu toto præsentis, ubi vero semen immaculatum ut in sequentibus.
- β, *vulgaris*, caule 3-6-pedali, foliis oppositis vel alternis, semine flavo-griseo venis reticulatis pallidioribus maculis nigris nullis vel paucis. — Spontanea locis supra cit.; spontanea? vel culta in India sept. (Roxb.! Wall. list n. 4665, Hook. f. et Thoms.!), spontanea ad Tong Dong Birmanorum (Wall.! cat. ined. itin. birm. n. 881), in Europa vulgo culta. C. sativa et C. Indica, auctorum supra cit. C. orientalis h. Noé 11. 62 in h. Boiss.! In speciminibus asiaticis folia sæpius alterna videntur, quæ in nostris cultis sæpius opposita, nunc alterna. Vires inebriantes pro copia glandularum infloresc. admodum variant. Semina in spec. sibiricis minora, in ipso specimine 1043 Kar. et Kir. herb. mei non maculata, vel (ut in α) nigro maculata; in indicis ut in cultis europæis magnit. vulgari fulvo-grisea, venis pallidioribus percursa, non præterea maculata. (v. v.)
- γ, *Pedemontana*, caule 10-20-pedali, cæteris ut in vulgari. — Colitur in pinguibus Europæ merid., ex gr. Bononiæ, Pedemontii, Aquitanix (Duch. in Bull. soc. bot. 13. p. 406).
- δ, *Chinensis*, caule 18-20-pedali, ramis diffusis, foliis sæpius alternis, segmentis 7-9 anguste lanceolatis, cæt. ut in vulgari. — E China in hortum vindob. anno 1827! Galliam merid. (Delile) et circa Parisios (Vilm. Bonjard.) introducta. C. Chinensis Del. et C. Koch ll. cc. Ex optimis specim. a me olim lectis in h. vind. semina die 5 sept. matura erant, vulgari viz minora, colore simili, Parisiis vero semina minora, plerumque sterilia fuerunt, tandem Berolini non maturaverunt et glandulis resinosis caruerunt. (v. v.)

## Ordo CLXXXIII. ULMACEÆ.

In futuro et ultimo volumine XVII<sup>o</sup> inquirendæ.

## Ordo CLXXXIV. ARTOCARPEÆ.

Cum Ulmaceis ad futurum volumen remissæ.

## Ordo CLXXXV. URTICACEÆ (1).

Castanearum sectionis genus *Adans. Fam. des pl.* 2. p. 376. — Scabridearum genera *Linn. Ordin. nat.* p. 593. — Urticarum genera *Juss. Gen. pl.* p. 400. — Urticearum genera *DC. Fl. fr.* 3. p. 321; *Agardh Aphor. bot.* p. 203; *Bartl. Ordin. nat.* p. 105; *Lindl. Nat. Syst. ed. 1.* p. 93; *Martius Conspect. regn. veg.* p. 13. — Urticeæ *R. Br. Obs. pl. Cong.* p. 35; *Brongn. Enum. gen. pl. cult. Mus. par.* p. 98; *Adr. de Juss. Cours élém. bot.* p. 592. — Urticearum tribus *Dumort. Anal. fam.* p. 17. — Urticeæ *Endlich. Prodr. fl. Norf.* p. 37; *ejusd. Gen. pl.* p. 282; *Meissner Pl. vasc. gen.* p. 348; *Lindley Veget. Kingd.* p. 260; *Wedd. Monogr. Urt. in Archiv. Mus. hist. nat. Par.* 9. p. 49.

Flores apetalæ, ut plurimum diclini monoicique v. dioici, perraro polygami (in gen. *Achudemia* et *Parietaria*), regulares aut irregulares. *Masc.* Perigonium viride rariusve coloratum (albidum v. rubens) et fere petaloideum, gamophyllum, 4-5- aut rarissime 2-v. 3-partitum, interdum monophyllum; segmentis æqualibus concavis, haud raro sub apice v. summo dorso mucronatis aut (sed minus frequenter) transversim cristatis, in præfloratione valvatis aut (saltem apice) imbricatis; alabastro depresso et subumbilicato aut acuminato vel mucronato. Stamina segmentis perigonii numero æqualia et opposita s. superposita, fundo perigonii inserta; filamentis crassiusculis, basi sæpe complanatis, superne plerumque attenuatis immo et antice sæpe transversim rugosis, juxta insertionem nonnunquam perigonio atque etiam inter se aliquantulum connatis, æstivatione inflexis, sub anthesi elastice patentibus; antheris bilocularibus, introrsis, fere medio dorso ope connectivi carnosuli adfixis, loculis contiguis sæpius oblongo-reniformibus basi et apice discretis rima longitudinali dehiscentibus; poline pulveraceo, e granulis sphaericis tenuissimis pallide luteis constante. Pistilli rudimentum lineari-conicum v. obovoideum v. clavatum v. cupuliforme, glabrum aut basi plus minus pilo-

---

(1) Auctore H. A. Weddell.

sum lanatumve. *Fem.* Perigonium calycinum tubulosum aut 3-5-lobum v. partitum (segmentis æqualibus inæqualibusve), post anthesin sæpissime accrescens, raro nullum. Ovarium liberum v. perigonio tubuloso plus minus adhærens connatumve, sessile v. breviter stipitatum, glaberrimum aut (deficiente simul perigonio) hispidum setulosumve, sæpius ovatum v. lanceolatum, imo loculo v. juxta fundum illius ope funiculi distincte erecti vel adscendentis superneque dilatati adfixum, apice ad basim styli v. stigmatis spectans summoque loculo persæpe cohærens; integumento duplici. Stylus nullus s. a stigmate ipso non distinctus v. brevis, rarissime (interdum in *Parietaria*) elongatus, simplex. Stigma polymorphum, capitatum v. oblongum v. spathulatum v. lineare, rectiusculum aut inflexum, sæpe unilateraliter papillosum, papillis brevibus prælongisve paucisque aut numerosis, nonnunquam fere discoideum aut plane penicilliforme, rarissime (in specie unica) brachiatum. Stamina rudimenta nulla v. (in *Procrideis*) squamiformia, inflexa, segmentis perigonii superposita. Fructus exsuccus (achænium) v. baccatus (drupa), modo nudus, scilicet perigonio subimmutato solum stipatus, modo eodem aucto sicco aut carnoso coloratoque et interdum suberoso sulcato aut varie appendiculato pro parte vel integre vestitus aut etiam cum illo connatus. Achænia sæpe plus minus compressa, ovoidea v. sphærica, recta v. obliqua, rarissime omnino asymmetrica, lævia v. punctulata v. tuberculata, nonnunquam anguste alata interdumque setosa aut (more seminis *Gossypii*) longissime lanata; pericarpio plerumque tenui submembranaceo v. fragili, rarius crustaceo, fuscescente v. atro aut etiam eburneo, opaco v. nitidissimo. Drupæ ut plurimum ovoideæ v. obovoideæ, in capitula densa moriformia vulgo aggregatæ coloremque luteum v. rubentem afferentes; endocarpio sæpius crustaceo, sarcocarpio plerumque e cellularum strato unico a perigonio distincto v. cum illo coalito constante. Semen ab endocarpio discretum eique conforme; testa tenuissima fusca, chalaza obscure fuscescente basi late notata. Albumen raro copiosum, sæpius parcum quandoque (sed raro) prorsus abortiens, oleosum, albidum. Embryo rectus, in albuminis axi antitropus ejusdemque fere longitudine, aut (deficiente albumine) cavitatem totam seminis explens; cotyledonibus carnosiss, applicatis, ovatis v. suborbicularibus extusque convexis; radícula cylindrica v. conoidea, supera; gemmula parum conspicua. — *Hermaphr.* Perigonium (post anthesin nonnunquam accretum) ut in flor. fem. aut masc. Stamina ut in flor. masc. Pistillum ut in flor. fem. — Herbæ annuæ perennesve v. suffrutices v. frutices (rarissime scandentes) v. interdum arbores, regionis fere nullæ omnino exsules, sed in tropicis et subtropicis totius orbis præsertim obviæ, odore (si ullo) herbaceo, succo aqueo v. rarissime (in *Neraudia*) lacteo; ramis junioribus nec non

caulibus herbaceis sæpissime angulatis; ligno molli sæpeque levissimo; cortice tenui, fibris tenacibus farcto; epidermidis cellulis permultis ampliatis singulisque concretione calcarea s. cystolitho parieti interno appensa expletis. Pili (si adsunt) simplices et innocui aut rarius (in *Urereis*) basi strato exteriore cellularum aucti urentesque. Gemmæ axillares, solitariae v. geminae v. ternae, aliæ floriferae, aliæ folia conduplicata stipulisque persæpe velata exhibentes. Folia decidua aut persistentia, alterna v. opposita, rarissime quaternata, interdum (etiam apud species oppositifolias) disticha ac inæquilatera et in eodem jugo (s. pare) haud raro etiam inæqualis magnitudinis, ut plurimum petiolata, simplicia, penni-sæpiusve tri-triplo-palmati-nervia (1), integra aut varie dentata rarissime palmatoloba. Stipulae ad basim petioli geminae, caulinae vel petiolares vel axillares, liberae v. inter se aut cum illis folii oppositi magis minus connatae. Flores sessiles v. pedicellati, constanter cymosi; cyma rarissime ad florem unicum redacta, nec sæpe e floribus pluribus in receptaculo communi convexo aut concavo cœgestis constante, vulgo multiflora et varie ramosa, interdum regulariter dichotoma s. bipara, sæpius autem unipara (ramis postgenitis axi primordiali plus minus perfecte substitutis (et tunc modo regulariter paniculiformi modoque plus minus divaricata, floribus nunc fere omnibus discretis et nonnunquam (præsertim in ramulis extremis) scorpioido-ordinatis, nunc (et frequentissime) in glomerulos s. capitula ad apices angulosque et sinus ramorum ramulorumque conglobatis; tandem aliquando glomerulis s. capitulis in axillis rami bracteati axi ipso cæterum nullo modo absimilis sessilibus inflorescentiamque mixtam atque vere indefinitam (2) conficientibus. Bractea interdum nullæ, plerumque parvæ persistentesque, sæpe scariosæ, quandoque in involucrium proprium v. commune gamo-v. di-polyphyllum coalitæ, — Quoad affinitates Urticearum confer ad procemium Monographiæ meæ ubi de his fusc disserterui primumque conatus sum in lucem proferre quo modo hic ordo cum Malvoideis Euphorbiaceisque connectitur.

(1) Quum nervi primarii (s. costulae) sunt omnes æqualiter patentes inferioresque sequentibus circiter æquilongi aut forte breviores, folium dicitur penninervium. Quum vero nervi primarii inferiores (s. basilares) cæteris multo longiores sunt et magis adscendentes (v. g. ad medium limbi vel ultra producti), folium dicitur trinervium aut triplinervium. In folio trinervio nervi basilares a nervo medio (s. costa) ad ipsam insertionem petioli, in folio triplinervio paulo supra hanc insertionem nascuntur.

(2) Clarissimus Aeh. Guillard, qui assiduo studio veram rationem inflorescentiæ enodare laboravit, inter inflorescentias indefinitas (s. progressivas) ducit illas cunctarum Urticearum, in unam confundens omnino omnes inflorescentias e caule aut ramo plantæ enatas (Bullet. Soc. bot. Fr. 4, p. 380 et 453).

Videas quoque quod in eodem exposui super geographicam distributionem Urticacearum, necnon de indole admodum polymorphica specierum, quam ob causam interdum difficillimum est eas rite circumscribere. — Species paucissimæ more herbarum oleracearum esculentæ sunt; fructus nonnullarum etiam comeduntur; complures tandem nervositate fibrarum corticalium insiguntur.

## CONSPECTUS TRIBUUM ET GENERUM.

A. — Flores masc. 4-5-andri rarissime 2-3-andri.

### Tribus I. UREREÆ.

Stirpes pilis urentibus s. stimulis armatæ, foliis decussatim oppositis aut spiraliter alternis, perigonio fl. fem. 4-partito v. 4-lobo, rariter bilobo v. bipartito aut tubuloso, semper libero.

\* Achænium rectum. Stipulæ laterales.

I. *Urtica*. Perigonium fl. fem. 4-partitum, foliolis exterioribus plerumque minoribus. Folia opposita.

II. *Hesperocnide*. Perigonium fl. fem. tubulosum. Folia opposita.

III. *Nanocnide*. Perigonium fl. fem. 4-partitum, foliolis exterioribus minoribus. Folia alterna.

III. \*\* Achænium obliquum. Stipulæ axillares. Folia alterna.

IV. *Obetia*. Perigonium fl. fem. tetraphyllum, fructiferum membranaceum achænium longe superans.

V. *Fleurya*. Perigonium fl. fem. inæqualiter 4-partitum, post anthesin membranaceum. Stigma lanceolatum v. filiforme. Achænium in utraque facie scrobiculatum sæpiusque tuberculatum. Plantæ annuæ.

VI. *Laportea*. Perigonium fl. fem. 4-partitum v. -lobum, post anthesin exsuccum. Stigma sæpius filiforme. Achænium discoideum lævigatum. Plantæ perennes s. lignosæ.

VII. *Urera*. Perigonium fl. fem. 4-lobum, fructiferum baccatum. Stigma sæpius capitatum.

VIII. *Scepcarpus*. Perigonium fl. fem. tubulosum ore contracto inconspicue denticulato, fructiferum baccatum. Stigma capitatum.

IX. *Gyrotænia*. Perigonium fl. fem. bilobum post anthesin subimmutatum. Stigma capitatum.

X. *Girardinia*. Perigonium fl. fem. bipartitum, segmento altero multo majore apice tridentato, altero lineari v. abortivo.

Tribus II. *PROCRIDEÆ*.

Stirpes inermes, foliis oppositis v. abortu alternis sæpeque distichis, perigonio fl. fem. libero, 3-rariusve 5-partito, stigmate penicillato.

\* Folia decussatim opposita.

XI. *Pilea*. Perigonium fl. fem. tripartitum, segmento uno majore sub apice gibboso. Flores in glomerulos laxè cymosos aut aggregatos dispositi.

XII. *Achudemia*. Perigonium fl. fem. subæqualiter 5-partitum. Glomeruli laxè cymosi.

XIII. *Lecanthus*. Flores in receptaculo discoideo v. concavo congesti.

\*\* Folia disticha, alterna v. opposita.

XIV. *Pellionia*. Perigonium fl. fem. 5-partitum, ovarium superans, segmentis sub apice mucronatis. Flores cymosi v. fasciculati.

XV. *Elatostema*. Perigonium fl. fem. vulgo 3-phyllum sæpiusque ovario brevius. Flores in receptaculo depresso v. concavo insidentes, capitulo sæpissime involucreto.

XVI. *Procris*. Perigonium fl. fem. 3-4-phyllum ovarium superans, segmentis cucullatis muticis. Flores in receptaculo globoso aggregati, involucreo nullo.

Tribus III. *BØEHMERIÆ*.

Stirpes inermes, foliis oppositis alternisve; perigonio fl. fem. libero aut ovario adnato v. cohærente, nunc sæpissimeque tubuloso, nunc brevissimo v. plane deficiente.

Subtribus I. *EUBØEHMERIÆ*. — Perigonium fl. fem. tubulosum, liberum vel ovario plus minus adhærens, fructiferum vulgo siccum s. membranaceum, ore conspicue 2-4-dentato.

\* Stigma in fructu persistens. Perig. fruct. nec alatum nec costatum.

XVII. *Bøehmeria*. Stigma lineare.

XVIII. *Chamabainia*. Stigma subcapitatum.

\*\* Stigma (lineare) cum apice ovarii articulatum et mox deciduum. Perigonium fructiferum sæpe costatum alatumve.

XIX. *Pouzolzia*. Perigonii fl. masc. segmenta dorso convexa. Nervi basiales foliorum ante apicem limbi in margine evanescentes.

XX. *Memoralis*. Perigonii fl. masc. segmenta summo dorso infracta et transversim cristata. Nervi basiales foliorum indivisi et usque ad apicem limbi producti.



Subtrib. II. *SARCOCHLAMYDEÆ*. — Perigonium fl. fem. liberum, fructiferum succulentum, ore varie dentato v. lobato.

\* Stigma in fructu persistens.

XXI. *Cypholophus*. Perigonium fl. fem. tubulosum, ore contracto. Stigma lineare superne incurvum. Flores in capitula sphæroidea sæpius dense congesta.

XXII. *Touchardia*. Perigonium fructiferum subcampanulatum 4-lobum. Stigma spathulatum longe ciliato-papillosum. Flores dense capitati.

XXIII. *Laurea*. Stigma? Flores in spicas simplices densissimas dispositi.

XXIV. *Sarcochlamys*. Perigonium fructiferum ventricosum, ore laterali contracto. Stigma capitatum. Flores densissime spicati, spicis simpliciter ramosis.

XXV. *Poikilospermum*. Perigonium fl. fem. subcampanulatum ovarium omnino vestiens, ore minutissime 4-denticulato. Stigma capitellato-peltatum. Flores cymosi.

\*\* Stigma cum apice ovarii articulatum et cito deciduum.

XXVI. *Neraudia*. Perigonium fructiferum ventricosum, ore valde contracto. Stigma lineare. Flores in glomerulos axillares paucifloros dispositi.

Subtrib. III. *VILLEBRUNEÆ*. — Perigonium fl. fem. ovario adnatum, limbo minimo dentato v. subintegro.

\* Stigma cum apice ovarii articulatum et valde caducum.

XXVII. *Pipturus*. Limbus perigonii fl. fem. minimus subdentatus. Stigma lineare.

\*\* Stigma in fructu persistens.

XXVIII. *Villebrunea*. Fructus siccus cupula carnosa suffultus. Stigma subpeltatum margine longe ciliatum.

XXIX. *Debregeasia*. Fructus baccatus. Stigma penicillato-capitatum.

Subtrib. IV. *MAOUTIÆ*. — Perigonium fl. fem. nullum v. brevissimum. Stigma persistens.

XXX. *Leucosyke*. Perigonium fl. fem. cupuliforme ovario multo brevius. Stigma capitatum. Flores dense capitati.

XXXI. *Maoutia*. Perigonium fl. fem. nullum. Stigma spathulato-capitatum. Achænium hispidum v. setosum. Flores in glomerulos laxè cymosos dispositi.

XXXII. *Myriocarpa*. Perigonium fl. fem. nullum; loco hujusce calyculus diphyllus ovarii stipitem amplectens. Achænium margine setosum. Flores in spicam longissimam digesti.

XXXIII. *Phenax*. Perigonium fl. fem. nullum. Stigma filiforme. Achænia minuta inter bracteas glomerulorum axillarium occulta.

#### Tribus IV. *PARIETARIÆ*.

Stirpes inermes, foliis semper alternis integerrimisque, floribus dielinis aut polygamis, inflorescentiis (saltem femineis) bracteis herbaceis 2-pluribus s. involucrio munitis, perigonio fl. fem. tubuloso libero.

XXXIV. *Parietaria*. Inflorescentia sæpius polygama 3-multi-flora. Stigma aspergilliforme.

XXXV. *Gesnouinia*. Inflorescentia androgyna 3-flora. Stigma lineare.

XXXVI. *Helxine*. Inflorescentia uniflora.

XXXVII. *Hemistylis*. Inflorescentia unisexualis v. androgyna, floribus masc. in glomerulos spicatos dispositis. Involucrum florum biflorum et diphyllum, foliolis basi utrinque lobulo lineari auctis.

XXXVIII. *Rousselia*. Inflorescentiæ unisexuales, floribus masc. cymulosis. Involueri florum fem. biflori et diphylli foliolis basi integerrimis.

B. — Flores masc. monandri v. (in specie unica) diandri.

#### Tribus V. *FORSKOHLEÆ*.

Stirpes inermes aut interdum pilis induratis aculeolatæ, foliis alternis oppositisve, floribus dielinis sæpe involucratia, perigonii fl. fem. tubuloso aut omnino deficiente.

Subtrib. I. *EUFORSKOHLEÆ*. — Flores in involucrio gamo-polyphyllo aggregati et plerumque densissime lanati.

XXXIX. *Forskohlea*. Involucrum di-poly-phylum.

XL. *Droguetia*. Involucrum gamophyllum.

Subtrib. II. *AUSTRALINÆ*. — Flores exinvolucrati lanaque destituti.

XLI. *Australina*. Flores masc. monandri. Perigonium fl. fem. tubulosum liberum. Stigma in fructu persistens.

XLII. *Didymodoxa*. Flores masc. monandri, feminei perigonio destituti. Stigma persistens.

XLIII. *Distemon*. Flores masc. diandri. Perigonium fl. fem. tubulosum, ovario adnatum. Stigma cum apice ovarii articulatum et mox deciduum.

Trib. I. *URUREÆ* Gaudich. Bot. Voy. Uran. p. 496; Wedd. Monogr. p. 53.

Herbæ fruticescæ (nonnunquam scandentes) vel arbores, pilis arentibus aculeisve (paucissimis exceptis) armati. Folia alterna (5 in singulis cyclis) aut decussatim opposita, homomorpha aut asymmetricè diversiformia, rarissime inæquilatera, cystolithis plerumque punctiformibus inspersa. Sûpulae caulinae, laterales vel axillares. Inflorescentiæ seu cymæ androgynæ aut unisexuales, modo vere dichotomæ sive biparæ, modo et frequentius uniparæ; floribus nunc discretis et in ramulis extremis subscorpioideo-ordinatis, nunc sæpissimeque glomerulatis, glomerulis late aut dense cymoso-paniculatis (1). Flores unisexuales. *Masc.* : perigonio 4-5-partito, segmentis apice membranaceis planiusculis muticisque præfloratione imbricatis; pistilli rudimento obovato vel cupuliformi glabro; alabastro umbilicato. *Fem.* : perigonio libero, inæqualiter 4-partito-lobato vel-dentato, raro bilobo aut (segmentis 3 coalescentibus) bipartito, post anthesin sæpissime accreto; stigmatibus vario; staminum rudimentis nullis. Fructus ut plurimum siccus, nunquam pubescens nec setosus. Embryo cotyledonibus vulgo orbicularibus basi et apice sæpe emarginatis, radícula conoidea longioribus; albumine parco.

I. *URTICA* Tourn. Instit. t. 308, et auct., exclus. spec.; Gaudich. Bot. Voy. Uran. p. 496; Wedd. Monogr. p. 55. — *Urtica*, a, Endlich. Gen. pl. p. 283.

Flores monoici aut dioici, semper glomerulati, glomerulis vulgo in axi seu rhachi indivisa vel ramosa (nunquam dichotoma) spicatis racemosisve vel paniculatis et interdum unilateralibus rarissime solitariis capituliformibusque, inflorescentiis (apud species monoicas androgynis unisexualibusve) in singulis axillis semper geminis, pedicellis florum masc. articulatis (2). *Masc.* Perigonium profunde 4-lobum, lobis ovatis intus concavis extus plus minus stimulosus-hispidis, alabastro in medio depresso. Stamina 4, antheris oblongo-reniformibus. Pistilli rudimentum cupuliforme, hyalinum. *Fem.* Perigonium vulgo 4-partitum s. fere 4-phyllum, segmentis inæqualibus, exterioribus fere semper minoribus plerumque oblongis et interdum subcarinatis patentibusque, interioribus erectis planiusculis aut rarissime (in *U. atrovirenti*) cucullatis. Ovarium rectum, ovoideum. Ovulum imo loculo adfixum, erectum, funiculo brevi. Stigma sessile v. rarius stylo brevi suffultum, penicillato-capitatum (papillis interdum valde elongatis), nunc deciduum, nunc diu persistens. Fructus omnino exsuccus, nempe achænium rectum, ovatum v. oblongum, compressum, læviusculum aut rarius plus minus verruculosum, perigonio aucto membranaceo v. rarissime carnosulo inclusum s. vestitum. Semen pericarpio fere conforme. Albumen parcum. Embryo coty-

(1) Hæc inflorescentias pro figura sua illam paniculæ definitæ admodum referentes paniculas sæpe nuncupavi, sed cymæ sunt revera in quibus rami laterales potentiores et continuata serie geniti axim principalem consumptum præcedunt.

(2) Vid. quoque quod de inflorescentia dicitur in charact. tribus.

ledonibus obcordato-rotundatis radículaque subconica donatus. — Stirpes herbaceæ, annuæ v. perennes, raro frutescentes, cosmopolitæ, in temperatis vero et subtemperatis utriusque orbis præcipue obviæ, stimulis obsitæ aut rarissime horumce abortu prorsus inermes, caulibus ramisque plerumque tetragonis. Folia opposita, varie dentata, raro incisolobata, rarissime integra, palmatim 5-7-v. rarissime, 3-nervia, cystolithis punctiformibus v. rarius oblongis linearibusve. Stipulæ laterales, cujusque paris foliorum nunc liberæ s. utrinque inter petiolos binæ, nunc utrinque in unam interpetiolarem solitariam coalitæ. Glomeruli basi bracteolati. Flores sæpissime ebracteati, utriusque sexus virentes, masculi mox decidui. Fructus diu persistentes.

§ 1. *Flores masc. et fem. in inflorescentiis omnibus aut rarius in intermediis solum intermixti* (1).

\* Inflorescentiæ petiolo breviores, sæpe glomeruliformes (2).

1. U. URENS (Linn. Spec. pl. p. 1396), herbacea, inter stimulos glabriuscula, foliis ovatis v. oblongo-ovatis inciso-serratis, dentibus triangulari-lanceolatis acutis obtusiusculisve (apicali lateralibus subæquali), stipulis inter petiolos utrinque binis, inflorescentiis androgynis petiolo brevioribus (rarissime longioribus), perigonii fructiferi segmentis majoribus pilo urente solitario sæpissime armatis. ☉ In cultis et incultis ruderalisque juxta habitationes hominum fere ubique occurrit, cis Cancrum et præsertim in Europa vulgatissima. in Asia orientali autem necnon inter et trans tropicos rarior. — Engl. bot. 18. t. 1236; Fl. dan. 5, t. 739; Reichb. Fl. germ. 12, t. 652, f. 1320; Wedd. Monogr. p. 58, t. 1, c. f. 12 et 13. U. minor Lam. Fl. fr. 2, p. 194. — Caulis erectus v. adscendens,  $1\frac{1}{2}$ -4 decim. altus, succulentus, subsimplex v. sæpius a basi ramosus, ramis caulem ipsum persæpe altitudine æmulantibus et sicut ille magis minusve stimulosis. Folia  $2\frac{1}{4}$ -8 centim. longa,  $1\frac{1}{2}$ -5 centim. lata, basi rotundata aut acuta v. nonnunquam subcordata sæpissimeque abrupte cuneata, dentibus raro acutis integris aut aliquando denticulis 1-2 instructis, dente apicali ut plurimum lateralibus æquilongo et æquilato vel etiam minore, pagina utraque stimulis sparsis armata cæterum glabra v. nervis folii junioris marginibusque subtiliter setulosis, petiolo quam limbus brevior v. ad nodos inferiores caulis illi æquali aut longiore stimulosi v. inermi. Stipulæ omnino liberæ, lanceolatæ, acutiusculæ, ciliatæ. Inflorescentiæ subsessiles, brevissime ramosæ, juniores congestifloræ, floribus masc. quam fem. paucioribus, perigonii lobis medio dorso setulosis. Paniculæ fructiferæ patulæ, laxifloræ, racemiformes, petiolum rariter æquantes, ramulis inermibus aut parce stimulosis, perigonii fem. segmentis interioribus s. majoribus planiusculis apicem versus et margine setuloso-hispidis necnon in dorso insuper pilo urenti majusculo instructis rariusve inermibus. Achænium ovatum,  $1\frac{1}{2}$ -2 millim. vulgo longum, obtusum, cinereo-rufescens, subtilissime scabriusculum, perigonium æquans. — Notam specificam optimam præbet exiguitas dentis apicalis foliorum. Minus constans est præsentia pili urentis in dorso segmentorum majorum perigonii fl. fem. Alabastri masculi quam in speciebus duabus sequentibus (quæ forsan ab eadem stirpe originem ducunt) triente v. dimidio majores sunt. (v. v. et s.)

β, *parvifolia*, caule et præsertim foliis multo minoribus quam in typo ① Passim. U. parvula Bl. Mus. bot. lugd.-bat. 2, p. 149. (v. s.)

γ, *inermis*, ex toto inermis ① Passim. U. inermis Forsk. Fl. Ægypt. p. 160. U. verticillata Vahl, Symbol. p. 76; Pers. Synops. 2, p. 252.

(1) Vid. etiam U. membranaceam (n. 29) atque species minus rite cognitæ sive dubiæ consanguinitatis ad calcem generis descriptas.

(2) Conf. etiam U. rupestrum cujus spicæ fem. petiolo sæpe breviores sunt, necnon interdum androgynæ.

2. *U. FLABELLATA* (Kunth in H. B. Nov. gen. et sp. 2, p. 40), herbacea, magis minusve stimulosā, cæterum fere glabra, foliis flabellato-incisis, laciniiis plerisque subtrilobis, lacinulis acutiusculis, stipulis inter petiolos utrinque binis, inflorescentiis androgynis densis petiolo brevioribus, perigonii fructiferi segmentis majoribus subinermibus v. sparsim stimulosohispidis. ① In Andibus Novogranatensibus et æquinocialibus; frequentissima ad muros villæ Querchu provinciae Pasto, alt. 2540 metr. (Humb. et Bonpl.); prope Hacienda de Antisana, alt. 3380 metr. et ad Cuenca (Hartweg, n. 1382); in pascuis alpinis prov. quitensis (Jameson), necnon in Peruvia (Pavon, Cl. Gay; Lechler, n. 1802) et in Bolivia (Mandon, n. 1100). — Wedd. Monogr. p. 65. — Caulis suberectus, a basi ramosus, plus minus stimulosus, cæterum glaber, ramis elongatis diffusisque. Folia sæpissime inæquilateralia, dimidio inferiore cuneata, laciniiis pinnatim 3-5-lobis rarius integerrimis, sinubus acutiusculis rotundatisve, utraque pagina stimulis longiusculis validis conspersa, nervis et margine insuper plus minus pilosis, petiolis foliorum rameorum limbo plerumque brevioribus caulinarum vero persæpe eo longioribus rarissime inermibus. Stipulæ parvæ. Glomeruli sessiles, ramulis inermibus. Flores masc. pauci, cito decidui, alabastris,  $\frac{3}{4}$  millim. vix latis, lobis sparsim stimulosohispidis. Perigonii fructiferi segmenta majora interdum fere inermia, at sæpius stimulis 1-5 prælongis atisque minoribus dorsalibus marginalibusve armata, achænio ovato  $1\frac{1}{2}$ - $1\frac{3}{4}$  millim. longo acuto obscure fusco-marmorato scabriusculoque æquilongæ. — Stimuli (teste Hartwegio) quam maxime urentes. (v. s.)
3. *U. SPATULATA* (Smith, in Rees' Cyclop. 37, n. 17), herbacea, inter stimulos fere omnino glabra, foliis parvis rotundatis incisis, dentibus lanceolatis acutis, petiolo quam limbus plerumque longiore, stipulis inter petiolos utrinque binis, inflorescentiis androgynis congestifloris petiolo brevioribus, perigonii fructiferi segmentis majoribus inermibus. ① Ubique in plateis circa urbem Montevideo (Commerson; etc.), in Brasilia meridionali (A. de St. Hilaire) — Wedd. Monogr. p. 60. *U. Scalpe* Commers. mscr. in hb. mus. par.; Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 198. *U. bonariensis* Pers. Synops. 2, p. 552. *U. urentissima* Commers. ex Pers. l. c. *U. urens* Miq. pro parte, in Mart. Fl. bras. fasc. 12. p. 195. — Caulis erectus v. ascendens,  $1\frac{1}{2}$ -3 decim. altus, basi ramosus, stimulis raris copiosisve armatus, cæterum glaber. Folia ambitu rotundata, 8-15 millim. lata, basi abrupte cuneata, dentibus subacuminatis 2-3 millim. longis (terminali et lateralibus fere æqualibus) integerrimis v. denticulo uno et altero instructis, sinubus acutis, pagina superiore stimulis longis et raris armata cæterum glabriuscula, inferiore sæpissime prorsus inermi nervis interdum subadpresse pilosulis, petiolo adulto rarissime quam limbus brevior sparse stimulosus. Stipulæ oblongo-lanceolatae,  $1\frac{1}{2}$ -2 millim. longæ, obtusiusculæ, ciliatæ. Glomeruli subsessiles, petiolo multo breviores, ramulis inermibus. Flores masc. pauci, cito decidui, alabaistro  $\frac{1}{4}$  millim. lato, lobis glaberrimis. Perigonii fructiferi segmenta majora inermia et glabriuscula, achænio oblongo-ovato  $1\frac{1}{2}$ - $1\frac{3}{4}$  millim. longo apiculato lutescente et sub lente minutissime scabriusculo æquilongæ. — *U. flabellatæ* affinium quam *U. urentis*, sed ab utraque bene distincta. Specimina vidi quæ ad quamdam formam sequentis speciei nonnihil accedere videntur. (v. s.)
4. *U. ECHINATA* (Benth. Pl. Hartweg. p. 252), herbacea, inter setas urentes sæpius crebre hispida, foliis ovatis grosse dentatis v. inciso-dentatis crenatisve dente apicali ut plurimum cæteris longiore, stipulis inter petiolos binis, inflorescentiis androgynis confertifloris petiolo brevioribus, segmentis majoribus perigonii fructiferi æqualiter hispidis. ①-②-4.
- α, foliis ovatis grosse dentatis, dentibus triangularibus acutis v. acuminatis (apicali cæteris longiore), petiolo quam limbus sæpius brevior stimulosus. ①-4 In regione alpina Peruviae et Boliviae secus montes et in pascuis rupium (d'Orbigny, Weddell, Mandon). *U. andicola* Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 198; Monogr. p. 60. t. I. C. f. 14-16. Radix in planta perennante crassa, carnosa. Caules sæpe numerosi, ramosi v. simplices, stimulis validis inter pilos breviores numerosissimosque conspersis. Folia  $1\frac{1}{2}$ -4 centim. longa, 1-3 centim. lata, basi rotundata v. cordata, dentibus vulgo integris, limbo plano aut interdum bullato-rugoso, petiolo

quam limbus circiter dimidio brevior rarius longior. Stipulæ oblongo-lanceolatæ, ciliatæ. Glomeruli sessiles, demum petiolo multo breviores, floribus masc. sæpius vix copiosis, lobis perigonii hispidis. Achænium oblongo-ovatum, submillimetrale, breviter apiculatum, læviusculum, fuscescens perigonium subæquans. — Caulis et rami in planta exsiccata sæpe lutescunt. Achænium in varietatibus omnibus fere dimidio minus quam in *U. urente* et in consimilibus. (v. s.)

β, *echinata*, foliis ovato-orbicularibus grosse crenatis v. inciso-dentatis (dentibus inæqualibus acutiusculis obtusiusculisve) subtus in nervis præsertim hirsutis, petiolo quam limbus brevior hirsuto. ①-② In Andibus quitensibus (Hartweg, n. 1383). *U. echinata* Wedd. Monogr. p. 64. (v. s.)

γ, *Berteroana*, caule inter stimulos glabro, foliis (quam in var. α et β vulgo minoribus) ovatis dentatis v. crenato-dentatis, dentibus ovatis acutis obtusisve (apicali majore) petiolo quam limbus sæpius longior stimulosus. ①-② In Andibus chilensibus, prov. Aconcagua (Philippi). *U. Berteroana* Philippi in Linnæa, ann. 1864, p. 235. — Primo intuitu simillima videtur hæc forma varietati *Orizabæ U. chamædryoidis*, sed configuratione cystolithorum facillime distinguitur. In *U. echinata* enim, sicut in *U. urente* et in affinis, hæc corpuscula luce transmissa folio inspecta constanter punctiformia sunt, quum in *U. chamædryoide* ut quoque in *U. magellanica* elongata sint sive oblonga. (v. s.)

δ, *trichantha*, caule elongato debili inter stimulos glabriusculo v. hispido foliis ut in forma præcedente sed basi haud raro late cuneatis, floribus longe denique hirsutis. ① In speluncis regionis alpinæ Andium boliviensium, prope Sorata (Mandon, n. 1102). — Forma valde insignis sed meo iudicio ab *U. echinata* non specificè distincta. (v. s.)

5. *U. MASAFUERÆ* (Philippi, in Annales de l'Univ. de Chili; et Ann. sc. nat. 4<sup>e</sup> sér. 7, p. 107), herbacea, subinermis et glabriuscula, foliis ovatis grosse inciso-crenatis v. crenato-serratis, crenis s. lobulis utrinque 2-4 (apicali majore), stipulis utrinque inter petiolos binis, inflorescentiis androgynis glomeruliformibus petiolo brevioribus, perigonii fructiferi segmentis majoribus æqualiter hispidis. ① In insula chilensi Masafuera maris Pacifici (Philippi). — Caulis 3 decim. altus, debilis, ramosissimus. fere omnino glaber, ramis gracillimis. Folia majora 2 centim. longa, 10-15 millim. lata, basi late cuneata, dentibus lobulisve obtusis v. acutiusculis, parce pilosula, superiora minora sæpe triloba, lamina exsiccata tenuiter membranacea cystolithis oblongis inspersa, petiolo tenui sæpe admodum elongato. Stipulæ parvæ, oblongo-lanceolatæ. Glomeruli sessiles petiolo multo breviores, floribus masc. paucis fugacibus, alabastro 1 millim. lato hispido. Achænium millimetrale, ovatum, pallidum, læviusculum (?), perigonium æquans. — Huc referenda sunt, ni fallor, specimina quædam a beato Hilario in sylvis umbræ prope Montevideo lecta et in hb. Mus. par. servata. Hæc olim pro varietatibus *U. magellanicæ* habui, et non erit mirandum si aliquando, iterum scrutata, ad eandem sententiam redirent botanici. Planta montevidensis multo elatior est quam masafuerensis foliisque majoribus donata, serraturis s. crenis obtusis utrinque 4-6. (v. s. typ.)

6. *U. CHAMÆDRYOIDES* (Pursh, Fl. Am. sept. 1. p. 113), herbacea, subinermis v. plus minus stimulososa, cæterum glabra, foliis lanceolatis ovatisve v. cordatis, superioribus sæpe multo minoribus, inferioribus præsertim grosse inciso-v. crenato-serratis, dentibus triangularibus ovatisve acutiusculis aut obtusis (apicali sæpius majore), stipulis utrinque inter petiolos binis, inflorescentiis androgynis, junioribus glomeruliformibus petiolo plerumque brevioribus, perigonii fructiferi segmentis majoribus inermibus aut stimulo dorsali armatis glabriusculisque vel magis minusve hispidis.

α, foliis plerisque ovato-lanceolatis, superioribus gradatim magnitudine decrescens augustioribusque inferioribus basi rotundatis aut vix cordatis. ① In Americæ septentrionalis provinciis australioribus frequens, usque ad altit. 3250 metr. in imperio mexicano obvia. — Wedd. Monogr. p. 61. *U. verna* Rafn. Atl. Journ. p. 16. *U. purpurascens* Nutt. Fl.

Arkans. p. 169. U. *Bovista* Riddell, Fl. West. st. U. *stachydfolia* Kunth et Bouché, pl. nov. et emend. h. b. berol. in. Ann. sc. nat. 3<sup>a</sup> ser. 7. p. 181. U. *propinqua* Liebmanna in Vidensk. Selsk. Skr. 5 R. nat. og math. Afd. 2. p. 293. U. *Berlandiera* Bl. Mus. bot. lugd.-bat. 2. p. 151 (Berlandier, n. 73). — Caulis erectus, 4-6 decim. altus, gracilis, simplex v. a basi parce ramosus (ramis sæpius simplicibus flaccidisque), pilis urentibus raris aut crebris armatus, cæterum glaberrimus, v. ad nodos apiceque inconspicue et brevissime pilosulus. Folia 3-5 centim. longa in parte inferiore plantæ, superiora multo minora angustaque lanceolata et magis minusve acuminata basi rotundata v. acuta, dentibus integris ad basim limbi minoribus, utraque pagina parce setuloso-hispida stimulis paucis majusculis hirsutis intermixtis, lamina exsiccata subtus pallidior sive subglaucescente cystolithis parvis plerisque oblongis cæteris punctiformibus inspersa, petiolo gracili quam limbus dimidio brevior subinermi mox glabrato. Stipulæ lineari-lanceolatæ, 3-5 millim. longæ, acutiusculæ, ciliolatæ. Inflorescentiæ juniores congestifloræ, subglobosæ, sessiles v. breviter pedunculatæ, petiolum æquantes eove breviores. Flores masc. cito decidui, lobis sparsim setulosus singulis stimulo solitario dorsali haud raro armatis, alabastro 1½ millim. lato. Paniculæ fructiferæ patulæ demumque reflexæ petiolo breviores, densæ, rhachi fere inermi, perigonio fruct. segmentis majoribus ciliolatis achænio ovato 1-1½ mill. long. acutiusculo sub lente punctulato æquilongis. — Subvar. *gracilescens*, foliis magis lanceolatis acutiuscule serratis. ④ Cum typo. U. *gracilis* Rafin. l. c. U. *gracilescens* Bl. l. c. (v. s.)

β, *latifolia*, foliis plerisque late cordato-ovatis (superioribus minoribus) grossiuscule et obtuse dentatis, dentibus interdum denticulo dorsali instructis. ④ Cum typo passim. (v. s.)

γ, *parvifolia*, caule quam in typo magis ramoso, foliis minoribus omnibusque fere æqualibus utrinque acutis serraturis parvis. ④ Texas, etc. (Wright, etc.) (v. s.)

δ, *Orizabæ*, caule gracili flexuoso irregulariter ramoso parce stimuloso, foliis parvis, superioribus et inferioribus subæqualibus late ovatis et plerisque subcordatis. ④ In montis Orizabæ imp. mexic. locis humidis obumbratisque (Liebmann) U. *Orizabæ* Liebm. l. c. p. 292. Folia 1-1½ centim. longa. (v. s.)

Varietates solum insigniores enumeravi, sed aliæ sunt et sat numerosæ quæ fortassis posterius notiscere mereantur; cavendum augeatur earum opespecierum debilius moles.

7. U. *GLANDULIFERA* (Liebmann, l. c. p. 293), herbacea, caule stimuloso, foliis ovatis basi cordatis incisio-serratis, dentibus falcatis obtusiusculis ciliatis, inflorescentiis androgynis glomeruliformibus petiolo brevioribus breviter pedunculatis. ④? In locis humidis umbrosisque regionis calidæ Mexici orientalis, prope Hacienda de Jovo prov. S. Cruz, etc. (Liebmann) — Wedd. Monogr. p. 62. — Caulis 4 decim. altus, a basi ramosus. Folia 2½ centim. longa et lata, petiolata superiora decrescens, anguste lanceolata, subsessilia, acuminata, pilosula, sparsim stimulosa. Stipulæ deciduæ. Glomeruli reflexi, floribus masc. et fem. intermixtis glandulisque clavatis albo-capitellatis numerosis interjectis. Achænia ovata, apice subobliqua, subtilissime granulata. (Descript. ex Liebmann, l. c.) — Species affinis dicitur præcedenti a cl. auctore, sed configuratione foliorum sequenti etiam affinis videtur. «Glandulæ clavatæ» a cl. Liebmannio in inflorescentia visæ sunt verisimiliter pedicelli florum marium delapsorum, nec ut ipse suspicatur florum abortivorum.

8. U. *SURINCISA* (Benth. Pl. Hartweg. p. 293), herbacea, inter stimulos sæpius densiuscule pubescenti-hispida, foliis late ovatis basi rotundatis cordatisve incisio-dentatis, dentibus triangulari-lanceolatis acutiusculis, dente apicali longiore, stipulis inter petiolos utrinque binis inflorescentiis androgynis petiolo (quam limbus interdum longiore) brevioribus rariusve longioribus, fructiferis laxis recemiformibus, perigonii fructiferi segmentis majoribus inæqualiter stimuloso-hispidis. ④ In imperio mexicano, prope Santa Rosa, ad sepes (Hart-

weg, n. 1620); Orizaba (Botteri); Santa Fé prope Mexico (Bourgeau, n. 641); locis umbrosis, prope Colipa (Liebmann); in ins. Porto Santo prope Madeira frequens (Moniz). — Wedd. Monogr. p. 62. U. falcicrenata Lieb. l. c. p. 293; Wedd. l. c. p. 63. — Caulis erectus 2-6 decim. altus, ramosus etiamque ramosissimus, crassitie admodum varia, stimulis copiosis armatus et hos inter (præsertim apice) pubescenti-hispidus. Folia 4½-6 centim. longa, 3½-4½ centim. lata (superiora minora), basi rotundata vel cordata, acuminata, dentibus sæpius integris, utrinque sparsim setigera subtusque pilosula, pagina superiore demum glabrata, dentibus integris vel denticulo hinc inde instructis acutis obtusiusculisve, apicali lineari-lanceolato quam laterales dimidio longiore, petiolo 4-6 centim. longo pilositate tenui patentique et stimulis longis densiuscule obsito. Stipulæ lanceolatæ, 3-5 millim. longæ, acutæ, pubescenti-pilosæ. Paniculæ floriferæ erectæ, petiolo breviores, raro longiores, brevissime pedunculatæ, pedunculo hispidio sparseque stimulosi. Flores masc. pauci, perigonii lobis pubescenti-hispidis. Paniculæ fructiferæ laxiusculæ, patulæ, pedicellis parce stimulosi, perigonii fructiferi laciniis achænio anguste ovato sesquimillimetrali vel nonnihil minore acutiusculo minute punctulato fusciscenti æquilongis. (v. s.)

β, *angustifolia*, foliis lanceolatis, dentibus horumce quam in typo angustioribus sinubusque magis inæqualibus. ① In imperio mexicano cum typo (Bourgeau, n. 641, pro parte) — (v. s.)

γ, *floribunda*, foliis minoribus quam in typo breviusque petiolatis, inferioribus ovatis, superioribus lanceolatis, inflorescentiis fructiferis folium ipsum nonnunquam superantibus. ① Cum typo in ins. Porto Santo (Moniz, Mandon. pl. mad. n. 227). — Caulis inter stimulos hispidus foliosus. merithallis a basi ad apicem plantæ pedetentim brevioribus. Folia 1½-3 centim. longa, superiora anguste lanceolata, basi rotundata v. subcordata, utrinque inter stimulos subtiliter hispida v. glabrata, petiolo quam limbos dimidio breviori. Paniculæ etiam floriferæ petiolum superantes et in speciminibus quibusdam apicem limbi superantibus, rhachi quam in typo sæpe magis stimulosa. Achænium vix millimetrale. (v. s.)

9. U. MEYERI (Wedd. Monogr. p. 64), herbacea, inter stimulos glabra v. pilosula, foliis amplis late ovato-cordatis acuminatis grosse subinciso-dentatis lobulatisve dente apicali s. acumine cæteris longiore, stipulis inter petiolos binis, inflorescentiis androgynis petiolo brevioribus rariusve longioribus pedunculatis, fructiferis laxis racemiformibus, segmentis majoribus perigonii fructiferi stimulis 1-5 dorsalibus armatis.

α, foliis late cordato-ovatis dentibus triangularibus plerumque obtusiusculis. ① In Africa australi (Drège, Ecklon) — U. lobata E. Mey. mscr. (nomen improprium), Bl. Mus. bot. lugd. bat. 2. p. 143. — Caulis erectus, 2-6 decim. altus, subsimplex v. breviter ramosus, dense stimulosus at cæterum fere glaber. Folia 5-10 centim. longa, 3½-8 centim. lata, basi sinu subrotundo parum profundo cordata atque nonnunquam juxta petioli insertionem abrupte cuneata, quasi ab ima basi dentata sparsim stimulosa subtusque et margine pilosa; dentibus inæqualibus (inferioribus ut plurimum minoribus) magis minusve angustis modo antrorsum modo extrorsum spectantibus integris vel denticulo hinc inde instructis, apicali quam laterales angustiore longioreque. Stipulæ minimæ, deciduæ. Paniculæ breviter pedunculatæ, pedunculo rhachique simplici glabriusculis puberulisve stimulisque sparsim armatis, sub anthesi oblongæ petiolumque subæquantes, dein interruptæ et petiolo accreto plerumque breviores. Flores masc. copiosi, longiuscule pedicellati, lobis perigonii sparsim stimulosi, alabastro 2 millim. lato. Perigonii fructiferi segmenta majora oblongo-ovata, piloso-hispida aut glabrata, ciliata stimulisque 1-5 gracilibus dorso munita, segmentis exterioribus interdum triplo longiora achænioque late ovato 1½-2 millim. longo obtuso scabriusculo fuscoque subduplo majora (v. s.)

β, *lobulata*, foliis cordato-ovatis rotundatisve inciso-lobulatis, lobulis subtridentatis. ① Cum typo (Drège) — U. lobulata E. Mey. mscr.; Bl. l. c.;



Wedd. Monogr. p. 84. — Caulis inter stimulos minute pilosulus. Folia utrinque inter setas subtiliter pubescentia. Inflorescentiæ fructiferæ quandoque longiuscule pedunculatæ. (v. s.)

\*\* Inflorescentiæ saltem fructiferæ petiolo longiores (1).

10. *U. HYPERBOREA* (Jacquemont mscr. Wedd. Monogr. p. 68), cæspitosa, herbacea aut basi lignescens, inter stimulos raros subtiliter pubescens, foliis ovatis cordatis argute serratis, stipulis inter petiolos utrinque binis, inflorescentiis parvis androgynis petiolo brevissimo longioribus, perigonii fructiferi segmentis majoribus inermibus achænium undique excedentibus. 4 In lapidosis alpinis Thibetæ occid. et orient. altit. 4500 metr. et ultra vulgarissima (Jacquemont; Strachey et Win'erbottom, n. 18; Hooker. fil. et Thomson). — Rhizoma crassum, lignescens, inferne sphacelatum radiculis,que validis tomentosis instructum, superius suberoso-squamosum. Caules robusti, densiuscule fasciculati, 2-3 decim. alti, simplices, basi denudati demumque in lignum indurati, cæterum subtiliter pubescentes pilisque urentibus paucis armati. Folia sinu lato cordata,  $1\frac{1}{2}$ -3 centim. longa, 1-2 centim. lata, rugosa, utrinque pilis urentibus conspersa, dentibus triangularibus subacuminatis omnibus fere æqualibus vel terminalibus minoribus, nervis paginæ inferioris pubescentibus glabrativæ, petiolo 3-4 millim. longo pubescente inermi vel setigero. Stipulæ oblongæ, petiolo fere æquilongæ, obtusissimæ, puberulæ, etiam ad nodos inferiores persistentes. Paniculæ vix multifloræ. laxifloræ, 1-2 centim. longæ. Flores masc. rari, alabastro 1 millim. lato, segmentis hispidis. Perigonii fem. segmenta interiora post anthesin quam in congeneribus magis ampliata, illa perigonii fructiferi orbicularia membranacea pilosulo-hispida achænium oblongo-ovatum 2 millim. circiter longum pallidum læviusculum undique excedentia. (v. s.)

11. *U. STACHYOIDES* (Webb, Phytogr. canar. 3. p. 259. t. 210), herbacea, caule ramisque præsertim superne copiose stimulosi, foliis ovatis v. elliptico-ovatis apice attenuatis crenatis vel crenato-serratis, dentibus parvis obtusissimis acutiusculisve, petiolo quam limbus brevioræ rarius hinc æquilongo, stipulis inter petiolos utrinque binis, inflorescentiis androgynis v. unisexualibus (masc. superioribus) sæpe spiciformibus petiolo longioribus brevioribusve, perigonii fructiferi segmentis majoribus hispidulis piloque urente dorsali majusculo persæpe instructis. ① In rupestribus, ad muros et sepes Teneriffæ meridionalis, secus viam inter Santa Cruz et oppidum Orotava, etc. (Webb, etc.) — Wedd. Monogr. p. 72. *U. stachyophylla* Webb, olim. — Caulis gracilis, ascendens vel prostratus, 2-6 decim. altus, valde ramosus, caule primario ut plurimum ramos excedente, pilis urentibus ad apices ramorum sæpe longioribus et crebrioribus. Folia  $1\frac{1}{4}$ -5 $\frac{1}{2}$  centim. longa, 8 millim. ad 3 $\frac{1}{2}$  centim. lata, basi acutiuscula rotundata aut cordata sæpe ab ipsa insertione petioli dentata, dentibus subæqualibus apud folia majora utrinque viginti v. numerosioribus, limbo planiusculo supra pilis urentibus gracilibus sparsim armato, subtus (in folio juniore præsertim) iisdem horrido, cæterum utraque pagina sed præsertim in nervis inferioris margineque plus minusve pubescenti-pilosulo, petiolo gracili longitudine variabili haud raro limbum æquante vel etiam superante pubescente sparsimque stimulosi. Stipulæ lineari-lanceolatæ, 2 millim. circiter longæ, longiuscule ciliatæ. Paniculæ spiciformes sive racemiformes, subsessiles, uni-bisexuales, masculæ et femineæ in ramis iisdem vel distinctis, masculæ superiores ac plerumque femineis longiores (nempe interdum fere decimetrales), rhachi haud raro subalata sparsimque setigera, femineæ densifloræ brevissimæ vel petiolum fere æquantes, rhachi magis minusve stimulosi. Perigonium masc. pubescenti-hispidum, alabastro 1 $\frac{1}{2}$  millim. lato. Perigonii fructiferi segmento interiore achænio ovato  $1\frac{1}{4}$ -1 $\frac{1}{2}$  millim. longo obtusiusculo cinereo-fuscescente oculoque armato distincte punctato subæquimagno. — Ad hanc speciem verisimiliter referenda est *U. rubricaulis* Hornem. mscr.; Bl. Mus. bot. lugd.-bat. 2. p. 150. (v. s.)

(1) Vide etiam species duas præcedentes, scil. *U. subincisam* et *U. Meyeri*.

12. *U. ATROVIRENS* (Req. in Loisel Nouv. not. p. 40, non Fl. gall.), herbacea aut lignescens, inter stimulos raras glabra, foliis cordato-ovatis anguste acuminatis argute inciso-dentatis serratisve petiolo quam limbus saepius brevior, stipulis inter petiolos utrinque binis, inflorescentiis androgynis petiolo longioribus vix ramosis, fructiferis ut plurimum cernuis, perigonii fructiferi segmentis cucullatis carnosulis conniventibus exterioribus interioribusque subaequimagnis, achænium laxissime includentibus. ① ♀ In Corsicæ et Ilvæ ruderatis, ad muros et secus vias, necnon in umbris montanis frequens. — Wedd. Monogr. p. 69. t. 1. c. f. 1-7; Mabile, herb. cors. n. 384. *U. grandidentata* Moris, Stirp. Sard. elench. fasc. 2. p. 9. *U. dioica*, β, *atrovirens* Gren. et Godr. Fl. fr. 3. p. 108. Cæspitosa, caulibus gracilibus erectis parce ramosis, annuis herbaceisque aut perennantibus et tunc basi lignosis, stimulis raris instructis, cæterum glabris. Folia plerumque late cordata, sinu baseos aperto, 2½-6 centim. longa, 2½-5 centim. lata, dentibus triangulari-lanceolatis acutissimis acumine brevioribus sæpissimeque latioribus, obscure virentia, juniora rugosa nitida, utrinque pilis urentibus validis hinc et inde armata, nervis paginæ inferioris margineque et interdum intervenio ipso superioris setulis brevissimis subadpressis vix conspicuis insuper conspersis, cystolithis punctiformibus, petiolo gracili dimidii limbi circiter longitudine rarius limbo subaequilongo setis raris iis caulibus conformibus prædito. Stipulae anguste lanceolatae, 4-6 millim. longæ, acutissimæ, omnino discretæ, glabriusculæ. Paniculae floriferæ rhachi simpliciter breviterque ramosa, juniores erectæ, petiolum vix superantes, glomerulis densissimis demum confluentibus. Flores minimi, masculi in singulis glomerulis 1-2 vel numerosiores (femineis interdum subnullis), segmentis perigonii sub apice hispidis stimuloque solitario in dorso armatis cæterum glabriusculis, alabastro ⅓ millim. vix lato. Paniculae fructiferæ patulae vel cernuæ, ramis brevibus fructibus undique dense obductis, perigoniorum segmentis crassis amplis valde concavis imbricatis conniventibus glaberrimis vel setulis raris subadpressis inspersis pallide virentibus achænium ellipticum 1 millim. circiter longum obtusiusculum læve pallidumque capsulae more continentibus. — Stirps structura perigonii ♀ quam maxime singularis nec ulli cognatum comparanda. (v. s. et v. cult.)

β, *floribunda*, paniculis laxè ramosis. ♀ In iisdem locis ac typus. (v. s.)

γ, *angustifolia*, foliis lanceolatis basi attenuatis. ♀ Cum typo. — *U. ilvensis angustifolia* urens foliis altius incisis Mich. pl. ital. et Germ. exsicc. n. 180. (v. s.)

Plantam vidi in hb. hort. bot. mospel quæ fortassis in varietatem quartam posterius constituenda erit. Paniculae speciminis omnino mascula sunt et dentes foliorum quam in typo magis inæquales denticulo dorsali sæpe instructi.

13. *U. SPITREALIS* (Bl. Mus. bot. lugd.-bat. 2. p. 152), herbacea, sparsim aut conferte stimulosa et inter setas puberula v. glabra, foliis inferioribus ovatis plus minus acuminatis basi rotundatis cordatisve serratis, superioribus gradatim decreescentibus lanceolatis anguste acuminatis, petiolo quam limbus dimidii brevior, stipulis inter petiolos utrinque binis, inflorescentiis spiciformibus pedunculatis petiolum superantibus patulis, inferioribus masculis, superioribus et intermediis omnibus androgynis v. supremis femineis, perigonii fructiferi segmentis interioribus quam exteriora multo majoribus hispidulis.

α, foliis inferioribus ovatis sensim et breviter acuminatis (acumine obtuso) basi rotundatis aut vix cordatis obtuse serratis. ①? In imperio mexicano, inter Tampico et Real del monte (Berlandier, n. 349). — Wedd. Monogr. p. 97. — Caulis suberectus, parce ramosus, inferne crassitie pennæ anatinæ, obtuse quadrangularis, ut petioli pube parca subtili setisque albidis urentibus sparsis vel (præsertim in ramis junioribus) confertissimis armatus aut illorum basibus crassioribus tuberculatus, internodiis inferioribus 4-6 centim. longis. Folia in specim. suppet. 1½-3 centim. longa sed nulla horumce ad partem inferiorem caulis pertinent, superiora anguste lanceolata basi integra acutaque et in triente superiore acuminata serraturis parvis subæqualibus. Spicæ folio breviores, breviter pedunculatae, pleræque interruptæ, rhachi sparsim stimulosa v. inermi parceque puberula.

Glomeruli masc. in spiram dispositi sec. cl. Blume, perigonii lobis hispidulis, alabastro 1-1½ millim. lato. Perigonii fructiferi segmenta majora demum fere glabrata, achænium submillimetrale ellipticum pallidum sequantia. (Descript. partim ex Bl. l. c. et partim ad specimen sat mancum in hb. Mus. par. servatum.)

β?, *ligulata*, foliis inferioribus ovatis basi cordatis apice in acumen elongatum liguliforme acutum protractis argute serratis. ①? In imperii mexicanæ regione frigida prov. Oaxaca, altit. 3000 metr. (Galeotti, n. 314). — Caulis erectus, elatus, simplex, obtuse angulatus, puberulus et sparsim setosus, basi forsan penna anserina crassior, internodiis etiam superioribus fere decimetralibus. Folia majora visa ampla, 15 centim. longa, 7 centim. lata, basi haud profunde cordata, acumine lineari 3 centim. longo integro, serraturis limbi triangularibus parum incurvis a basi ad apicem limbi magnitudine crescentibus, utraque pagina hispida sparseque stimulosa, petiolo gracili 2-3-centimetrali. Stipulæ anguste lanceolatae 5-8 millim. longæ, acuminatæ, hispidulæ. Spicæ graciles 3-5 centim. longæ, pedunculo 10-15-millimetrali rhachique subtiliter pubescentibus et sparsim setigeris, superiores omnino femineæ. Floris masc. alabastrum 1½-2 millim. latum. (v. s.)

14. U. MAGELLANICA (Poir. in Lam. Encycl. suppl. 4, p. 221), herbacea, parce stimulosa aut inermis, caule ramisque sæpissime omnino glabris, foliis ovatis v. lanceolato-ovatis et haud raro basi cordatis subacuminatis plerumque grosse crenato-serratis dentatisve, stipulis inter petiolos utrinque binis, inflorescentiis spiciformibus petiolo longioribus patulis, perigonii fructiferi segmentis interioribus quam exteriora multo majoribus hispidulis. 2 In America andina et insulis magellaniceis sat frequens: in temperatis sylvestribus Novæ Granatæ (Humb. et Bonpl.) in pascuis alpinis regionis quitensis, altit. 3600-3950 metr. (Jameson); in regione alpstri Peruviae et Boliviae (Cl. Gay; Mardon n. 1098; Weddell, etc.); ad sepes et in sylvis chilensibus præcipue australibus et in ins. Chiloe et Chonos (d'Urville, Bertero, Cl. Gay, Hook. fil. etc.); prope Port Galant (Commerson). — Wedd. Monogr. p. 70. U. leptophylla Kunth in H. B. Nov. gen. et sp. 2, p. 39. U. Darwinii Hook. fil. Fl. Antarct. p. 343; Cl. Gay, Fl. chil. 5. p. 359. Caules graciles, 5-8 decim. alti, flaccidi, simplices v. parce ramosi, ut plurimum setis sparsis raro copiosis armati, sub apice et ad nodos pilis subadpressis et vix conspicuis etiam frequenter conspersi. Folia inferiora plerumque ovata (superiora sæpe angustiora), basi obtusa v. rotundata v. subcordata, dentibus late triangularibus acutis obtusisve prope basim limbi sæpissime multo minoribus vel quidem subnullis, dente apicali s. acumine lanceolato, lamina in utraque pagina et præcipue in inferiore subtiliter et subadpresse pilosa cæterum subnervi v. sparsim stimulosa, exsiccata obscure viridi cystolithisque punctiformibus oblongisve inerspersa, petiolo gracili dimidii limbi circiter longitudine rarius longiore interdum breviora pilosulo stimulisque raris armato. Stipulæ oblongæ, 3-4 millim. longæ, obtusiusculæ, ciliatæ. Inflorescentiæ breviter pedunculatæ, rhachi fere simplici pubescente juxtaque glomerulos crebriusculæ stimulosæ, juniores congestifloræ erectæ petiolo breviores. Flores masc. numerosi, majusculi femineos minimos initio tegentes v. pauciores, alabastro 2 millim. circiter lato, lobis subtiliter hispidis. Paniculæ fructiferæ patulæ, racemos graciles interruptos petiolo longiores fingentes, perigonii fract. segmentis majoribus planiusculis dorso hispidulis achænio late ovato 1½ millim. longo pallideque lutescente æquimagnis. — Stirps natura valde polymorpha necnon pro ætate sat variabilis. Ineunte anthesi flores masc. interdum adeo sunt numerosi ut, femineis sub hinc latentibus, planta dioica videatur. — Subvar. *pubescens*, caulibus foliisque pubescentibus. — Cum typo. (v. v. et s.)

β, *glomeruliflora*, foliis quam in typo minoribus, inflorescentiis in glomerulos petiolum vix superantes aut etiam breviores redactis. 2? In Brasilia meridionali (A. de St. Hilaire). — Cum U. Masafueræ conferenda. (v. s. in hb. Mus. par.)

γ, *bracteata*, foliis cordato-ovatis lanceolatisve, dentibus plerisque subacuminatis, stipulis inter petiolos utrinque plus minus coalitis, racemis in-

ferioribus masculis, superioribus femineis, intermediis solum androgynis. 4 In Chile prope Rancagua (Bertero, Bridges, etc.). — U. bracteata Steud. in Bot. Zeit 33. p. 258; Bl. Mus. bot. lugd.-bat. 2. p. 139; Wedd. Monogr. p. 82. — Stipulis inter petiolos subcoalitis ad speciem sequentem spectat, atque inflorescentiis subunisexualibus ad præcedentem.

15. U. *BALLOTEFOLIA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 197; Monogr. p. 73), herbacea parce stimulosae etiamque subinermis, pubescens v. glabrata, foliis ovatis subcordatis subduplicato-crenatis rarius simpliciter dentatis, petiolo quam limbus multo brevior, stipulis inter petiolos utrinque solitariis, inflorescentiis androgynis spiciformibus pendulis petiolo multo longioribus. ④? In Nova Granata, prope Bogota (Goudot; Triana, etc.); in Peruvia (Pavon). — U. morifolia Smith in Rees' Cyclop. 37. n. 20, non Poir. — Caulis erectus 4-6 decim. et ultra altus, subsimplex, pilis urentibus vulgo paucis armatus. Folia apice attenuata v. breviter aut longiuscule acuminata, 3-11 centim. longa, 2-6 centim. lata, dentibus obtusissimis vel acutiusculis integris aut denticulo uno et altero dorso instructis, acumine subintegro obtusiusque v. denticulato et acuto, pagina superiore pilis inæqualibus conspersa, inferiore ubique sed præsertim in nervis magis minusve pubescenti-hispida, cystolithis punctiformibus, petiolo villosa inermi aut breviter stimulosa. Stipulæ ovato-oblongæ (singulis e duabus coalitis constantes), 4-5 millim. longæ, apice emarginato-bifidæ v. subtrilobæ, hispidæ, ciliatæ, sæpe reflexæ. Spicæ cylindricæ, densifloræ, longitudine variabiles, minores 2-3-centimetrales, floribus subunilateralibus, rhachi pubescenti-stimulosa. Flores masc. cito decidui, lobis perigonii puberulis hispidisve, alabastris 1½ millim. lato. Perigonii fructiferi segmenta majora hispidula, ciliata, demum glabrata, achænio ovato 1 millim. longo apiculato lævisculo fusciscenti æquilongo. (v. s. in hb. Smith nunc Soc. Linn., Mus. par. et Boiss.)

β, *macrostachya*, foliis longiuscule acuminatis spicas gracillimas non aut vix æquantibus. ④? Prope San Antonio Novo-Granatensium, loco dicto Boqueron de la Mesa (Goudot). — (v. s. in hb. Mus. par.)

## § 2. Inflorescentiæ unisexuales (1).

\* Stipulæ inter petiolos utrinque binæ, omnino liberæ aut basi vix connatæ (2).

16. U. *PILULIFERA* (Lin. Spec. pl. p. 1395), monoica, herbacea, plus minus stimulosae, cæterum, glabra vel pilosula, foliis ovatis sæpeque cordatis incisedentatis, dentibus obtusiusculis, stipulis inter petiolos utrinque binis, inflorescentiis unisexuales, glomerulis femineis globosis s. capituliformibus sæpe in pedunculo solitariis. ①-② In ruderatis et cultis Angliæ Europæ mediæ et mediterraneæ necnon Africæ borealis, ins. St. Helenæ et Asiæ australis. — Engl. bot. 3. t. 148; Lam. Encycl. Illustr. t. 761; Reichb. Fl. germ. 12. t. 653; Wedd. Monogr. p. 74. t. 1. C. p. 11 et 12. Caulis erectus v. adscendens, ramosus, 2-6 decim. altus vel procerior, ad nodos sæpe pilosulus. Folia 3-8 centim. longa, 2½-6 centim. lata, basi cordata v. truncata rariusve cuneato-attenuata, apice plus minus acuminata, dentibus majusculis triangularibus lanceolatisve integris aut denticulo uno aut altero instructis, apicali longiore et angustiore, limbo utrinque setis urentibus sparsim consperso sæpissimeque (præsertim in nervis inferioris paginæ) pilosulo et magis minusve ciliolato, petiolo quam limbus brevior v. longiore stimulosae rariusve subinermi. Stipulæ oblongæ v. ovatæ v. lanceolatæ, magnitudine variabiles (2-8 millim.), omnino discretæ v. inter petiolos nonnihil connatæ, ciliatæ. Inflorescentiæ masc. et fem. in iisdem axillis v. inferiores omnes mascule. Paniculæ masc. petiolo longiores, plerumque inæqualiter bifurcæ, paucifloræ s. e glomerulis 6-8 discretis constantes, ramulis gracilibus stimulosis, lobis perigonii

(1) Vide etiam U. stachydifoliam et U. magellanicam inter species inflorescentiis androgynis, ut quoque species dubiæ sedis ad calcem generis descriptas.

(2) Exceptis var. γ et δ U. dioicæ.

stimulis tenuibus longiusculisque armatis. Capitula fem. breviter aut longiuscule pedunculata (pedunculo stimuloso), nunc (et frequentissime) solitaria, nunc didyma, sub anthesi parva s. glomerulum masculum magnitudine æmulantia at densius setosa, dein accreta, sphaeroidea, diametro subcentimetræli v. minoris s. pisi ciceris circiter magnitudine, obtuse crenulata pilisque urentibus horrida. Perigonii fructiferi segmenta interiora maxima, 3-4 millim. longa, obcordato-rotundata, cucullata, conniventia, infra apicem stimuliferum adpresse hispidula, ciliata. Achænium late ovatum, 2½-2¾ millim. longum, 2 millim. latum, obtusum, fuscum, nitidum, Lini semen referens, perigonio vesiculosulo inclusum. (v. v. et s.)

Subvar. *Balearica*, vegetior, foliis profundius cordatis magisque rugosis quam in typo, stipulis latioribus ①-②. Cum typo. — U. Balearica Linn. l. c.; Gussone, Fl. Sic. synops. 2. p. 581. — Caulis et petioli (teste Gussone) sæpius atroviolacei. (v. s. et v. cult.)

Subvar. *Chesneyana*, parce stimulosa etiamque glabrata, omnibus partibus multo minoribus quam in typo, foliis brevius dentatis, capitulis femineis depauperatis ①. Passim. — U. Chesneyana Bl. Mus. bot. lugd.-bat. 2. p. 152.

β, *Dodartii*, foliis plerumque ovatis integerrimis sinuatisve aut hinc inde dentatis. ① Passim? et in hortis botanicis frequentissime culta. — U. Dodartii Linn. l. c. et auct.; Reichb. Fl. germ. 12. t. 653; Wedd. Monogr. p. 76. t. 1. C. f. 8-10. U. integrifolia Poin. Encycl. 4. p. 637. U. convexa Hornem. mscr.; Spreng. Syst. veg. 3. 839. — Hujus varietatis insignis nulla specimina nisi ex hortis botanicis excerpta videre licuit.

17. U. CANNABINA (Linn. Hort. Ups. p. 282; Spec. pl. p. 1396), monoica, herba-cea, caule fere omnino glabro et inermi, foliis tripartitis lobis pinnati-bipinnati-partitis, stipulis inter petiolos utrinque binis, inflorescentiis unisexualibus, fructiferis spiciformibus petiolo longioribus erecto-patulis. ¼ In Sibiria uralensi, altaica, baikalensi et orientali, necnon in Dahuria (Ledebour) et in Persia prope Teheran secus flumen Chahrout (Aucher Eloi, n. 5314). — Ledeb. Fl. ross. 3. p. 638; Wedd. Monogr. p. 76. — Caulis gracilis, elatus (1-2-metralis), subsimplex aut ramosus, glaberrimus vel subtilissime pilosulus, inermis, folia 6-15 centim. longa, latitudine varia, basi sæpius obtusata, lobis magis minusve productis, intermedio semper majore æquilaterali pinnati-vel bipinnati-partito, lacinii hujusce inæqualibus lanceolatis acutis, terminali inferioribus æquimagna, lobis lateralibus sive inferioribus ut plurimum inæquilateralibus, horum lacinii exterioribus majoribus, sinubus nunc acutis nunc at rarius obtusissimis, limbi utraque pagina setis urentibus rarissimis conspersa cæterum glaberrima vel margine inconspicue ciliolata, petiolo gracili quam limbus multo brevior plus minusve stimuloso vel inermi. Stipulæ lineari-oblongæ, 2-6 millim. longæ, obtusæ vel acutiusculæ. Paniculæ breviter pedunculatæ, pedunculo rhachique pubescentibus glabratisve et pilis urentibus sparsim armatis, glomerulis confluentibus; masculæ ex axillis inferioribus ortæ breviter et diffuse ramosæ, florum perigonio sparsim breviterque hispido, alabastro 1½-2 millim. lato; femineæ in axillis superioribus caulibus ramorumve nascentes, initio erectæ dein patulæ; fructiferæ spiciformes crasse cylindricæ. Perigonium fructiferum glabriusculum ciliolatumve, sic-cum cystolithis prominentibus persæpe asperatum, segmentis majoribus inermibus vel stimulis 1-3 dorso armatis quam exteriora dimidio circiter longioribus achæniisque ovatum 2 millim. longum obtusum apiculatumve minute punctulatum fuscenscensque paulo superantibus. (v. v. cult. et s.)

18. U. RUPESTRIS (Guss. Cat. h. reg. in Boccad. p. 66 et p. 83; Fl. Sic. synops. p. 580, non Lowe), monoica aut interdum dioica, suffruticosa, subinermis et glabriuscula, foliis ovatis v. ovato-ellipticis basi late cuneatis rotundatisve grosse serratis, dentibus acutiusculis v. obtusis, stipulis inter petiolos utrinque binis, inflorescentiis spiciformibus erectiusculis sessilibus, masculis petiolo longioribus femineis brevioribus, perigonii fructiferi segmentis majoribus glabriusculis et fere inermibus. ¼-½ Inter saxa et rupes umbrosos vallibus di Noto Siciliæ (Gussone; Huet du Pavillon, n. 177). — Spreng. Syst. veg. 3. p. 580; Wedd. Monogr. p. 83. — Ramosa, caulibus erectis 3-9 decim. altis cras-

sitie calami scriptorii nonnunquam tortuosis basi albidis, ramis gracilibus glabriusculis et inermibus lævibus scabriusculisve. Folia  $1\frac{1}{2}$ –10 centim. longa, 1–6 centim. lata, apice acuminata, basi interdum fere truncata, dentibus integris, glabra vel pilis sparsis crassiusculis et subinnocuis obsita, obscure viridia, nitidiuscula, lamina exsiccata in pagina superiore cystolithis oblongis conspersa, petiolo gracili  $1\frac{1}{2}$ –4 $\frac{1}{2}$  centim. longo sparsim stimulosi. Stipulæ angustissimæ, vix bimillimetrales vel longiores, demum deciduæ. Spicæ masc. in axillis superioribus ramorum v. in ramis distinctis sessiles, lineares, 3–5-centimetrales, rarissime breviter ramosæ, erectiusculæ, rhachi fere undique floribus tecta vel altero latere nuda et subcompressa stimulis nonnullis instructa vel inermi, floribus numerosis, alabastris  $1\frac{1}{2}$  millim. latis, perigonii segmentis dorso brevissime setulosis. Spicæ fem. quam masculæ breviores (an semper?), initio erectiusculæ, demum patulæ, simplices, rhachi parum stimulosi. Perigonii fructiferi segmenta interiora quam exteriora duplo longiora, achænio ovato 1–1 $\frac{1}{2}$  millim. longo obtusiusculo granuloso æquilonga. — Habitu U. morifoliam nonnihil refert, sed ab illa et cæteris omnibus congeneribus distinctissima est. Inflorescentiæ, teste Gussone, interdum androgynæ. (v. s.)

19. U. *DIÖICA* (Linn. Spec. pl. p. 1396), monoica aut dioica, herbacea v. rarissime basi suffrutescens, cæspitosa, rhizomatibus plus minus elongatis, caulibus elatis varie pilosis stimulosisque aut omnino glabris, foliis ovatis oblongisve v. lanceolatis sæpius ab ima basi serratis, stipulis inter petiolos utrinque binis (hisce rarissime in unam interpetiolarem coalitis), inflorescentiis unisexualibus sæpius ad basim plus minus ramosis, fructiferis pendulis.

α, *vulgaris*, dioica, herbacea, caulibus simplicibus pubescenti-hispidis, foliis cordato-ovatis (superioribus angustioribus) utrinque subtiliter pubescentibus et supra sparsim stimulosi grosse dentato-serratis. ♀ Ad sepes, in fossis ruderalisque et secus muros subhumidos per totam fere Europam (polari excepta) admodum frequens, in Africa, Asia occidentali etiamque orientali et in America boreali temperata et subtropica quoque obvia, in America tropica autem rarissima atque ab Oceanis insulis (Nova Hollandia excepta) ut videtur adhucdum exul. — Lam. Encycl. t. 761; Fl. dan. 5. t. 746; Engl. bot. 25. t. 1750; Reichb. Fl. germ. 12, t. 654; Wedd. Monogr. p. 77. U. major Kanitz, in Bot. Zeit. (1863), p. 51. Caulis erecti v. adscendentes basi que radicales, 4–12 decim. alti elatiorese, raro ramosi sæpius inter stimulos undique pube brevi tenui patulæque vestiti. Folia vulgo majuscula, nempe 5–15 centim. longa, 2 $\frac{1}{2}$ –10 centim. lata, inferiora ut plurimum basi sinu lato cordata, superiora minora haud raro oblonga et subacuminata, dentibus plerumque integerrimis, apicali acutiusculo lateralibus sæpe duplo longiore, petiolo quam limbus duplo triplove breviori pubescenti-setuloso. Stipulæ lineari-oblongæ vel oblongo-lanceolatæ, 4–8 millim. longæ, acutæ aut obtusæ, puberulæ ciliolatæque. Inflorescentiæ masc. sub anthesi erectiusculæ v. patulæ, glomerulis approximatis, perigonii pubescenti-hispidis; femineæ initio patulæ sæpius basi longiuscule et diffusæ ramosæ, ramulis villosis et sparsim stimulosi, glomerulis discretis; fructiferæ nutantes vel pendulæ, vulgo subintricatæ, glomerulis confluentibus. Perigonii fructiferi segmenta interiora quam exteriora plerumque dimidio longiora, achænium ovatum uni-sesqui-millimetrale obsolete punctatum v. læve paulo superantia. — Inter flores masculos exstant interdum nonnulli feminei quorum perigonium majus est et minus hispidum quam in inflorescentiis plantæ femineæ. Rarissime occurrunt specimina inflorescentiis aliis masculis aliisque femineis in eodem caule, masculis tunc (sicut in varietatibus pluribus monoicis infra designatis) ex axillis inferioribus femineisque e superioribus enatis. (v. v.)

Subvar. *umbrosa*, caulibus folisque subtiliter pubescentibus vix stimulosi. ♀ In sylvis Europæ passim. (v. v.)

Subvar. *hispida*, caulibus folisque et ramulis inflorescentiæ crebre stimulosi, perigonii fructiferi segmentis valde hispidis. ♀ In Italia, Corsica et Gallia meridionali frequens, nec non in Afghanistan (Grif-

- 5th). *U. hispida* DC. Fl. fr. 5. p. 355; Duby, Bot. gall. 1. p. 418; Guss. Fl. Sic. synops. 2. p. 581. Folia sæpe latiora quam in typo. (v. s.)
- Subvar. *horrida*, caulibus foliorumque pagina superiore et segmentis perigonii fructiferi præsertim horridis. ♀ Cum præcedente passim.
- Subvar. *duplicato-serrata*, foliis (vix cordatis) caulibusque crebre stimulosi-hispidis, serraturis foliorum utrinque denticulo instructis. (v. s.)
- Subvar. *glabrata*, caulibus foliisque sparsim stimuloso-hispidis nec pubescentibus. ♀ In regione alpina Valesiæ, juxta monticularum domos, altit. 2400 metr. (J. Gay); in Dalmatia (Visiani), in Cache-myria (Jacquemont, n. 1159). *U. glabrata* Clementi, in Vis. Fl. Dalm. 1. p. 217. (v. s.).
- β, *Eckloniana*, monoica, caulibus fistulosis setis urentibus densissime hirsutis, foliis cordato-ovatis subtus v. utrinque pubescentibus et magis minusve stimulosis. ♀ In Africa australi (Drège n. 2844; Zeyher n. 1547). — *U. Eckloniana* Bl. Mus. bot. lugd.-bat. p. 142.
- γ, *platyphylla*, monoica, caulibus undique et præsertim ad nodos stimuloso-hispidis pubescentibusve sed stimulis veris ut videtur parcissime armatis, foliis amplis cordato-ovatis inæqualiter serratis supra stimulis minimis sparsis exasperata, stipulis utrinque solitariis integris (duabus nempe in unam interpetiolarem ex toto connatis), perigonio fructifero hispidulo. ♀ In Asia boreali-occidentali, juxta sinum Awatschka (Seemann n. 1293). *U. platyphylla* Wedd. Monogr. p. 86. Folia 15–20 centim. longa. Stipulæ ovalæ, fere sesquicentimetrales. (v. s. in hb. Kew.)
- δ, *Kioviensis*, monoica, caule setis urentibus obsito cæterum pube destituto, foliis ovatis, stipulis foliorum superiorum inter petiolos sæpissime plus minus alte connatis, perigonio fructifero glaberrimo. ♀ Prope Kiew in Rossia australi (Bogowitch); ad S. Georgen in com. Posoniensi Hungariæ (rev. Botta). — *U. kioviensis* Bogowitch, in Bull. Soc. Nat. Mosc. (1843) n. 2. p. 324. *U. dioica* var. *monoica* Trautvetter, in Bull. phys. math. S. Pet. 13. p. 189. *U. radicans* Botta, in Verhandl. des Vereins für Naturk. in Presburg, 1 (1856) Abhandl. p. 6–7. SB. p. 24. *U. major* & *kioviensis* Kanitz, in Bot. Zeit. (1863) p. 51.
- ε, *galeopsisifolia* (Kanitz, in litt.), foliis ovato-lanceolatis supra pilosis subtus caulibusque tomentosis, ramis inflorescentiæ spiciformibus petiolum subæquantibus pedunculis tomentosis. ♀ In Hungaria, ad lacum Peiso (Wierzbicki). — *U. galeopsisifolia* Wierzb. in Gr. Anth. Herb. n. 41, ex Kanitz l. c.; Bl. Mus. bot. lugd.-bat. p. 145, non Jacq. fil.; Wedd. Monogr. p. 100. — Similitudine chirographica delusus (teste cl. Kanitz) b. Blumius hanc plantam false nomini Jacquini fil. addixit. Specimen a celeberrimo auctore in herb. cæs. vindob. visum a Wierzbickio in Hungaria ann. 1824 lectum fuit schedulaque ipsius manu scripta instructum est. — Forma sequens cum hac forte consocianda est.
- ζ, *pubescens* (Trautvetter in Bull. phys. math. S. Pet. 13. p. 188), dioica, caule foliisque stimulis fere prorsus destitutis dense molliterque pubescentibus, perigonio fructifero puberulo, cæt. ut in typo. ♀ In Rossia australi (Ledebour, Trautvetter). — *U. pubescens* Ledeb. Fl. ross. 3, p. 638.
- η, *mollis*, monoica, caule foliisque pube molli densaque obtectis et sparsim stimulosi, inflorescentiis masc. et fem. intermixtis (Steudel), ramis spiciformibus ad caulem subverticillatis. ♀ In Andibus chiliensibus prope Quillota et Rancagua (Bertero); in imperio mexicano, altit. 2600 metr. (Heller).
- θ, *diplothricha*, dioica, caule vix ramoso, foliis ovato-lanceolatis (dente apicali lineari producto) cauleque pilis brevibus mollibus appressis pubescentibus et setis urentibus hispidulis, ramis inflorescentiarum masc. (femineis non visis) ad caulem subverticillatis interrupte glomeruliferis

folio brevioribus. ♀ In Andibus chilensibus prov. Aconcagua (Philippi). U. diplotricha Philippi in Linnæa 33 (1864) p. 234. — Teste cl. auctore, glomeruli præter setas filis crassioribus apice truncatis (nonne pedicellis exsiccatibus?) longitudine pilos calycesque æquantibus cincti sunt.

1, *pycnantha*, monoica, inflorescentiis fem. contractis spiciformibusque, glomerulis confluentibus, cæt. ut in typo. ♀ In Bolivia Andibus prov. Larecaja (Mandon, n. 1098) et passim.

x, *Steudelii*, dioica, caule ramuloso, foliis lanceolato-oblongis (saltem superioribus) basi obtusis crenato-dentatis cauleque setoso-hispidis, ramis inflorescentiæ spiciformibus verticillum mentientibus, perigonii fructiferi segmentis breviter hispidis. ♀ Ad sepes secus vias Rancaguæ in Chile. U. pseudo-dioica Steud. in Bot. Zeit. 23. p. 257; Wedd. Monogr. p. 99.

λ, *angustifolia* (Ledebour, Fl. alt. 6. p. 240; Fl. ross. 3. p. 637), dioica, caulibus elatis puberulis glabrativæ sæpius inermibus, foliis ovato-vel oblongo-lanceolatis lanceolatisve basi ætutis aut rotundatis grosse et argute serratis v. subincisis utrinque at præsertim subtus scabriusculis hispidis rarissimeque stimulosi, perigonio fructifero hispidulo. ♀ In Sibiria præsertim boreali et in Rossia australi, in imperio sinensi circa Pekin (Rev. P. David), nec non passim et rarissima in America boreali temperata (Liebmann, cæt.). — U. angustifolia Fisch. Hort. Gorenk.; Horn. Hort. Hafn. Suppl. p. 107. U. aquatica Liebm. l. c. p. 291; Wedd. Monogr. p. 80. Caules nonnunquam 3-metrales. (v. s. et v. cult.)

μ, *subniscia*, dioica, caule foliisque ovatis inciso-serratis glabrativæ atque fere inermibus, cæt. ut in var. præced. ♀ In Syria. (v. s.)

v, *procera*, monoica, cæt. fere ut in var. λ. ♀ In America boreali fere tota; in ditone mexicana usque ad altit. 2400 metr. frequens (Berlandier, n. 340 et 432; Galeotti, n. 334; Schiede, n. 27; Bourgeau, n. 188, etc.); in Mauritania prope Blidah (Barrat et Durando). — U. gracilis Ait. Hort. Kew. ed. 1. p. 341; Hook. Fl. bor. am. 2. p. 141. U. procera Muhlenb. in litt. ad Willd.; Willd. Spec. pl. 4. p. 353; Pursh, Fl. Am. sept. 1. p. 113. U. mexicana Bl. Mus. bot. lugd.-bat. 2. p. 140 non Liebm. U. Serra Bl. l. c. U. dioica β angustifolia Schlechtend. in Linnæa, 7. p. 141. (v. s.)

Subvar. *duplicato-serrata*, dentibus foliorum 1-2-denticulatis. ♀ In Kentucky America borealis.

ξ, *Sicula*, dioica, caulibus basi suffrutescentibus gracilibus ramosis piloso-hispidis, foliis quam in formis præcedentibus minoribus basi rotundatis argute serratis hispidiusculis, cæt. ut in typo. ♀ In Sicilia prope Palermo et Polini (Gasparrini). U. sicula Ga-parr. in Guss. l. c. p. 580. Habitus foliorum fere U. rupestris (sec. cl. Gussone), sed quoad flores U. dioicæ et U. hispidæ affinis; tamen ab ipsis caulibus basi suffrutescentibus, foliis minoribus minus profunde serratis et habitu gracili sui proprio aliena.

20. U. INCISA (Poir. Encycl. Suppl. 4. p. 223), monoica v. dioica?, caulibus inter stimulos glabris, foliis oblongis linearibusve vel lanceolatis aut triangulari-ovatis dentatis vel inciso-serratis. serraturis sinubusque acutis, stipulis inter petiolos utrinque binis, inflorescentiis unisexualibus ut plurimum racemiformibus petiolo longioribus v. brevioribus, perigonio fructifero glabro.

α, foliis lanceolatis v. lineari-lanceolatis basi acutis. ♀ In locis umbrosis Australiæ felicis et Tasmaniæ. — Wedd. Monogr. p. 81. U. lucifuga β linearifolia Hook. fil. in Lond. Journ. of bot. 6. p. 285; Walp. Ann. bot. 1. p. 644. — Caules graciles 6-10 decim. alti, simplices v. breviter ramosi, pilis urentibus densiuscule aut parce conspersi, cæterum glabri. Folia 6-15 centim. longa, 5-10 millim. lata, dentibus triangularibus acutis frequenter acuminatis et vulgo integris, apicali lanceolato, pagina utraque limbi stimulis sparsis armata, cæterum glaberrima v. subtilissime



pilosula, petiolo gracili quam limbus brevior parum setigero. Stipulæ oblongæ v. lineari-lanceolatæ, 3-5 millim. longæ, acutæ, glabræ. Inflorescentiæ plerumque geminæ interdum ramosæ, masculæ et feminæ in distinctis ramis aut plantis? vel in eodem ramo ortæ, masculæ sæpe superiores, patulæ pendulæque, rhachi setosa v. inermi. Perigonii masc. lobi hispiduli v. glabrati, alabastro 2 millim. lato. Perigonium fructiferum glaberrimum et inerme, segmentis majoribus minora dimidia longitudine excedentibus achænioque late ovato  $1\frac{1}{4}$  millim. longo apiculato læviusculo pallido æquilongis. — A varietatibus nonnullis glabrioribus *U. dioicæ* haud multum recedit. Flores masculi quam in omnibus semper majores. — In herbario Hookeriano formam vidi monstruosam perigonio fem. tubuloso inæqualiterque 4-dentato ut in gen. *Hesperocnidi*. (v. s.)

β, foliis oblongo-ovatis basi truncatis, petiolo quam limbus brevior. γ Cum typo et in Nova Zelandia. (v. s.)

γ, foliis triangulari-ovatis basi cordatis, petiolo quam limbus sæpe longiore. γ In Tasmania præsertim et Nova Zelandia. — Folia 3-6 centim. longa, 2-4  $\frac{1}{2}$  centim. lata. (v. s.)

\*\* Stipulæ inter petiolos utrinque solitariæ (s. duæ in unam interpetiolarem coalitæ), integerrimæ v. apice bifidæ (1).

21. *U. AUSTRALIS* (Hook. fil. Fl. antarct. p. 68), monoica? herbacea pilosula v. glabrata, foliis late ovato-cordatis grosse dentatis, dentibus acuminatis sinubusque acutis, stipulis utrinque inter petiolos solitariis, inflorescentiis unisexualibus, femineis racemiformibus erectis petiolo plerumque brevioribus, achænio lævi. γ? In ins. Auckland et in Novæ Zelandiæ insula septentrionali (Hook. fil., cæt.). — Wedd. Monogr. p. 88. — Caulis elatus, basi prostratus et radicans, validus, ad nodos crebriusculæ stimulis. Folia ampla, 10-14 centim. longa, 7-12 centim. lata, opposita aut superiora ternata, basi sinu lato cordata atque e medio sinu brevissime cuneata, apice attenuata aut subacuminata sæpe cochlearia, ab ipsa basi dentata, dentibus triangularibus ovatisve sæpissime mucronulato-acuminatis integerrimis v. denticulo hinc inde instructis apicali angustiore lanceolato, limbo carnosulo 5-7-nervio utraque pagina sparsim pilosula superiore utem demum omnino glabrata inferioreque pilis paucis urentibus insuper in nervis conspersa, petiolo crasso limbum æquante v. multo brevior, primum subtiliter pilosulo et juxta limbum stimulosulo demumque glaberrimo. Stipulæ interpetiolares ovatæ, 8-12 millim. longæ, ad medium usque v. etiam profundius bifidæ, lobis oblongis acutis in nervis et margine pilosulis. Racemi densiflori, breviter pedunculati, rhachi subinermi. Perigonium fructiferum carnosulum, glaberrimum, segmentis interioribus quam exteriora dimidio longioribus, achænium elliptico-ovatum,  $1\frac{1}{4}$ - $1\frac{1}{2}$  millim. longum paulo superantibus. — *U. dioicæ* affinis. (v. s.)

22. *U. AUCKLANDICA* (Hook. fil. Fl. antarct. p. 68), monoica?, herbacea, tota pubescens, foliis late cordato-ovatis acutis grosse serrato-dentatis, dentibus acuminatis sinubusque acutis, stipulis inter petiolos sæpius solitariis bifidisque aut bipartitis, inflorescentiis unisexualibus, masculis breviusculis. γ? In insulis Auckland, ad oras sylvarum necnon in arenosis maritimis (Hooker fil.). — Walp. Ann. bot. p. 643; Wedd. Monogr. p. 91. — Caulis erectus, robustus, angulatus, crassitie pennæ anserinæ, 3 decim. altus, ad nodos incrassatus et setosus. Folia majuscula, 6-10 centim. longa, 4  $\frac{1}{4}$ -7  $\frac{1}{4}$  centim. lata, omnia opposita, basi sæpius cordata, cochlearia, coriacea, stimulosula, siccitate nigricantia, juniora fusco-pubentia, petiolo 3 centim. circiter longo valido sulcato. Stipulæ interpetiolares majusculæ, foliaceæ, late ovatæ, raro usque ad basim fissæ. Inflorescentiæ masculæ juniores (solum visæ) floribus pedicellatis aggregatis, perigonii lobis dorso setosis. — Species nondum satis cognita. *U. dioicæ* etiam sat affinis, ut videtur. (v. s.)

(1) Vide etiam varietates γ et δ *U. dioicæ*.

23. *U. SIMENSIS* (Hochstetter, in Ach. Rich. Tent. fl. Abys. 2, p. 260), dioica, herbacea, caule crasso fistuloso pubescente v. glabrato stimulisque undique hirsuto, foliis cordatis v. triangulari-ovatis serratis, serraturis integris denticulatisve, sinubus acutis, stipulis inter petiolos solitariis, inflorescentiis racemiformibus erectis petiolo longioribus, achænio majusculo lævi. 2? In Abyssiniæ montibus simensibus, altit. 3500 metr. (Schimper). — Wedd. Monogr. p. 87. — Caulis erectus v. adscendens, basi interdum radicans, 4-12 decim. altus, digiti nonnunquam crassitudine, raro ramosus, sæpe rubens, inferne plerumque denudatus, pilis urentibus undique plus minus hirsutus et inter illos subtiliter pubescens aut demum glabratus. Folia 3-7 centim. longa, 2-5½ centim. lata, a basi lata magisque aut minus cordata, rarius rotundata, sæpius sensim attenuata, interdum subacuminata, dentibus triangularibus vel triangulari-lanceolatis, inæqualibus, juxta petiolum minoribus nonnunquam acuminatis integris aut dorso vel utrinque denticulis 1-3 instructis, apicali magnitudine varia, limbo utrinque sed præsertim subtus puberulo stimulisque brevibus consperso inconspicue ciliolato; petiolo quam limbus dimidiis vulgo brevior pilis urentibus obsito. Stipulæ interpetiolares, ovato-rotundatæ, 4-5 millim. longæ, integræ, obtusæ, pubescenti-ciliolatæ, dorso glabriusculæ. Inflorescentiæ masculæ sub anthesi erectæ aut patentes, rhachi tereti puberula, perigonii masculi lobis glabriusculis pilo urente solitario sub apice armatis; alabastro 1½-2 millim. lato; fructiferæ patentes vel cernuæ, densifloræ, subcylindricæ. Perigonii fructiferi segmenta glabrata, inermia v. sæpius pilis urentibus 1-3 dorsalibus instructa, interiora exterioribus lineari-oblongis plus quam dimidio longiora achænioque ovato 2-2½ millim. longo-obtusiusculo flavescenti æquimagna. (v. s.)

24. *U. MORIFOLIA* (Poiret, in Lam. Encycl. Suppl. 4. p. 222, non Smith), monoica, frutescens, caule ramisque mox glaberrimis, foliis ovatis basi cordatis rariusve rotundatis crenatis dentatisve, inflorescentiis unisexualibus racemiformibus patulis pendulisve petiolo longioribus, achænio tuberculato.

α, foliis grosse crenatis, crenis mucronulatis. ̄ In regione sylvarum Teneriffæ et Canariæ ad margines torrentium, altit. 1000 metr. et infra, in rupibus convallium nec non in apricis et ad sepes vulgaris (Webb; Bourgeau, pl. can. n. 199 et 902 sub U. membranacea, cæt.); in rupestribus montis Pico dos Ariedros Maderæ (Mandon, n. 226); in sylvis et vineis editioribus insulæ Pico rarius (Hochstetter, pl. azor. n. 126). — Webb. Phytogr. can. 3. p. 258. t. 211; Wedd. Monogr. p. 91. — Frutex metralis vel altior, diffuse ramosus, ramis elongatis teretibus aut obsolete angulatis, annotinis glaberrimis epidermide rubente lævi demumque sublacera vestitis, novellis sparsim hispidis stimulisque brevibus ad nodos interdum crebriusculis (ut plurimum tamen parce) armatis. Folia 5-10 centim. longa, 3-7 centim. lata, basi fere semper cordata, plerumque acuminata, ab ipso baseos sinu dentata, dentibus inferioribus minoribus, superioribus latioribus nunc obtusissimis nunc acutiusculis et brevissime mucronulatis, integerrimis vel interdum extus denticulo uno alterove instructis, apicali cæteris longiore et sæpius lanceolato; utraque limbi pagina setis urentibus conspersa, superiore pilis tenuissimis adpressis insuper obducta et inferiore plus minusve pubescente; petiolo gracili, dimidium limbum circiter æquante, villosulo glabratove, plus minusve setigero. Stipulæ interpetiolares, lanceolatæ, 4-6 millim. longæ, apice bifidæ, glabriusculæ aut ciliolatæ, mox deciduæ. Racemi graciles, masculi et feminei in eodem ramo (masculis superioribus) vel e ramis distinctis orti; masculi petiolo multo longiores, folium interdum subæquantes, breviter pedunculati, simplices aut ramesi, pedunculo rhachique glabriusculis sparsimque setigeris, glomerulis subconfluentibus, perigonio masc. hispido, alabastro 1-1½ millim. lato; feminei sub anthesi erecto-patuli, demum subcylindrici, petiolum vix superantes illoque haud infrequenter breviores, longe pedunculati, pedunculo filiformi glabro, glomerulis subdiscretis. Perigonii fructiferi segmenta majora margine et in nervo prope apicem hispida setisque 1-2 dorsalibus frequentissime armata, achænio late ovato 1½-2 millim. longo obtuso insigniter tuberculato pallido æquitonga.

β, *elevata*, dentibus foliorum triangularibus acutis. ♂ In insula Madeira (Lowe, Moniz, Mandon). U. *elevata* Lowe, Primit. faun. et fl. Mader. et Port. Sanct. p. 15. U. *Lowe* Seub. Fl. azor. p. 28. (v. s.)

25. U. *PARVIFLORA* (Roxb. Fl. Ind. 3. p. 581), monoica, herbacea, glabra pubescente, foliis ovatis acuminatis basi sæptissime subcordatis subduplicato-crenatis serratisve, stipulis inter petiolos utrinque solitariis, paniculis floribundis erecto-patulis, superioribus femineis persæpe (foliis abortientibus) inflorescentiam unicam terminalem mentientibus, perigonii fructiferi segmentis majoribus fere rotundatis, achænio læviusculo. ♀? In Indiæ orientalis montibus Himalaya, regione subtropica et temperata, et Assam (Wallich, Jacquemont, Strachey et Winterbottom, Wight, Hooker fil., Thomson etc.). — Wight, Icon. pl. Ind. 2. t. 690; Wedd. Monogr. p. 85. U. *virulenta* Wall. Cat. n. 4586. U. *ardens* Lamk. U. *himalayensis* Kunth et Bouché Sp. nov. h. berol. in Ann. sc. nat. 3<sup>e</sup> sér. 7, p. 182. — Caulis erecti, metrales elatioresve, graciles, ramosi, ramis obsolete angulatis et haud raro (fortassis siccitate) fistulosi, modo pube tenui brevissima cinereaue undique obductis, modo glabratis, sparsim aut dense stimulosi. Folia 3-15 centim. longa, 1½-7 centim. lata, basi rotundata vel subcordata, ut plurimum anguste acuminata, superiora interdum in bracteis abeuntia, dentibus magis minusve inæqualibus ad basim et apicem limbi sæpe minoribus et juxta petiolum nonnunquam prorsus nullis triangularibus rariusve sublancoelatis argutis obtusiusculisve ut plurimum dorso et interdum utrinque denticulo uno alterove instructis, nervis basalibus medium limbum vix attingentibus, lamina tota utrinque stimulis raris armata, nervis primariis subtus nunc horridis, nunc fere inermibus, pagina superiore præterea sæpe inconspicue pilosula inferioreque subtiliter pubescente, cystolithis aliis punctiformibus aliis oblongis, petiolo gracili longitudine varia puberulo vel glabrato et plus minus stimulosi. Stipulæ utrinque in unam interpetiolarem oblongo-ovata 4-12 millim. longam integram obtusissimamque puberulam concretæ. Inflorescentiæ breviter pedunculatæ, pedunculo ramisque puberulis inermibus sparsimve stimulosi, inferiores masculæ foliis plerumque breviores et haud raro fere simplices, superiores femineæ præcedentibus vulgo multo longiores, ramulis gracilibus laxè patulis, glomerulis junioribus discretis at demum confluentibus. Perigonium fl. masc. hispidulum, alabastro 1½-2 millim. lato. Perigonium fructiferum æqualiter hispidum, segmentis interioribus quam exteriora plus quam dimidio longioribus achænioque elliptico-rotundato submillimetrali obtusiusculo vel apiculato pallido lævisculoque aut sub lente minute (ad marginem præsertim) scabro æquilongis. (v. s.)

β, foliis ovatis lanceolatisve, dentibus s. serraturis integris. ♀? In iisdem locis ac typus.

26. U. *THUNBERGIANA* (Sieb. et Zucc. in Abhandl. d. math.-phys. Kl. d. kgl. baier. Akad. d. Wiss. 4. 3. p. 214), dioica, herbacea, inter stimulos pubescens, foliis e basi cordata ovatis acuminatis (acumine lineari-subfalcato) grosse inciso-serratis, stipulis utrinque inter petiolos solitariis, inflorescentiis masculis spiciformibus simplicibus v. parce ramosis elongatis patulis pendulisve, femineis... ♀? In Japonia (Siebold) — Walp. Ann. bot. 1. p. 643; Wedd. Monogr. p. 80. — Caulis elati?, simplices, pilis decumbentibus obsiti et stimulis longis insuper armati, internodiis elongatis. Folia illis U. *pituliferæ* similia, 6-10 centim. longa, 5-7 centim. lata, acumine 15-20-millimetrali, serraturis triangulari-subfalcatis acutis integris v. extus ad basim aut utrinque unidentatis, pagina superiore parce stimulosa, inferiore in nervis præsertim hispidopubescente, petiolo quam limbus breviorè gracili puberulo-hispido et parce stimulosi. Stipulæ interpetiolares, oblongæ, 6-8 millim. metientes, obtusæ v. emarginato-bifidæ. Spicæ masc. folia superantes, graciles, laxæ s. interruptæ, simplices v. inferne ramosæ, rhachi pubescente vix stimulosa. Perigonium subtiliter hispidulum v. glabrum. (v. s. in hb. Monac. et Mart.)
27. U. *GRANDIDENTATA* (Miq. Pl. Junghuhn. 1, p. 27; non Moris), dioica, in caule præsertim horrida, foliis rotundato-ovatis acute duplicato-serratis, stipulis magnis inter petiolos solitariis, inflorescentiis femineis præsertim diffuse ramosis patulis petiolo longioribus, achænio lævi. ♀? In ins. Java

(Leschenault; Zollinger, n. 2539). — Wedd. Monogr. p. 90. U. bullata Bl. Mus. bot. lugd.-bat. 2. p. 145. — Caulis erectus, robustus, subsimplex, pilis urentibus horridus et hos inter pubescenti-hispidus. Folia 8-12 centim. longa, 7-11 centim. lata, basi latiuscule cordata, a medio ad apicem sensim attenuato-acuminata, dentibus magnis triangularibus acutissimis utrinque serraturis argutis 1-3 instructis rarius integris ornata, acumine ipso lanceolato, supra sparsim stimulosæ cæterum glabra, subtus in nervis stimulis paucis etiam armata intervenio pubescente v. glabrato, petiolo limbum subæquante puberulo stimulisque horrido. Stipulæ interpetiolares, magnitudine insignes (nempe nonnunquam bicentimetrales), late ovatæ, integræ, nervosæ, ciliolatæ dorsoque puberulæ v. glabratae. Paniculæ masc. petiolo longiores, erectæ, patule ramosæ, rhachi puberula et stimulosæ, floribus subsessilibus, perigonii lobis hispidis, alabastro 1½ millim. lato; femineæ majusculæ pedunculatæ, pedunculo rhachique ramis elongatis ramulisque patentibus dense et brevissime hispidis, glomerulis parvis confluentibus. Perigonium fructiferum hispidulum, segmentis interioribus rotundatis quam exteriora dimidio circiter longioribus, achænium elliptico-rotundatum 1½ millim. longum et sub lente minutissime punctulatum æquantibus. — Habitu Girardiniam quandam mire mentitur. (v. s.)

28. U. FEROX (Forst. Prodr. n. 346), dioica, fruticosa, stimulis paucis elongatis armata, foliis triangulari-oblongis lanceolatisve basi cordatis aut rotundatis et interdum utrinque lobatis sæpius grosse dentatis (dentibus acuminatis sinubus rotundatis sejunctis) rariusve sinuatis v. subintegris, stipulis utrinque inter petiolos solitariis, inflorescentiis masc. racemiformibus quam petioli brevioribus, femineis... ♀ In ruderatis insularum septentrionalis et mediæ Novæ Zelandiæ. — Hook. fil. Fl. Nov. Zel. p. 224; Wedd. Monogr. p. 89. — Frutex 2-metralis, diffuse ramosus, ramulis teretibus striatis, novellis pubescentibus canescentibusque et pilis urentibus deciduis conspersis. Folia vulgo fere hastata, 4-12 centim. longa, latitudine variabili, juxta basim nunc lobo unico et asymmetrico, nunc lobis duobus oppositis brevibus ovatis triangularibusve et sicut reliquum limbum dentatis instructa, superne sensim attenuata, dentibus subæqualibus vel inferioribus minoribus ab ipsa basi sensim acuminatis et in pilum longiusculum productis, lamina (sicca tenuiter membranacea) supra sparssime hispidula nervo medio pilis paucis urentibus longeque bulbosis armato, subtus (in nervis præsertim) stimulis similibus instructa subtiliterque pubescente. Stipulæ lineari-oblongæ, 4-6 millim. longæ, integræ, obtusæ, margine pubescenti-pilosæ. Racemi masc. teretiusculi, rhachi pubescente, floribus breviter pedicellatis sessilibusve. perigonio densiuscule pubescenti-hispido, alabastro submillimetræli. Flores feminei.... — Species ab omnibus congeneribus distinctissima. Stimuli bulbo protracto valde insignes et, teste Colenso, quam maxime urentes, dolore etenim ex illorum ictu orto per dies quatuor haud raro permanente. (v. s.)

29. U. MEMBRANACEA (Poir. Encycl. 4. p. 638), monoica rariusve dioica, herbacea, caule nisi ad nodos inter stimulos glabro, foliis ovatis v. elliptico-ovatis inciso-serratis v. grosse dentatis crenatisve, stipulis utrinque inter petiolos solitariis, inflorescentiis unisexualibus v. rarissime androgynis spiciformibus, superioribus petiolorum superantibus, rhachi mascularum dorso utrinque membranaceo-dilatata et altera facie tantum florifera.

α, caule parce stimulosæ, foliis acute dentatis, spicis unisexualibus, rhachi mascularum membranaceo-dilatata. ○ In cultis et ruderatis Europæ mediæ et præsertim australioris, ins. Azoricis, Madeiraque et Africa boreali inclus. ins. Canariis; in Gallia usque in Armorica secus oras oceani atlantici obvia; in ins. Borbonia olim a Commersonio etiam lecta at verisimiliter non vere indigena. — Desf. Flor. atlant. 2. p. 340; DC. Fl. fr. 5. p. 355; Guss. Fl. Sic. synops. 2. p. 579; Reichb. Fl. germ. 12. t. 652; Wedd. Monogr. p. 93. U. dioica β Linn. Spec. pl. p. 1396. U. dubia Forsk. Fl. Ægypt. Arab. p. 121. U. caudata Vahl Symbol. 2. p. 96; Brot. Phytogr. Lusit. t. 151, non Poir. Nonnunquam biennis, caule erecto v. adscendente, 2-10 decim. alto, simplice vel breviter ramoso, pilis urentibus parce armato rariusve subinermi, ad nodos et præcipue ad

ipsam basim petiolorum magis minusve pubescenti-piloso, cæterum glabro. Folia 2-8 centim. longa, 12 millim. - 6 centim. lata, basi plerumque rotundata aut late cordata sed in ipso sinu brevissime cuneata, apice attenuata vel subacuminata, dentibus triangularibus persæpe acutiusculis integris aut rarius hinc v. inde unidenticulatis, ad basim limbi minoribus vel etiam prorsus deficientibus prope apicem vero ut plurimum majoribus et haud raro margine exteriori convexis apicali cæteris subæquali; limbo utrinque sparsim setigero ac plus minusve hispido et ciliato; cystolithis parvis punctiformibus aut sæpissime oblongis linearibusve sparsis vel juxta nervos crebrioribus, petiolo gracili longitudine variabili sæpius elongato s. limbum subæquante inermi v. setigero. Stipulæ oblongo-ovatæ, 2-6 millim. longæ, integerrimæ, ciliatæ. Spicæ masc. fem. in eadem aut rarius in distinctis plantis; masculæ plerumque longiores ex axillis superioribus ortæ, folia (nonnunquam minima) sæpissime longe superantes, floribus e medio faciei superioris rhacheos nec ex ejus aliis enatis, perigonio masc. pubescenti-hispido, alabastro 1 millim. lato; femineæ pedunculo gracili suffultæ, petiolo longiores brevioresve, nonnunquam (præsertim inferiores) minimæ, subcylindricæ v. ovatæ, continuæ v. subinterruptæ, glomerulis densifloris subsecundis, rhachi acute angulata necnon interdum subalata sparsim setigera. Perigonii fructiferi segmenta majora hispida v. setuloso-hispida, achænio ovato 1 millim. longo obtusiusculo sæpe fusco-punctulato et medio dorso interdum impresso æquimagna. — Ab omnibus congeneribus distincta conformatione spicæ masculæ. (v. v. et s.)

β, *crenata*, foliis multo minoribus quam in typo, dentibus numerosissimis subæqualibus obtusiusculisque. ☉ Passim.

γ, *horrida* (Wilkomm et Lange, Prodr. fl. hisp. p. 251, robustior, stimulus longis crebrisque horrida, spicis superioribus petiolo longioribus floribus paucis fem. cum masc. intermixtis, rhachi parum dilatata undique florigera. ☉ In Hispania (Wilkomm et Lange).

δ, *neglecta*, minus robusta quam typus, foliis sæpius ovato-ellipticis basi cuneatis et acute serratis, spicis omnibus similibus more spicarum fem. typi confectis sed (teste Gussone) floribus fem. et masc. intermixtis. ☉ Passim. In iisdem locis ac typus et præsertim in Sicilia et insulis vicinis (Gussone, cat.). U. neglecta Guss. ind. sem. h. reg. Boccad. ann. 1825, p. 11; Fl. Sic. synops. p. 579. U. azorica Seubert, Fl. azor. p. 28. U. atlantica et U. Pittoni Bl. Mus. bot. lugd. bat. p. 149 et 147. — Specimina quæ vidi sub nomine U. neglectæ Guss. sunt individua feminea depauperataque typi.

### *Species dubiæ affinitatis.*

30. U. GLOMERULIFLORA<sup>1</sup> (Steud. in Bot. Zeit. 33. p. 257), dioica?, fruticosa?, undique hispida et aspera, foliis elliptico-ovatis acuminatis inæqualiter serratis nervis subtus pilosulis, petiolo brevissimo, stipulis utrinque binis, floribus in glomerulos dense globosos congestis, perigonii fructiferi segmentis majoribus medio dorso setulosis. ♂ In sylvis humidiusculis montium insulæ Juan Fernandez (Bertero). — Blume, Mus. bot. lugd. bat. 2. p. 154; Wedd. Monogr. p. 94. Boehmeria fernandeziana Cl. Gay, Fl. chil. 5. p. 360. — Frutex? v. suffrutex, inermis (saltem sic videtur), caule erecto semimetrali v. elatiore rigido basi teretiusculo simpliciter ramoso (ramis virgatis) undique hispido asperoque v. inferne lævigato. Folia basi attenuata,  $1\frac{1}{2}$  -  $3\frac{1}{4}$  centim. longa, 5-20 millim. lata, dentibus acutis obtusiusculisque integris v. denticulo hinc et inde instructis, dente apicali angustiore, limbo pilis brevibus subadpressis rigidisque utrinque exasperato, exsiccato rigido, petiolo crassiusculo 2-3 millim. longo pubescenti-hispido. Stipulæ lanceolatæ, 2-3 millim. longæ, ciliolatæ. Inflorescentiæ fructiferæ (floriferæ non suppetebant) densissime glomeratæ, subglobosæ, pisi circiter magnitudine, in ramis propter merithallorum brevi-

tatem valde approximatae petiolumque aliquantulum superantes. Perigonii fem. segmenta interiora dorso setosula, achænio millimetrali oblongo apiculato ferrugineo lævisculoque paulo breviora. — Specimina mea sunt omnia fructifera; affinitates veræ itaque dubiæ; ob defectum pedicellorum florum ♂ delapsorum forsitan tamen melius inter species dioicas collocanda quam inter androgynas ad quas ipsa facie accedere videtur. Planta fructifera habitu rigido Boehmeriam quamdam glomerulifloram etiam nonnihil refert. (v. s.)

31. U. TRACHYCARPA (Wedd. Monogr. p. 95), dioica?, ramis tomentosis, foliis lanceolatis acuminatis serratis subtus pubescenti-tomentosis et subincanis, petiolo quam limbus dimidius breviora, stipulis inter petiolos utrinque binis, inflorescentiis unisexualibus (an semper?) densis spiciformibus pendulis, perigonio fructifero dense hispido. ♀ In California (Beechey). — Urticæ spec. Hooker et Arn. Bot. Beech. voy. p. 159. — Suffrutex? ramulis teretibus tomentosis pilisque urentibus gracillimis conspersis. Folia (in specimine examinato) 2-3½ centim. longa, 6-10 millim. lata; dentibus parvis triangularibus integris acutiusculis vel obtusis, apicali lineari, laminæ (exsiccatae coriaceæ) pagina superiore pubescente stimulisque raris conspersa, inferiore mollior tomentosa inermi, petiolo tomentoso setigeroque. Stipulæ lanceolatae, 2-3 millim. longæ, pubescentes. Inflorescentiæ fructiferæ (floriferæ mihi ignotæ) densæ, spiciformes, petiolo duplo vel triplo longiores, pendulæ. Perigonii fructiferi segmenta interiora exteriora fere dimidio excedentia, achænio elliptico submillimetrali utrinque acuto sub lente scabro cinereo-fusce-cente paulo longiora. — Fragmentum suppetens aliquanto mancum; species autem ut videtur distinctissima. (v. s. in hb. Kew. et De Lessert.)
32. U. NICARAGUENSIS (Liebmann, in Vidensk. Selsk. Skr. 5 R. nat. og math. Afd. 2. p. 292), monoica, herbacea, caule sparsim setoso et reflexe pilosulo, foliis ovatis acuminatis basi rotundatis grosse serratis, serraturis repandenticulatis, stipulis inter petiolos utrinque binis?, inflorescentiis androgynis spiciformibus petiolum æquantibus eove brevioribus pendulis. ♀? In monte ignivomo Irasu, prope S. Juan de Nicaragua; altit. 2900 metr. (Orsted). — Wedd. Monogr. p. 72. — Caulis simplex. Folia 6-8 centim. longa, 3-3½ centim. lata, utrinque pilosula et supra stimulis sparsim obsita, petiolo 1½-3 centim. longo pilosulo. Stipulæ ovatae, 4 millim. longæ, subacutæ, ciliolatae, reflexæ. Spicæ 15 millim. longæ, floribus sessilibus. Perigonium fl. masc. pilosulum; feminei segmenta majora rotundata pilosula, achænio rotundato-ovato apiculato subtilissime punctulato fusco circiter æquilonga (Descript. ex Liebmann, l. c.). Conferenda cum U. Magellanica.
33. U. MEXICANA (Liebmann, l. c. p. 291), monoica, herbacea, stimulosa et pubescenti-pilosa, foliis cordato-ovatis grosse crenatis (crenis obtusiusculis) longe petiolatis, inflorescentiis androgynis petiolo sæpe brevioribus, floribus masculis plerumque inferioribus, spicis fructiferis pendulis. ♀? In locis humidis imperii mexicani prope vicum Chinantla, prov. Puebla, altit. 2200 m.-tr. (Liebmann). — Wedd. Monogr. p. 67. — Caulis erectus, 6-10 decim. altus, tetragonus, angulis rotundatus, lateribus canaliculatis, ramosus v. subsimplex. Folia 7-10 centim. longa, 5-7 centim. lata, acuminata, petiolo 2½-5 centim. longo. Inflorescentiæ spiciformes, sessiles, laxæ, patulæ, 2½ centim. mentientes longioresve. Perigonium fl. masc. puberulum, floris feminei segmentis majoribus pilosis achænio ovato subtilissime punctulato flavescens subæquilongis. — Specimen nullum vidi; U. chamædryoidi propinqua videtur.
34. U. MICRANTHA (Kunth et Bouché, Spec. nov. h. bot. berol. in Ann. sc. nat. 3<sup>e</sup> sér. 7. p. 182), dioica, glabriuscula, foliis ovatis acuminatis basi late rotundatis grosse serratis, inflorescentiis fem. petiolo brevioribus. ♀ Patria ignota. — Wedd. Monogr. p. 98. Suffruticosa, caulibus erectis teretibusculis. Folia subtus glabriuscula. Inflorescentiæ masc. non visæ. In horto cum U. rupestri Guss. confusa; longe differt præsertim inflorescentia (Kunth et Bouché).
35. U. FOLIOSA (Blume, Mus. bot. lugd. bat. 2. p. 142), dioica, setis urentibus hispida, foliis lanceolatis basi rotundatis v. subcordatis acutis v. acuminatis serratis supra punctato-scabris subtus ad nervos strigosulis, stipulis

utrinque inter petiolos binis, spicis masc. simplicibus petiohū æquantibus. 3 In Japonia. — Wedd. Monogr. p. 100. — Caulis suffruticosus, erectus, serangularis, circiter metralis, ramis brevibus decussatis, foliosus, fistulosus, inferne asper setisque destitutus, superne et præsertim rami tenuiores setis urentibus hirtulus. Folia  $2\frac{1}{2}$ -7 centim. longa,  $1\frac{1}{4}$ -3 centim. lata, serraturis acutis interdum majusculis, in nervis subtus magis minusve strigosula et interdum setulis nonnullis conspersa. Stipulæ breves, acutæ, strigosulæ, apice cito emorientes. Spicæ masc. ex axillis foliorum superiorum breviter pedunculatæ, 12-15 millim. longæ, simplices, glomerulifloræ (Ex Bl. l. c.).

36. U. FASTIGIATA (Blume, Mus. bot. lugd. bat. 2. p. 151), monoica, herbacea, caule albido-setoso, foliis subcordato-ovatis obtusis grosse obtuseque crenato-serratis, stipulis utrinque inter petiolos binis, spicis sessilibus androgynis petiolo brevioribus. 4? In Chile (Cuming). — Wedd. Monogr. p. 98. — Caulis ascendens fastigiato-ramosus. Folia utrinque obsolete puberula et setis raris adpersa. Stipulæ parvæ, acutæ. Flores glomerulati (Descript. ex Bl. l. c.). — A cl. auctore affinis dicitur U. Magellanica, conferenda cum forma brachystachya hujus stirpis.

37. U. STIPULACEA (Bertol. Miscel. 2. p. 24), monoica, herbacea, nudiuscula, foliis ovatis grosse et acute serratis sparse strigoso-pilosis, stipulis floralium membranaceis ciliatis, spicis androgynis simplicibus axillaribus pedunculatis fasciculatis laxifloris folio multo brevioribus. ☉ E ruinis, prope castrum Sedjim Kala (Pl. sicc. Euphr. n. 61). — Proxima U. urenti, secundum cl. auctorem, sed certe diversa dentibus foliorum latioribus, stipulis fol. flor., spicis pedunculatis, cæt.

### *Species exclusæ.*

- U. ACERIFOLIA Zenker = Girardinia palmata.
- U. ACUMINATA Poir = Urera acuminata.
- U. ADOENSIS Hochst. = Girardinia condensata.
- U. ÆSTUANS Linn. = Fleurya æstuans.
- U. AFFINIS Hook. et Arn. = Fleurya interrupta.
- U. ALBA Blume = Leucosyke capitellata.
- U. ALBIDO-PUNCTATÄ Steud. =? Pipturus argenteus.
- U. ALCEÆFOLIA Poir. = Urera Caracasana.
- U. ALIENATA Linn. = Pouzolzia indica.
- U. ? AMENA Wall. list, n. 4626 = alius ordinis.
- U. ANGULATA Blume = Pilea angulata.
- U. ANGUSTATA Blume = Debregeasia longifolia.
- U. ANGUSTIFOLIA Hamilt. = Boehmeria macrophylla.
- U. ANISOPHYLLA Wall. = Pilea anisophylla.
- U. APPENDICULATA Wall. = Villebrunea appendiculata.
- U. AQUATICA Moon = Memorialis aquatica.
- U. ARBOREA Linn. fil. = Gesnouinia arborea.
- U. ARBORESCENS Poir. =? Leucosykes spec.
- U. ARDENS Blume =? Laportea microstigma.
- U. ARGENTEA Forst. = Pipturus argenteus.
- U. ARGUTA Kunth. = Pilea arguta.
- U. ARMIGERA Presl = Urera baccifera.

- U. ARTICULATA Juss. = *Pilea articulata*.
- U. ARVENSIS Poepp. = *Pilea hyalina*.
- U. ATROX Leschen. = *Laportea peltata*.
- U. BACCIFERA Linn. = *Urera baccifera*.
- U. BACCIFERA Blanco Fl. Filip. ed. 1. p. 695 (in ed. 2. sub *U. arborescens*) = ?
- U. BETULÆFOLIA Swartz = *Pilea hederacea*.
- U. BICOLOR Roxb. = ? *Debregeasia* spec.
- U. BIFIDA Zolling. = *Boehmeria biloba*.
- U. BILOBA hort. = *Boehmeria biloba*.
- U. BLANDA Wall. = *Boehmeria platyphylla*, var.
- U. BLITUM Salz. = *Phenax Sonneratii*.
- U. BULBIFERA S. et Zucc. = *Laportea bulbifera*.
- U. BULLOSA Hochst. = *Girardinia bulbosa*.
- U. CÆRULEA Blume = *Cypholophus cæruleus*.
- U. CAFFRA Thunb. = ? *Fleurya peduncularis*.
- U. CALLITRICHIOIDES Kunth. = *Pilea microphylla*.
- U. CALOPHYLLA Schlechtend. = ?
- U. CANADENSIS Linn. = *Laportea Canadensis*.
- U. CANDICANS Burm., Blume = ? *Boehmeria nivea*.
- U. CANDIDISSIMA Blume = *Leucosyke candidissima*.
- U. CAPENSIS Linn. fil. = *Leidesia Capensis* Mull. arg. in Prodr. 15. s. 2. p. 793.
- U. CAPENSIS E. Mey. = *Fleurya peduncularis*.
- U. CAPITATA Linn. = *Boehmeria cylindrica*.
- U. CAPITELLATA Poir. = *Leucosyke capitellata*.
- U. CARACASANA Jacq. = *Urera Caracasana*.
- U. CARAVELLANA Schrank = *Fleurya æstuans*.
- U. CATARACTÆ Steud. = *Pilea trinervia*.
- U. CAUDATA Burm. = *Boehmeria platyphylla*, var.
- U. CAUDATA Poir. = *Boehmeria platyphylla*, var.
- U. CAUDATA Wall. = *Memoralis caudata*.
- U. CAUDIGERA Wall. = *Boehmeria platyphylla*, var.
- U. CELEBICA Blume = *Boehmeria Celebica*.
- U. CELLULOSA Spreng. = *Pilea ciliaris*, var.
- U. CHIPLE Hamilt. = *Pouzolzia viminea*.
- U. CHURTA Hamilt. = *Laportea crenulata*.
- U. CILIARIS Linn. = *Pilea ciliaris*.
- U. CILIATA Swartz = *Pilea Swartzii*.
- U. CINERASCENS Blume = *Pipturus argenteus*.
- U. CINERASCENS Poir. dict. suppl. 4, p. 226 = ?
- U. CINERASCENS Wall. = *Boehmeria platyphylla*, var.
- U. COLLINA Bertero = *Pilea radicans*.
- U. COMOSA Hamilt. = *Boehmeria comosa*.
- U. CONDENSATA Hochst. = *Girardinia condensata*.
- U. CONGLOBATA Blume = *Pouzolzia conglobata*.



- U. CONGLOBATA Pav. = *Phenacis* spec.
- U. CONTRACTA Blume = ? *Villebruneæ* spec.
- U. CORALLINA Liebm. = *Urera* *Caracasana*.
- U. CORDATA Steud. = *Fleurya æstuans*.
- U. CORYLIFOLIA Juss. = *Fleurya æstuans*.
- U. CORYMBOSA Poir. = *Pilea corymbosa*.
- U. CRASSIFOLIA Willd. = *Pilea crassifolia*.
- U. CRENULATA Roxb. = *Laportea crenulata*.
- U. CYLINDRICA Swartz = *Pilea cuneifolia*.
- U. CUNEATA A. Rich. = *Fleurya cuneata*.
- U. CUNEIFORMIS Poir. = *Pilea cuneiformis*.
- U. CUSPIDATA Willd. = *Pilea lucens*.
- U. CYMOSA Hassk. = *Fleurya cymosa*.
- U. CYLINDRICA Linn. = *Boehmeria cylindrica*.
- U. CYLINDRICA Wall. = *Boehmeria platyphylla*, var
- U. DASYPHYLLA Poepp. = ? *Pilea globosa*.
- U. DAUCIODORA Pav. = *Pilea dauciodora*.
- U. DEBILIS Endl. = *Parietaria debilis*.
- U. DECUMANA Rumph. = *Laportea decumana*.
- U. DEPRESSA Swartz = *Pilea depressa*.
- U. DICHOTOMA Blume = *Debregeasia longifolia*.
- U. DICHROA Poepp. = *Pilea nummularifolia*.
- U. DIFFUSA Swartz = *Pilea diffusa*.
- U. DISTACHYA Spreng. = ? *Boehmeria cylindrica*.
- U. DIVARICATA Forsk. = ? *Fleurya æstuans*.
- U. DIVARICATA Linn. = *Laportea Canadensis*.
- U. DIVERGENS Meyer = *Fleurya æstuans*.
- U. DIVERSIFOLIA Blume = *Maoutia diversifolia*.
- U. DIVERSIFOLIA Link = *Girardinia heterophylla*.
- U. ELATA Swartz = *Urera elata*.
- U. ELONGATA Blume = *Pilea trinervia*.
- U. EVITATA Wall. = *Laportea evita*a.
- U. EXCELSA Bert. = *Boehmeria excelsa*.
- U. FASCICULATA Poir. = *Pilea pumila*.
- U. FATUA Burm. = *Fleurya interrupta*.
- U. FERNANDEZIANA Steud. = *Parietaria debilis*.
- U. FRUTESCENS Poir. dict. enc. suppl. 4. p. 226. = ?
- U. FEROCISSIMA Sweet = *Girardinia heterophylla*.
- U. FICIFOLIA Poir. = *Obetia ficifolia*.
- U. FILIFORMIS Walt. = *Boehmeria platyphylla*, var.
- U. ? FINLAYSONIANA Wall. list n. 4644 = ?
- U. FLORIGUNDA Kunth. = *Pilea multiflora*.
- U. FOLIOLOSA Wall. = *Pouzolziæ* spec.
- U. FREIREÆFORMIS Steud. = *Parietaria debilis*.

- U. MOROIDES Cunnigh. = *Urera* ? *moroides*.  
 U. MULTIFLORA Poir. = *Pilea multiflora*.  
 U. MURALIS Vahl Symb. 1. p. 77 (U. *parasitica* Forsk.) = ?  
 U. MURICATA Heyne = *Debregeasia longifolia*.  
 U. MUTISIANA Spreng. = *Pilea multiflora*.  
 U. NANA Ham. = *Pouzolzia indica*.  
 U. NAUCLIFLORA Roxb. Fl. ind. ed. 1832. v. 3. p. 592 = *alius ordinis*.  
 U. NEMOROSA Kunth. = *Fleurya æstuans*.  
 U. NITIDA Vell. = *Urera baccifera*.  
 U. NIVEA Linn. = *Boehmeria nivea*.  
 U. NUDICAULIS Swartz = *Pilea nudicaulis*.  
 U. NUMMULARIFOLIA Sw. = *Pilea nummularifolia*.  
 U. NUTANS Poepp. = *Pilea nutans*.  
 U. OBEA Wall. = *Pilea umbrosa*.  
 U. OBTUSATA Vahl = *Pilea obtusata*.  
 U. ORBICULATA Rich. = ? *Pileæ spec.*  
 U. OREOPHILA Miq. = *Pilea trinervia*.  
 U. OVALIFOLIA Blume Bijdr. p. 504 = ?  
 U. OVALIFOLIA Buching. = *Fleurya peduncularis*.  
 U. PALMATA Forsk. = ? *Girardinia palmata*.  
 U. PALMATA Leschen. = *Girardinia palmata*.  
 U. PALUSTRIS Juss. = *Boehmeria cylindrica*.  
 U. PANICULATA Steud. = *Fleurya ruderalis*.  
 U. PANICULATA Roxb. Fl. ind. ed. 1832. v. 3. p. 589 (e Molucc.) = ?  
 U. PARASITICA Blume = ? *Pellionia Javanica*.  
 U. PARASITICA Forsk. Descr. p. 160 (U. *muralis* Vahl) = ?  
 U. PARIETARIA Poir. = *Pilea articulata*.  
 U. PARIETARIA Linn. = *Pilea ciliaris* ?  
 U. PARVA Blume = *Pouzolzia indica*.  
 U. PARVIFOLIA Hamilt. = *Pouzolzia viminea*.  
 U. PARVIFLORA Roxb. = *Memorialis parvifolia*.  
 U. PEDUNCULARIS E. Mey. = *Fleurya peduncularis*.  
 U. PEDUNCULARIS Smith = *Pilea trinervia*.  
 U. PELLUCIDA Labill. = *Pipturus incanus*.  
 U. PELTATA Blume = *Laportea peltata*.  
 U. PENDULA Willd. = *Pilea rupipendia*.  
 U. PENDULIFLORA Wall. = *Boehmeria macrophylla*.  
 U. PENTANDRA Roxb. = *Memorialis pentandra*.  
 U. PEPOIDES Steud. = *Pilea peploides*.  
 U. PETIOLARIS Sieb. et Zucc. = *Pilea petiolaris*.  
 U. PETIOLATA Steud. = *Fleurya æstuans*.  
 U. PHOTINIPHYLLA Cunnigh. = *Laportea photiniphylla*.  
 U. PILOSA Loureir. = *Acalypha Poiratii* Spreng. ex Prodr. 15. s. 2. p. 880 (ubi  
 errore sub U. *hispida* citatur).

- U. PILOSIUSCULA Blume = *Boehmeria platyphylla*, var.  
 U. PILULIFERA Burm. = *Fleurya interrupta*.  
 U. POLYSTACHYA Wall. = *Boehmeria platyphylla*, var.  
 U. PORTULACINA Spr. = ? *Pilea microphylla*, var.  
 U. PORTULACOIDES Spr. = *Pilea microphylla*.  
 U. PUBESCENS Wall. list, 4614, ex h. Roxb. (*U. Roxburghii* Steud.) = ?  
 U. PULCHELLA Link Enum. h. ber. alt. part. 2 p. 385. = ?  
 U. PULCHERRIMA Roxb. = *Sarcochlamys pulcherrima*.  
 U. PULEGIFOLIA Poir. = *Pilea pulegifolia*.  
 U. PULUS Steud. = ? *Laportea microstigma*.  
 U. PUMILA Linn. = *Pilea pumila*.  
 U. PUNCTATA Ham. = *Pouzolzia viminea*.  
 U. PUNCTATA Kunth. = *Pilea punctata*.  
 U. PUSILLA Poir. = *Australina pusilla*.  
 U. PUSTULATA Liebm. = *Laportea pustulata*.  
 U. PUTA Hamilt. = *Maoutia Puya*.  
 U. QUINQUENERVIS Wall. = *Memoralis hirta* et M. 5-nervis.  
 U. RACEMOSA Plum. = *Fleurya æstuans*.  
 U. RADICANS Swartz = *Pilea radicans*.  
 U. RADICANS Wight = *Pilea Wightii*.  
 U. RADULA E. Mey. = ? *Didymodoxæ spec.*  
 U. RECLINATA hort. = *Pilea crassifolia*.  
 U. REPANDA Blume = *Pipturus repandus*.  
 U. REPENS Swartz = *Pilea repens*.  
 U. REPENS Wall. = *Pouzolzia indica*.  
 U. RETICULATA Sw. = *Pilea reticulata*.  
 U. RHOMBEA Linn. = *Pilea rhombea*.  
 U. RIEDLEI Steud. = *Pilea lucens*.  
 U. RIVULARIS Schlecht. = *Droguetia pauciflora*.  
 U. ROTUNDIFOLIA Ham. = *Boehmeria platyphylla*.  
 U. ROTUNDIFOLIA Poir. = *Pipturus rotundifolius*.  
 U. ROXBURGHII Steud. nom. (*U. pubescens* Wall. list, 4614) = ?  
 U. RUBESCENS Blume = *Villebrunea rubescens*.  
 U. RUDERALIS Forst. = *Fleurya ruderalis*.  
 U. RUFA Swartz = *Pilea rufa*.  
 U. RUGOSA Swartz = *Pilea rugosa*.  
 U. RUGOSISSIMA Blume = *Boehmeria platyphylla*, var.  
 U. RUGULOSA Wall. = *Boehmeria rugulosa*.  
 U. RUPIPENDIA Poir. = *Pilea rupipendia*.  
 U. SALICIFOLIA Juss. = *Procris peduncularis*.  
 U. SANDWICENSIS Wedd. = *Hesperocnide Sandwicensis*.  
 U. SANGUINEA Blume = *Pouzolzia viminea*.  
 U. SCABRA Blume Bijdr. p. 504 = *alius ordinis*.  
 U. SCABRELLA Hornem. (*Ildefonsia* Mart. in lit. ad Schrank, *U. Ildefonsia* Steud.) = ?

- U. SCABRELLA Roxb. = *Boehmeria platyphylla*, var.  
 U. SCHIMPERIANA Hochstett. = *Fleurya restuans*.  
 U. SCRIPTA Hamilt. = *Pilea scripta*.  
 U. SEMIDENTATA Poir. = *Pilea semidentata*.  
 U. SEMISERRATA Smith = *Phenax Sonneratii*.  
 U. SERPYLLACEA Kunth. = *Pilea serpyllifolia*.  
 U. SERRULATA Swartz = *Pilea serrulata*.  
 U. SESSILIFLORA Swartz = *Pilea sessiliflora*.  
 U. SESSILIFOLIA Poir. = *Pilea sessilifolia*.  
 U. SESSILIFOLIA Blanco = ? *Boehmeria spec.*  
 U. SINUATA Blume = *Laportea crenulata*.  
 U. SMILACIFOLIA Wall. = *Pilea smilacifolia*.  
 U. SPHÆROCEPHALA Roxb. Fl. ind. ed. 1832 v. 3 p. 589 = ?  
 U. SPICATA Blume = *Boehmeria platyphylla*, var.  
 U. SPICIGERA Steud. = *Fleurya interrupta*.  
 U. ? SPINOSA Blume Bijdr. p. 504 = alius ordinis.  
 U. SPIRALIS Domb. = *Myriocarpus densiflorus*.  
 U. SQUAMIGERA Wall. = *Chamabainia squamigera*.  
 U. STIMULANS Linn. fil. = *Laportea stimulans*.  
 U. STIPULOSA Miq. = *Pilea angulata*.  
 U. STOLONIFERA Sw. = *Pilea stolonifera*.  
 U. STRIGOSA Blume = *Pouzolzia indica*.  
 U. SUBCORDATA Commers. = *Pilea lucens*.  
 U. SUBPERFORATA Wall. = *Boehmeria subperforata*.  
 U. SUFFRUTICOSA Roxb., Hamilt. = *Pouzolzia indica*.  
 U. SUCCULENTA Salzm. = *Pilea hyalina*?  
 U. ? SUPERBA Wall. list, n. 4625 = alius ordinis.  
 U. SYCOPHYLLA Bory = *Obetia ficifolia*.  
 U. SYLVATICA Blume = *Villebrunea sylvatica*.  
 U. TASMANICA Müller = *Australina Muellieri*.  
 U. TENACISSIMA Roxb. = *Boehmeria nivea*.  
 U. TENUIS Steud. = *Pouzolzia indica*.  
 U. TERNATA Heyne = *Memorialis concinna*.  
 U. TETRAPHYLLA Hochst. = *Pilea tetraphylla*.  
 U. THYMIFOLIA Kunth. = *Pilea globosa*.  
 U. TILIÆFOLIA Kunth. = *Urera Caracasana*.  
 U. TOMENTOSA Juss. = *Boehmeria platyphylla*, var.  
 U. TRIANDRA Blume = *Memorialis parvifolia*.  
 U. TRIANTHEMOIDES Sw. = *Pilea trianthemoides*.  
 U. TRILOBATA Poir. = *Pilea trilobata*.  
 U. TRINERVIA Roxb. = *Pilea trinervia*.  
 U. TRIPLINERVIA Poir. = ? *Pilea lucens*.  
 U. TRIPLIVERVIA Rottl. = *Pouzolzia indica*.  
 U. TRIPLINERVIA Hamilt. = *Pilea scripta*.

- U. TRIPLINERVIS Juss., Pers. = *Pilea lacens*.  
 U. TRIPLINERVIS Wall. = *Pouzolzia indica*.  
 U. TUBEROSA Roxb. = *Pouzolzia tuberosa*.  
 U. TUBERCULATA Anderson = *Fleurya æstuaus*.  
 U. ULMIFOLIA Kunth. = *Urera Caracasana*.  
 U. ULMIFOLIA Vahl = *Phenax Sonneratii*.  
 U. UMBELLATA Blanco Fl. de Filip. ed. 1 p. 696 (sub nom. *U. ferax* in ed. 2 =  
 U. UMBELLATA Bory = *Pilea umbellata*.  
 U. UMBROSA Wall. = *Pilea umbrosa*.  
 U. URAGERA Steud. = *Boehmeria platyphylla*, var.  
 U. UROPHYLLA Wall. = *Villebrunea appendiculata*.  
 U. VARIEGATA Spreng. = *Pilea sessiliflora*.  
 U. VENOSA Wall. = *Boehmeria rugulosa*.  
 U. VERBASCIFOLIA Poir. = *Pilea verbascifolia*.  
 U. VERONICEFOLIA Dietr. ex Steud. nom. = ?  
 U. VERRUCOSA Liebm. = *Urera Caracasana*.  
 U. VERRUCOSA Moon = *Memoralis aquatica*.  
 U. VESICARIA Roxb. = *Pouzolzia vesicaria*.  
 U. VILLOSA Thunb. Fl. Jap. p. 70 = ?  
 U. VILLOSA Salzm. = *Fleurya æstuaus*.  
 U. VIMINEA Wall. = *Pouzolzia viminea*.  
 U. VIRGATA Forst. = *Boehmeria platyphylla*, var.  
 U. WALLICHIANA Steud. nom. (*U. paniculata* Roxb.) = ?  
 U. WHITLOWI Muehlenbg = *Laportea Canadensis*.  
 U. ZEYLANICA Burm. = *Girardinia Zeylanica*.

II. HESPEROCNIDE Torr. et A. Gr. in Bot. of Whipple's exp. p. 83. —  
 Urticæ spec. Wedd. olim.

Flores monoici, glomerulati, glomerulis axillaribus androgynis, pedicellis articulatis. *Masc.* Perigonium 4-partitum, segmentis elliptico-ovatis, intus concavis, extus setuloso-hispidis, alabastro in medio depresso. Stamina 4, antheris reniformibus. Pistilli rudimentum urceolato-obovoideum, hyalinum. *Fem.* Perigonium tubulosum, oblongo-ovatum, ore minuto bi-quadri-dentato, dentibus duobus majoribus, extus pilis hamatis undique hispidum. Ovarium rectum, oblongo-ovatum, sessile. Ovulum.... Stigma sessile, penicillato-capitatum, persistens. Achenium rectum, late ovatum, lenticulari-compressum, apiculatum, perigonio aucto membranaceo arcte vestitum. Semen pericarpio conforme. Albumen parcum. Embryo cotyledonibus transverse elliptico-rotundatis, basi et apice emarginatis, radícula conoidea subæquilongis. — Herbæ annuæ, altera oceanica altera californica, pilis urentibus donatæ; foliis oppositis, inciso-vel crenato-serratis, tri-quinque-nerviis, cystolithis oblongis, punctiformibusve; stipulis lateralibus liberis, floribus ebracteatis. — Genus a cl. aut. propter perigonium ♀ tubulosum, ad *Boehmerias* et præsertim ad *Pouzolzias* adductum, sed absque dubio inter *Ureras* collocandum. Conformatio floris feminei ab illa typi normalis tribus certo sat discre-

pat, et cæteri omnes characteres et imprimis facies cum illis *Urticarum* genuinarum egregie congruunt; nec prætermittendum exstare speciem *Urticæ* (v. *U. incisam*) cujus flores, quando anomali, illos gen. *Hesperocnide* omnino mentiuntur.

1. *H. TENELLA* (Torr. et A. Gr. l. c.), simplex aut vix ramosa, parce stimulosa, cæterum glabriuscula, foliis ovatis obtuse dentato-serratis, glomerulis laxiusculis. ☉ Ad scopulos obumbratos vallis Napa Californiæ. — Caulis gracilis 1-2-decimetralis. Folia latiuscule ovata, 1-2-centimetralia, dentibus apice rotundatis, utrinque sparsim stimulosa, subtiliter ciliata. Petiolus limbi plusquam dimidio brevior. Stipulæ ovatæ parum conspicuæ. Glomeruli 15-20-flori, floribus masculis inter femineos 1-2 mox deciduis. Perigonium fructiferum 1-2 millim. longum, ovatum, compressum, pilis brevibus hamatis undique obsitum. Achænium conforme apiculatum utrinque obscure punctatum.
2. *H. SANDWICENSIS*, ramosissima, setoso-hispida pubescensque, foliis ovato-lanceolatis inciso-serratis, dentibus sæpius obtusis, glomerulis densifloris. ☉ In montibus Mouna-Roa et Kaak insulæ Owhyhee (Macrae). — *Urtica Sandwicensis* Wedd. Monogr. p. 67. — Caulis adscendens, in speciminibus suppetentibus circiter 4-decimetralis, ramosissimus, ramis diffusis, aliis axi fere æquilongis, aliis brevioribus, omnibus undique subæqualiter breviterque stimuloso-hispidis. Folia majora vix sesquicentimetralia, 6 millim. circiter lata, at pleraque subcentimetralia, apicem versus attenuata vel subacuminata, basi obtusata, utrinque sed præsertim subtus hispidio-pubescentia, dentibus illis foliorum *Urticæ* urentis fere similibus sed pro limbi magnitudine latioribus paucioribusque, apicali elliptico; pilis inferioris paginæ in nervis primariis longioribus crebrioribusque; nervis secundariis tertiariisque haud perspicuis; lamina exsiccata supra obscure viridi, cystolithis minimis punctiformes et oblongos inter pilos exhibente, subtus nonnihil cinerascens; petiolo 1-2-millimetrali. Stipulæ parvæ, oblongæ, obtusæ, hispidulæ. Glomeruli sessiles, ii cujusque nodi confluentes, etiam fructiferi vix piso dimidio æquimagni; floribus masc. paucis, perigonio parum hispidio, alabastro vix millimetrali. Flores fem. subsessiles, perigonio urceolato minimo 2-4-denticulato, fructifero ovato, achænium conforme 1½ millim. longum læve pallideque fuscum arcte involvente. (v. s. in hb. Lindley et DC.)

II. NANOCNIDE *Blume*, *Mus. bot. lugd.-bat.* (1) 2. p. 154; *Wedd. Monogr.* p. 289.

Flores monoici, glomerulati, glomerulis in singulis axillis solitariis geminisve unisexualibus, masculis laxiusculis pedunculatis, femineis densis sessilibus, pedicellis florum ♂ articulatis. *Masc.* Perigonium 4-vel 5-partitum, segmentis ovatis intus concavis extus setulosis glabrisve, alabastro in medio depresso. Stamina 4-5, antheris reniformibus. Pistilli rudimentum late obovoideum, hyalinum. *Fem.* Perigonium 4-partitum, segmentis inæqualibus erectis, 2 majoribus exterioribus carinatis, 2 interioribus angustioribus planiusculisque. Ovarium rectum, oblongum. Ovulum imo loculo insertum, erectum, funiculo brevi. Stigma sessile, penicillato-capitatum, persistens. Achænium rectum, late ovato-ellipticum, compressum, læviusculum, perigonio aucto membranaceo omnino inclusum. Semen pericarpio fere conforme. Albumen parcum. Embryo cotyledonibus ellipticis, radícula conica. — Herbæ annuæ, insulas sinenses et Japoniam incolentes, inermes (?) caulibus gracilibus diffusis vel prostratis. Folia alterna, grosse crenata, tri-quinque-nervia, membranacea, cystolithis oblongis linearibusve. Stipulæ laterales, li-

---

(1) Vide quod dicitur in monographia mea (pp. 54, 90 et 588) de temporibus in quibus datum est opus Blumeanum.

beræ. Flores basi bracteati, masculi decidui, feminei persistentes. — Hoc genus a cl. Blumeo prope Pellioniam inter Prærideas collocatur; at, meo iudicio, ejus affinitates veræ sunt cum Urereis, quibus conformatione perigonii et stigmatis, necnon defectu staminodiorum et ipso habitu bene congruit. Plantæ exsiccatae stimulis veris expertes videntur, sed vivæ forsân urunt.

1. *N. JAPONICA* (Bl. l. c. t. 17), foliis late ovatis aut flabelliformibus grosse crenato-dentatis, junioribus margine et in nervis subtus pubescentibus, segmentis perigonii masc. 5-partiti glabris extusque plica seu cristula transversali auctis, illis feminei sub apice setula instructis (Blume). ☉ In Japonia ubi vocatur Katen-sô; Corea (Oldham n. 766, Wilford); Formosa (Swinhoe); Loo-Choo (Wright n. 301). — Wedd. l. c. t. 9 B. — Herba circiter bidecimetralis, e basi ramosa, ramis subfastigiatis. Folia 2 centim. longa, 1½ centim. lata, obtusa basi subcordata vel truncata aut vage cuneata integraque, tri-tripli aut quintupli-nervia, crenis acutiusculis obtusisve in utroque latere limbi 5-7, lamina exsiccata supra striolata (Blume); petiolis inferioribus longissimis, superioribus limbo brevioribus illumve æquantibus. Flores masc. umbellati (Blume), pedunculo petiolum superante. (v. s.)

2. *N. LOBATA*, foliis ambitu late ovatis rotundatisve in dimidia parte superiore 3-5-lobatis rarius hic et illic omnino integris utrinque sparsim pubescenti-pilosis, perigonii masculi 4-partiti segmentis extus convexis sicutique illa perigonii feminei densiuscule setuloso-hispidis. ☉ In insulis sinensibus Loo-Choo (Wright). — Herba, caulibus fasciculatis gracilibus prostratis 1-4 decim. longis retrorsum hispidis aut glabratiss, internodiis elongatis. Folia 6-15 millim. longa et lata (saltem in specim. obviis), obtusissima; basi ad insertionem petioli sæpius breviter cuneata, trinervia, lobis obtusis subæqualibusque vel exterioribus paulo minoribus, pilis in nervis paginae inferioris numerosioribus; lamina exsiccata tenuiter membranacea læte viridi et fere concolore, utrinque cystolithis oblongis sparsis notata; petiolo limbum æquante vel breviori. Stipulæ ovatae, haud millimetrales, obtusæ, ciliatae. Cymulæ masc. parvæ, in iisdem axillis ac femineæ aut in axillis distinctis nascentes, pedunculo gracili petiolum æquante setuloso, pedicellis apice incrassatis et cum basi perigonii articulatis, alabastro semimillimetrali. Glomeruli fem. in axillis ultimis (nonnunquam aphyllis) caulium præcipue obviis, 3-5 millim. lati, floribus basi bracteis hyalinis fimbriatis stipatis; perigonio fructifero millimetro nonnihil longiore, segmentis exterioribus admodum carinatis laterales subspathulatos fere omnino involventibus achæniumque ellipticum compressumque arcte includentibus. (v. s.)

IV. *OBTETIA* Gaudich, Bot. Voy. Bonite, t. 82; Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 178; Monogr. p. 106. t. 1 B. — Urticæ spec. auct.

Flores dioici, in glomerulos laxè cymoso-paniculatos dispositi. *Masc.* Perigonium 5-partitum, lobis ovatis obtusis puberulo-hispidis. Stamina 5. Pistilli rudimentum globosum. *Fem.* Perigonium 4-phyllum, foliolis inæqualibus ovatis rotundatisve glabris. Ovarium primo erectiusculum, mox obliquum. Ovulum erectum. Stigma subsessile, ovatum villosumque s. penicillato-capitatum, persistens, denique inflexum. Achænium oblique ovatum, compressum, margine acutatum et in utraque facie scrobiculato-concavum læviusculumque, pedicello recte insidens perigonioque valde aucto et tenuiter membranaceo laxè involucrium. Semen pericarpio subconforme. Albumen parvum. Embryo cotyledonibus transverse oblongo-rotundatis, basi et apice emarginatis, radícula conica paulo longioribus. — Frutices madagascarienses, pilis urticantibus parce armati; folia alterna, lobata v. grosse crenata, penninervia aut subpalmatinervia, cystolithis punctiformibus. Stipulæ liberæ, laterali-axil-

lares, subpersistentes. Flores bracteis linearibus intermixti, perigonii fructiferis diu persistentibus, pedicellis teretibus.

1. O. FICIFOLIA (Gaudich. Bonite, t. 82), foliis polymorphis ut plurimum sub-7-lobatis, lobis (raro nullis) crenulatis dentatisve, subtus pubescentibus sæpiusque canescentibus, paniculis ramosissimis.  $\text{\textcircled{J}}$  In ins. Bourbon, præsertim in lapidosis (Commerson, etc.). — Wedd. Monogr. p. 107. *Urtica ficifolia* Poir., Encycl. 4. p. 638. *U. sycophylla* Bory, voy. Iles afr. I. 281. *Urtica?* *ficifolia* Gaudich. Uran. p. 496. — Subinermis. Ramuli crassi, basi denudati, crebre cicatricosi epidermideque cinerea vestiti, apice foliosi breviterque pubescenti-tomentosi. Folia in apice ramulorum magis minusve approximata, 6-20 cm. longa et lata, basi late cordata, palminervia et fere pinnato-lobata, lobis plus minus profundis sæpius 7 rarissime 3 vel paucioribus aut fere nullis nonnunquam 8-9 aut pluribus (*Urt. heracleifolia* Juss.), plerumque lanceolatis ovatisve, apice acuminatis acutisque, rarissime linearibus, inferiore (majore) et apicali persæpe hinc inde lobulis nonnullis minoribus instructis, sinubus angustis fundoque rotundatis, crenis sæpius obtusis, lamina exsiccata submembranacea nervis subtus prominulis, folio juniore supra sparsim hispido rugosuloque mox glabrato subtus molliter incano-puberulo vel pubescente, nervis primariis breviter sparsimque stimulosis, petiolo limbo æquilongo vel breviori puberulo stimulosi aut inermi. Stipulæ fere liberæ, ovatae, 1-2-centimetricæ, puberulæ, ciliatæ. Paniculæ masc. et fem. ex axillis foliosis nascentes, longiuscule pedunculatæ, petiolos superantes, erecto-patulæ, ramosissimæ, rhachi ramulisque patulis subtiliter puberulis, glomerulis masc. fere discretis vix numerosis pisi circiter magnitudine, floribus sessilibus lobis hispidulis, alabastris bimillimetralibus; glomerulis fem. parvis numerosissimis, perigonii fructiferi folioliis achænio oblique ovato vix millimetrali duplo triplo longioribus. — Negat Bory de St-Vincent contactum hujus arboris esse noxium, et idem asserit de cæteris *Urticis* insulas Mascarenas incolentibus; verumtamen non dubium est viatorem quoad nonnullas in errorem inductum fuisse (v. s.)

2. O. MADAGASCARIENSIS (Wedd. in Ann. sc. nat., sér. 4. I. p. 178), foliis oblongo vel elliptico-lanceolatis obtuse crenatis glabriusculis, paniculis parce ramosis.  $\text{\textcircled{J}}$  In ins. Madagascar (Commerson). — Wedd. Monogr. p. 108. t. I. B. f. 1. *Urtica madagascariensis* Juss. mscr. *Urtica?* *madagascariensis* Gaudich. Uran. p. 496. Frutescens, ramis teretibus epidermide obscure cinereo-luscescente subtiliterque pubescente vestitis. Folia in ramulis brevissimis lateralibus approximatique fasciculata, 5-10 centim. longa, 1  $\frac{1}{2}$ -3 centim. lata, a medio sæpius utrinque attenuata, basi obtusata vel haud raro ante petioli insertionem emarginata, apice obtuse acuminata, crenis grossiusculis subæqualibus obtusissimis, penninervia, nervis primariis utrinsecus 8-10 tenuibus subtus vix prominentibus intra marginem reticulatim anastomosantibus, supra glaberrima, costa subtus sparsim retrorsumque stimulosi-hispida vel inermi et haud raro puberula, nervis folii junioris pilosulo-hispidis, lamina exsiccata membranacea, petiolo subcentimetrica breviter pubescente. Stipulæ fere liberæ, lanceolatæ, petiolo paulo breviores, glabriusculæ, foliis delapsis remanentes, ramulos squamarum instar imbricatim tegentes. Paniculæ masc. non visæ. Paniculæ fem. ex ramulis aphyllis (?) vel inter folia nascentes iisque dimidio circiter breviores, oblongæ, patulæ, brevissime pedunculatæ, rhachi gracili ramulisque puberulis, floribus in apice ramulorum glomerulatis, perigonii fructiferi laciniiis achænio oblique ovato 1 millim. longo subdimidio longioribus (v. s. in hb. Mus. par., Juss. et Lessert.).

V. FLEURYA Gaudich. Bot. Voy. Uran. p. 497; Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 204; Monogr. p. 109. t. 1. A; Miq. in Mart. Fl. bras. fasc. 12. p. 195. — *Urticæ* spec. auct. — *Schychowskia* Endl. in Annal. des Wien. Mus. 1. p. 187. t. 13. — *Urtica*, d. *Fleurya* Endl. Gen. pl. p. 283.

Flores monoici v. dioici, in cymulas dichotomo-scorpioideas aut in glomerulos modo paniculatos modo subdistiche racemosos digesti, mas-



culi et feminei in eadem aut in distinctis inflorescentiis nascentes, paniculis racemisve in singulis axillis solitariis, pedicellis florum utriusque sexus articulatis. *Masc.* Perigonium 4-5-partitum, segmentis ovatis lanceolatisve glabris aut pubescentibus v. sub apice stimuligeris, alabastro umbilicato. Stamina 4 v. 5. Pistilli rudimentum globosum v. clavatum v. sublobatum. *Fem.* Perigonium 4-partitum v. 4-lobum, segmentis s. lobis interdum fere æqualibus at sæpius inæqualibus, interioribus (s. lateralibus) tunc quam exteriores multo majoribus oblique ovatis planiusculis, exteriorumque altero (superiore) rotundato cucullato inermi v. pillo urente calcarato, altero (inferiore) lanceolato v. ovato vix perspicuo. Ovarium initio rectum, dein magis minusve obliquum, ovoideum. Ovulum juxta fundum loculi adfixum, obliquum, funiculo gracili. Stigma sessile, ovatum v. lanceolatum v. lineare, persistens et denique uncinatum reflexum, aut stigmata 2 sæpiusve 3 conniventia, uno mox elongato et solum bene conspicuo, binis aliis ad hujus basim velut stigmata adventitia s. rudimentaria persistentibus. Achænium oblique ovatum rotundatumve, compressum, margine abrupte acutatum interdumque anguste membranaceo-alatum, in utraque facie sæpissime scrobiculato-concavum tuberculatumque, pedicello oblique v. sæpius obliquissime insidens, perigonio parum aucto pro parte vestitum et cum illo ad maturitatem e pedicello desarticulato mox cadens. Semen endocarpio conforme. Albumen nisi juxta apicem seminis parvissimum. Embryo cotyledonibus transverse elliptico-rotundatis basi et apice parum emarginatis, radícula conica. — Herbæ annuæ, inter tropicos utriusque orbis diffusæ, stimulis armatæ aut fere inermes. Folia alterna, serrata v. crenato-dentata, trinervia, cystolithis linearibus. Stipulæ axillares, profundè bifidæ. Flores sæpissime ebracteati, masc. albidî v. rosei, feminei virentes, pedicellis horumce uni-bi-articulatis cylindricis rariusve sub perigonio lateraliter compressis. — Genus admodum naturale, a cæteris bene distinctum conformatione achæniorum necnon horumce cito lapsu.

1. *F. æSTUANS* (Gaudich. Uran. p. 196, addit. spec.), pilosa vel stimulosâ aut fere omnino inermis et glabra, foliis ovatis vel rotundato-ovatis, basi cordatis vel rotundatis cuneatisve grosse dentatis rariusve crenato-dentatis, paniculis androgynis aut interdum prorsus unisexualibus androgynis femineisque fastigiatis, florum fem. perigonio valde inæqualiter 4-partito, pedicello haud dilatato, stigmate brevissimo. ① In Americæ septentrionalis et præsertim meridionalis Antillarumque locis calidis cultis et incultis, juxta sepes et ad oras sylvarum frequens; in Africa orientali et occidentali, in ins. mascarenis necnon in ins. indicis etiâ obvia. — Grisebach Fl. Br. W. Ind. p. 154. Caulis erectus vel adscendens, 2-10 decim. altus, vulgo subsimplex. Folia 5-25 centim. lata, breviter acuminata, acumine triangulari vel lanceolato acuto aut obtusiusculo, limbo ima basi integro aut in toto margine dentato, dentibus baseos minoribus et magis approximatis, cæterum subæqualibus; lamina exsiccata tenuiter membranacea, subtus pallidior; petiolo gracili, longitudine varia, limbum sæpissime sequante. Stipulæ circiter ad medium vel altius bifidæ, laciniis lineari-subulatis ciliatis glabrativæ. Flores masc. cum femineis in fisdem cymulis intermixti et mox decidui aut in pedunculis distinctis glomerulos pisiformes conficientes, perigonio 4-vel 5-partito; cymulis panicularum androgynum et fem. alternis, breviter pedunculatis, laxifloris, bis terve dichotomis deinceps fastigiato-vel fasciculato-scorpioideis, explanatis, racemum compositum subdistiche ramosum pedunculatumque efformantibus. Flores fem. initio rectiusculi, sed in pedicellum subcylindricum mox incurvi demumque sub angulo acutissimo reflexi; perigonii fructiferi aucti segmentis interioribus sive lateralibus quam exteriores multo majoribus, ovatis rotundatisve, margine glanduloso-ciliatis, aliorum altero superiore pileato stimulo longiusculo dorsali frequentissime armato, altero inferiore lineari. Achænium ovatum, 1-1½ millim. longum, præsertim versus apicem anguste membranaceo-alatum,

perigonii segmentis majoribus haud dimidio longius, stigmate ovato vel lanceolato valde recurvo apice uncinatum et in utraque facie area impressa tuberculataque ovata v. rotundata ornatum. (v. s.).

α, *Linnaæana*, caule imprimis superne petiolisque et pedunculis piloso-hirsutis, stimulis diaphanis cum pilis aliis longioribus innocuis mixtis, foliis basi rotundatis cordatisve in toto margine grosse crenato-dentatis, dentibus late triangularibus vel triangulari-semirotundatis juxta petiolum minoribus acutiusculis obtusisve, limbo utrinque sparsim stimulosos nervis subtus magis hirsutis, floribus monoicis (raro dioicis), paniculis plerumque ovatis androgynis raro unisexualibus, masculis tunc ut in var. β dispositis, perigonii masc. sæpius 4-partiti segmentis hispidis, achænio frequenter millimetrum vix metiente scrobiculis lateralibus rotundatis. — In Antillis (Cl. Richard), Fernando Po (Mann, n. 154), Monbaze (Boivin), Abyssinia (Schimper, Quartin-Dillon et Petit), Java (Zollinger, n. 1878). — *Urtica racemosa* Herb. Burm. U. æstuans Linn. Spec. pl. p. 1397; Jacq. Hort. Schoenbr. 3. t. 388. ? U. divaricata Forsk. Descr. pl. Egypt. et Arab. cent. 6. p. 160. ? U. hirsuta Vahl Symbol. bot. 1. p. 77. U. Schimperiana Hochstet. in Bot. Zeit. 33. p. 259; Walp. Ann. 3. p. 412. *Fleurya æstuans* Gaudich. Uran. p. 196; Bl. Mus. bot. Lugd. Bot. 2. f. 11, Wedd. Monogr. p. 112, exclus. var. β; Miq. Fl. Ind. bot. 1. pars 2. p. 288. F. ingrata Miq. in Zolling. System. Verzeichn. p. 106.

β, *glandulosa*, caule petiolisque et pedunculis glanduloso-pilosis, paniculis ut in typo androgynis sæpiusve unisexualibus, muscularum glomerulis paucis vel numerosis capituliformibus pisi circiter mole in apice ramulorum pedunculi bis ter quaterve dichotomi solitarii v. laxè aggregatis, pedunculis in singulis axillis solitariis vel geminis raro omnino simplicibus, perigonio masc. vulgo 5-partito, panicularum configuratione i. androgynarum simili, cæt. ut in var. α. — In Brasilia orientali (Martius, hb. fl. bras. n. 84); Peruvia (Pavon), necnon in littore orientali Africæ ad Zanzibar (Boivin). *Fleurya glandulosa* Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 205. F. æstuans β glandulosa ejusd. Monogr. p. 112. (v. s. in hb. Mus. par., DC. et Boiss.)

γ, *racemosa*, caule petiolisque et pedunculis glabriusculis vel glaberrimis cæterum plus minus stimulosus aut fere inermibus, foliis vulgo amplis cordatisque sed interdum ad insertionem petioli rotundatis cuneatisve breviter acuminatis ima basi integris aut sæpius cum reliquo limbo dentatis (dentibus sæpe subacuminatis raro obtusis) utrinque sparsim stimulosus, paniculis fere pyramidalis, floribus masc. mox deciduis et fem. vulgo in iisdem cymulis intermixtis, muscularum perigonio ut plurimum 4-partito, segmentis sæpiissime apice piliferis, achænio 1½-1¼ millim. longo scrobiculis lateralibus ovatis. — Frequens in insulis Indiæ occidentalis; Nicaragua (Orsted), Venezuela (Fendler n. 1242), Cumana Funcke), Nova Granata (Humb. et Bonpl. Goudot, etc.), Peruvia (Pœppig, n. 2086), Guiana (Kegel, Schomburgk, etc.), Brasilia prope Bahia et Para (Salzmänn), Pernambuco (Gardner), prov. Rio et Minarum (Martius), Caravellas (Max. Neuwied), etc. — *Urtica racemosa* Plum. Cat. pl. Am. p. 11. U. caravellana Schrank, Pl. rar. hort. monac. t. 82. U. latifolia Rich. in Act. Soc. hist. nat. par. 1. p. 113; Persoon, Synops. 2. p. 554. U. corylifolia Juss. herb. U. nemorosa Kunth in nov. gen. et sp. 2. p. 42. U. divergens Meyer, Primit. fl. Esseq. p. 264. U. cordata Steud. Nomencl. 2. p. 734. *Fleurya cordata* Gaudich. Uran. p. 497, Bonite t. 83; Wedd. Mon. p. 110; Miq. Fl. Ind. bot. 1 (2) p. 227. F. caravellana Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 204. F. æstuans Miq. in Mart. Fl. bras. fasc. 12. p. 196. ? *Urtica Gaudichaudiana* Henslow, in Ann. of nat. hist. 1. p. 341. t. 12. (v. s.)

δ, *tuberculata*, caule tuberculato-hispido, petiolis pedunculisque hispidis, cæt. ut in præced. — In ins. Galapagos (Andersson). — U. tuberculata Andersson, Fl. Galapag.

ε, *petiolata*, caule petiolisque et nervis paginæ inferioris foliorum peduncu-

lisque creberrime breviusculeque stimulosi, cæt. ut in typo. — In ins. mascarenis (Commerson, Gaudich.), et Timor (Riedlé). *Fleurya petiolata* Decsne Herb. Timor p. 162.

2. *F. cymosa* (Wedd. Monogr. p. 113), caule pilis minutis patentibus puberulo aliisque longioribus robustioribusque insuper instructo, foliis late ovatis acuminatis basi subcordato-peltatis grosse dentatis utrinque puberulis, inflorescentiis androgynis fastigiatis, superioribus quam folia longioribus, perigonii fem. inæqualiter 4-partiti pedicello cylindrico, stigmate crasso elongato. ④ In ruderalis et hortis prope Bataviam frequentissima vulgoque a Malalis Saroni dicta. — Miq. Fl. ind. bat. (2) p. 228. *Urtica* (*Fleurya*) *cymosa* Hassk. Pl. jav. rar. p. 200. Herba stricta, robusta, 6-12 decim. alta, caule simplice aut subramoso, pilis majoribus apice subclavato-incrassatis. Folia 5-16 centim. longa, 3-13 cm. lata, 3-nervia, utrinque puberula, supra atro-viridia, subtus pallida. Stipulæ a basi ad medium connatæ. Inflorescentiæ in eodem cum foliis plano explanatæ, superiores sæpe 2-decimetrales foliisque longiores, inferiores multo minores, pedunculo 2-3-chotomo petiolis longiore ramisque et pedicellis puberulis; floribus umbellatis aut subcapitatis in apice ramorum brevium aggregatis, masculis præcocius inter femineos immixtis, perigonii albidii 5-lobi lobis in juventute pilo erecto apice clavato instructis. Flores fem. perigonio 4-partito, segmentis exterioribus anguste linearibus concavis subfalcatis, superiore brevior, duobus interioribus s. lateralibus subrotundis planis. Achænium securidiforme, anguste marginatum (margine inferiore convexo) tenuissimeque subalatum, compressum, tuberculatum, stigmate reflexo coronatum. (Descript. ex Hasskarl, l. c.) — A *F. æstuante* præsertim stigmate elongato differre videtur.

3. *F. cuneata* (1), magis minusve stimulosa, cæterum glaberrima, foliis rhomboideo-ovatis acutis inferne late cuneatis vel rotundato-truncatis excepta basi grossiuscule dentato-serratis (serraturis utrinque 4-6), inflorescentiis corymbiformibus androgynis petiolo brevioribus, pedicello floribus fem. cylindrico, perigonio ejusdem inæqualiter 4-partito, stigmate brevissimo. ④ In cultis ruderalis et sylvis insulæ Cubæ (Poeppig, Despreaux, Ramon de la Sagra, Wright, etc.). — *Urtica cuneata* A. Rich. in Ram. de la Sagra Hist. ins. Cub. Bot. 2. p. 224. *Fleurya umbellata* Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 205; Monogr. p. 114. — Caulis erectus, 1-3 decim. altus, succulentus, simplex v. a basi ramosus, striatus, stimulis longis conspersus, exsiccatus cystolithis oblongis creberrime albo-punctatus. Folia 1-5 centim. longa, 2-3  $\frac{1}{2}$  centim. lata, paulo infra medium late cuneato-attenuata v. rotundata, rarius truncata, sursum sensim attenuata serraturisque 4-6 triangularibus acutis utrinque instructa, utraque pagina stimulis longis et raris armata, lamina exsiccata tenuiter membranacea, cystolithis linearibus sparsis juxta nervos vero numerosioribus, petiolo gracili limbum longitudine superante vel æquante sparsim stimulosus. Stipulæ in planta exsiccata vix conspicuæ. Cymulæ solitariae, axillares, corymbiformes (umbellas referentes), subsessiles petiolisque breviores, rhachi bis terque dichotoma, ramulis pedicellisque subfasciculatis. Flores masc. ad basin cymularum subsolitarii et mox labentes, perigonii 4-partiti segmentis glabriusculis; fem. ut in *F. æstuanti*, achænio 1 millim. vix longo. — Species optima, characteribus essentialibus constantissimis. (v. s.)

β, *grossa*, caule parce ramoso, foliis ovatis 2-4 centim. longis, petiolo limbum sæpe æquante eove longiore, serraturis grossis utrinque 3-4. In ins. Cuba, cum typo (Wright, n. 1453 pro parte, ann. 1860-1864).

γ, *ovata*, caule a basi valde ramoso, stimulis quam in typo crebrioribus armato necnon inter stimulos hispido, foliis 1-1 $\frac{1}{2}$  centim. longis late ovatis basi rotundatis supra basin integram dentato-serratis. Cum præced. (Wright, n. 1453, pro parte.)

δ, *subintegra*, fere omnino inermis, caule gracili præsertim superne ra-

---

(1) Nomini specifico a me huic plantæ olim dato regulæ 57 codicis botanici contrario habui necesse alium, prioritatis causa dignius, subjicere. Item, etsi non semper libente animo, alia sat numerosis in pag. seq. occurrentis.

β, *mitis*, caule glabrato aut apice pubescente, inermi. ④ In Africa australis (Drège). *Urtica mitis* E. Meyer, l. c. *Fleurya mitis* Wedd. in Ann. sc. nat. l. c.

7. F. GROSSA (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 183), sparsim stimulosā, cæterum undique glaberrima, foliis deltoideo-ovatis acuminatis basi subtruncatis grosse acuteque dentatis, paniculis unisexualibus vel rarissime androgynis laxiuscule pyramidalis petiolo brevioribus, floribus femineis perigonio demum inæqualiter 4-partito, pedicello cylindrico, stigmate lanceolato brevi. ④ In Promontorio Bonæ spei (Drège; Zeyher, n. 3865). — Wedd. Monogr. p. 119. t. 1. A. f. 1-3. *Urtica grossa* E. Meyer, mscr. in Sched. exs. Dreg. Caulis erectus, 2-6 dm. altus, subsimplex, stimulis longiuscule pedicellatis robustisque hic et illic armatus. Folia 3-8 centim. longa, 2½-7 centim. lata, basi rectiuscule latissimeque truncata vel sæpius prope petiolum nonnihil rotundata obsoleteque peltata, acumine anguste lanceolato, dentibus utroque latere 5-7 triangulari-acuminatis acutissimis, in nervis stimulis paucis majusculis dispersa, lamina exsiccata tenuiter membranacea supra subtusque concolore, cystolithis punctiformibus parum conspicuis, petiolo gracili quam limbus brevior vel illi subæquilongus stimulis robustis retrorsis armato. Stipulæ minimæ, angustæ, haud profunde bifidæ. Flores sæpius monoici. Inflorescentiæ masc. inferiores, breviter pedunculatæ, inermes, floribus vix numerosis in rhachi communi vel in ramulis lateralibus brevibus incurvisque glomerulatis brevissimeque pedicellatis, perigonio 5-partito. Inflorescentiæ fem. ex axillis superioribus ortæ, masculas forma referentes, floribus initio erectiusculis sed mox in pedicellum incurvis demumque omnino reflexis, perigonio inæqualiter 4-lobo, fructiferi accreti lobis interioribus majoribus ovato-rotundatis, exteriorum altero superiore cucullato inermi, altero lineari-lanceolato. Achænium ovatum, 1½-1¼ millim. longum, margine membranaceo prorsus carens, perigonii segmentis circiter dimidio longius, scrobiculis lateralibus lævibus, stigmate valde incurvo apice uncinatum. (v. s.)

8. F. PODOCARPA, non aut vix stimulosā, fere undique hispido-pilosula, foliis ovatis acuminatis basi subtruncatis acutiuscule serratis, inflorescentiis parvis unisexualibus, femineis (e glomerulo unico constantibus) longe pedunculatis, perigonio fructifero campanulato 4-lobo pedicello haud dilatato, stigmate subulato basi brachiato, achænio ad insertionem valde angustato. ④ In Africa occidentali, prope Angiama (Barter). — Caulis gracilis, erectus, 3-4-decimetralis, simplex v. ramosus, demum fere glabratus. Folia 3-6 centim. longa, 2-4½ centim. lata v. majora, acumine lineari 5-10-millimetrali acuto, limbo ima basi integro et ad insertionem petioli sæpe breviter cuneato, serraturis subæqualibus, lamina exsiccata membranacea, petiolo gracili quam limbus brevior magis minusve hispido. Stipulæ parvæ, laciniis lanceolato-subulatis. Cymæ masculæ in axillis superioribus præsertim nascentes, pedunculo erecto petiolum dimidia fere longitudine sæpe superante, e glomerulis 2-3 pisiformibus approximatis confertæ, floribus subsessilibus vel breviter pedicellatis. Cymæ femineæ ex axillis inferioribus ut videtur præcipue nascentes; fructiferæ valde depauperatæ, pedunculo gracillimo petiolum superante (saltem in specimine viso), pedicellis 1-2 millim. longis, perigonio campanulato bimillimetrali paulo infra medium 4-lobo, lobis 3 majoribus triangularibus subæqualibus (intermedio carinato), quarto brevior oblongo concavo. Achænium oblongo-ovatum, 4 millim. longum, basi longiuscule angustatum s. pediculatum, undique acute marginatum et in utraque facie area ovata angusta vix tuberculata notatum, stigmate (s. stigmatis reliquo?) vix millimetrali basi incurvata appendice brevi simplice v. bifido adpresso aucta. (v. s. in hb. Mus. par. et Kew.)

β, *Mannii*, caule ramisque et petiolis glabriusculis, foliis multo majoribus crenato-serratis basi rotundatis v. subtruncatis. ④ In montibus Cameroun insulæ Fernando Po, altit. 750 metr. (G. Mann, n. 1950). — Herba plus quam metralis, ramis gracilibus internodiis elongatis. Folia 12-15 centim. longa, 6-8 centim. lata, acumine lineari crenis breviter acuminatis ad basim limbi minoribus, limbo utrinque sparsim hispido, petiolo gracili quam dimidius limbus longiore. Stipulæ sesquicentimetales, lineares,

circiter ad medium bifidæ. Cymæ masculæ....; femineæ e glomerulo parvo 6-10-flore constantes, floribus sessilibus v. breviter pedicellatis, perigonii 4-partiti segmentis tribus majoribus ovatis acuminatis, quarto oblongo-lanceolato dimidio brevior. Ovarium oblongo-ovatum, sessile, circiter millimetrale. Stigma subulatum, 4 millim. longum, basi appendice stigmatiformi bi-tri-millimetrali ad medium bifido auctum. (v. s. in hb. Kew.)

*Species non satis nota.*

F. LURIDA (Blume, Mus. lugd. bot. 2. f. XI, absque descript.) Patria? — Fructus in op. cit. delineatus illum F. æstuantis bene refert.

*Species exclusæ.*

F. GLECHOMÆFOLIA Miq.: alius ordinis.

F. GLOBULIFERA Miq.: alius ordinis.

F. HUMULIFOLIA Kunth = Girardinia heterophylla.

F. PHOTINIFOLIA Kunth = Laportea photinifolia

F. SCABRA Miq.: alius ordinis (1).

VI. LAPORTEA Gaudich. Bot. Voy. Uran. p. 498; Wedd. Monogr. p. 121; Miq. Fl. Ind. bat. 1 (2), p. 230; Seem. Fl. Vit. p. 238. — Urticæ spec. auct. — Dendrocide Miq. Pl. Jungh. p. 29. — Sclepsion Rafn. mscr. — Discocarpus Liebm. (non Klotzsch. nec Pfeiff.)

Flores dioici v. monoici, glomerulati, glomerulis in inflorescentias paniculiformes semperque unisexuales digestis, paniculis fem. et masc. apud species monoicas ex axillis distinctis ortis, masc. superioribus, pedicellis florum masc. articulatis. Masc. Perigonium 4-v. 5-partitum, segmentis ovatis glabrisque aut magis minus hispidis apice membranaceis, alabastro in medio depresso. Stamina 5. Pistilli rudimentum globosum. Fem. Perigonium 4-partitum v.-lobum, segmentis s. lobis fere æqualibus aut interdum admodum inæqualibus, nempe interioribus multo majoribus et plerumque ovatis rotundatisve glabris v. dorso et margine pilosulis exteriorumque altero (denique superiore) latiore rotundato et sæpius cucullato, altero lanceolato multo brevior. Ovarium junius rectum, sed mox obliquum, ovoideum. Ovulum erectum v. adscendens, funiculo brevi aut longiusculo suffultum. Stigma sessile, lineare elongatumque aut rarissime breve et oblongum, villosulum, in fructu persistens. Achænium oblique ovatum rotundatumve, raro ventricosum (pericarpio tunc nonnihil succulento), plerumque compressum et haud raro discoideum necnon juxta insertionem abruptissime angustatum, margine æquali vel parum incrassato faciebus læviusculis aut rarius granulosus pedicello plus minus oblique insidens perigonioque membranaceo subimmutato vestitum. Semen pericarpio conforme. Albumen tenue. Embryo cotyledonibus rotundatis apice subtruncatis basi emarginatis, radícula conica brevi. — Herbæ perennes fruticesve vel arbusculæ aut interdum arbores elatæ, in America boreali, Asia tropica et Oceania indigeni, stimulis raris aut crebris armati. Lignum cum exstat album succosum. Folia alterna, varie dentata aut integer-

(1) Planta sub nomine *Fleuryæ*? *rugosissimæ* a cl. Miquelio (Fl. Ind. Bot. suppl. prim. p. 412) descripta sine ullo dubio non est *Fleurya* et forsân cum *F. globulifera* et *F. scabra* ejusdem auctoris ad genus *Fatouma* Gaudich. inter *Artocarpeas* adscribenda.

rima, penninervia, cystolithis minutis punctiformibus inspersa. Stipulæ axillares, binerviæ, integræ v. bifidæ, deciduæ. Cymæ a. paniculæ sæpe distiche ramosæ. Flores haud raro ebracteati, ut plurimum virentes, pedicellis femineorum cylindricis aut varie dilatatis incrassatisve interdumque subfastigiato-coadunatis. Fructus diu persistens. — Genus nonnihil polymorphum propter varietates perigonii feminei, necnon habitus. Sectio prima ad Fleuryam characteribus quibusdam spectat.

Sect. I. *SCLEPSION* Wedd. monogr. — Flores ♀ perigonio 4-partito, segmentis valde inæqualibus, pedicellis sæpissime alatis.

1. *L. CANADENSIS* (Gaudich. Uran. p. 498), herbacea, foliis late ovatis acumina-  
tis basi subcordatis serratis parce stimulosus inermibusve supra plerumque  
asperulis, inflorescentiis femineis subterminalibus, pedicellis alatis, perigonii  
fructiferi segmentis lateralibus ovatis, achænio sub lente granuloso.  $\frac{2}{4}$  In Ame-  
ricæ septentrionalis fere totius regionibus temperatis et subtemperatis fre-  
quens. — Wedd. Monogr. p. 123. t. 2. c. f. 2-4; Asa Gr. Man. bot. Un. St.  
Urtica canadensis et U. divaricata Linn. Spec. pl. p. 1397 et auct; Schkahr  
Bot. Handb. t. 289. U. Whitlowi Muehlenb. in Green Cat. pl. N. York. Sclep-  
sion divaricatum Rafinesque mscr. Fleurya canadensis Benth. in Hook.  
Nig. Fl. p. 517. — Herba cæspitosa, caulibus metralibus altioribusve, erectis,  
subsimplificibus, magis minusve stimulosus demumque inermibus et (persis-  
tente basi stimulorum) interdum asperatis, inter pilos urentes glaberrimis,  
internodiis superioribus nonnunquam valde abbreviatis. Folia 6-16 centim.  
longa, 3-11 centim. lata, circiter a medio ad apicem late rotundata vel subcor-  
data, rarissime plane cordata, apice sensim vel subabrupte acuminata, acu-  
mine angusto acutiusculoque, serraturis foliorum etiam ejusdem rami parvis  
v. grossiusculis sæpius acutis nunc subacuminatis nunc extus vel utrinque  
convexis juxta petiolum sæpissime nullis, nervis primariis utrinsecus 7-8  
subtus perum prominentibus jugo inferiore omnino basilari vel subbasilari,  
secundariis tenuissimis rectiuscule vel reticulatim anastomosantibus, sicca  
membranacea, juniora supra sparsim breviterque stimulosa sed mox glabrata,  
limbo subtus in nervis (in costa præsertim) pilis urentibus numerosioribus  
et sæpe validioribus armato, petiolo gracili quam limbus breviori sparsim  
stimuloso vel subinermi. Stipulæ oblongæ, apice bifidæ, glabriusculæ. Flores  
monoici, cymis masculis et femineis plerumque in eodem caule. Inflorescen-  
tiæ masc. pedunculatæ, erecto-patulæ, foliis breviores, ovatæ, divaricato-  
ramosissimæ, floribundæ, ramulis pedunculoque puberulis stimulosisque aut  
inermibus, floribus laxiuscule glomerulatis mox deciduis subsessilibus pedi-  
cellatisve bracteolis fuscis stipatis, perigonii 5-partiti segmentis parce hispi-  
dis. Inflorescentiæ fem. ex axillis superioribus ortæ, folia apicalia approxi-  
mata vulgo superantes, pyramidatæ, longiuscule pedunculatæ, subsimplificiter  
distiche ramosæ, ramulis patulis pubescentibus stimulosisque, floribus in  
glomerulos parvos laterales breviterque pedunculatos digestis initio erectius-  
culis sed mox in pedicellum alatum incurvis demumque reflexis, perigonii  
inæqualiter 4-partiti segmentis lateralibus multo majoribus et in medio his-  
pidulis, superiore pileato hispido ciliatoque, inferiore lanceolato minimo.  
Achæmium oblique orbiculatum, 2-2½ millim. latum, valde compressum, bre-  
viter pedicellatum, sub lente fusco-marmoratum minutissimeque granulo-  
sum, perigonio fere duplo longius. (v. v. cult. et s.)

β, foliis basi rotundatis. — In hisdem locis ac typus (v. s.)

γ, foliis multo minoribus quam in typo, superioribus subverticillatis. —  
Cum præcedentibus. (v. s.)

- 2, *pustulata*, foliis basi rotundatis vel subcordatis serrato-dentatis, nervis  
paginæ inferioris petiolisque et ramulis inflorescentiæ sicut rami ipsi  
densiuscule stimulosi. In provincia mexicana Puebla, prope vicum Chi-  
nantla, circiter ad altit. 2200 metr. (Eratid?) — Urtica (Laportea) pustu-  
lata Liebmann, l. c. p. 294. Laportea pustulata Wedd. Monogr. p. 140.

— Folia, teste cl. Liebmanno, supra pustulata, sed illa speciminis a me visi haud asperiora sunt quam in typo. (v. s.)

2. *L. EVITATA*, herbacea, foliis oblongo-vel lanceolato-ovatis ovatisve acuminatis basi obtusis rotundatisve aut subcordatis argute serratis (dentibus apice setigeris) crebriuscule stimulosis utrinque sed praesertim supra asperis, inflorescentiis femineis subterminalibus, pedicellis alatis, perigonii fructiferi segmentis lateralibus ovatis oblongisve achenio suborbiculari laeviusculo plusquam dimidio brevioribus.  $\frac{1}{2}$  In sylvis montium temp. peninsulae Indiae orientalis et Himalayae humidioribus, alt. 1200-2400 metr. a Sikkim ad Kumaon, et in montibus Nilghiri, necnon in Japonia. — *L. terminalis* Wight, Ic. pl. Ind. 6. t. 1972; Wedd. Monogr. p. 125. t. 2. c. f. 1. *Urtica evitata* Wallich, Cat. n. 4588 (In herb. Wall. sub eodem nomine exstant specimina nonnulla Pileae umbrosae quae omnino innocua est). — Caulis erectus, 3-8 decim. altus, stimulis sparsim vel densiuscule conspersis, basi demum glabreatus, magis minusve asperatus, inter stimulos superne saepe hispidulus. Folia 4-20 centim. longa, 2-10 centim. lata, a medio apicem versus sensim attenuato-acuminata, acumine lanceolato v. lineari-acuto, serraturis triangularibus subaequalibus singulis stimulo cuspidatis, nervis primariis utrinsecus 5-7 (jugo inferiore basali vel subbasali dimidium limbum superante), secundariis tenuibus plerisque transverse inosculantibus, lamina (siccitate rigidula) supra plus minus asperato-hispida et utrinque (at subtus in nervis praesertim) stimulosa, stimulis paginae superioris saepius sparsis, petiolo gracili elongato brevi aut limbi nonnunquam longitudinem aequante plus minus stimulos. Stipulae lanceolatae, subcentimetales, integrae vel apice bifidae, dorso sparsim stimulosi-hispidae, deciduae rariusve persistentes. Flores monoici aut nonnunquam (?) dioici. Cymae masc. ut in specie praecedente, perigonii 5-partiti laciniis hispidulis glabriusculisve. Cymae fem. plerumque longius pedunculatae, pedunculo et ramis undique vel tantum uno latere densiuscule hispidulis stimulosque crebre sparsimve hirsutis, perigonii fructiferi segmentis lateralibus ovatis oblongisve obtusis. Achenium oblique rotundatum vel ovatum, 2-2 $\frac{1}{2}$  millim. longum, valde compressum, fusco-marmoratum interdumque minute granulosum; caetera ut in *L. canadense*. (v. s.)

$\beta$ , foliis utrinque laevibus, caet. ut in typo. (v. s.)

$\gamma$ , foliis ovatis breviter stimulosis supra fere laevibus, subtus asperulis, perigonii fructiferi segmentis longiuscule stimulosis, achenii fere ovati dimidium superantibus. (v. s.)

$\delta$ , foliis ovatis utrinque fere laevibus, perigonii fructiferi segmentis lateralibus rotundatis adpresse hispidis (nec stimulosis), achenio rubromarginato subaequimagnis. — In iisdem locis ac praecedentes et typus. (v. s.)

3. *L. ALATIPES* (Hook. fil. in Journ. Linn. Soc. 7, p. 215), herbacea, grosse setigera, foliis ovatis acuminatis basi rotundatis v. subcordatis acute dentato-serratis, inflorescentiis axillaribus, femineis quam folia longioribus, pedicellis alatis flabellatim connexis, perigonii fructiferi segmentis lateralibus oblique ovatis achenio margine incrassato dimidio brevioribus.  $\frac{1}{2}$  In montibus Caneoron ins. Fernando Po, altit. 2150 metr. (G. Mann, n. 1973.) — Caulis metralis, subsimplex, angulato-sulcatus, setis rigidis undique horridus et inter setas hispidus. Folia 10-15 centim. longa, 5-7 centim. lata, sensim acuminata, ab ima basi serrata, serraturis inferioribus minoribus, nervis primariis utrinsecus 5-6, jugo inferiore omnino basilari; lamina supra sparsim et inaequaliter stimulosa, pagina inferiore in venis venulisque omnibus hispida sed in costa nervisque majoribus solum stimulis veris armata, exsiccata membranacea, petiolo gracili 3-5 centim. longo horrido. Stipulae lanceolatae, breviter bifidae, dorso hispidulae. Flores dioici? Inflorescentiae masc. petiolos non aut vix aequantes, breviter pedunculatae, parce ramosae, patulae, e glomerulis vix numerosis discretis v. subapproximatis constantes, floribus pedicellatis et cito deciduis, perigonio 4-partito parce hispidulo. Inflorescentiae fem. longe pedunculatae ramis paniculatis subsecundis, ramulis divaricatis, pedicellis 8 millim. longis in laminas flabelliformes connatis, floribus in apice pedicellorum oblique impositis, perigonio post anthesin haud accreto. Achenium

- 3 millim. circiter latum, compressum, suborbiculare, stylo brevi filiformi coronatum, marginibus late incrassatis, faciebus planis vix granulatis. — Duabus præcedentibus et præsertim priori affinis, sed ab utraque bene distincta conformatione pedicellorum fl. fem., stigmatis brevitate, cæt. (v. s. in hb. Kew.)
4. *L. DECUMANA* (Wedd. Monogr. p. 127), herbacea, foliis ovatis vel elliptico-ovatis acuminatis basi cordatis subinæqualiter dentatis, dentibus acutis interdumque ipsis denticulatis, utrinque at præsertim in nervis paginæ inferioris crebriuscule stimulosis, inflorescentiis distiche subfastigiato-ramosis, floribus femineis sessilibus, stigmate filiformi quam ovarium multo longiore.  $\frac{2}{3}$ ? In ins. Moluccis ad oras sylvarum et juxta sepes, necnon in hortis olim et forsân etiam hodie culta (Zollinger, n. 1149). — Miq. Fl. Ind. bat. 1, pars 2. p. 230. *Urtica decumana* Rumph. Herb. Amb. 6. t. 20. f. 1; Roxb. Fl. Ind. 3. p. 587; Wight, Icon. fl. Ind. or. 2. p. 689. *Boehmeria macrothyrsa* Miq. in Zolling. Syst. Verz. 2. p. 103; Fl. Ind. bot. 1 (2). p. 254; Wedd. Monogr. p. 377. — Rami crebre stimulosi. Folia 1-2½ decim. longa, 6-15 centim. lata, ab ipsa basi dentata, anguste cordata, sensim acuminata, acumine etiam dentato, dentibus modo integris triangularibus acutis modo denticulo uno alterove hinc v. inde instructis, sinibus acutis aut obtusis, nervis fere ut in *L. terminali*, pagina superiore stimulis brevibus crebriuscule conspersa sed vix aut non asperata, nervis inferioris pilis urentibus tenuibus dense obsitis. Inflorescentiæ masc. non visæ. Inflorescentiæ fem. interdum maximæ (nempe una in specimine examinato cum pedunculo plus quam 4-decimetrali), distiche et alterne ramosæ, ramis inferioribus elongatis sequentibus sensim abbreviatis superioribus brevissimis, floribus in glomerulos parvos congestis initioque sessilibus, glomerulis breviter et æqualiter pedunculatis secus ramulos secundarios distiche racemosis, pedunculo ramis ramulisque hispido-stimulosis atque siccitate cystolithis minutis oblongis inter pilos punctulatis, perigonii 4-partiti segmentis valde inæqualibus ovatis, lateralibus multo majoribus ovarium superantibus anguste acuminatis exsiccatisque fasciculis raphidum crasse albo-lineatis, superiore cucullato piloso, ovario valde obliquo quam stigma subulatum villosulumque circiter triplo brevior. — Specimen unicum vidi femineum et absque fructu ex hb. Reinwardtiano benevole communicatum. Flores omnino sessiles sunt, sed fieri potest ut pedicelli rudimentarii post anthesin magis minusve elongati sint. Folia, teste Rumphio (Herb. Amb. 6. p. 47), in quotidiano usu sunt apud Indos ins. Moluccarum ad fricandum corpus ægrotorum, quo facto pars fricta mox rubet vesiculæque magna copia intumescunt, quodam modo sicut apud nos Sinapios ope aut Cantharidis. Planta dicitur, malice : *Danu gattal besaer*; hoc est : « Foliis pruritus magnus. » (v. s. in hb. Reinw. Lugd. bat.).
5. *L. PELTATA* (Gaudich. Uran. p. 498), arborea ? foliis late ovatis breviter acuminatis paulo supra basim rotundatam peltatis acute vel obtuse denticulatis supra magis minusve hispidis glabrativè subtus stimulosopilosis vel canescenti-pubescentibus, inflorescentiis femineis lateralibus, rami apicem haud superantibus, pedicellis sæpius alatis, perigonii fructiferi segmentis lateralibus late ovatis, achænio æqualiter tuberculato.  $\frac{3}{4}$  In insulis Java et Timor (Leschenault, Riedlé, Junghuhn). — Decsne Herb. Timor. p. 162; Wedd. Monogr. p. 126; Miq. Fl. Ind. bot. 1. c. *Urtica peltata* Blume, Bijdr. p. 496. *U. atrox* Leschen. Dendrocnide peltata Miq. Pl. Jungh. p. 30. — Ramuli herbacei aut lignescentes, vix crassi, stimulis brevibus gracillimisque hirsuti, foliosi. Folia in specim. visis 7-15 centim. longa, 5½-12 centim. lata, a medio ad basim late rotundata rariusve plus minus cordata, manifeste peltata, superne sensim attenuata apiceque acuminata, acumine acuto, denticulis triangularibus acutis acuminate aut obtusissimis, sinibus acutis vel obtusis, nervis primariis tenuibus utrinsecus 7-8 intra marginem arcuatim anastomosantibus, subtus vix prominentibus, jugo inferiore ex petioli insertionem orto limbi midium superante, nervis secundariis tenuissimis transverse connexis, pagina superiore sparsim stimulosopilosis, inferiore subtiliter pilosula v. canescenti-pubescente stimulisque gracilibus sæpe conspersa, lamina exsiccata tenuiter membranacea, petiolo gracili quam limbus longiore v. breviter puberulo retrorsumque stimulosopiloso. Stipulæ ovatæ, 6-8 millim. longæ, pubescentes.



Flores dioici (?). Cymæ masc. non visæ. Cymæ fem. ex axillis inferioribus ramulorum ortæ, pedunculatæ, subcorymbosæ, patulæ, ramosæ, ramulis gracilibus retrorsum stimulosi, glomerulis longiuscule pedunculatis, pedicellis inæqualibus plerisque late et crassiuscule alatis stimulisque tenuibus obstitis. Perigonium fructiferum in pedicellum sub angulo valde acuto inflexum, inæqualiter 4-lobum s. partitum, segmentis lateralibus late ovatis cum superiore nonnihil brevioribus magis minusve coalitis hispidisque, inferiore minimo lanceolato. Achænium ovatum, compressum,  $2\frac{1}{2}$  millim. longum, æqualiter tuberculatum, stylo incurvo coronatum, perigonio vix dimidio longius. (v. s.)

6. LAXIFLORA, herbacea ? foliis ovatis aut fere rotundatis breviter acuminatis supra basim rotundatam peltatis integerrimis supra puberulis glabrativse subtus pubescentibus, inflorescentiis femineis petiolo longioribus rami apicem superantibus ?, pedicellis crassis angulatis vel subalatis, perigonii fructiferi segmentis lateralibus lanceolatis, achænio læviusculo. ♀ In insula Java (Zollinger, it. sec. n. 3136). — Ramuli subherbacei, crassiusculi, puberuli, nodis approximatis. Folia suppetentia 7-12 centim. longa, 5-8½ centim. lata, acumine acuto, basi late rotundata, nervis primariis tenuibus utrinsecus 8-10 in margine desinentibus, subtus parum prominentibus, jugo inferiore dimidium limbi haud attingente, nervis secundariis tenuissimis transversim inter primarios anastomosatis; stimulis in utraque pagina ut videtur fere nullis; lamina exsiccata membranacea viridi; petiolo supra basim limbi adulti circiter centimetrum orto illoque longiore, puberulo vel glabriusculo et fere inermi, siccitate gracillimo. Stipulæ ovato-lanceolatæ, acuminatæ, puberulæ. Flores dioici ? Inflorescentiæ fructiferae (solum visæ) solitariae, folia superantes, paniculiformes, pedunculo quam petiolus brevioris suffultæ, ramulis gracilibus erecto-patulis hispidulis et sparsim retrorsum stimulosi, glomerulis breviter pedunculatis pedicellis plerisque brevibus angulatis subinermibusque. Perigonium fructiferum profunde inæqualiter 4-partitum, segmentis lateralibus (majoribus) et inferiore lanceolatis denticulatis, superiore (inferiore subæquilongis) ovato. Achænium oblique rotundatum,  $1\frac{1}{2}$  - 2 millim. longum, compressum, faciebz planis læviusculisque, margine nonnihil incrassatum, stigmatibus filiformi valde incurvo coronatum, ima basi (ut in præcedentibus) abruptissime angustatum perigonioque circiter duplo longius. (v. s. in hb. DC.).

7. L. VRISEIANA, arborea ?, foliis rotundato-ovatis breviter et argute acuminatis supra basim rotundatam peltatis integerrimis coriaceis supra glabrativse subtus velutinis, inflorescentiis femineis petiolo brevioribus ramulis dense pubescentibus stimulisque retrorso-hispidis. ♀ ? In ins. Java (de Vriese). — Ramulus unicus suppetens crassus. ligneus, pubescens, internodiis brevibus. Folia 16 centim. longa, 13 centim. lata, acumine brevi subulato, nervis primariis circiter 10 in margine desinentibus secundariisque subparallele transversis bene distinctis jugo inferiore medium limbum haud attingente, lamina exsiccata coriacea fusca, petiolo supra basim limbi altiuscule (circiter 2 centim.) enato 8 centim. longo puberulo inermi. Stipulæ . . . . . Flores dioici. Inflorescentiæ femineæ (floriferis solum visis) in apice ramuli approximata breviter pedunculatæ, floribus in glomerulos minimos discretos digestis subsessilibusque, perigonii segmentis ovatis acutis acuminativse puberulis, lateralibus cæteris duplo longioribus, ovario suboblique ovato quam stigma paulo brevior. — Præcedenti ut videtur a specimine sat manco suppetenti affinis. (v. s. in hb. Mus. par. ex hb. Lugd. bat.).

- 8 ? L. REPENS, herbacea, caule gracili elongato repente, foliis rotundato-ovatis breviter obtuseque acuminatis basi sinu profundo obtuseque cordatis longæ petiolatis toto margine subobsoleto crenatis tri-quingue-nerviis supra glabriusculis v. sparsim stimulosi nervis subtus subtiliter puberulis, cymis masculis longissime pedunculatis laxifloris. ♀ In insula Fernando Po (G. Mann). Habitu ab omnibus congeneribus distinctissima. Caulis subherbacei, glabri, pennam corvinam crassitudine æmulantes, ad nodos præsertim radicantes, internodiis 3-4-centimetralibus. Folia 6-12 centim. longa et lata, subabrupte acuminata, sinu basilari ad petioli insertionem rotundato, nervis primariis utrinque 5-6 subtus parum prominentibus sparsisque stimuligeris, jugo infe-

riore in margine juxta limbi medium desinente, lamina exsiccata membranacea supra circa basim stimulatorum et subtus secus venulas minores reticulatas cystolithis minimis oblongis linearibusve ornata, petiolo gracili 10-15-centimetroali puberulo sparsimque et breviter stimulosi. Stipulæ breves, ovatae, acuminatae. Inflorescentiæ subcorymbiformes et nodis defoliatis ortæ, pedunculo petiolos circiter æquante gracillimo subtiliter puberulo et sparsissime stimuligeri, floribus in apice ramulorum 3-5 subumbellatis, pedicellis bi-tri-millimetralibus glabriusculis basi bracteolatis, perigonii 5-partiti segmentis oblongo-lanceolatis acutiusculis, alabastro sesquimillimetrali. (v. s. in hb. Kew.). — Species sedis nonnihil incertæ ob defectum florum fem.

9. *L. BULBIFERA* (Wedd. Monogr. p. 139), herbacea, foliis ovatis vel ovato-ellipticis acuminatis basi rotundatis cuneatisve serrato-dentatis parce setosis supraque asperulis, inflorescentia feminea solitaria subterminali longiuscule pedunculata vix multiflora, pedicellis haud (?) alatis, stigmatate elongato cymis masculis ex axillis inferioribus una cum bulbillis 1-3 ovatis nascentibus. 4 ? In Japonia (Siebold). — *Urtica* (Laportea) bulbifera Sieb. et Zucc. Abhandl. der math. physik. Kl. der Baier. Akad. 4 (3). p. 214. — Caulis glaber vel setulis parvis obsitus. Folia 6-14 centim. longa, 3-7 centim. lata, longe petiolata. Stipulæ linearilanceolatae, siccitate membranaceæ. Bulbilli ad basim pedunculorum ♂ solitarii vel 2-3 glomerati, ovato-globosi, apiculati, magnitudine circiter seminis Cannabis. Inflorescentiæ masc. plures, solitarie e singulis axillis cum bulbillis 1-3 enatæ aut interdum geminæ absque bulbillis; perigonii 4-partiti segmentis acutis. Florum fem. perigonium valde inæqualiter 4-partitum, segmentis lateralibus multo majoribus post anthesin oblique orbiculari-obovatis, glabris. Achæmium (stigmatate brevius) orbiculari-ovatum, substipitatum, valde compressum, læviusculum. (v. s. in hb. Monac.).
10. *L. MILNEI* (Seemann, Fl. Vit. p. 238), arborea, foliis late ovatis acuminatis triplinerviis repando-dentatis stimulosi demum glabratis, inflorescentiis fem. axillaribus, pedicellis alatis, achænio oblique ovato acuto. ♂. In ins. Viti Lævu (Milne). — Folia 1½ decim. longa, 6-8 centim. lata, carnosula, costulis utrinque 9. Flores non visi. Achæmium fere magnitudine lentis.

Sect. II. *SARCOPIUS* Wedd. Monogr. — Flores fem. perigonio 4-partito, segmentis valde inæqualibus. Pedicelli fructiferi botuliformi-incrassati.

11. *L. GIGAS* (Wedd. Monogr. p. 129. t. 3 et 4. f. 1-9), arborea, foliis magnis late ovatis rotundatisve apice sæpius abrupte attenuatis breviterve acuminatis basi cordatis subintegris vel inæqualiter sinuato-crenatis serrulatisve tri-vel subquinque-nerviis supra pilosulis sparsimque stimulosi aut glabratis subtus mollior pubescentibus, inflorescentiis quam folia brevioribus, pedicellis fructiferis botuliformibus, stigmatate subulato. ♂ In sylvis umbrosis Novæ Hollandiæ orientalis, a fluvio Clarence River usque ad limites districtus Illawarra præcipue obvia (Cunningham, Leichard, Verreaux n. 559, Macarthur, F. Mueller). — *Urtica* Gigas All. Cunningham, mscr. Urtica excelsa et U. rotundifolia Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 178. — Arbor 15-30-metralis vel etiam multo excelsior trunco erecto basi processibus aliformibus singulariter corroborato dein sensim attenuato atque interdum ad altitudinem 40 metr. attingente antequam in comam expandatur, cortice tenui parce rimoso pallidoque et in apice ramulorum pubescente necnon cicatricibus latis post lapsum foliorum notato. Folia 13-35 centim. longa, 10-30 centim. lata, ima basi sinu profundo angustoque cordata et haud raro subpeltata, acumine obtuso vel acutiusculo, in dimidio inferiore sæpe integra, superne autem sæpius repanda vel (etiam in toto margine) sinuato-dentata, rarissime omnino integra, nervis primariis utrinsecus 6-10 subtus parum prominentibus, jugo inferiore (folii trinervi) dimidium limbum paululo superante sub margine cum sequentibus arcuatum connexo, nervis secundariis transverse rectiusculeque vel reticulatim anastomosantibus, lamina exsiccata membranacea vel coriacea subtus plus minus canescente, petiolo longitudine varia interdum limbo fere æquilongo tomentoso vel pubescente exsiccato gracili. Folia plantæ junioris quam adultæ etiam ampliora et stimulis validis armata. Stipulæ mox deciduæ, ovato-lanceolatae, 2-4 centim. longæ, subacuminatæ apiceque brevi-

ter bifidæ, extus pubescentes. Flores dioici. Inflorescentiæ petiolum vix æquantes v. superantes, pedunculatæ, patule ramosæ, ramis ramulisque magis minusve tomentosis et nonnunquam densiuscule stimulosi, glomerulis parvis discretis approximativse. Flores masc. subsessiles vel breviter pedicellati, perigonii 4-partiti segmentis puberulis. Flores fem. pedicello brevi tereti crassiusculo insidentes, initio erectiusculi sed mox incurvi, perigonio inæqualiter 4-partito haud millimetrâli quam pedicellus fere dimidio longiore ovariumque ovatum fere æquante, stigmatibus quam flos ipse duplo triplove longiore. Glomeruli fructiferi sæpe confluentes e pedicellis accretis carnosius et plus minus botuliformi-cylindraceis aut mutua compressione varie difformibus sparce stimulosi maxima ex parte constantes; perigonii fructiferi in pedicellum omnino incurvi necnon in fossula laterali apicis ejusdem partim recepti segmentis lateralibus majoribus ovatis fimbriato-denticulatis, superiore cucullato quam inferius lanceolatum dimidio longiore. Achænium oblique rotundatum,  $1\frac{1}{2}$ -2 millim. longum et latum compressum granulato-scabrum perigonii duplâ longitudinem æquans (1). (v. s.)

12. L. PHOTINIPHYLLA (Wedd. Monogr. p. 138), arbor adulta parce stimulosa aut subinermis, foliis parum amplis oblongo-ellipticis ovatisve acuminatis basi acutatis rotundatisve vel subcordatis inæqualiter sinuato-crenulatis aut integris triplinerviis utrinque glaberrimis, inflorescentiis paniculiformibus folio brevioribus, pedicellis fructiferis botuliformi-incrassatis, stigmatibus subulato. § In sylvis littoralibus Novæ Hollandiæ orientalis tropicæ et subtropicæ sat frequens (All. Cunningham, Leichard, F. Mueller); in ins. Lifa Novæ Caledoniæ (Thiebault); ins. Viti s. Fidji (Seemann). — L. vitiensis Seem. Acc. gov. miss. Viti Isl. p. 427; Fl. Vit. p. 239. t. 60. — Urtica photiniphylla All. Cunningham. mscr. Fleurya photiniphylla Kunth, Ind. sem. h. b. berol. (1846) p. 11. — Arbor, trunco 12-16 metr. alto (teste F. Mueller), ramulis gracilibus vix succulentis inermibus vel aculeis parvis conicis obsitis foliosis exsiccatis epidermide fusca vestitis. Folia 7-20 centim. longa, 3-9 centim. lata, a medio versus apicem deorsumque sensim attenuata sensimque aut abrupte acuminata, acumine obtusiusculo, ad ipsam petioli insertionem sæpe acute angustissimeque cordata, vulgo infra medium vel prope basim tantum integra, cæterum magis minusve profunde crenulato-sinuata aut obtusissime crenata; nervis primariis subsecus plus minus patentibus prope marginem arcuatim reticulatimque anastomosantibus, subtus vix prominentibus, jugo inferiore suprabasilarî limbum medium superante, secundariis tenuissimis reticulatim connexis, lamina exsiccata membranacea, petiolo gracili, 3-5 centim. longo, glaberrimo, hic et illic (ut ipse limbus) stimulis tenuibus armato vel omnino inermi. Stipulæ triangulari-ovatæ, parvæ, glabræ. Flores dioici. Inflorescentiæ paucæ axillis superioribus ramorum ortæ, pedunculatæ, petiolum æquantes superantesve, laxè subdivaricato-ramosæ, ramis ramulisque fere inermibus, glomerulis parvis discretis. Flores masculi perigonio 4-partito, segmentis glabris. Flores feminei erectiusculi aut in pedicellum flore brevioribus parum incurvi; pedicello post anthesin valde accreto albidocarnoso botuliformi, figura autem, sicut apud speciem præcedentem, secundum in glomerulo situm atque pedicellorum vicinorum compressionem sæpe plus minus mutata, flore ipso mox in apicem pedicelli incurvum et denique omnino laterali. Perigonium sub anthesi minimum et vix inæqualiter 4-lobum vel partitum, segmentis lateralibus inde cum ovario accretis ovato-rotundatis minutissime denticulato-ciliatis superioreque fere cucullato et inferiore ovato dimidio longioribus. Ovarium quam stigma duplo triplove brevius. Achænium oblique rotundato-ovatum,  $2\frac{1}{2}$  millim. longum, basi cum perigonio semimillimetrâli in scrobiculo laterali pedicelli receptum, stigmatibus persistente et omnino incurvo nonnihil longius, siccitate fuscum et sub lente granulatum. — Cum primum ex speciminibus omnibus floriferis femineisque hancce stirpem describere conatus sum, affinitates mihi haud innotuere; sed interea fructus et flores ♂ qui tunc non suppeditabant mihi benigne misit Ferd.

(1) Videas quoque quod de hac egregia arbore scripsi in Monographia mea necnon figuram trunci ex notis cl. Macarthur delineatam.

Mueller melburniensis simul cum floribus utriusque sexus *L. Gigantis*. Planta junior (teste eodem) admodum urens est. (v. v. in hort. Kew. et s.)

13. *L. HARVEYI* (Seem. l. c.; Fl. Vit. p. 238. t. 59), arborea, adulta subinermis et fere omnino glabra, foliis amplis late cordato-ovatis rotundatisve acuminatis inæqualiter crenato-dentatis sinuatisve tripli-v. subquintuplinerviis utrinque glaberrimis, inflorescentiis paniculiformibus folio brevioribus, pedicellis fructiferis botuliformi-incrassatis, stigmate subulato. 5 In ins. Viti (Seemann n. 426, Harvey). — Arbor 12-metralis, habitu præcedentis cui affinis. Folia 15-20 centim. longa et lata, apice sensim acuminata, basi sinu lato sed vix profundo cordata, nervis primariis subsenis subtus parum prominentibus, secundariis distinctis inter præcedentes arcuatim connexis, lamina exsiccata membranacea viridi, petiolo 4-6 centim. longo. Stipulæ triangulari-ovatae, sesquicentimetrales, acuminatae, dorso puberulae, cito deciduae. Flores dioici. Inflorescentiæ masculæ ut in specie præcedente sed ramulis inermibus subtilissime puberulis. Flores feminei non visi. Perigonium fructiferum minimum, in apice aut fere in apice excavato? pedicelli incrassati sessile, segmentis 4 subæqualibus denticulatis. Achænium oblique ovatum, vix bimillimetrale, stigmate persistente et valde incurvo circiter dimidio longius, sub lente minute granulosum. (v. s. in hb. Kew. et Mus. par.)

Sect. III. DISCOCARPUS *Wedd. Monogr.* — Flores fem. perigonio 4-partito, segmentis valde inæqualibus. Pedicelli fructiferi nec alati nec carnos.

14. *L. MEXICANA*, frutescens, foliis ovatis apice attenuatis basi obtusatis repandis vel inæqualiter sinuato-denticulatis subtus præsertim stimulosi-hispidis, floribus dioicis, inflorescentiis folia æquantibus superantibusve, pedicellis fructiferis haud incrassatis cylindricis, achænio disciformi. 5 In imperio mexicano circa Cordova et Orizaba, altit. 300-1500 metr. frequens (Bourgeau); prope Oaxaca et in monte San Felipe, ad altit. circiter 2000 metr. (Galeotti n. 7128; Liebmann). — *L. platycarpa* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 181; *Monogr.* p. 132. t. 2. C. f. 7. *Urera platycarpa* Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 60. *Discocarpus mexicanus* Liebmann, in kön. dansk. Vid. Selsk. Skr. (1851) p. 294. — Frutex 1-3-metralis, truncus sat crasso parce ramoso, ramulis succulentis desiccatione plicatis cicatrices foliorum lapsorum monstrantibus et epidermide pallida glabraque vestitis. Folia in ipso apice ramulorum valde approximata, 5-10 centim. longa, 2½-6 centim. lata, juniora in utraque pagina sed in inferiore præsertim stimulosi-hispida (pilis costæ interdum retrorsis), supra demum glabrata; petiolo gracili brevi densiuscule stimulosi. Stipulæ oblongæ, 6-8 millim. longæ, integræ, puberulae. Flores dioici. Cymæ ex axillis foliosis ortæ, longiuscule pedunculatæ, folia æquantibus iisve longiores, patule ramosæ, ramis pedunculoque sæpissime retrorsum stimulosi, floribus masc. perigonio 5-partito, laciniis parce stimulosi-hispidis. Cymæ fem. floribundæ, 10-15 centim. longæ, ovatae, floribus breviter pedicellatis laxiuscule glomeratis primo erectiusculis et denique in pedicellum magis minusve inflexis, perigonii (violacei) inæqualiter 4-partiti segmentis laterilibus late ovatis acuminatis glabris, aliorum brevissimorum altero (demum superiore) late rotundato ciliato altero ovato. Achænium orbiculare, disciforme, 4 millim. latum, quam maxime compressum, membranaceum, margine incrassato, læve, pellucide virescens, perigonio vix aucto multoties longius. (v. s.)
15. *L. NICARAGUENSIS*, arborescens, foliis ovatis acutiusculis basi obtusis crenatis supra adpresse pilosulis subtus dense cano-tomentosis, floribus dioicis, inflorescentiis valde ramosis folia superantibus, pedicellis fructiferis haud incrassatis cylindricis, achænio disciformi. 5 In Nicaragua prov. Segovia, prope Matajalpa (Orsted). — *L. Liebmannii* Wedd. *Monogr.* p. 133. *Discocarpus Nicaraguensis* Liebmann l. c. — Arbor mediocris, ramulis patulis crassis succulentis cicatricosis epidermide tenui glabra vestitis. Folia 4-5 centim. longa, 2½-4 centim. lata, crassa, petiolo 1½-2½ centim. longo incano-hispido et retrorsum stimulosi. Flores dioici. Planta masc. aphylla florens, cymis 5-8 centim. longis irregulariter repetite ramosis, ramulis hispidis retrorsumque stimulosi, floribus glomerulatis, perigonii 5-partiti laciniis hispida.

Planta fem. foliifera florens, cymis 8-13 centim. longis suberectis supra decomposite ramosis, ramulis hispidis, achænio ut in specie præcedente. (Descript. ex Liebmann l. c.)

Sect. IV. DENDROCNIDE *Wedd. Monogr.* — Flores fem. perigonio 4-loba, lobis subæqualibus. Pedicelli fructiferi haud incrassati cylindrici aut nulli. Ramuli extremi inflorescentiæ interdum fastigiato-dilatati.

16. *L. CRENULATA* (Gaudich. Bot. Voy. Uran. p. 498), arborescens, ramis inermibus, foliis ellipticis ovatisve acuminatis basi cordatis emarginatisve in dimidio apicali inæqualiter sinuato-crenulatis rariusve integris penninerviis utrinque glabris costa subtus breviter stimulosi, floribus dioicis, inflorescentiis pedunculatis petiolum superantibus, femineis laxifloris ramulis haud dilatatis, pedicellis fructiferis cylindricis, perigonio subæqualiter 4-loba, stigmate subulato. ♂ In provinciis humidioribus Indiæ orientalis: Bengal, Assam, Silhet, Chittagong et Concan, nec non in regione tropica Himalayæ orientalis ad Sikkim, alt. 0-1500 metr.; in ins. Zeylaniam et Java. — *Wedd. Monogr.* p. 133. t. 2. C. f. 5; *Miq. Fl. Ind. bat. l. c.* p. 230. *L. gigantea* Gaudich. l. c. *L. latifolia* ejusd. Bonite t. 81. *U. gigantea* Poir.? *Encycl. suppl.* 4. p. 224. *U. sinuata* Bl.? *Bijdr.* p. 605. *U. crenulata* Roxb. *Fl. ind.* 3. p. 591; *Wight Ic. Fl. Ind.* 2. t. 686. *U. Churta* Hamilt. in *Wall. Cat. n.* 4608. *Urera javaensis* et *U. gigantea* Gaudich. *Uran.* p. 496. *U. crenulata* et *U. Commersoniana* *Wedd.* in *Ann. sc. nat.* 3<sup>e</sup> sér. 13. p. 104 et 105. *Dendrocniide crenulata* *Miq. Pl. Jungh.* 1. p. 31. — *Arbuscula* 3-6-metralis, trunco longe nudo, ramis patentibus, ramulis crassis lævibus glabris. Folia 10-35 centim. longa, 5-14 centim. lata, ut plurimum a medio utrinque attenuata basi que ante ipsam petioli insertionem emarginata vel inferne latiuscule rotundata cordataque apice acuminata (acumine acutiusculo), nervis primariis utrinsecus 12-15 ante marginem arcuatum anastomosantibus subtus prominulis, jugo inferiore dimidium limbum haud attingente, secundariis tenuissimis rectiuscule v. reticulatim connexis, limbo juniore utrinque sparsissime stimulosi-pilosulo mox glabrato, stimulis vix nisi in costa inferioris paginæ persistentibus, petiolo valido 2-6 centim. longo breviter stimulosi minutissimeque puberulo rarius glabro et inermi. Stipulæ ovato-lanceolatæ, 1-2-centimetrales, nonnunquam acuminatæ, puberulæ v. glabræ. Inflorescentiæ in axillis foliatis erectiusculæ v. patentes, petiolo longiores, divaricato-ramosæ, pedunculo 1-2-centimetrali v. longiore ramulisque puberulis glabrativis et stimulis tenuissimis crebre armatis, glomerulis paucifloris discretis, floribus sessilibus v. breviter pedicellatis. Perigonium masc. puberulum et magis minusve stimulosum, 4-partitum; fem. rectiusculum, fere campanulatum, subæqualiter 4-lobum s.-dentatum, dentibus acutis, crebriuscule stimulosum. Achænium late ovatum, vix obliquum, 5-6 millim. longum, ventricosum (pericarpio aliquantum succulento), acuminatum, læve, perigonio minimo crateriformi et membranaceo insidens, stigmate incurvo persistente coronatum. — Quod ad vires nocuus stimulorum attinet vid. monographiam meam, l. c. (v. s.)

β, *Luzonensis*, foliis anguste ellipticis oblongisve basi acutis aut obtusis integris glaberrimis inermibusque. — In insulis Philippinis, prope Manillam, etc. (Callery; Cuming n. 522). (v. s.)

17. *L. MICROSTIGMA* (Gaudichaud mscr. in hb. Mus. par.), arborea?, foliis oblongis acuminatis basi rotundatis integerrimis penninerviis glabriusculis, floribus dioicis, inflorescentiis fem. longe pedunculatis, floribus sessilibus in ramulis distichis apice incrassatis dense glomeratis, perigonio 4-loba, stigmate minimo ovato-oblongo. ♂ In ins. Java (Leschenault). — *Wedd. Monogr.* p. 138. t. 2. C. f. 6; *Miq. Fl. Ind. bat. l. c.* p. 232. ? *Urtica ardens* Blume *Bijdr.* p. 504. *Urera microstigma* *Wedd.* in *Ann. sc. nat.* 4<sup>e</sup> sér. 1. p. 177. — Ramuli crassi, sicci obscure cinerei, apice et præsertim ad nodos pubescentes subtiliterque stimulosi. Folia 10-25 centim. longa, 5-10 centim. lata, a medio deorsum sensim attenuata, ima basi anguste rotundata et interdum quasi subcordato-peltatæ, acumine apicali brevi angusto obtusiusculoque, nervis primariis utrinsecus 10-12 subtus parum prominulis, cæteris parum perspicuis,

limbo utrinque glabro costa subtus prominula tenuiterque piloso-stimulosa, exsiccato rigido, petiolo robusto 1½-5 centim. longo pubescenti-piloso cinereoque. Stipulæ ovatæ, 2-2½ centim. longæ, acuminatæ, integræ, dorso pubescentes, ad marginem glabratae. Inflorescentiæ utriusque sexus in axillis foliatis longe pedunculatæ, pedunculis petiolo longioribus puberulis pubescentibusve, compressæ, ambitu triangulari-ovatæ, decomposite ramosæ, folium tri-quadri-pinnatum quodammodo referentes, ramis ramulisque horizontaliter patentibus puberulis subinermibus, glomerulis in ramulis extremis confluentibus, floribus sessilibus, masc. perigonio 4-partito, laciniiis extus puberulis, femineis minimis valde obliquis, perigonii 4-lobi lobis æqualibus inæqualibusve; perigonio fructifero glabro quam achænium rotundatum 1-1½ millim. latum compressum læviusculum subdimidio brevius. — Stigmatis exiguitate stirps admodum in genere insignis, Laporteam Ureramque aliquatenus connectens. (v. s. in hb. Mus. par.)

18. *L. GAUDICHAUDIANA* (Wedd. Monogr. p. 137), frutescens vel arborea, foliis magnis late ovato-oblongis sæpius abrupte acuminatis cordatis integerrimis longe petiolatis penninerviis supra subtiliter pilosulis subtus breviter pubescenti-tomentosis, floribus dioicis, inflorescentiis femineis longiuscule pedunculatis petiolo multo longioribus bracteis majusculas ad basim ramulorum gerentibus, floribus in apice subfastigiato horumce dense glomeratis, perigonio profunde 4-loba, stigmatate elongato-filiformi. ♀ In insulis Philippinis, ubi *Lipa* vocatur (Gaudichaud, Cuming). — Miq. l. c. p. 232. Utera Gaudichaudiana Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. l. p. 177, non Henslow. — Ramuli succulentii, exsiccati etiam crassitie digiti minoris, epidermide pallida glabraque vestiti cicatricibusque latissimis foliorum delapsorum notati. Folia 18-36 centim. longa, 10-20 centim. lata, basi late rotundata sinuque vix profundo cordata, a medio apicem versus magis minusve attenuata, acumine brevi acuto, nervis primariis utrinsecus 12-14 subtus prominulis, jugo inferiore subbasilari, secundariis tenuibus transversim linea recta anastomosantibus, pagina inferiore tota molliter subincano-tomentosa, lamina exsiccata rigida, petiolo succulento siccitate gracili elongato puberulo. Stipulæ..... Flores dioici. Cymæ masc. haud visæ. Cymæ fem. longiuscule pedunculatæ, laxæ et distiche ramosæ, in specimine suppetente fere bidecimetrales, pedunculo ramisque densiuscule stimulosi, bracteis ad basim ramulorum majusculis (6-10 millimetralibus) late ovatis, extus puberulis, floribus numerosis sessilibus in glomerulos discretos ramulos apice spathulato-vel subfastigiato-dilatatos terminantes congestis; perigonio profunde 4-loba lobis parce stimulosi, ovarium æquante styloque filiformi semicentimetrali multoties brevius. Achænium haud visum. — Sequenti similis quoad ordinationem florum fem., sed ab illa et ab omnibus congeneribus magnitudine bractearum inflorescentiæ facillime distinguenda. (v. s. in hb. Mus. par.)

19. *L. STIMULANS* (Miq. in Zolling. Cat. p. 103), arborescens, foliis obovato-lanceolatis abrupte breviterque acuminatis deorsum angustatis et ima basi rotundatis integris (sicine semper?) supra glabris nervis subtus stimulosopilosis, inflorescentiis longe pedunculatis petiolumque longe superantibus, bracteis ad basim ramulorum vix conspicuis, floribus dioicis, femineis in apice fastigiato-dilatato ramulorum dense glomeratis sessilibusque perigonio 3-4-loba, stigmatate subulato. ♀ In insulæ Javæ sylvis umbrosis, prope Tjikoya (Zollinger, n. 510). — Blume, mus. lugd. bot. 2; Wedd. Monogr. p. 135; Miq. Fl. Ind. bat. l. c. L. sinuata Bl. ? l. c. *Urtica stimulans* Linn. fil. Suppl. 418 et auct. U. Meyeniana Walp. ? in Nov. act. ac. Leup. Car. 16. suppl. 2. pl. 422. — Ramuli crassi, puberuli, exsiccati epidermide rubente vestiti, apice foliiferi, infra apicem cicatricibus triangulari-rotundatis foliorum lapsorum signatis. Folia 12-20 centim. longa, 3-6 centim. lata, a triente superiore deorsum sensim attenuata, ima basi angustissime rotundata et ante ipsam insertionem petioli emarginata, acumine angusto acuto, nervis primariis utrinsecus 10-12 ante marginem arcuatim anastomosantibus evanescentibusve subtus parum prominentibus (jugo inferiore suprabasiliari brevissimoque), secundariis tenuissimis reticulatim connexis, pilis costæ tenuissimis patulisque, lamina exsiccata rigida glaucescente pagina inferiore cystolithis reticulata, petiolo 2-4-centimetrali puberulo sicco albicante. Stipulæ triangula-

res, 6-10 millim. longæ, acuminatæ, nervis pilosis. Inflorescentiæ erectæ, masc. longe fem. longissime pedunculatæ, ramis paucis gracilibus pedunculoque patule vel reflexe stimulosi-pubescentibus, bracteolis vix millimetralibus, floribus masc. sessilibus pedicellatisve, in glomerulos discretos laterales vel terminales digestis, perigonii 4-partiti laciniis hispidulis. Flores fem. sessiles, perigonio profunde 3-4-lobi, lobis triangularibus stimulosi, fructifero quam achænium elliptico-rotundatum 2-2½ millim. latum valde compressum marginatum granuloso multoties brevior. — Habitu *L. Microstigma* refert, sed characteribus inflorescentiæ et cæt. primo intuitu longe distat. (v. s.)

20. *L. COSTATA* (Miq. Fl. Ind. bat. 1. pars 2. p. 231), arborea, foliis e basi obtusa subobovato-oblonga breviter acuminata, a medio versus apicem crenato-repandis valde costulatis, in costa utrinque appressissime setulosis, inflorescentiis paniculiformibus pedunculatis alterne ramosis bracteatis hirtellis ramulis ultimis apice dilatato-floridis, stigmatibus subfiliforme quam ovarium longiore. ♂ In insula Java (Junghuhn). Dendrocide costata Miq. Pl. Jungh. 1. p. 30. Rami crassi. Folia conferta, 20-25 centim. longa, chartacea. Stipulæ ovatæ, coriæ. Flores dioici, feminei sessiles; perigonii lobis 4 breviter ovatis puberis, lateralibus majoribus. — Hanc plantam ad præcedentem retuleram in Monographia mea et revera paucis notis ab illa discrepare videtur. Descriptionem juxta illam cl. Miquelii scripsi.

β, *contracta* (Miq. l. c.), foliis brevioribus et latioribus valide costatis. In ins. Celebes. *Laportea contracta* Blume, herb.

21. *L. PTEROSTIGMA*, arborea? foliis elliptico-oblongis sensim acuminatis basi rotundata v. subcordata excepta minute denticulatis penninerviis supra pubescentibus subtus tomentosis, inflorescentiis (femineis) pedunculatis petiolumque longe superantibus, bracteis ad basim ramulorum parvis, floribus in apice fastigiato horumce glomerulatis sessilibusque, perigonio 4-partito, stigmatibus lineari compresso-subalato. ♀? In insulæ sinensis Formosæ regione austro-occidentali (Swinhoe). — Folia nonnulla avulsa et inflorescentiæ quædam femineæ solum suppetunt. Folia (haud adulta?) 6-10 centim. longa, 3½-4½ centim. lata, acumine acuto, ad insertionem petioli interdum vix profunde emarginata, nervis primariis circiter 12 juxta marginem arcuatim anastomosatis evanidisve subtus parum prominentibus, jugo inferiore sequentibus breviori, venulis inter præcedentes parallele arcuatis, tomento s. pubescentia nervorum breviori quam in intervenio, lamina exsiccata subcoriacea subtus albicante, petiolo 2-2½ centim. longo sat robusto canescenti-pubescente. Inflorescentiæ femineæ 10-15 centim. longæ, paniculiformes, laxè patuleque ramosæ, ramis gracilibus puberulis stimulisque parvis sparsim armatis, bracteis ovatis lanceolatisve. Glomeruli discreti, e floribus 6-10 constantes, perigonii segmentis ovatis, altero superiore minore. Ovarium ¼ millim. circiter longum, ovatum, apice cum stigmatibus sesquimillimetrali tæniæformi et plus minus incurvo continuum. (v. s.)

*Species incertæ consanguinitatis v. omnino dubiæ.*

22. *L. AMPLISSIMA* (Miq. Fl. Ind. bat. 1. pars 2. p. 232), foliis maximis usque bipedalibus elliptico-oblongis acutiusculis vel obtusis coriaceis multicostulatis novellis in costis subpuberis mox glaberrimis lævibus, cymis femineis dichotome ramosis, ramulis ultimis dilatatis floriferis. ♀? In insula Java (Teysm.). Inflorescentiæ in nulla specie generis *Laportæ* mihi cognita dichotomæ sunt; forsitan eadem ac *Uvera amplissima* Blume, Mus. lugd. bat. t. 22 (fig. anal.) absque descript. quæ etiam *Laportæ* referenda est.

23. *L. OLERACEA* (Wedd. Monogr. p. 141), herbacea, foliis amplis late ovatis acuminatis basi rotundatis obtusatisve in toto margine acute grossiusculeque serratis utrinque sed præsertim supra sparsim breviterque stimulosi lævibus in planta viva aliquantulum succulentis, inflorescentiis masculis subsessilibus densiuscule multifloris quam petioli multoties brevioribus, femineis ignotis. ♀? In sylvis Himalayæ orientalis temperatæ, prov. Sikkim, alt. 2730-3040 metr. (Hook. fil.). — Caulis circiter bimetralis, debilis, angulato-sulcatus,

stimulis gracilibus longeque bulbosis sparsim armatus et inter illos superne hispidulus. Folia in specim. visis 12-18 centim. longa, 8-10 centim. lata, acumine s. dente apicali lineari uni-sesqui-centimetrali, dentibus cæteris triangularibus apice piliferis subæqualibus inæqualibusve juxtaque petiolum minoribus, nervis basilaribus circiter usque ad trientem limbi superiorem productis, lamina exsiccata tenuiter membranacea oculo armato subtus minutissime creberrimeque badio-punctata, petiolo fere decimetrali subinermi. Stipulæ lanceolatæ, apice bifidæ, persistentes. Cymæ masculæ juniores ovatæ vel rotundatæ, ramosissimæ, pedunculo brevissimo breviter bracteato ramulisque hispidulis; floribus parvis, pedicellatis, perigonio 5-lobato hispidulo. Cætera desiderantur. (v. s.)

24. *L. MOROIDES* (Wedd. Monogr. p. 142), arborea? ramulis apice tenuissime stimulosus, foliis latiuscule cordato-ovatis acuminatis peltatis in toto margine dentatis trinerviis tenuiter piloso-pubescentibus stimulosisque. ♀ In Australia, juxta Endeavour river (Cunningham). — *Urtica moroides* Cunningham. mser. — Ramulus in specimine unico viso vix crassus, foliatus, internodiis brevibus, stimulis tenuissimis fere bimillimetralibus crebriusculè obsitus. Folia 7-8 centim. longa et lata, circiter a medio ad apicem attenuata, breviuscule acuminata, basi truncata v. cordata manifesteque peltata, dentibus parvis inæqualibus acutis, sinubus obtusis, acumine longitudine varia raro centimetrali integro v. dentato acuto, nervis primariis utrinque 6-7 tenuibus ante marginem arcuatim anastomosantibus, jugo inferiore supra limbi basim ex insertione petioli orto dimidiumque limbum superante, lamina exsiccata tenuiter membranacea subtus glaucescente et molliuscule pubescente, pagina superiore in nervis præsertim stimulosa, petiolo gracili limbum subæquante tenuissime retrorsum stimuloso. Stipulæ ovatæ semicentimetales, pubescentes. Cætera ignota. — Ad hanc stirpem forte adscribenda erit planta inflorescentiis pulchre violaceis cujus exstat in Musæo britannico speciosa picta imago mihi a beato R. Brownio olim benigne monstrata.

25. *L. ORLONGATA* (Miq. in Zolling. Cat. p. 103), foliis alternis breviter et longiter petiolatis ellipticis vel sæpe elliptico-vel obovato-oblongis acuminatis supra basim subobtusam dense crenato-dentatis subcoriaceis, costulis utrinque circiter 8, supra glabris subtus in nervis et petiolo scabriusculè pubescentibus, 4½-8½ poll. longis, stipulis linearibus hirsutis, inflorescentiis ex axillis nudis vel foliatis paniculatis pubescentibus, floribus femineis in ramulis dense et interrupte subunilateraliter spicatis, achæniis punctatis. — In Javæ sylvis humidis prope Tjikoia (Zollinger, n. 581). — Miq. Fl. Ind. bat. l. c. p. 231; Wedd. Monogr. p. 140. — Prope *L. crenulatam* (Miq. l. c.). — Specimen a me visum in herbario Zollingeriano (num. cit.) non est stirpis hujus generis sed *Villebrunneæ* (V. rubescens.)

VII. *URERA* *Gaudich. Bot. Voy. Uran. p. 496; Miquel, in Mart. Fl. bras. fasc. XII, p. 194; Fl. Ind. bat. l. pars. 2. p. 232; Wedd. monogr. p. 143. t. 2. A; Griseb. Fl. Br. W. Ind. p. 154. — Urticæ spec. auct.*

Flores dioici rariusve monoici, modo discreti, modo glomerulati (glomerulis interdum capituliformibus), floribus glomerulisque cymam biparam (seu dichotomam) vel uniparam (sive non vere dichotomam) efformantes; inflorescentiis in singulis axillis solitariis unisexualibusque, pedicellis florum ♂ articulatis. *Masc.* Perigonium 4-v. 5-partitum, segmentis ovatis apice tenuiter membranaceis, alabastro fere umbilicato. Stamina 4 v. 5, antheris reniformibus. Pistilli rudimentum globosum v. depressum v. subcupuliforme. *Fem.* Perigonium 4-lobum, lobis sæpius inæqualibus, interioribus s. lateralibus ut plurimum majoribus, exteriorum altero nonnunquam minimo ad rarissime prorsus deficiente, vulgo glaberrimis. Ovarium rectum v. obliquum, ovoideum. Ovulum imo loculo affixum, erectum aut vix adscendens, funiculo brevi suffulsum. Stigma subsessile, penicillato-capitatum rotundatumque v. rarissime lanceolatum denseque villosopapillosum, persistens. Achænium



fere exsuccum, recte v. oblique ovatum rotundatumve, compressum aut utrinque convexum, lævisculum v. magis minusve tuberculatum; perigonio accreto baccato et colorato vestitum. Semen pericarpio subconforme. Albumen parciissimum. Embryo cotyledonibus rotundatis, apice et præsertim basi emarginatis, radícula plerumque multo longioribus. Fructes aut arbusculæ vel rarissime suffrutices erecti aut interdum scandentes, inter tropicos Novi Orbis necnon Africae et Oceaniæ provenientes, rarissime extratropici, stimulis aculeisque copiose aut parce armati, iisque ut videtur nonnunquam prorsus carentes. Folia alterna, integerrima vel crenulata aut varie dentata v. incisa, palmatinervia vel penninervia, cystolithis numerosissimis punctiformibus vel rarius fusiformibus oblongisque. Stipulæ axillares, bicarinatæ apiceque bifidæ aut integræ, deciduæ. Flores sæpissime ebracteati, colorati. — Genus in tribu insigni perigonio fructifero baccato.

### § 1. — *Cymæ biparæ sive vere dichotomæ.*

#### A. — *Species neogææ.*

1. U. CARACASANA (Griseb. Fl. Brit. W. Ind. p. 154), arborea v. fruticosa, foliis late ovatis acuminatis basi sinu lato et apertio cordatis crenato-dentatis supra scabro-hispidis subtus plus minus pubescentibus, cymis regulariter dichotomis, floribus femineis in apice ramulorum discretis ternisve, perigonio fructifero rubente vel albidio? 3-5 in silvis Americæ tropicæ et subtropicæ: in imperio mexicano (Botteri, Bourgeau, etc.), insulis Indiæ occidentalis, Nova Granata, Peruvia? Guyana et Brasilia. — U. alceæfolia Gaudich. Uran. p. 496. U. Jacquini Wedd. in Ann. sc. nat. 3<sup>e</sup> ser. 18. p. 200; Monogr. p. 144, exclus. var. Urtica caracasana Jacq. Hort. schönbr. 3. p. 71. t. 386. Urt. alceæfolia Poir. Encycl. suppl. 4. p. 224. Ramuli puberuli pubescentesve, stimulis tenuibus insuper armati, rarius inermes. Folia magnitudine varia, majora 20-25 centim. lata, 25-30 centim. longa, in eodem ramo nonnunquam diversiformia, alia scilicet cordata aliaque basi rotundata, dentibus sive crenis nunc obtusissimis nunc subacutis sæpissime mucronulatis æqualibus inæqualibusve, nervis primariis secundum limbi magnitudinem numero vario (6-12) ante marginem reticulatim vel subarcuatim anastomosantibus subtus parum prominulis (jugo inferiore basilari ultra medium limbum protracto), secundariis tenuissimis persæpe linea recta connexis parallelisque, pagina superiore plus minus scabra et haud raro (in folio juniore præsertim) verruculosa, verruculis apice pilis brevibus rigidulisque tandem caducis instructis, nervis utrinque puberulis aut densiuscule piloso-pubescentibus necnon primariis subtus patentim aut retrorsum stimulosi, stimulis sæpius parum numerosis longiusculis brevissimisve, intervenio paginæ inferioris varie pubescente rarissime omnino glabrata, lamina exsiccata submembranacea, petiolo gracili quam limbus interdum longiore at sæpius brevior glabrato aut pubescente stimulosi vel inermi. Stipulæ ovatæ, apice persæpe breviter bifidæ, hispidulæ vel pubescentes. Flores dioici vel nonnunquam monoici (anne fortuito?), floribus masculis tunc ex axillis superioribus femineisque ex axillis inferioribus (foliis demum lapsis) ramuli ortis. Cymæ masc. erecto-patulæ, petiolis longiores, pedunculatæ, quater-sexies dichotomæ, ramulis divaricatis pedunculoque 2-3-centimetrali vel longiore puberulis pubescentibusve stimulosi aut etiam inermibus, floribus subsessilibus aut pedicellatis, in capitula pisi circiter magnitudinis et coloris rosei digestis, perigonii 4-partiti lacinii hispidis. Cymæ fem. breviter pedunculatæ v. subsessiles, multoties dichotomæ, patulæ, ramulis (plerisque ebracteatis) quam muscularum brevioribus sed magis stimulosi demumque glabratis et inermibus, pilis interdum subunilateralibus, floribus pedicellatis solitariis vel subternis, stigmate rotundato; perigonio rectiusculo 4-lobo, lobis lateralibus demum majoribus, fructifero oblique valde succulento stramineo siccitate cystolithis majusculis albo-punctato achænium millimetrale obliquum late ovatum ventricosum sub lente punctulatum scabriusculumve ex integro vestiente.

- β, *subpeltata*, foliis sinu angusto cordatis, lobis basilariibus haud raro imbricatis. — In Brasilia amazonica (Spruce, n. 1573), etc. — *Urera subpeltata* Miq. in Mart. FL. bras. fasc. 12. p. 189. t. 66. *Urtica tiliaefolia* Kunth? in H. et B. Nov. gen. et sp. 2. p. 141. *Cymæ* ♀ plantæ *Kunthianæ* non visæ, synonymon ergo dubium. (v. s. in hb. Mus. par. Boiss. et Mart.)
- γ, *mitis*, inermis, foliis (quam in præcedenti minoribus) ovatis oblongisve vel ellipticis basi rotundatis aut vix cordatis dentatis v. crenato-dentatis. — In ins. Martinica (Plée), in Nova Granata (Pavon) et in Brasilia. — *Urera mitis* Miq.? l. c. U. Jacquinii, var. *ulmifolia* Wedd. Monogr. p. 145. *Urtica ulmifolia* H. B. Kunth? l. c. *Cymas* femineas plantarum a Kunthio et Miquelio descriptas non vidi; synonyma itaque dubia. (v. s. in hb. Mus. par., Boiss. et Mart.)
- δ, *Miquelii*, foliis quam in typo magis acuminatis basi rotundatis vel ad petiolum subtruncatis paucinerviis, cæteris charact. similibus. — In Brasilia fluminensi et Peruvia (Cl. Gay). *Urera acuminata* Miq. l. c. p. 190, non Gaudich.
- ε, *tomentosa*, foliorum pagina inferiore petiolisque apice ramulorum et ramis inflorescentiarum dense hirtotomentosis, cymis masc. quam in typo minus diffusis. — In monte ignivomo Irasú ditionis Costaricae altit. 2500 metr. (Orsted); in Nova Granata, prope Ibaque (Goudot), In Brasilia. *Urera densiflora* Miq. l. c. p. 190. *Urtica verrucosa* Liebm. l. c. p. 265. *Cymas* femineas non vidi. (v. s.)
2. U. CORALLINA, arborea, foliis late ovatis basi leviter cordatis acutis late denticulatis supra sparse setosis punctatis elevatis asperis ad costam nervosque adpresse pilosulis, cymis in axillis foliorum delapsorum subsessilibus intricato-ramosissimis ramis divaricato-dichotomis puberulis. ♀ In ditione Costa Rica, prope Aguacate (Orsted). — *Urtica* (*Urera*) *corallina* Liebm. l. c. p. 295. *Urera* Jacquinii, ? *corallina* Wedd. Monogr. p. 145. — Rami teretes, flexuosi, rubentes, sursum versus pubescentes. Folia 12-16 centim. longa, 9-12 centim. lata, longe petiolata, trinervia, supra saturate viridia, subtus glaucescentia, petiolo pilosulo. Flores dioici. *Cymæ* 3-4 centim. longæ, 8 centim. latæ; stigmatate rotundato; perigonio fem. post anthesim miniato achænium late ovatum parum compressum apiculatum læve flavescens includente.
3. U. SIMPLEX, frutescens, inermis, caule simplici, foliis oblongis, utrinque sensim attenuatis apiceque anguste et breviter acuminatis basi obtusis acutiuscule serrato-dentatis triplinerviis subpenninerviis supra densiuscule verruculoso-rugosis hirsutisque subtus dense pubescenti-hirsutis, cymis femineis subsessilibus petiolos non aut vix superantibus, dichotomis ramis hirtis, floribus in apice ramulorum solitariis v. ternis. ♀ In provincia Bogota Novæ Granatæ ad Salto de Tequendama (Triana). Caulis bimetralis, gracilis, inferne denudatus, apice foliatus denseque hirsutus, pilis urentibus destitutus. Folia 10-20 centim. longa, 5-8 centim. lata, ima basi laxius dentata, dentibus subæqualibus raro obtusatis, nervis primariis 6-8, intra marginem arcuatim anastomosatis; jugo inferiore paulo supra basim limbi nato, limbo exsiccatocoriaceo subtus minute alveolato, nervis nervulisque præsertim hirsutis, petiolo foliorum inferiorum 5-6 centim. longo sicut caulis glabrato. Flores dioici. *Cymæ* masculæ.... *Cymæ* fem. ex axillis foliatis vel denudatis ortis densiusculis, 3-4 centim. latis, fere rotundatis, repetite dichotomis, stigmatate rotundato. Perigonium fructiferum ut in præcedente miniatum achænium late ovatum vix compressum læviusculumque fere omnino vestiente. Habitu formaque foliorum et indumento a præcedente facillime distinguenda, sed illi nihilominus sat affinis. Caule omnino simplici sequente etiam congruens. (v. s.)
4. U. AURANTIACA (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 201; Monogr. p. 147), frutescens, caule simplici, foliis cordato-ovatis acuminatis crenato-dentatis, dentibus plerumque obtusis, trinerviis, supra sparsim exasperato-hispidis, subtus mollior pubescenti-tomentosis inermibus, cymis pedunculatis repetite dichotomis, floribus fem. in apice ramulorum subsessilibus pedicellatisve subternis s. laxè glomerulatis. ♀ In Brasilia meridionali, ad ripas inundatas flum. Mondego et Paraguay (Wedd.). — Caulis metralis, modice crassus, basi longè

nudus, internodiis elongatis, exsiccatus epidermide cinereo-fuscescente pubescente vel glabrata obectus, apice tomentosus. Folia 7-12 centim. longa, 4-9 centim. lata, sinu profundiusculo aperto cordata, superne sensim acuminata, ab ipsa basi usque ad apicem subæqualiter crenata, nervis primariis 6-8 ante marginem arcuatim anastomosantibus subtus prominulis (jugo inferiore dimidium limbum vix attingente), secundariis tenuissimis transversim rectiuscule vel subreticulatim connexis, pagina superiore pilis brevibus rigidisque et demum labentibus conspersa, costa prope petiolum densiuscule pilosa, nervis cæteris persæpe magis minusve pilosulis, lamina exsiccata rigida cystolithis minimis inæqualibus punctiformibus oblongisque creberrime inspersa, petiolo 3-5 centim. longo pubescenti-tomentoso inermi. Stipulæ oblongo-ovatæ, apice bifidæ, pilosæ. Flores dioici. Cymæ ex axillis foliatis natæ, petiolis breviores, patentes, divaricato-ramosæ, ramis pedunculoque sesquicentimetrali pubescentibus stimulosisque, masc. ter quaterve dichotomæ, floribus sessilibus aut pedicellatis in capitula pisi circiter magnitudine læteque aurantiaca digestis, perigonii 4-partiti lacinii piloso-hispidis; fem. pluries dichotomæ, floribus pedicellatis, perigonio 4-dentato, fructifero haud viso. (v. v.)

5. U. ELATA (Griseb. Fl. Brit. W. Ind. p. 154), scandens (an semper?), fere inermis, foliis oblongo-vel elliptico-ovatis obovatisve breviter acuminatis basi rotundatis aut subcordatis in dimidio apicali sinuato-crenulatis vel integris trinerviis supra glaberrimis vel parciissime hispido-stimulosis, costa subtus adpresse inconspicueque pilosula, cymis repetite dichotomis, floribus fem. discretis. ☉ In sylvis ins. Jamaica (Swartz, Wilson) et isthmi Panama (Sutton Haynes, n. 750). — U. sinuata Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 201; Monogr. p. 148. *Urtica elata* Swartz, Fl. Ind. occid. 1. p. 322; Poir. Encycl. suppl. 4. p. 224. — Truncus scandens 6-10 metralis (teste Sutton Haynes), ramulis crassiusculis, siccis epidermide lævi fusco-rubente v. cinerea vestitis cicatricibusque foliorum lapsorum notatis. Folia 7-18 centim. longa, 5-8 centim. lata, raro subcordata, acumine acuto, nervis primariis tenuibus utrinsecus 5-8 ante marginem arcuatim anastomosantibus subtus parum prominulis, secundariis tenuissimis inter primarios transverse parallelis reticulatisque denticulis interdum ciliolatis, lamina exsiccata tenuiter membranacea, supra cystolithis minimis creberrimisque sub lente albo-punctulata, iisdem corpusculis in nervis paginæ inferioris etiam perspicuis pubemque brevem quodammodo mentientibus, petiolo gracili 1-2 centim. longo puberulo. Stipulæ triangulari-ovatæ, 6-8 millim. longæ, inconspicue pubescentes. Flores dioici. Cymæ masc. ex axillis defoliatis horizontaliter patentes, pedunculo bi-tri-centimetrali, 5-ies dichotomæ, ramulis gracilibus subcentimetralibus, angulis apertis vacuis, floribus sessilibus in glomerulos pisiformes terminales congestis, perigonii 4-partiti segmentis rotundato-ovatis glabris, pistilli rudimento obconico depresso obsolete dentato. Cymæ fem. etiam in axillis foliorum lapsorum breviter pedunculatæ, patentes, divaricato-ramosissimæ, 6-8-ies dichotomæ, confluentes, ramulis gracillimis intricatis demum glaberrimis, floribus breviter pedicellatis rectiusculis, stigmatibus rotundato; perigonio fructifero 4-dentato, achæniis valde obliquum late ovatum ventricosum sub lente punctulatum omnino vestiente. — A cl. Grisebach dicitur arborea. (v. s.)
6. U. EXPANSA (Griseb. l. c. p. 155), arborea?, foliis ellipticis vel elliptico-oblongis obtusiusculis acutiusculisve basi vulgo emarginatis crenatis rariusve sinuato-repandis, cymis globosis axillaribus sive lateralibus pedunculatis repetite divaricato-dichotomis, floribus fem. longe pedicellatis, achænio.... ☉ In insulæ Jamaica ditione australi (Swartz, etc.). *Urtica expansa* Swartz (sec. Griseb.). Folia 7-18 centim. longa, 3-7 centim. lata. — Præcedenti affinis. (Griseb. l. c.)
7. U. CAPTATA (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 199; Monogr. p. 154), arborescens, plus minus stimulosa interdumque fere inermis, foliis ovatis acuminatis basi sinu aperto cordatis raro rotundatis crenato-dentatis, trinerviis supra sparsim scabro-hispidis, subtus pubescentibus, cymis pedunculatis dichotomis, floribus fem. in apice ramulorum demum conglomerato-capitatis, capitulis discretis didymisve, perigonio fructifero atro? ☉ In Venezuela, inter Caracas et La Guayra (Fendler, n. 1274); in sylvis humidis provinciæ Yungas

(Wedd.). — Arbor 3-4-metralis, trunco ramoso, ramulis crassiusculis apice pubescentibus et magis minus stimulosus (siccis epidermide cinerascente vestitis), demum fere omnino glabris, cicatricibus foliorum delapsorum signatis. Folia 10-15 centim. longa, 6  $\frac{1}{2}$ -10 centim. lata, apice sensim acuminata, dentibus inæqualibus obtusis vel acutiusculis paucis etiam mucronulatis, nervis primariis 5-8 juxta marginem reticulatim connexis vel in illo evanescentibus subtus parum prominentibus (jugo inferiore dimidium limbum superante), secundariis tenuibus transversim inosculantibus, lamina exsiccata rigidula, venis utrinque sed præsertim subtus intervenioque pubescentibus pilosulisve, costa petioloque gracili et plus minus elongato stimulis tenuibus retrorsisque sparsim aut dense armatis. Stipulæ ovatæ, bicuspidatæ, pubescentes. Flores dioici. Cymæ ex axillis foliatis at demum nudis ortæ, masc. vulgo majores laxioresque, fem. regulares vel irregulares, ramulis pubescentibus et densiuscule retrorsum stimulosus demumque magis minusve glabris, perigonio masc. subtiliter puberulo vel glabrato; glomerulis fructiferis capituliformibus ex apice ramuli dilatato receptaculiformi et perigoniis baccatis constantibus pisi circiter magnitudine, stigmatibus rotundato, achænio ovato submillimetrali læviusculo perigonio incluso. (v. v.)

β, *Pavonii*, fruticosa vel arborescens, subinermis, foliis basi rotundatis aut sinu aperto et haud profundo cordatis quam in typo sæpe minoribus obtuse crenatis subtus in nervis præsertim puberulis, floribus femineis in apice ramulorum 7-15 subsessilibus glomerulos parvos sive capitella formantibus. γ In Nova Granata et Peruvia (Pavon) necnon in Venezuela, valle amnis Tui, altit. 1200-1500 metr. (Fendler, n. 1275). Rami apice breviter densiusculeque pubescentes stimulis destituti. Folia nunc ovato-cordata cæterumque quoad formam et magnitudinem illis typi similia, nunc elliptico-vel oblongo-ovata basi que rotundata 6-15 centim. longa, 3-8 centim. lata, crenis parvis vel latiusculis depressisque subæqualibus juxta petiolum ut in typo plerumque minoribus etiamque nullis, rarissime mucronulatis, nervis omnibus præsertim subtus sicut petiolus puberulis et ut videtur inermibus. Cymæ fem. densiusculæ, ramulis plus minus divaricatis puberulisque et parce vel parcissime stimulosus, bracteis ut apud typum vix conspicuis, glomerulis fructiferis quam in eodem minoribus, achænio millimetrali, late ovato, vix compresso. — Stirps, velut plures præcedentium non satis cognita, quoad fructus atque in vivo denuo examinanda. Specimen alterum fructiferum a me huc adlatum fructibus atris gaudet (teste inventore Fendler), caractere cujus ope facillime distinguenda erit hæc species ab illis quorum perigonium femineum ad maturitatem rubrum fit aut albidum. (v. s. in hb. Boiss. et DC.)

#### B. — *Gerontogea*.

8. *U. SANDWICENSIS* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 177; Monogr. p. 150), arborescens, foliis plerumque oblongis breviter acuminatis basi obtusis acutatisve in dimidio superiore inæqualiter laxè et obtuse crenato-serratis rarius omnino integris penninerviis supra glabris subtus in nervis stimulosopilosulis, cymis dichotomis, floribus femineis in apice ramulorum discretis sive laxiuscule glomerulatis. δ In insulis Sandwicensibus (Gaudichaud, Remy, etc.). — Villebrunea crenulata Gaudich. Bot. Voy. Bon. t. 92. — Ramuli vix crassi, teretiusculi, glabrati, foliosi aut cicatricibus foliorum lapsorum notati et dessiccatione nodosi, epidermide castanea vel fusca vestiti. Folia 12-20 centim. longa, 5-8 centim. lata, a medio utrinque attenuata, apice acuta v. sæpius abrupte acuminata, acumine triangulari vel lanceolato et plerumque integro, nervis primariis utrinsecus 12-15 apice subdiscretis v. juxta marginem inter se anastomosantibus subtus prominulis (jugo inferiore quam sequentes multo breviores), nervis secundariis tenuibus transversim linea recta vel reticulatim inter se connexis, lamina exsiccata cystolithis minimis punctiformibus vix conspicuis dispersa, petiolo gracili 1  $\frac{1}{2}$ -4 centim. longo stimulosopilosulo. Stipulæ lanceolatæ, 2-3 centim. longæ, integræ, in nervis sparsim hispida. Flores dioici. Cymæ masc. ex axillis foliatis aut nudatis ortæ, patentes, pedunculo 1-2-centimetrali suffultæ, 4-5-ies dichotomæ, ramis semi-

centimetralibus patentibus sparsim brevissimeque setulosis, floribus in glomerulos pisiformes terminales congestis, perigonii 5-partiti segmentis ovatis glabris. Cymæ fem. pedunculatæ, corymbiformes, petiolos superantes, ramulis divaricatis hispidis sed demum glabris, floribus rectiusculis breviter pedicellatis subternis, stigmate rotundato; perigonio fructifero fere æqualiter 3-4-dentato v. denticulato valde carnoso; achænio suboblique ovato-rotundato,  $1\frac{1}{2}$ - $1\frac{1}{2}$  millim. lato, abrupte et obtuse acuminato, compresso, facie utraque tuberculata, perigonio undique vestito. (v. s. in hb. Mus. par., Lessert., Kew. etc.)

β, *glabra*, foliis ovatis inæqualiter crenulato-serratis subtus in axillis nervorum hirtellis. ♂ vel ♀. In ins. Oahu Sandwicensium (Beechey, Remy). — *Urera glabra* Wedd. Monogr. p. 149. *Procris glabra* Hook. et Arn. Bot. Voy. Beech. p. 96. — Folia paulo minora quam in typo (saltem in specim. visis), 7-12 centim. longa, 4-7 $\frac{1}{2}$  centim. lata, basi integra, nervis primariis 8-12 subtus vix prominulis, jugo inferiore paulo supra basin nato sequentibus æquali, achænio minute granuloso. (v. s. in hb. Mus. par., Lessert et Kew.)

γ, *mollis*, foliis membranaceis profundius serratis subtus in nervis et intertenuo molliter breviterque pubescentibus. — In insula Sandwicensi Owhihee, ad montem Kaah (Macrae). (v. s. in hb. Kew.)

## § 2. — *Cymæ uniparæ sive non vere dichotomæ.*

A. — *Species neogææ.*

\* *Stigma lanceolatum.*

9. *U. LACINIATA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> ser. 18, p. 203; Monogr. p. 160, t. 2, A. f. 10), arborea, aculeata, foliis inciso-lobatis, cymis uniparis floribundis, floribus femineis in apice ramulorum laxè glomerulatis, stigmate lanceolato. ♂ In sylvis Novæ Granatæ prope Maracaybo (Plée); prope Quindiu (Goudot); necnon in isthmo Panamæ (Seemann); in Peruvia orientali (Pavon, Spruce, n. 4140). — *U. girardinoides* Seemann, Bot. of Herald, n. 494, p. 194. *Urtica laciniata* Goudot, mscr. Arbor 3-4-metralis (teste Goudot), ramulis apice herbaceis siccitate valde angulato-plicatis aculeis elongatis cartilagineis (stimulis induratis) horridis et inter illos pubescentibus. Folia 2-3 decim. longa et lata, basi alte cordata, longe acuminata, profunde inciso-lobata, lobis lanceolatis utrinque 4 (inferiore multo majore extus laciniis 4-5 inciso, lobis reliquis versus apicem limbi decrescentibus), interdum falcatis integrisque v. dorso dentibus 1-4 magnis et inæqualibus instructis, terminali lateralibus conformi, nervis primariis utrinsecus numero loborum discretis subtus parum prominentibus, secundariis reticulatim vel linea recta connexis, lamina utrinque sparsissime hispidula, costa et nervis primariis aculeis (iis ramulorum similibus) parce obsitis, exsiccata submembranacea, petiolo gracili decimetrali vel breviori puberulo glabratove longe et crebriuscule aculeolato. Stipulæ ovatæ, bicuspidatæ, pubescentes. Flores dioici. Cymæ valde multifloræ, in axillis foliorum delapsorum (?) pedunculatæ, petiolos æquantes vel superantes, laxè ramosæ, ramisque et pedunculo puberulis haud stimulosi. Cymæ masc. ramulis elongatis diffusis, floribus sessilibus in glomerulos discretos laterales sessilesque digestis, perigonio 5-partito glabriusculo. Cymæ fem. erectiusculæ, diffuse ramosæ, glomerulis paucifloris sessilibus, lateralibus terminalibusque fere discretis, floribus subsessilibus, perigonio inæqualiter 4-lobato, segmentis lateralibus majoribus ovarium subæquantibus, fructifero quam achæmium rotundato-ovatum seu lenticuliforme  $1\frac{1}{2}$  millim. longum triente brevior. — A cæteris conformatione stigmatis distinctissima. Habitu autem sequentem quodammodo refert atque sicut illa vernacule apud Novogranatenses Pringamosa dicitur vel Papaya colorada. (v. s.)

\*\* *Stigma rotundatum.*

10. *U. BACCIFERA* (Gaudich. Bot. Voy. Uran. p. 497), fruticosa vel arborea, caule seu trunco ramisque persæpe aculeato, foliis late ovatis obovatisve vel

floribus in apice ramulorum cymæ fere discretis, stigmatibus rotundato.  $\zeta$  In ins. Madagascar (Chapelier, Meller). — Caules e rhizomate repente orti, pauci, erecti, simplices, 1-4 decim. alti, basi teretiusculi, rugosi, rubentes, prope apicem sub-4-anguli stimulisque brevibus horridi. Folia 10-26 centim. longa, 2-3½ centim. lata, fere a basi ad apicem sensim attenuata, in toto margine dentata, dentibus æqualibus inæqualibusve acutis sinibusque angulatis vel sæpius rotundatis sejunctis, acumine ipsius limbi etiam dentato acutoque; nervis primariis utrinque 10-18 horizontaliter patentibus, pagina superiore secus costam et nervos majores præcipue stimulosæ cæterum glaberrima, inferiore subtiliter pubescente, nervis eminentibus densiusculeque stimulis, petiolo 1-2-centimetro horrido. Stipulæ ovatæ, acuminatæ, in nervis hispidæ. Flores dioici? Cymæ fem. petiolis vix longiores, parce breviterque ramulosæ, compressiusculæ (teste Chapelier), erecto-patulæ, pedunculo brevissimo ramisque hispidis et sparsim stimulosis, floribus sessilibus valde obliquis, perigonii inæqualiter 4-lobi lobis glanduloso-ciliatis, interioribus sive lateralibus ovatis quam exteriores multo majoribus, lobo inferiore minimo acuto, superiore gibboso. Achæmium oblique ovatum, 1 millim. vix longum, compressiusculum, angustissime marginatum, facie utraque medio læviuscula cæterquin punctato-scabra. — Specimen unicum vidi floribus omnibus femineis; tamen planta ab inventore dicitur monoica. Stimuli testimonio ejusdem vehementer urentes sunt. Perigonium fem. illud specierum quarundam sectionis prioris gen. Laportæ refert. (v. s. in hb. Mus. par. et Kew.)

\*\* Frutices scandentes.

15. U. ACUMINATA (Gaudich. Bot. Voy. Uran. 496, non Miq.), scandens, ramis inermibus, foliis ellipticis vel elliptico-rotundatis abrupte acuminatis basi obtusatis rotundatisve integerrimis trinerviis glabris, cymis uniparis, floribus in apice ramulorum discretis vel laxè glomerulatis, stigmatibus rotundato.  $\cup$  In insula Mauritio (Commerson, du Petit Thouars. l. c.) et Timor (Riedlé). — Dcne Herb. Timor. p. 162; Wedd. Monogr. p. 157; Miq. Fl. Ind. bat. 1 (2). p. 233. Urtica acuminata Poir. Encycl. suppl. 4. p. 224. — Rami lignosi, alii foliferi graciles elongati, alii breviores crassioresque floriferi, epidermide rubente glaberrimaque vestiti. Folia 5-10 centim. longa, 3½-6 centim. lata, acumine brevi angusto obtusaque, nervis primariis utrinsecus 2-3, secundariis inter illos transverse anastomosantibus, membranacea, utrinque glaberrima vel juniora in costa nervisque parum eminentibus subtus puberulis stimulisque rarissimis brevibusque armatis, pagina superiore laminæ exsiccata cystolithis oblongo-linearibus creberrime ornata, petiolo gracili mox glabrato quam dimidius limbus brevior, exsiccato cystolithis subtilissime punctulato. Stipulæ lanceolatæ, 4-5 millim. longæ, apice sæpius bifidæ, glabriusculæ, cito deciduæ. Flores dioici. Cymæ in axillis foliorum delapsorum breviter pedunculatæ, patentes, petiolis multo longiores, divaricato-ramosæ, ramulis glaberrimis aut pilis urentibus hic et illic armatis, floribus masc. breviter pedicellatis, perigonio 5-partito, singulis laciniis stimulo gracili dorsali plerumque instructis cæterum glaberrimis; floribus fem. subsessilibus magis minusve obliquis, perigonio sub-4-dentato vel 4-lobato, lobis glaberrimis, lateralibus interdum multo majoribus « fructifero seminis milliæci magnitudine exquisitè coccineo-rubente apiculo obtuso (achæni apice) nigricante » (Commerson). Achæmium ovatum, 2 millim. circiter longum, ventricosum, subtilissime granulosum, perigonio vix longius. — Vulgo Liane du voyageur vocatur in Mauritio necnon Liane blanche. (v. s. in hb. Mus. par., Juss. et Lessert.)
16. U. HYPSELODENDRON (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 203), scandens, foliis ovatis vel rotundato-ovatis acuminatis obtusis vel subcordatis minute crenulatis subtriplinerviis glabriusculis, cymis uniparis, floribus femineis in apice ramulorum laxiuscule glomerulatis, stigmatibus rotundato.  $\cup$  In locis humidis umbrosissimis montium Abyssiniæ, alt. 2000-3000 metr. (Schimper). — Wedd. Monogr. p. 158. Urtica Hypselodendron Hochstett. msc. in A. Rich. Tent. fl. Abyss. 3. p. 260. Hypselodendron Schimper Hochstett. in sched. exs. Schimp. n. 1136. — Caulis lignosus, frutices arboresce etiamque proceras scandens, ramis extremis pendulis, ramulis epidermide lævi palli-

deque castanea glaberrima vel puberula vestitis. Folia 6-15 centim. longa, 3-12 centim. lata, apice plerumque abrupte acuminata, acumine angusto obtusiusculo, inferne integra, cæterum crenulis obtusis subæqualibus usque ad apicem ornata, supra glaberrima, lamina exsiccata cystolithis punctiformibus creberrime inspersa subtus nervis transversis eleganter reticulata, glabrata vel costa venisque primariis puberulis sparsimve pilosulis cystolithosque raros hic illic exhibentibus, petiolo gracili puberulo vel glabrato longitudine varia. Stipulæ lanceolatæ, fusco-scariosæ, apice bifidæ, nervis et interdum margine pilosulis, cito deciduæ. Flores dioici. Cymæ masc. petiolis breviores vel paulo longiores, vix multifloræ, pedunculo brevi ramisque (haud numerosis) puberulis, floribus brevissime pedicellatis, perigonii 4-partiti segmentis subtilissime puberulis. Cymæ femineæ divaricato-ramosæ, ramis minutissime puberulis, floribus subsessilibus valde obliquis, perigonii inæqualiter 4-lobi vel subbilobi (lobis exterioribus subnullis), lobis interioribus achænium oblique ovatum  $1\frac{1}{2}$  millim. circiter longum subventricosum subtiliter granulosum subæquantibus. (v. s.).

17. *U. OBOVATA* (Benth. in Hook. Niger fl. p. 515), scandens, ramis novellis stimulis crassis sive aculeis horridis, foliis ellipticis vel oblongo-ellipticis obovatisve abrupte acuminatis basi obtusis rotundatisve plerisque crenatis trinerviis glaberrimis inermibusque vel in costa et nervis subtus parce breviterque stimulosus, cymis uniparis, floribus femineis in apice et ad latera ramulorum inflorescentiæ laxiuscule glomeratis, stigmatibus rotundato. ☉ In Sierra Leone (Vogel) et Fernando Po (Mann). — Wedd. Monogr. p. 159. Caulis fruticosus. Rami elongati, teretes, exsiccati epidermide obscure rubente vestiti. Folia 5-12 centim. longa,  $2\frac{1}{2}$ -6 centim. lata, acumine subcentimetro angusto obtuso et plerumque integro, nervis primariis utrinsecus 5-6 subtus parum prominentibus secus marginem arcuatim connexis (jugo inferiore medium limbum attingente interdumque superante), secundariis linea recta aut reticulatim inter se anastomosantibus, lamina exsiccata submembranacea aut rigidula supra cystolithis minimis crebriuscule inspersa, subtus hæc corpuscula fere linearia secus nervos præsertim exhibente, petiolo gracili longitudine variabili nunc vix centimetro nunc limbo æquilongo inermi rariusve stimulosus, exsiccato sæpe cystolithis minutis punctato. Stipulæ. . . . . Flores dioici. Cymæ in ramulis foliosis numerosæ, pedunculatæ, foliis breviores rariusve illis æquilongæ, pedunculo 1-3-centimetro erecto-patulo, ramis subdivaricatis glabris et fere inermibus, ramulis extremis vero seu floriferis demum (in cyma feminea) stimulis longiusculis subfasciculatisque horridis. Flores masc. breviter pedicellati, in glomerulos parvos discretos digesti, perigonio 5-partito glabro, alabastro sesquimillimetro sphaeroideo, pistilli rudimento obconico depresso obsolete 5-lobato. Flores fem. subsessiles, in glomerulos numerosos ad apices ramulorum fere confluentes congesti, stimulis quasi involucrati, perigonio tubuloso ovarium fere æquante, ore inæqualiter denticulato, fructifero baccato rubro-aurantiaco quam achænium oblique ovatum bimillimetrale paulo brevior. (v. s. in hb. Kew.)

♀, *longifolia*, ramulis fere omnino inermibus lævibus et glaberrimis, foliis quam in typo minoribus sæpiusque subintegrè serrulatisve, petiolo elongato. ☉ In iisdem locis ac typus (Vogel). — *Urera oblongifolia* Benth. l. c. p. 516. (v. s. in hb. Kew.).

18. *U. CAMEROONENSIS*, scandens, fere glaberrimus et (ut videtur) omnino inermis, foliis oblongis anguste acuminatis basi angustatis et ad insertionem petioli subemarginatis integerrimis tri-tripli-nerviis, nervis primariis paucis, cymis uniparis, floribus fem. . . . . ☉ In montibus Cameroon ins. Fernando Po, altit. 1500 metr. (G. Mann, n. 2173). — Frutex 5-metralis, ramis teretibus siccitate fuscis, internodiis superioribus 1-2-centimetro. Folia 8-12 centim. longa, 3-4 centim. lata, acumine lineari obtusiusculo, nervis primariis in utroque latere limbi 2-3 arcuatis in margine evanescentibus folii junioris subtilissime puberulis, jugo inferiore medium limbum superante subtusque parum prominente, venulis parum conspicuis subparallele transversis, lamina exsiccata coriacea supra minutissime fusco-punctulata, cystolithis in pagina inferiore prorsus inconspicuis, petiolo 2-4 centim. longo. Stipulæ triangulari-ovatæ, vix centimetro cito deciduæ. Cymæ masc. (solum visæ) breviter

pedunculatæ, folio breviores, ramis paucis patulis, glomerulis subdiscretis, floribus subsessilibus, perigonii 4-partiti alabastro bimillimetrali (v. s. in hb. Kew.).

*Species exclusæ.*

- U. AMPLISSIMA Bl. = *Laportea amplissima*?
- U. COMMERSONIANA Wedd. = *Laportea crenulata*.
- U. CRENULATA Wedd. = *Laportea crenulata*.
- U. EXCELSA Wedd. = *Laportea Gigas*.
- U. GAUDICHAUDIANA Henslow = *Fleurya æstuans*.
- U. JAVAENSIS Gaudich. = *Laportea crenulata*.
- U. LAMIFOLIA Gaudich. = ? *Boehmeria nivea*.
- U. LEICHARDIANA Wedd. = *Laportea photiniphylla*.
- U. MADAGASCARIENSIS Gaudich. = *Obetia Madagascariensis*.
- U. MICROSTIGMA Wedd. = *Laportea microstigma*.
- U. PALMATA Gaudich. = *Girardinia palmata*.
- U. PARIETARIFOLIA Gaudich. = ?
- U. ROTUNDIFOLIA Wedd. = *Laportea Gigas*.
- U. STIMULANS Wedd. = *Laportea microstigma*.

VIII. SCEPOCARPUS (1).

Flores dioici?, glomerulati, glomerulis in cymam uniparam s. non vere dichotomam digestis, inflorescentiis axillaribus solitariis. *Masc.* (non visi). — *Fem.* Perigonium tubulosum, ore contracto inconspicue denticulato, ovarium omnino vestiens. Ovarium rectum, oblongo-ovatum. Ovulum erectum, in fundo loculi subsessile. Stigma sessile, penicillato-capitatum rotundatumque, persistens. Achænium oblique ovatum, utrinque convexum, læviusculum, perigonio baccato et colorato arcte vestitum. Semen ut in *Urera*. — Frutex scandens, insulæ Fernando Po Africa occid. incola, pilis urentibus parce armatus. Folia alterna, dentata, trinervia, cystolithis minimis subpunctiformibus. Stipulæ axillares, integræ, deciduæ. Flores feminei ebracteati? Fructus sessiles sive pedicello v. ramulo incrassato oblique insidentes.

S. MANNII, scandens, foliis elliptico-ovatis abrupte breviterque acuminatis basi subcordatis toto margine crenato-dentatis utrinque hispido-pubentibus. ☉ In ins. Fernando Po (G. Mann). — Caulis 3-6-metralis, ramis apice pubescentibus foliiferisque cæterum ut videtur glabris nudatisque. Folia in specim. viso 8-12 centim. longa, 5½-8 centim. lata, acumine vix centimetrali, dentibus versus petioli insertionem minoribus, nervis primariis in utroque latere limbi 5, jugo inferiore dimidium limbum vix superante, lamina exsiccata membranacea obscure viridi, petiolo 2-4-centimetrali pubescente atque (velut limbus ipse) stimulis parvis insperso. Stipulæ semicentimetales ovatae, et longiuscule acuminatæ, dorso pubentes. Cymæ fructiferæ (solum visæ) in ramo foliis denudato ovatae v. rotundatæ, pedunculo 3-centimetrali patulo suffultæ, patulo v. divaricato-ramosæ, ramis pedunculoque puberulis, ramulis ad insertionem perigoniorum incrassatis stimulis brevibus horridis. Perigonium fructiferum ovatum, ventricosum, 1¼-1½ millim. longum (in sicco), in planta viva (teste inventore) corollam luteam mentiente. Achænium millimetrale, subtiliter granulatum. — Habitus *Ureræ* cujusdam. Indagatio accurata partium fructus in sicco sat difficilis. Perigonium in segmenta duo (valvarum

(1) E verbis græcis *σκηπε*, vestio et *καρπός*, fructus.



more) in herbario frequenter dehiscit. Basis fructus vivi forsan in excavatione ramuli cymæ incrassati recepta, marginibus brunneis hujus excavationis a cl. Mannio (in sched.) verisimiliter pro calyce acceptis.

IX. GYROTÆNIA Griseb. in *Mem. Acad. Am. sc. et art.*, N. sér. 8. (1860) p. 174.  
— Ureræ spec. Wedd. *Monogr.*

Flores dioici, spicati aut capitati (rhachi s. receptaculo femineorum carnosus), inflorescentiis axillaribus solitariisque. *Masc.* Perigonium 4-partitum, segmentis orbicularibus, alabastro subumbilicato. Stamina 4, antheris reniformibus. Pistilli rudimentum globosum. *Fem.* Perigonium parvum bilobum, lobis oppositis æqualibus margine ciliato-denticulatis. Ovarium rectum, ovoideo-oblongum. Ovulum imo loculo affixum, erectum. Stigma sessile, penicillato-capitatum rotundatumque. Achænium subsuccum, vix oblique ovatum, subcompressum, stigmatem persistente coronatum basique perigonio subimmutato stipatum. Semen pericarpio conforme. Albumen... Embryo... — Frutices sive arbusculæ antillanæ, stimulis veris forsan carentes. Folia alterna, serrato-dentata vel subintegra, cystolithis punctiformibus oblongisque inspersa. Stipulæ axillares, binerves, apice integræ. Flores fem. ebracteati. Cl. Grisebachii sententia hoc genus inter Procrideas collocandum est, genus Pileam cum Myriocarpa (gen. Boehmeriearum) connectens; at me iudice locus ejus legitimus est inter Ureæas cum quibus structura et indole omnium fere partium optime congruit. A genere Ureræ non essentialiter discrepat nisi perigonio bilobo post anthesin parum aucto.

1. G. MYRIOCARPA (Griseb. l. c. Cat. pl. cub. p. 60), foliis lanceolatis vel elliptico-lanceolatis acuminatis in toto margine dentato-serratis subtus pubescentibus, floribus femineis in glomerulos spicatos dispositis, rhachi demum carnosâ irregulariter complanato-dilatata et sinuato-convoluta. ♀ In sylvis Cubæ orientalis prope villam Monte-Verde (Wright n. 531 et 531'). — Arbor parva, ramulis epidermide rubente hirtopubescenti demumque glabra vestitis. Folia 4-7 centim. longa, 1½-3 centim. lata, basi acuta vel obtusiuscula, plerumque sensim acuminata, acumine ipso serrato, triplinervia, nervis basilaribus ultra medium limbi productis, pagina superiore folii junioris sparsim pilosa, situ pilorum in folio seniore exsiccato et submembranaceo nonnihil prominente et albicante, petiolo 1-1½ centim. longo patulo sicut nervi primarii densiuscule hirtopubente. Stipulæ ovatæ, 5 millim. circiter longæ, acutæ v. subacuminatæ, extus adpresse pubentes, persistentes. Inflorescentiæ utriusque sexus 3 centim. circiter longæ, patulæ, rhachi junioris breviter hirta. Spica masc. gracilis, floribus bracteatis, alabastro 1½ millim. lato. Spica fem. e glomerulis, discretis vel plus minus confluentibus interdumque subunilateralibus confecta, floribus haud semimillimetralis numerosissimis in lobulis rhacheos sessilibus perigonii lobis ovatis margine breviter glanduloso-ciliatis, ovario recto stigmatem rotundato papilloso coronato. Inflorescentia fructifera valde aucta, achæniis ¼ - ½ millim. longis, recte vel suboblique ovatis nonnihil compressis basi perigonio vix aucto stipatis atque (teste cl. Wrightio) in rhachi pulposa subpeltata albidâ plus minus lobata nidulantibus.
2. G. SPICATA, foliis suboblique ellipticis oblongisve acuminatis integris vel obsolete sinuatis glaberrimis, floribus femineis in spicas axillares rhachi rectiuscula carnosâ congestis. ♀? In insula Jamaica (Swartz). Ureræ spicata Wedd. *Monogr.* p. 155; Griseb. *Fl. Br. W. Ind.* p. 155. — Ramuli teretiusculi, circiter crassitie pennæ anserinæ, epidermide exsiccata pallide rubente glabra lævisque. Folia 6-12 centim. longa, 2½ centim. lata, in triente superiore sensim acuminata, acumine obtusiusculo, juxta petioli subcentimetralis insertionem nonnihil emarginata, trinervia, nervis basilaribus didimidum limbi plerumque superantibus et cum primariis sequentibus prope marginem arcuatim anastomosatis, secundariis paucis laxè reticulatis, lamina exsiccata

rigidula. Stipulæ anguste triangulares, semicentimetrales, puberulæ. Inflorescentiæ fem. (solum visæ) 1-1½ centim. longæ, glomerulis confluentibus rhachim carnosam simplicem obtegentibus. Perigonii lobi rotundati vel ovati, ciliolati. Achænium ovatum, ¾ millim. longum, ventricosum. (v. s. in hb. Mus. brit., Soc. Linn. et Kew.).

3. *G. TRINERVATA*, foliis ellipticis obtusiusculis prope apicem serrato-dentatis cæterum integris subtus subtiliter tomentosis albicantibusque, floribus fem. in receptaculis globosis breviter pedunculatis sessilibus. ¾ In insula S. Domingo (Bertero). — *Procris trinervata* Poir. *Encycl. illustr.* t. 763, f. 2. *Boehmeria trinervata* Pers. *Synops.* 2. p. 556. — Ramulus unicus visus gracilis, epidermide rubente adpresse puberula vestitus. Folia 3-4 centim. longa, 2 centim. circiter lata, basi rotundata, sub apice utrinque dentibus 3-4 acutis obtusiusculisve notata, tri-tripli-nervia, nervis basilaribus ad trientem superiorem limbi protensis et cum nervis primariis sequentibus arcuatim anastomosatis, intervenio subtus albicante et distincte reticulato; lamina exsiccata coriacea supra rugosula et nitida. Stipulæ ovata, binerviæ, caducæ. Inflorescentiæ fem. (solum visæ) in singulis axillis ut videtur solitariæ, petiolo breviores, receptaculis fructiferis globulosis 4-5 millim. latis in apice peduncululi solitarii vel geminis. Achænia ½-¾ millim. longa, ovata, stigmatibus rotundato persistente coronata basi perigonio bilobo s. fere naviculari stipato. (v. s. in hb. Monac.)

- X. *GIRARDINIA* Gaudich. *Bot. Voy. Uran.* p. 498; *Decsne. in Jacquem. Voy.* 4. p. 151; *Wedd. Monogr.* p. 163. t. 2. B. — *Urticæ spec. auct.* — *Urtica*, e, *Girardinia* Endlich. *Gen. pl.* p. 283.

Flores dioici vel monoici, masculi in glomerulos racemum simplicem vel furcatum aut paniculam efformantes congesti, feminei cymoso-paniculati et in ramulis extremis scorpioideo-ordinati, cymis fructiferis densissimis stimulisque horridis, inflorescentiis (semper unisexualibus) in singulis axillis solitarii geminisve, pedicellis florum masculorum articulatis. *Masc.* Perigonium 4-5-partitum, segmentis ovatis hispidis. Stamina 4-5. Ovarii rudimentum globosum v. cupuliforme. *Fem.* Perigonium bipartitum, segmento altero sæpius multo majore fere tubuloso sed hinc ad basim usque fisso necnon apice denticulato, altero minimo lineari parum conspicuo, rarissime late ovato aut prorsus abortiente. Ovarium ovoideo-lanceolatum. Ovulum sæpius adscendens, funiculo gracili sustentatum. Stigma elongato-filiforme, sæpius subtilissime papillosum, subpersistens. Achænium oblique rotundatum, compressum, facie utraque convexiuscula lævi v. subtiliter granulosa rarissime setulosa, perigonio persistente patenteque calceato. Albumen parvissimum. Embryo cotyledonibus rotundatis utrinque emarginatis, radícula parva oblonga. — *Herbæ* elatæ, annuæ aut perennantes et plus minus lignosæ, asiaticæ et africanæ, erectæ, pilis aculeisve urentibus ut plurimum crebre armatæ. Folia alterna, grosse serrata aut varie lobata, trinervia, cystolithis punctiformibus inspersa. Stipulæ sæpe majusculæ, utriusque folii in unam axillarem integram v. apice bifidam coalitæ, haud persistentes. Flores virescentes, masculi cito decidui bracteis paucis minimis stipati, feminei bracteolati aut fere ebracteati inter setas creberrimas achænia maturautes diuque persistentes; ramulis cymæ fem. demum valde incurvatis latereque convexo præsertim horridis.

1. *G. HETEROPHYLLA* (Decne in Jacquem. *Voy.* 4. p. 151. t. 153), foliis ovatis vel tri-quinque-lobis basi cordatis breviter petiolatis acute serratis, lobis lanceolatis acuminatis, impari præsertim nonnunquam lobulo laterali instructo, supra magis minusve pilosis venis venulisque subtus hispido-pubescentibus et sparsim stimulosi, stipulis vix amplius oblongis ovatisve, cymis fructiferis

elongatis, pedunculis petiolo longioribus. ④ In Indiæ orientalis regione temperata Himalayæ, alt. 1200-3500 metr. a Sikkim ad Marri, in sylvis ruderalisque frequens, nec non in montibus Khasia, Ava? (Wall.), Assam, Kashmir et in Arabia (Forsk.). — Blume, Mus. lugd. bat. 2. p. 158; Wedd. Monogr. p. 164. *Urtica palmata* Forsk. Fl. Ægypt. arab. p. 159. *U. heterophylla* Vahl. Symb. bot. 1. p. 76; Willd. Sp. pl. 4. p. 362. *U. armata* Kunth. *U. diversifolia* Link. Enum. 2. p. 385. *U. horrida* ejusd. 1. c. Caulis basi sæpe digitum crassus, demum ligneus, ramulis gracilibus striato-sulcatis pubescenti-hispidis ad nodos sæpe hirsutis stimulisque raris et sæpe majusculis insuper conspersis. Folia 6-8 centim. longa, lobis subæqualibus inæqualibusve, lobis inferioribus etiam interdum extus lobulo instructis, serraturis parvis triangularibus, acumine lineari acutissimo, petiolo quam limbus dimidius multo brevior pubescenti-hirto et sparsim stimulosus. Stipulæ oblongæ, pro genere parvæ, bifidæ, pubescenti-hispidæ, caducæ. Flores dioici, inflorescentiis solitariis. Racemi masculi graciles, laxè paniculati, paniculis quam folia brevioribus sed petiolis superantibus, floribus subsessilibus, perigonio hispidulo. Paniculæ fructiferæ solitariae, elongatæ, lobulatæ, basi nonnunquam interruptæ, cum pedunculo (3-4-centimetrali) 1-1½ dm. longæ, rhachi pedunculoque pubescenti-hispidis aut inermibus, ramulis pedicellisque contra stipulis validis horridis et inter illos hispidis. Perigonii segmentum superius oblongo-ovatum, inferiore lineari dimidio longius, dorso hispidum, dentibus apicalibus triangularibus. Achænium late ovatum, nonnunquam subcordatum, 2-2½ millim. longum, minute punctatum, nigrescens, stigmatè persistente duplo triplove longius. — Herba robusta adspectu quidem formidabili, sed ictu (teste Hookerio) haud periculoso, in Indiæ orientalis regionibus temperatis (circa mayseta præsertim) maxima copia pullulescens. (v. s.)

2. *G. PALMATA*, foliis 3-5-lobatis basi late cordatis (lobis oblongo-lanceolatis grosseque et acute inciso-serratis) longiuscule petiolatis utrinque sed præsertim subtus hirsuto-pilosis sparsimque stimulosis, stipulis sæpe majusculis lateque cordatis rarius ovatis, cymis fructiferis elongatis quam pedunculi sui multo longioribus. ④ In Indiæ orientalis regione temperata montium Nilghiri (Leschenault, Hooker et Thomson, etc.). — G. Leschenaultiana Decsne l. c. p. 152; Wedd. Monogr. p. 165. *G. palmata* Gaudich. Uran. p. 498; Bl. l. c. *Urtica palmata* Leschenault, non Forsk. *U. acerifolia* Zenker, Pl. Ind. Schmid. dec. 1. n. 3 et 4. Herba elata, ramis ramulisque sulcatis adpresse pilosis et stimulis longis validisque horridis. Folia in specim. visis 18-25 centim. lata, sinubus loborum magis minusve profundis nunquam vero ipsam costam attingentibus, lobis inferioribus extus lobulo unico majusculo v. 2-3 parvis instructis, serraturis magnis triangulari-acutis, dente terminali triangulari-lanceolato, petiolo limbum longitudine subæquante pubescenti-piloso, costaque et nervis primariis subtus plus minus stimulosis. Stipulæ late ovato-cordatæ, 1½-3-centimetrales, extus sparsim et adpresse pilosæ. Flores dioici. Racemi masc. solitarii, simplices v. bifurci v. simpliciter ramosi, pedunculati, pedunculo 2-3-centimetrali hirsuto et parce stimulosus, floribus stimulosopilosis. Cymæ fructiferæ solitariae, elongatæ, cum pedunculo (3-centimetrali) 1-2 dm. longæ longioresve, prope basim v. hic et illic interruptæ, rhachi pilosa, ramulis brevibus incurvis pedicellisque pilosis et stimulis longis validisque dense obsitis. Perigonii segmentum superius oblongo-ovatum, inferiore vix longius, dorso sparsim stimulosopilosum, dentibus apicalibus triangulari-lanceolatis. Achænium ovatum, 2½-3 millim. longum, stigmatè persistente subæquilongum. — A præcedente cui affinis differt stipulis plerumque majoribus, dentibus foliorum, etc. (v. s.)
3. *G. ZEYLANICA* (Decsne l. c. p. 152), foliis pinnatifido-5-lobatis basi late cordatis lobis magis minusve profundis inciso-serratis lobulatis longe petiolatis supra sparsim hispidis aut hirsutis stimulosisque vel inermibus, venulis subtus adpresse pubescentibus vel hirto-pubescentibus, nervis stimulosis, stipulis late cordatis, cymis fructiferis fere reniformibus pedunculo longioribus. ④ In montibus temperatis Peninsulæ Indiæ orientalis Nilghiri et Concan, necnon in Zeylan et Java. — Wedd. Monogr. p. 167; Miq. Fl. Ind. bat. 1 (2). p. 233. *G. hibiscifolia* Miq. Pl. Jungh. 32; Bl. l. c. *Urtica Zeylanica* Burm. Thes. zeylan. p. 232. *U. heterophylla* Wight, Icon. pl. Ind. 2. t. 687, non Vahl.

Rami sulcati, adpresse pubescentes et stimulis gracilibus brevibusque conspersi. Folia (in var.  $\alpha$ ) deltoideo-rotundata, 12-16 centim. lata, raro ad costam usque partita, lobis inferioribus extus lobulo majusculo instructis, serraturis triangularibus acutiusculis obtusiusculisve, dente terminali lanceolato, petiolo quam limbus haud raro longiore adpresse pubescente, costa nervisque primariis paginæ inferioris stimulis gracilibus magis minusve horridis. Stipulæ late cordatæ, subcentimetricæ, apice acuminatæ angusteq. bifidæ, extis adpresse pubescentes. Flores monoici. Racemi masc. in axillis inferioribus ramulorum simplices vel bifurci, raro decimetricæ, pedunculo pubescente 1-3 centim. longo suffulti, floribus hispidis. Cymæ fem. ut plurimum solitariae, pedunculatæ, sub anthesi pyramidatæ, ramulis subdistichis laxiuscule patentibus pedunculoque 2-3-centimetrici pubescentibus et stimulosi; fructiferae (in var.  $\alpha$ ) 3-4 centim. longæ, pedunculis suis paulo longiores, densiuscule hispidæ stimulisque gracilibus hirsutissimæ. Perigonii segmentum superius oblongum, inferiori lineari æquilongum, dorso sparsim hispidopilosum, dentibus apicalibus lanceolatis. Achænium late ovatum, 2-2½ millim. longum stylo persistente paulo longius. — A duabus præcedentibus prima facie distinguitur cyma fructifera brevi et coarctata stimulisque ejusdem multo gracilioribus. (v. s.)

$\beta$ , *vitifolia*, foliis magis minusve profunde trilobatis vel cordato-ovatis lobis nullis aut vix ullis, stipulis ovato-oblongis, pedunculis quam in typo longioribus. ① In ins. Java (Zollinger, n. 2212). G. vitifolia Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 181. (v. s.)

4. G. *EROSA* (Decsn. l. c. p. 151 et 152), foliis pinnatifido-subseptemlobatis (lobis inciso-serratis lobulatisve) basi truncatis vel late cordatis longiuscule petiolatis supra sparsim hispidis stimulosisque subtus adpresse pubescentibus nervis horridis, stipulis ovatis, cymis fructiferis pedunculo multoties longioribus. ① In Madagascarie provincia Emirna juxta prædium Ra-Massina (Bojer) et prope Tananarivo in fossis (Goudot). — Wedd. Monogr. p. 167, t. 2, B, f. 8 et 9. — Stirpis 1-2-metralis, ramis gracilibus sparsim et sub-adpresse hispidopilosulis stimulisque longis horridis. Folia rotundata, 10-15 centim. lata, non raro prope usque ad costam partita, lobis inferioribus (quam cæteri multo majoribus) extus plerumque 2-3-lobulatis, serraturis triangularibus lanceolatisve (terminalibus s. acuminibus angustissimis), petiolo 4-6 centim. longo adpresse v. patentim hispido costaque et nervis primariis stimulis longis armatis. Stipulæ ovatæ, 1-2-centimetricæ, apice bifidæ, extus subtiliter hispidæ. Flores monoici. Racemi masc. simplices, breviter pedunculati, petiolis breviores, floribus perigonio sparsim hispidulo. Cymæ fem. solitariae v. in iisdem axillis ac racemi masculi nascentes, oblongæ; fructiferae stimulis tenuibus numerosissimisque horridæ, perigonii segmento superiore ovato v. oblongo inferiori lineari-subulato æquilongio dorso sparsim setulosos dentibus apicalibus lanceolatis, stylo quam ovarium multo longiore. (v. s. in hb. Mus. par. et Delessert).

5. G. *BULLOSA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 203, et 4<sup>e</sup> sér. 1. p. 181; Monogr. p. 168. t. 2. B. f. 5-7), foliis rotundato-ovatis duplicato-serratis sublobulatis basi truncatis cordatisve breviter acuminatis longiuscule petiolatis utrinque sparsim stimulosi subtusque subtiliter pubescentibus, stipulis ovatis, cymis fructiferis plus minus elongatis brevissime pedunculatis. ① In Abyssiniæ præruptis, prope Aosa altit. circiter 2500 metr. (Schimper). — Urtica bulbosa Hochstett. in Bot. Zeit. 33. p. 259; A. Rich. Tent. Fl. Abyss. 2. p. 262. Herbacea v. demum lignescens, præalta, ramis crassis fistulosis pubescenti-hispidis, stimulis brevibus longiusculisve et interdum retrorsis albicantibus basi que obscuris dense obsitis. Folia ampla, 1½-4 decim. longa et lata, basi plerumque cordata, subacuminata, dentibus lobuliformibus inæqualibus utrinque subacute serratis, interdum bullato-rugosa, petiolo quam limbus dimidius brevior densiuscule brevissimeque pubescente v. tomentoso et stimulis longiusculis consperso rarius subinermi. Stipulæ 2 centim. longæ, integræ v. brevissime bifidæ, extus subtilissime puberulæ. Flores dioici, inflorescentiis solitariis geminisve. Racemi masc.... Cymæ fem. lobulatæ, subinterruptæ, rariusve longiter ramosæ, petiolo breviores longioresve, fructiferae demum densissimæ, ramulis pedicellisque creberrime stimulosi, perigonii

segmento superiore late oblongo s. quadrato quam inferius subulatum subduplo longiore dorso crasse carinato et infra apicem tridentatum gibbosum pilisque quibusdam rigidis subadpressis insuper instructo dentibus apicalibus triangularibus. Achænium rotundatum, 3-3½ millim. longum, minute granulatum, nigrum, stylo persistente haud dimidio longius. (v. s.)

6. *G. CONDENSATA* (Wedd. Monogr. p. 169. t. 2. B. f. 1-5), foliis plerumque late ovatis rarius bi-tri-lobis basi obtusatis vel truncatis subcordatisve acuminatis ut plurimum grosse serratis utrinque stimulosi supra sparsim hispidis subtus breviter pilosulis, petiolo quam limbus dimidius sæpius brevior, stipulis ovatis, cymis fructiferis rotundatis vel oblongis subsessilibus. ① In Abyssiniæ locis umbrosis ad altit. circiter 2000 metr. (Schimper), necnon in Indiæ himalayensis reg. bor. occid. temperata ad altit. 2200 metr. (J. Thomson). *G. condensata* et *G. adoensis* Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18, p. 203; et 4<sup>e</sup> sér. 1. p. 181. *Urtica condensata* et *U. adoensis* Hochstett. l. c.; A. Rich. l. c. p. 263. Herbacea, caule 6-10 decim. alto, erecto, subsimplice, striato-sulcato, subsucculento, hispidulo-pubescente, basi glabrato, stimulis validis consperso vel subinermi. Folia magnitudine maxime variabili, 6-25 centim. longa, 4-25 centim. lata, raro subæqualiter 3-lobata (lobis plus minus profundius acuminatis extus interdum lobulo parvo instructis), sæpius regulariter ovata, basi subcordata et apicem versus sensim acuminata, acumine lanceolato, nonnulla etiam in eodem caule hinc vel inde sublobata, serraturis æqualibus inæqualibusve acutis vel subacuminatis, petiolo adpresse pubescente stimulisque validis (in planta exsiccata basi rufescentibus) armato. Stipulæ 5-10 millim. longæ, extus sparsim pubescenti-hispidæ, apice bifidæ, mox deciduæ. Flores monoici vel dioici, inflorescentiis brevissime pedunculatis solitariis geminisve. Racemi masc. simplices vel bifurcati, petiolo plerumque breviores, floribus perigonio hispido. Cymæ fem. in iisdem axillis cum racemis masculis interdum commixtæ, petiolis brevioribus, in specimenibus quibusdam vix centimetrales, ramulis brevibus et pedicellis hispidulis stimulisque albidis dense obsitis. Perigonii segmentum superius oblongum, hispidum, dentibus apicalibus minimis vix distinctis, inferius nullum. Achænium ovatum vel subcordatum, 2½-3 millim. longum, stylo persistente paulo longius. (v. s.)

7. *G. CUSPIDATA*, foliis late ellipticis basi sæpius rotundatis apice abruptissime longissimeque cuspidatis grosse acuteque inciso-dentatis dentibus superioribus cæteris majoribus utrinque hispido-pubescentibus pilosulisve sparsimque stimulosi, petiolo limbum dimidium æquante, stipulis vix conspicuis, inflorescentiis masc. laxis paniculiformibus, fem. graciliformibus spiciformibus. ② Ad rivulos montis Occ. prope Pekin, in imperio sinensi (R. P. David). — Herbacea, caule circiter metrali subsimplice inferne crassitie pennæ cycnæ et plus minus angulato-sulcato hispidulo et sparsim stimuloso, internodiis elongatis. Folia 9-15 centim. longa, 6-10 centim. lata, basi obtusa v. rotundata v. subtruncata, cuspidate lineari-lanceolata 3-5 centim. longa basi sæpe angustata e fundo sinus obtusi inter dentes duos apicales magis elongatos limbi orta, dentibus cæteris ab apice ad basim limbi gradatim magnitudinæ diminutis inferioribusque sæpe minimis, petiolo gracili puberulo stimulisque sparsim obsito. Stipulæ triangulares bifidæ? Flores monoici. Inflorescentiæ masc. inferiores, solitariae aut geminae (altera minore), pedunculatæ, folio nonnihil breviores, ramulis laxis erecto-patulis hispidulis et parce stimulosi, glomerulis paucifloris, perigonio quinque-partito hispidulo. Spicæ femineæ ex axillis superioribus ortæ, initio filiformes foliumque superantes, glomerulis discretis, rhachi juxta glomerulos præsertim stimulosa; perigonii segmentis circiter æquimagnis, superiore elliptico apice obsolete bidentato dorso carinato crasseque pilosulo, inferiore ovato subacuminato etiamque dorso pilosulo. Ovarium ellipticum, perigonium superans et utrinque inter hujus segmenta crassiuscule pilosulum, stylo complanato altero latere papilloso subæquilongum. — Species insignis a congeneribus omnibus valde discrepans. (v. s.)

*Species exclusa.*

*G. FURIALIS* BL. = *Obetia ficifolia*.

Trib. II. *PROCRIDEÆ* Wedd. *Monogr.* p. 170. — *Elatostemæ* Gaudich. *Bot. Voy. Uran.* p. 496.

Stirpes fere omnes herbaceæ, paucissimæ suffrutescentes, inermes. Folia decussatim opposita v. abortu alterna et tunc sæpe disticha, in singulis jugis s. paribus homomorpha ac plerumque inæquimagna v. symmetrica heteromorpha et inæquilatera, cystolithis frequentissime linearibus in planta exsiccata pilos adnatos mire mentientibus. Stipulæ caulinae utriusque folii in unam axillarem integerrimam binervem coadunatæ. Flores unisexuales aut rarissime (in Achudemia) polygami. *Masc.* perigonio sæpius 4-partito, segmentis sub apice sæpissime mucronatis in præfloratione imbricatis (saltem apice), alabastro sæpe mucronato; pistilli rudimento conico. *Fem.* perigonio libero, 3 vel rarissime 5-partito, segmentis (interdum minimis) subæqualibus planiusculisque aut altero rariusve omnibus gibbo v. mucrone dorsali aut subapicali instructis post anthesin ut plurimum parum mutatis, stigmate penicillato, staminum rudimentis squamiformibus. Fructus siccus v. rarissime carnosulus, glaber. Embryo cotyledonibus ovatis orbicularibusve, radícula conica crassa brevi, albumine parco v. subnullo. *Hermaphr.* perigonio et pistillo ut in flor. fem., staminibus ut in flor. masc.

XI. *PILEA* Lindl. *Collectan. t.* 4; *Endlich. Gen. Pl.* p. 284; *Wedd. Monogr.* p. 172. t. 6. C. 7 et 8. — *Dubruleilia* Gaudich. *Bot. Voy. Uran.* p. 495. — *Urticæ* et *Parietaræ* spec. auct.

Flores monoici (inflorescentiis androgynis unisexualibusve) aut dioici glomerulati, glomerulis confertim v. laxè paniculato-cymosis rarius solitariis v. geminis, pedicellis florum masc. articulatis. *Masc.* Perigonium 4-partitum aut rarissime 2-v. 3-partitum, segmentis concavis carnosulis sæpissime albidis v. roseis sub apice membranaceo vulgo crasse mucronatis, mucronibus in alabastro conniventibus discretisve. Stamina tot quot segmenta perigonii. Pistilli rudimentum conicum. *Fem.* Perigonium 3-partitum, segmentis plerumque inæqualibus carnosulis virentibus rubescentibusve, lateralibus sæpius minoribus planiusculis, intermedio ut plurimum dorso gibboso seu cucullato. Staminum rudimenta squamiformia, inflexa, segmentis perigonii opposita. Ovarium rectum, compressiusculum. Ovulum in fundo loculi suberectum, funiculo obliquo gracili. Stigma breviter penicillatum, sessile. Achænium suboblique ovatum rotundatumve, compressum, læve aut scabriusculum, subnudum v. perigonio parum aucto incomplete vestitum. Semen pericarpio conforme. Embryo fere semper exalbuminosus aut subexalbuminosus, cotyledonibus ovatis rotundatisve radícula conica longioribus. — Herbæ annuæ v. perennes (rariusve suffrutices), pleræque tropicæ et subtropicæ, neogæ et gerontogæ, erectæ v. humi prostratæ et tunc sæpe radicanter. Folia opposita, ut plurimum petiolata, in eodem jugo s. pare sæpissime inæquimagna et interdum heteromorpha, raro æquilateralia, integerrima aut varie dentata, vulgo trinervia v. triplinervia, raro penninervia, limbo nonnunquam carnosulo nervisque omnino immersis, glabra aut varie pilosa cystolithisque fusifor-

mibus linearibusve v. punctiformibus v. etiam stellatis ut aliæ partes plantæ sæpius farcta. Stipulæ utrinque in unam intra-axillarem integram deciduam v. persistentem connatæ. Cymæ s. paniculæ sæpe distiche ramosæ. Flores bracteis parvis perigonio brevioribus (rarissime majusculis) muniti, masculi mox decidui. — Genus admodum naturale, perigonio femineo tripartito et irregulari facillime distinguendum.

*Conspectus distributionis specierum.*

**INTEGRIFOLIÆ**

Folia penninervia vel subenervia. . . . .	Confer ad sp.	1
Folia distincte trinervia vel triplinervia.		
Gerontogæ. . . . .		10
Neogæ. . . . .		13

**HETEROPHYLLÆ**, sive foliis in eodem jugo diversiformibus aut valde inæquimagnis.

Gerontogæ. . . . .	30
Neogæ. . . . .	33

**DENTATÆ**, sive foliis nec integris nec dimorphis nec in eodem jugo valde inæquimagnis.

**Gerontogæ.**

Glabræ, sive caule foliisque pilis omnino destitutis.

Longipedunculatæ, sive pedunculis (saltem cymarum fructiferarum) quam petioli plerumque longioribus.	
Stipulæ breves, triangulares, persistentes . . . . .	49
Stipulæ majusculæ, ovatæ vel lanceolatæ. . . . .	55

Brevipedunculatæ, sive pedunculis (saltem inflorescentiarum fructiferarum) quam petioli sæpius brevioribus.

Stipulæ breves, triangulares. . . . .	57
Stipulæ majusculæ, ovatæ, vel oblongæ. . . . .	67

Pubescentes, sive caule foliisque puberulis vel pubescentibus vel pilosis.

Longipedunculatæ, sive pedunculis (saltem cymarum fructiferarum) quam petioli plerumque longioribus. . . . .	75
Brevipedunculatæ. . . . .	79

**Neogæ.**

Glabræ.

Longipedunculatæ.	
Stipulæ breves, triangulares. . . . .	84
Stipulæ majusculæ, ovatæ oblongæ vel rotundatæ.	
Nervi basilares folii apicem limbi haud attingentes. . . . .	88
Nervi basilares usque ad apicem limbi producti. . . . .	98
Brevipedunculatæ. . . . .	102

Pubescentes, sive caule foliisque puberulis pubescentibus vel pilosis.

Longipedunculatæ. . . . .	117
Brevipedunculatæ. . . . .	144

§ 1. — *Integrifoliæ.*

A. — *Folia penninervia vel subenervia* (1).

1. P. MICROPHYLLA (Liebm. Vidensk. Selsk. Skr. 5 Række, naturv Og math. Afd. 2. p. 296), vulgo monoica, herbacea, glaberrima, caulibus ramosissimis diffusis, foliis parvis ellipticis oblongisve vel obovatis rarius ovatis ut plurimum obtusis basi sæpe cuneato-attenuatis integerrimis penninerviis vel sub-

(1) Vid. quoque P. subcrenata inter *Dentatas*.

enerviis, cymulis sæpius e glomerulo unico constantibus sessilibus pedunculatisve folio superatis androgynis aut unisexualibus. ①-② Fereper omniæm Americam tropicam diffuse loca humida et umbrosa præsertim incolens, usque ad 2000 metr. supra Oceani ripas obvia: in imperio mexicano, ins. Antillis, Nova Granata, Æquatore, Peruvia, Guyana, Brasilia, etc. — Griseb. Fl. Br. W. Ind. p. 155. *P. muscosa* Lindl. Collectan. t. 4; Wedd. Monogr. p. 174. *P. muscosa*, *P. callitrichioides* et *P. microphylla* Schlecht. in Linnæa, 27. p. 494 et seq. *Parietaria microphylla* Linn. Sp. pl. p. 1492; Poir. Encycl. 5. p. 16. *Urtica microphylla* Swartz, in Act. Holm (1787) p. 61; ejusd. Fl. Ind. occid. p. 305. *U. callitrichioides* Kunth in H. B. Nov. gen. et sp. 2. p. 40. *Dubreuilia microphylla* Gaudich. Uran. p. 495. Caules debiles filiformes aut plus minus succulenti, longitudine varia, ramique valde foliosi. Folia plerumque in quoque pare admodum inæquimagna,  $1\frac{1}{2}$ –5 millim. longa, nervis subimmersis et vix aut non conspicuis, exsiccata membranacea et nonnunquam pellucida, pagina superiore cystolithis immersis linearibus transversim lineolata, inferiore his corpusculis nisi in nervo medio destituta, petiolo brevissimo vel 1–5 millim. longo pulvino prominente sæpe suffulso. Stipulæ plerumque omnino inconspicue et fortassis interdum omnino deficientes. Cymæ minimæ fere ex omnibus axillis nascentes, solitarie aut geminæ, sæpissime sessiles (pedunculo quum extat quam folium semper brevior), androgynæ aut unisexuales, floribus vulgo breviter pedicellatis masculis paucioribus, perigonii segmentis breviter et crasse mucronatis, alabastro 1 millim. circiter longo, perigonii fructiferi segmento intermedio cucullato lateralibus vix longiore achænioque ovato semimillimetrali scarbiusculo paulo brevior. (v. s. et v. cult.)

β, *herniarioides*, caulibus filiformibus parum ramosis, foliis obovato-vel rotundato-spathulatis. ① In iisdem locis ac typus. — *Pilea herniarioides* Lindl. l. c.; Schlecht. l. c. p. 488. *P. peperomiæfolia* Liebm. l. c. p. 297; Wedd. Monogr. p. 179. *P. tenerrima* Miq. in Linnæa ann. 1853. p. 219. *P. herniarifolia* Bl. Mus. lugd. bot. 2. p. 44. *Urtica herniarioides* Swartz Act. Holm. (1787) p. 65. t. 2. f. 1; ejusd. Fl. Ind. occid. p. 309; Poir. Encycl. suppl. 4. p. 220. *U. herniarifolia* Willd. Sp. pl. 1. 359. — Caules interdum breves et erectiusculi, at sæpius prostrati vel repentes. Folia 5–10 millim. longa sæpiusque in petiolum gracilem cuneato-attenuata, exsiccata tenuiter membranacea. Cymulæ fere semper sessiles. (v. s.)

γ, *longifolia*, caulibus vix elongatis erectiusculis ramosissimis, foliis (cum petiolo gracili sæpe sesquicentimetralibus) oblongo-spathulatis, lamina exsiccata rigidula subtus sicut caulis et rami glaucescente, cymulis plerisque unisexualibus, fem. sessilibus vel breviter pedunculatis, masculis pedunculatis petiolum plerumque superantibus. ① In imperio mexicano (Sumichrast). — Habitu et cymulis vulgo longius pedunculatis. *P. serpyllaceam* nonnihil refert. (v. s. in hb. DC.)

δ, *portulacoides*, caulibus prælongis prostratis divaricato-ramosis crassis, foliis majusculis obovatis longiuscule petiolatis. ①? In Brasilia meridionali (Aug. St-Hilaire) et passim cum typo? — *Pilea Portula* Liebm.? l. c. 297. *P. portulacina* Bl.? l. c. *P. succulenta* Hook. fil.? Trans. linn. soc. 20. p. 152. *Urtica portulacina* Spr.? Syst. 3. p. 838. — Caules robustiores ramique et folia magis succulenti quam apud typum. Folia in speciminibus brasiliensibus 10–15 millim. longa, 4–6 millim. lata, in petiolum 2–3-millimetrali cuneato-attenuata. — Planta fortassis ulterius specificè distinguenda; a typo habitu sat diversa. (v. s. in hb. Mus. par.)

2. *P. TRIANTHEMOIDES* (Lindl. Collectan. t. 4), dioica, glaberrima, demum suffrutescens, foliis parvis obovatis breviter petiolatis integerimis carnosis enerviis, cymulis unisexualibus folio brevioribus sessilibus aut breviter pedunculatis. ②? In collibus rupibusque secus rivulos insulæ S. Domingo (Swartz, L. Cl. Richard, etc.), in Jamaica (Sloane), in ins. Cubi (Linden, n. 1951, Wright, n. 538, pro parte). — Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 207; Schlecht. l. c.; Bl. l. c. p. 44. *P. muscosa* γ trianthemoides Wedd. Monogr. p. 175; Griseb. Fl. Br. W. Ind. p. 155. *Herniaria lucida* aquatica Sloane, Voy. 1. t. 93.



- f. 2. *Urtica trianthemoides* Swartz, Act. Holm. ann. 1787. p. 68; ejusd. Fl. Ind. occid. p. 307; Smith, in Rees' Cyclop. 36, Urt. n. 45; Poir. Encycl. suppl. 4. p. 220. — Caulis admodum succulentus demumque lignescens, vulgo prostratus seu declinans, 3-6 decim. longus, cylindricus, viridissimus, ramis plerisque alternis distiche plano-patentibus ad nodos inflatis. Folia 5-15 millim. lata, subspathulata, obtusissima, petiolo 1-3-millimetrali suffulta, convexa, in vivo lucida, exsiccata autem opaca subtusque lacunosa sive rugosa et supra cystolithis parvis numerosissimisque oblique striolata. Stipulæ inconspicuæ. Cymæ nunquam androgynæ, cæterum floresque et achænia ut in *P. muscosa*. (v. s.)
- β, *microphylla*, caule foliisque minoribus quam in typo. — In ins. Cuba (Wright, n. 2230). *P. microphylla* Griseb. Cat. pl. cub. p. 59, pro parte. (v. s.)
3. *P. SERPYLLIFOLIA*, dioica, plerumque herbacea, glaberrima, foliis parvis obovato-orbiculatis cuneatisve obtusissimis integerrimis penninerviis, cymulis pedunculatis folia vulgo superantibus capitelliformibus. (1) In sylvis mexicanis, usque ad altitudinem 1500 metr. prope Xalapa (Schiele; Galeotti, n. 7084); in Nova Granata (Humb. et Bonpl.), prov. Pasto, alt. 2500 metr. (Triana); in Costa-Rica (Orsted). *P. serpyllacea* Hook. et Arn. Bot. Voy. Beech. p. 96; Wedd. Monogr. p. 177. t. 6. c. f. 10. *P. muscosa* hort. ex part. *P. serpyllifolia* hort. *P. callitrichioides* Kunth, Ind. sem. bot. berol. (1846) p. 12. *P. peplidifolia* Schlecht. in Linnæa, 27. p. 496. *Parietaria serpyllifolia* Poir. Encycl. 5. p. 16. *Urtica serpyllacea*. Kunth in H. B. Nov. gen. et sp. 2. p. 37; Cham, et Schlecht. in Linnæa, 6. p. 356. Caulis herbaceus, succulentus aut demum basi lignescens, erectus, 2-4 decim. altus, ramosissimus, valde foliosus. Folia 3-8 millim. longa, 2½-4 millim. lata, crassiuscula, nervis immersis, lamina exsiccata membranacea, pagina superiore cystolithis numerosissimis linearibus transversim lineolata, inferiore iisdem nisi in costa et margine destituta, petiolo 1-4 millim. longo. Stipulæ inconspicuæ (fortassis nullæ). Cymæ singulis axillis solitariae vel geminae, e glomerulo unico seminis cannabini circiter magnitudine constantes, pedicello filiformi 1½-10 millim. longo suffultæ, floribus masc. pedicellatis, albo-roseis, lobis perigonii breviter et crasse mucronatis, alabastro 1 millim. circiter longo; fem. subsessilibus, perigonii fructiferi segmento intermedio cucullato laciniis lateralibus achænioque ovato semimillimetrali lævi vel scabriusculo æquilongo. (v. s. et v. cult.)
4. *P. GLOBOSA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 208; Monogr. p. 178), monoica?, glaberrima, habitu suffruticuloso, foliis parvis admodum succulentis gibbis sæpeque subglobosis integerrimis enerviis, cymulis unisexuilibus capitelliformibus pedunculatis folia sæpius superantibus. ♀? In cultis, prope urbem Pasto Novo-Granatensium et villam Teindala, alt. 2600 metr. (Humb. et Bonpl.); in Peruvia (Pavon). — *P. thymifolia* Bl. Mus. bot. iugd. bat. 2. p. 44. *Urtica globosa* Pavon, mscr. *U. dasyphylla* Poepp. mscr. *U. thymifolia* Kunth in H. B. Nov. gen. et sp. 2. p. 37 (nomen omnino improprium). A basi ramosissima, ramis diffusis inferne denudatis carnosius, ramulis valde foliosis. Folia 2-3 millim. vix lata, exsiccata rigide membranacea margine sæpe undulata, pagina superiore cystolithis crassiusculis fusiformibus creberrime ornata, inferiore rugoso-lacunosa iisdem corpusculis prorsus destituta, petiolo uni-sesqui-millimetrali. Cæt. ut in *P. serpyllacea*, sed cymulæ paulo minores et haud raro brevius pedunculatæ. — Planta dicitur a Kunthio esse monoica, sed specimina a me visa sunt omnia unisexuala. (v. s.)
5. *P. PENDULA* (Griseb. Cat. pl. cub. p. 59, non Blume), dioica, herbacea, glaberrima, foliis rhomboideo-ovatis acute acutiusculis basi acutis obtusisve integerrimis subuninerviis (cæteris immersis), cymis pedunculatis foliaque superantibus glomerulis laxè racemoso-paniculatis. ♀ In ins. Cuba (Wright, n. 2232). — Caulis ramique elongati, graciles, e rupibus arboribusque ut videtur penduli, in specim. suppet. 3-4 decim. longi, exsiccati striati pallide virentes cystolithisque oblongis creberrime variegati. Folia pleraque reflexa, in singulis jugis altero plerumque minore breviusque petiolato, 6-12 millim. longa, 3-8 millim. lata, penninervia, lamina exsiccata rigida pallida margi-

nata, pagina superiore cystolithis linearibus numerosissimis eximieque intricatis ornata, inferiore iisdem munita sed minus conspicuis et longitudinaliter ordinatis, petiolo folii majoris 3 millim. circiter longo. Stipulæ breves, triangulares. Cymæ ex axillis ultimis ramorum nascentes, paucæ, solitariae; masc. non visæ; fem. pedunculo gracillimo 15-20 centim. longo suffultæ, parce ramosæ, glomerulis minimis discretis, floribus sessilibus, perigonii segmento intermedio oblongo breviter cucullato lateralibus longiore; achænio.... (v. s.)

6. P. FRANQUEVILLEANA, fruticulosa, foliis parvis oblongo-vel subpathulato-lanceolatis obtusis inferne in petiolum brevem attenuatis penninerviis margine sparsim breviterque spinuloso-ciliatis cæterum glabris, cymis minimis glomeruliformibus androgynis aut abortu unisexualibus subsessilibus petiolos vix superantibus. ♂ In ins. S. Domingo (Cl. Richard). Caulis elongatus, prostratus et sæpe radicans, demum adscendens, ramosissimus, ramis valde foliosis. Folia 5-8 millim. longa, 1½-2 millim. lata, nervis lateralibus parum conspicuis lamina exsiccata rigidula (lucis transmissa) hyalinis; cystolithis in eadem vix aut non distinctis linearibus; petiolo 1-2 millim. longo. Stipulæ breves, triangulares, persistentes. Cymulæ numerosæ aut breviter pedunculatæ, e glomerulo unico sæpius constantes; floribus masc. paucissimis breviter pedicellatis, perigonii lobis sub apice mucrone brevi instructis; floribus fem. sessilibus, perigonii segmentis subæqualibus. — Species distinctissima, habitu ab omnibus congeneribus (sequente excepto) recedens. (v. s. in hb. Franqueville.)
7. P. GNIDIODES (Griseb. Pl. Wr. in Mem. am. Acad. sc. et art. nov. ser. 8. 1860. p. 173; Cat. pl. cub. p. 59), dioica, fruticulosa, glabra, foliis crebris parvis lanceolatis utrinque acutis integris subuninerviis, cymulis pedunculatis e glomerulis paucis confectis folium superantibus. ♂ In regione orientali ins. Cubæ, prope villam Monte Verde (Wright, n. 1451). — Caules in specim. visis 10-15 centim. alti, castitiosi, graciles, lignosi, brunnei, basi denudati, superne parce ramosi denseque foliosi, glabri vel lente hispiduli, internodiis brevissimis. Folia 4-8 millim. longa, 1-1½ millim. lata, in singulis jugis subæquimagnis vel altero paulo minore, patula, breviter petiolata, lamina exsiccata rigida fusco-viridi subtus pallidiore margine incrassato subrevoluta, pagina superiore cystolithis linearibus elongatis oblique ordinatis alifisque punctiformibus adamussim interjectis eximie ornata, pagina inferiore his corpusculis etiam crebre notata sed homomorphis et longitudinaliter dispositis, petiolo subtilissime hispidulo. Stipulæ brevissimæ, triangulares. Cymæ fem. (masculæ haud visæ) in singulis axillis solitariae, prope apicem ramorum nascentes, simplices vel parce ramosæ, glomerulis parvis discretis, pedunculo tenuissimo 5-10 millim. longo, floribus breviter pedicellatis, perigonii fructiferi segmento intermedio oblongo breviter cucullato lateralibus dimidio longiori achænio semimillimetrali scabriusculo triente brevior. (v. s.)
8. P. DELTOIDEA (Liebm. l. c. p. 298), dioica vel monoica, herbacea, pusillima, caulibus tenuissimis humifusis, foliis exiguis rotundato-deltoides apice rotundatis juxta petiolum cuneatis integerrimis subnerviis supra sparsim pilosis, margine longiuscule distanter ciliatis, floribus in axillis superioribus vel inter folia terminalia paucis glomerulatis, perigonio femineo subnullo. ♂ In Costa-Rica (Osted), necnon prope villam Monte Verde dictam Cubæ orientalis (Wright, n. 1458). — Wedd. Monogr. p. 178. P. microphylla, var. perexigua Griseb. Pl. Wr. in Mem. am. Acad. sc. et art. nov. ser. 8 (1860) p. 173; Cat. pl. cub. p. 59. Caules 1-3 centim. longi, radicanes, pulvinato intricati, glaberrimi, subhyalini, apice præsertim foliosi, indivisi vel ramosi. Folia 1-2 millim. longa lataque, terminalia rosulata, nervis omnino inconspicuis, exsiccata membranacea obscure viridia subtus lacunoso-cellulosa, supra cystolithis linearibus obliquis transversalibusque ornata, petiolo quam limbus dimidius breviori glabro. Stipulæ breviter ovatæ, hyalinæ. Cymulæ minimæ, sessiles sive e floribus paucis sessilibus pedicellatisve et subfasciculatis confectæ, in rosula terminali præsertim obviæ foliisque breviores, unisexuales et æpissime in distinctis plantis natæ, perigonio masc. glabro aut sparsissime piloso segmentis sub apice mucrone conico breviusculo instructis, alabastro ½ millim. longo, fem. minimo sive e segmentis triangularibus subæqualibus quam

staminum rudimenta brevioribus constante. Achænium ovatum, semimillimetrale, oculo armato subtilissime granulato. — A speciebus præcedentibus optime distincta. Perigonium femineum rudimentarium, illum floris fem. Elastematis referens, in nullo congenere notatur. (v. s.)

β, *Domingensis*, foliis triplo majoribus quam in typo, sub lente distincte trinerviis, pagina superiore pilis longis diaphanis conspersa. — In ins. S. Domingæ (L. Cl. Richard). Caulis 3-6 centim. longus, merithallis inferioribus valde elongatis. Folia 3-6 millim. lata, nervis basilariibus tenuissimis dimidium limbi vix superantibus, lamina exsiccata subhyalina. Staminodia fl. fem. achænio delapso valde accreta lineari-oblonga inflexa et in apice pedicelli elongati segmenta perigonii majuscula mentientia. (v. s.)

9. *P. SARCOPHYLLA*, dioica, herbacea, glaberrima, caule crasso erecto, foliis succulentis, inferioribus ellipticis, superioribus (majoribus) oblongo-lanceolatis sensim acuminatis utrinque acutis integerrimis vel (in sicco) vage muricatis tripli-penninerviis nervis etiam in planta exsiccata fere omnino immersis, cymis masc. laxiusculis longe pedunculatis. ½? In sylv. montosis saxosisque ins. Antigua (Wulschlägel). — Caulis 2-4 decim. altus, angulato-sulcatus, internodiis 2-4-centimetralibus. Folia inferiora circiter 2 centim. longa, superiora (magis approximata) 5 centim. longa, 1½-2½ centim. lata, breviter petiolata, lamina exsiccata in dimidia parte superiore undulato-crenata nervis oculo armato etiam ægre perspicuis, cystolithis nullis aut minimis punctiformibus. Stipulæ.... Cymæ masc. (solum visæ) ex axillis superioribus natæ, floribus pedicellatis, perigonii segmentis sub apice longiuscule mucronatis. (v. s. in hb. monac.)

B. — *Folia distincte trinervia vel triplinervia.*

1° Gerontogææ.

10. *P. PEPOIDES* (Hook. et Arn. Bot. Voy. Beech. p. 96), monoica, glaberrima, foliis parvis rotundato-rhombeis utrinque obtusatis integerrimis trinerviis, cymis sessilibus quam petioli brevioribus, floribus masc. et fem. in iisdem glomerulis commixtis. ① In ins. Sandwich (Beechey, Gaudichaud, etc.), Gallapagos (Hook. fil.) et Java (Zollinger n. 2961). — Wedd. Monogr. p. 179; Miq. Fl. Ind. bat. 1 (2). p. 237. *P. pygmæa* ejusd. in Zoll. Syst. Verzeichn. p. 106. *Dubruëllia peploides* Gaudich. Uran. p. 495. — Caulis herbaceus, a basi ramosus, ramis brevibus vel elongatis gracillimis et prostratis subdecimetralibus, merithallis brevibus, nodis plerisque floriferis. Folia 3-6 millim. longa et lata, nervis lateralibus parum conspicuis dimidium limbum paulo superantibus, lamina exsiccata membranacea pagina superiore cystolithis linearibus inerspersa, inferiore iisdem destituta sed haud raro fusco-punctata, petiolo gracili quam limbus ut plurimum longiore. Stipulæ inconspicuæ. Cymæ parvæ, corymbiformes, multifloræ, floribus breviter pedicellatis, masculis paucis, lobis breviter mucronatis, alabastro ¼ millim. longo, femineis numerosissimis, perigonii fructiferi segmento intermedio anguste oblongo lateralibus triplo longiore achænioque late ovato semimillimetrali lævi paulo brevioribus.

β, *major*, caule decumbente vel adscendente succulento, foliis multo majoribus quam in typo (7-13 millim. latis) in dimidia parte superiore crenatodentatis, nervis conspicuis. — In ins. Sandwichensibus Oahu et Maui (Remy); Birma Ind. or. (Griffith).

11. *P. ARTICULATA* (Wedd. Monogr. p. 180), dioica, glaberrima, foliis ovato-lanceolatis acuminatis basi rotundatis integerrimis trinerviis, cymis masculis geminis capituliformibus, pedunculis petiolum vix superantibus. ½ In ins. Mauritio (Commerson, Du Petit-Thouars). — U. *Parietaria* β 1 Poir. Encycl. 4. 639. *U. articulata* Herb. Juss. — Caulis erectus, 4 dm. altus, simplex, crassitie circiter pennæ anserinæ, basi longe nudus et cylindricus, merithallis superioribus brevibus, cæteris magis minusve elongatis, epidermide exsiccata albicante cystolithisque numerosissimis nonnunquam punctulata ad nodos

- β, *latifolia*, foliis oblongo-lanceolatis 12-15 millim. latis subtriplinerviis (petiolo 5-15 millim. longo) exsiccatis concoloribus. — In ins. Cuba (Wright, n. 2233). (v. s.)
16. P. *VIRGATA*, dioica, glaberrima, foliis oblongo-lanceolatis acuminatis acutis ad insertionem petioli emarginatis integerrimis tri-triplinerviis, nervis basilaribus a margine limbi subdistantibus, stipulis brevissimis truncatis, cymis parvis subsessilibus petiolo brevioribus, achænio fere rotundato. ♂ ? In ins. Jamaica (Alexander). — Caulis (s. ramus) in specimine unico viso gracilis, virgatus, basi lignosus, circiter 4-decimetralis, simplex, epidermide exsiccato obscuro et undique cystolithis punctulato, internodiis circiter bicentimetralibus subæqualibusque. Folia in singulis nodis nonnihil inæquimagna, 4-6 centim. longa, 14-18 millim. lata, in  $\frac{1}{2}$  superiore acuminata, nervis basilaribus ad basim acuminis in margine evanidis, secundariis subvis bene conspicuis, lamina exsiccata subcoriacea obscura cystolithis punctiformibus minimis utrinque crebre obsita, pagina inferiore nonnihil pallidior. Petioli in utroque nodo valde inæquilongi, majore internodium circiter æquante. Stipulæ persistentes sed, ob exiguitatem, vix discernendæ. Cymæ femineæ (solum visæ) petiolo minore sæpius breviores, vix multifloræ, perigonii fructiferi segmento intermedio late oblongo apice truncato vel subtrilobo lobo intermedio sive cucullo gibbosulo, segmentis lateralibus orbiculari-ovatis intermedio breviori dimidiumque achæni parum superantibus, achænio millimetrali lævi maturo marmorato. (v. s. in hb. Kew.)
17. P. *STELLULIGERA* (Wedd. Monogr. p. 184. t. 6. f. 9), monoica, glaberrima, foliis rotundato-rhomboides utrinque obtusatis integerrimis trinerviis, stipulis brevibus ovatis, cymis masculis longe pedunculatis folia superantibus. ♀ ? In insula S. Domingo (Nectoux). — P. rhombea Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 17. p. 211, non Liebm. — Caulis herbaceus, in specimine unico suppetente 4 decim. longus, suberectus, breviter ramosus, merithallii elongatis, nodis inferioribus denudatis, siccitate albicans cystolithisque minimis obsolete punctulatus. Folia 10-20 millim. longa lataque, nervis basilaribus valde arcuatis apicem limbi fere pertingentibus, lateralibus intra arcum conspicuis, lamina exsiccata firmula; cystolithis in pagina superiore (extremo margine excepto) nullis, in pagina inferiore vero numerosissimis, stellato-ramosis; petiolo gracili, 4-8 millim. longo. Stipulæ parvæ. Cymæ masc. e glomerulis paucis constantes, pedunculis filiformibus limbo folii longioribus suffultæ, floribus sessilibus, perigonii lobis breviter mucronatis, alabastris 1 millim. circiter longis. Cymæ fem. non visæ. — P. rhombea refert; hujusce tamen folia acuta sunt et ciliata cystolithique admodum diversi. (v. s. in hb. Mus. par.)
18. P. *CONSANGUINEA*, glaberrima, foliis late rhomboideo-ovatis obtusis basi abrupte cuneatis integerrimis trinerviis, stipulis minimis truncatis, cymis.... ♀ In ins. S. Domingo (Bertero). — Caulis gracilis, elongatus, repens, ramis demum adscendentibus apice foliosis. Folia subcentimetralia, 8 millim. circiter lata, nervis basilaribus ut in specie præcedente, limbo ima basi seu in ipsa insertione petioli subemarginato, lamina exsiccata rigide membranacea supra cystolithis minimis linearibus creberrime obsita, margine ipso iisdem numerosioribus incrassato, pagina inferiore purpurascente cystolithisque destituta, petiolo circiter limbi dimidii longitudine. Cæt. ignota. — Præcedenti habitu arcte affinis, sed foliis basi abrupte cuneatis et forma cystolithorum facillime distinguenda. (v. s. in hb. Mus. par.)
19. P. *RHOMBEA* (Liebm. l. c. 305), monoica, herbacea, diffuse ramosa, foliis elliptico-rhomboides utrinque acutiusculis integerrimis subtriplinerviis supra glaberrimis nervis paginæ inferioris margineque sparsissime ciliatis, stipulis brevibus ovatis, cymis parvulis androgynis vel unisexualibus, pedunculis petiolo brevioribus. ♀ ? In Nova Granata (Mutis). — Wedd. Monogr. p. 191. *Urtica rhombea* Linn. fl. Suppl. p. 417; Willd. Sp. pl. 4. p. 357; Smith in Rees Cyclop. *Urtica*, n. 36. — Caulis striatus, ramis gracillimis, internodiis elongatis. Folia 3 centim. circiter longa, 1½-1 centim. lata, nervis basilaribus conspicuis fere ad apicem limbi usque pertingentibus, lamina exsiccata teniter membranacea, pagina superiore cystolithis linearibus crebre ornata,

inferiore iisdem prorsus destituta, petiolo gracili limbo æquilongo vel brevior. Stipulæ 1-2-millimetricæ. Cymæ e glomerulis 1-4 paucifloris approximatis subsessilibusque constantes, florum masc. lobis longiuscule acuminato-mucronatis, alabastro fere millimetrici, fem. perigonii segmento intermedio breviter cucullato lateralibus longiore achænioque fere æquilongo. — Forma foliorum et nervatione necnon brevitate pedunculorum et cæt. a sequente bene distincta.

20. *P. WRIGHTIANA*, monoica, caule parce ramoso, basi radicante, foliis late ovato-vel obovato-rotundatis utrinque rotundatis obtusisve integerrimis trinerviis margine petioloque longiusculo sparsim pilosulis sive ciliolatis cæterum glabris, stipulis brevibus late rotundatis, cymis parvis (e glomerulo unico vel paucis constantibus) unisexualibus, pedunculis quam petioli longioribus, segmentis perigonii fl. masc. longe mucronatis. ♀ In ins. Cuba (Wright, n. 2233, ex parte). — *P. stelluligera* Griseb. Cat. pl. cub. p. 59, non Wedd. Caulis herbaceus, 3-5 decim. altus, adscendens, sat robustus, internodiis elongatis, exsiccatò albicante, ramis paucis brevibus gracilibusque. Folia 1½-3 centim. longa et lata, nervis basilaribus dimidium limbi parum superantibus nec ad apicem ejusdem protractis, lamina exsiccata membranacea cystolithis parvis fusiformibus in pagina superiore crebre munita, inferiore iisdem corpusculis linearibus sub lente etiam crebrius ornata, petiolo gracili limbum æquante vel paulo brevior. Stipulæ bi-tri-millimetricæ, obtusissimæ, membranaceæ, glabræ, Cymæ 5-10 centim. latæ, pedunculis gracilibus 2-4-centimetricis glabris, masc. e glomerulo unico seu fasciculo confectis, floribus longiuscule pedicellatis, segmentis perigonii 2 millim. longis, mucrone subapicali cylindrico segmentum ipsum fere æquante et plus minus hispido cymis fem. parce ramosis, glomerulis parvis subdiscretis, perigonii fructiferi segmento intermedio oblongo breviter cucullato lateralibus dimidio longiore achænioque ovato haud millimetrici læviusculo aut circinnatim minute granuloso parum brevior. (v. s.)

β, *microphylla*, pusilla, monoica v. dioica, caulibus filiformibus repentibus, ramis adscendentibus glabris puberulisve, foliis parvis basi sæpe subcuneatis margine subciliatis, petiolo et nervis puberulis, cymis utriusque sexus e glomerulo unico constantibus, perigonii masc. segmentis quam in typo brevius mucronatis alabastro millimetrici. ♀ Prope Monte Verde, in regione orientali ins. Cubæ (Wright, n. 1456 a). — *P. microphylla*, var. *herniarioides* Griseb. Pl. Wr. in Mem. Acad. Am. sc. et art. nov. ser. 8 (1860) p. 173. — Folia 3-5 millim. longa et lata, petiolo quam limbus sæpe longiore, cystolithis paginæ superioris linearibus ad marginem crebrioribus.

γ, *intermedia* quoad staturam et habitum inter var. β et typum *intermedia*, monoica vel dioica, inflorescentiis ut in var. præcedente, segmentis perigonii masc. breviter mucronatis. ♀ In Cuba orientali (Wright, n. 1680 et n. 2238). — *P. rotundata*, var. foliis subintegerrimis, Griseb. l. c. et Cat. pl. cub. p. 60. — Folia pleraque in speciminibus visis 1 centim. longa et lata, cystolithis ut in var. β, alabastra florum masc. 1-1½ millim. longa, mucronibus conicis glabris dimidiam partem segmenti vix æquantibus. Achænia late ovata, ¼ millim. longa.

21. *P. CEPHALANTHA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 209; Monogr. p. 191), dioica, interdum suffrutescens, foliis ovatis vel elliptico-ovatis acuminatis basi obtusatis aut obsolete cordatis integris (exsiccatis margine undulatis) trinerviis glabris vel nervis subtus puberulis, stipulis majusculis rotundatis, cymis masc. capituliformibus, pedunculis petiolos superantibus. ♀ In ins. S. Domingo (Nectoux). — Herba, caulibus raro suffrutescentibus, basi sæpe repentibus, 2-4 decim. longis, prostrato-diffusis, ramosis, pennæ corvinæ circiter crassitie, exsiccatis obsolete sulcatis cystolithisque minutissime punctulatis, merithallis inferioribus brevibus, nodis aphyllis. Folia 1½-3 centim. longa, 7-15 millim. lata, acumine brevi acuto, nervis basilaribus dimidium limbi superantibus, lamina exsiccata firmulæ, pagina superiore cystolithis punctiformibus minutissimisque inspersa, inferiore iisdem ut videtur destituta, petiolo 2-4 millim. longo. Stipulæ petiolo multo longiores, rotundatæ,

patulæ, superiores internodia proxima superantes, in stirpe exsiccata ferrugineæ, glabræ, diu persistentes. Capitula masc. valde densiflora, globosa, pisi circiter magnitudine, pedunculo subcentimetro glabro suffulta, floribus bracteis superantibus brevissime pedicellatis, alabastris  $1\frac{1}{2}$  millim. longis, segmentis perigonii mucronatis. (v. s. in hb. Mus. par. et DC.)

22. *P. RUBIÆFOLIA* (Bl. Mus. bot. lugd.-bat. 2. p. 49), herbacea vel inferne suffrutescens, ramosissima, foliis ovatis v. subrotundo-ovatis rarissime obovatis acutis vel obtusis basi rotundatis vel oblique obtusis vel retusis integerrimis trinerviis subciliatis cæterum glabris, stipulis brevibus ovatis, glomerulis racemosis vel paniculatis.  $\frac{1}{4}$  In Guatemala, ad rivos. — Wedd. Monogr. p. 192. — Folia (alterna paulo minora vel brevius petiolata) 8-26 millim. longa et lata, striolata. (Descript. ex Bl.). — Sequentibus affinis videtur, ab illis habitu præsertim formaque cymarum distinguenda, diagnosi autem nonnihil manca.

23. *P. NITIDA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 211; Monogr. p. 185), monoica, glaberrima, foliis elliptico-ovatis utrinque obtusis integerrimis trinerviis, stipulis minimis, cymis sessilibus quam petioli brevioribus androgynis. (1) In Peruvia (Dombey). — Herba, caule erectiusculo 6-10 centim. alto gracili, merithallo inferiore prælongo. Folia 5-15 millim. longa, 3-8 millim. lata, nervis lateralibus parum perspicuis medium limbi vix superantibus, laminæ exsiccatae membranaceæ pagina superiore cystolithis linearibus subtransversis creberrime obsita, inferiore iisdem destituta aut magis minusve lacunosa, petiolo dimidio limbo subæquilongo. Stipulæ vix perspicuæ, pellucidæ. Cymæ suborbiculares, congestifloræ, floribus masc. paucis, perigonii lobis breviter mucronatis, alabastro  $\frac{1}{2}$  millim. longo, femineis numerosissimis, perigonii fructiferi segmento intermedio cucullato lateralibus dimidio longiore achænioque  $\frac{1}{2}$  millim. longo lævique æquilongo. (v. s. in hb. Mus. par.)

24. *P. CILIARIS* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 209; Monogr. p. 186), monoica vel dioica, foliis ovatis superne sensim attenuatis vel acuminatis acutis basi obtusis rotundatisve aut subcordatis integerrimis trinerviis supra pilis diaphanis conspersis subtus glabris vel in costa nervisque pilosulis margine longiuscule ciliatis rarissime omnino glabris, stipulis abbreviatis triangularibus vel triangulari-acuminatis, cymis folio brevioribus rarius longioribus diffuse racemosis, floribus masc. et fœm. in iisdem glomerulis intermixtis aut in cymis distinctis, masc. bractea sua longioribus.  $\frac{1}{4}$  In insulis Antillis frequens. — Parietaria racemosa, foliis ad oras villosis Plum. sp. p. 10. t. 120. f. 2. *Urtica ciliaris* Linn. Sp. pl. p. 1396; Poir. Encycl. 4. p. 642. *Dubreuilia ciliaris* Gaudich. l. c. 495. Herba, rhizomate subhorizontali, caulibus diffusis, 2-5 decim. altis simplicibus ramosisve crassitie varia striatis glaberrimis, merithallis inferioribus elongatis, nodis plerisque foliigeris, exsiccatis minute punctulatis. Folia magnitudine admodum variabili, majora 10-12 centim. longa, 4-5 centim. lata, a triente inferiore vel solum a medio apicem versus attenuata vel sensim acuminata, acutiuscula, nonnunquam inæquilateralia, raro basi acutata, nervis basilaribus apicem limbi fere attingentibus, laminæ exsiccatae tenuiter membranaceæ pagina superiore cystolithis minutissimis et solum oculo armato manifestis obsita, inferiore iisdem corpusculis linearibus fusiformibusve inerspersa, petiolis ejusdem paris foliorum sæpe sicut limbi ipsi valde inæquilongis, hisce plerumque dimidio brevioribus rarissime longioribus glabrisculis. Stipulæ vulgo breves, a basi semiamplexicauli acuminato-atenuatæ. Cymæ in singulis axillis solitariae rariusve geminae, pedunculatæ, glabræ, glomerulis discretis, floribus subsessilibus; alabastris florum masc. millimetralibus, segmentis perigonii sub apice acute mucronatis; perigonii fructiferi segmento intermedio breviter cucullato quam lateralia dimidio longiore achænioque elliptico-ovato vix semimillimetr. lævi paulo brevior. (v. s.)

β, *Rivoriæ*, caulibus basi lignescentibus robustioribus et magis ramosis quam in typo, foliis vulgo oblique cordatis, floribus masc. dimidio majoribus. In insula Martinica (D<sup>r</sup> Rivoire), et Jamaica (Bauer, Wilson, etc.). — *Pilea Rivoriæ* Wedd. in Ann. sc. nat. l. c. p. 210; Monogr. p. 187. *P. Parietaria* Bl. Mus. lugd. bat. 2. p. 48; *Urtica Parietaria* Linn.? Sp. pl. p. 1397. — Cymæ ut videtur plerumque unisexuales, masc. ex axillis su-

perioribus ortæ. Alabastra fl. masc. sesquimillimetralia. Cystolithi in utraque pagina foliorum punctiformes.

γ, *Wydleri*, caulibus ramisque gracilioribus diffusioribusque quam in præced., foliis multo minoribus ovatis basi obtusatis rotundatisve aut rarius subcordatis, cystolithis in utraque pagina limbi punctiformibus, cymis ad glomerulum unicum vel paucos spicatos redactis. — In ins. Portorico (Wydler, n. 199); Cuba (Wright, n. 535); Martinica (Sieber, n. 51) etc. — *Pilea Wydleri* Bl. l. c. p. 49; Wedd. Monogr. p. 190; Griseb. Cat. pl. cub. p. 535. — Folia interdum vix centimetrum longa, cystolithis in pagina inferiore sicut in superiore punctiformibus. — Ad hanc formam pertinere videtur *Urtica cellulosa* Spreng.

ε, *tumida*, glaberrima caulibus carnosius ad nodos tumidis parce ramosis basi longe denudatis, foliis ovatis vel lanceolato-ovatis subacuminatis, cymis laxè ramosis. — In ins. S. Domingo (L. Cl. Richard). *Pilea puncticaulis* Wedd. Monogr. p. 181. — Caulis basi prostrati ramique adscendentes, siccitate cystolithis brevibus fusiformibus conspicue albo-punctatis. Folia subtus purpurascens, exsiccata in pagina inferiore cystolithis linearibus delicatulis oculo nudo conspicuis conspersa.

25. *P. ALSINIFOLIA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 211; Monogr. p. 189), dioica, foliis ovatis obtusiusculisve vix acuminatis basi obtusatis aut obsolete cordatis integerrimis trinerviis ciliatis utrinque pubescenti-pilosis, stipulis ovato-rotundatis, cymis femineis pedunculatis quam folia brevioribus. ¾ In Nova Granata, prope Bogota, locis muscosis humidisque (Goudot). Herba caulibus herbaceis v. basi lignescens, 3-4 decim. altis, ramosis, crassius pennæ corvinæ, glabris, exsiccatis teretiusculis pallide castaneis albicanibusve cystolithis punctulatis, nodis inferioribus denudatis. Folia 1½-2½ centim. longa, 8-15 millim. lata, a medio apicem versus sensim attenuata, nervis basilariibus medium limbi paulo superantibus subtus vix prominentibus, lamina exsiccata tenuiter membranacea pagina superiore cystolithis minutissimis brevissimeque fusiformibus crebre obsita, pagina inferiore iisdem corpusculis destituta, petiolo gracili 3-5 millim. longo glabriusculo. Stipulæ a basi semiamplexicauli rotundatæ, glabræ. Cymæ fem. pedunculis glabriusculis quam petioli longioribus suffultæ, parvæ, laxiuscule ramosæ, glomerulis approximatis, perigonii fructiferi segmento intermedio oblongo cucullato segmenta lateralia achæniūque ovatum ¼ millim. longum læve æquante. — Cymæ masc. ignotæ. Species nonnihil dubia, stipulis majusculis ad *P. Lindenianam* accedens. (v. s. in hb. Mus. par.)

26. *P. PROPINQUA* (Wedd. Monogr. p. 190), dioica vel monoica, foliis ellipticis brevissime apiculato-acuminatis acutis basi rotundatis rarius subcordatis integerrimis trinerviis ciliatis utrinque pubescentibus, stipulis triangularibus, cymis unisexualibus pedunculatis folia subæquantibus, bracteis glomerulorum masc. quam flores brevioribus. ¾ In ins. S. Domingo (Poiteau). Præcedentibus affinis. Caulis herbacei, erecti, subcæspitiosi, in specim. viso 2-3 decim. alti, simplices, exsiccati grisei cystolithis crebre punctulati, superne subtiliter pubescentes, merithallis sesqui-bi-centimetralibus, nodis inferioribus mox denudatis. Folia 2-4 centim. longa, 1½-2 centim. lata, nervis basilariibus usque ad limb. apicem haud protractis, lamina exsiccata rigidula subconcolore, cystolithis in utraque pagina creberrimis, in superiore autem minimis fere punctiformibus et in inferiore fusiformibus confusisque, petiolo subcentimetrali pubescente. Stipulæ late triangulares, a basi semiamplexicauli sensim attenuatæ, sparsim pubescentes. Cymæ laxè ramosæ v. racemiformes, pedunculis glabriusculis quam petioli longioribus suffultæ, glomerulis discretis, florum masc. perigonio pubescente, lobis sub apice breviter mucronatis, alabastro millimetrali, perigonii fructiferi segmento intermedio late ovato pileato lateralibus achæniisque subæquilongis. (v. s.)

27. *P. INTEGRIFOLIA* (Liebm. l. c. p. 298), monoica, foliis ovatis acutiusculis basi breviter emarginatis integris trinerviis ciliatis et utrinque paleaceo-setosis, stipulis lineari-lanceolatis, cymis androgynis quam petioli brevioribus, glomerulis discretis. ¾? In provincia mexicana Oajaca (Liebmann). — Wedd. Monogr. p. 188. — Caulis herbaceus, erectus, 1½-2 decim. altus, indivisus,

tetragonus, glaber, striatus. Folia 2-3 centim. longa, 10-15 millim. lata, supra fusco-vidua subtilissime punctulata nervis impressis, subtus glaucescentia, petiolo subcentimetro glabro. Cymæ geminae, patulae, floribus minutis, perigonis tuberculatis. (Descr. ex Liebm. l. c.) — Videtur affinis nonnullis præcedentium a quibus stipularum forma imprimis distinguenda.

28. P. LINDENIANA (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 210; Monogr. p. 188), monoica vel dioica, foliis ovatis vel oblongo-ovatis sensim acuminatis acutis basi oblique rotundatis subcordatisve integerrimis trinerviis ciliatis utrinque pubescentibus, stipulis majusculis ovatis oblongisve, cymis pedunculatis folia sæpe superantibus unisexualibus, bracteis glomerulorum masc. pisiformium flores superantibus. ? In humidis Novæ Granatæ, loco dicto Salto de Tequendama, prope Bogotâ (Goudot; Linden, n. 799. — P. Lindenii Bl. Mus. bot. lugd. bat. 2. p. 49. — Herba, caule subrecto, 3-5 decim. alto, simplice, crassitie pennæ anserinæ, exsiccatō fistuloso angulato-sulcato cystolithis punctulato, merithallis inferioribus valde elongatis sæpeque aphyllis. Folia 3½-7½ centim. longa, 1½-3 centim. lata, nonnunquam subpeltata, nervis basilariibus apicem limbi fere attingentibus, laminæ exsiccatæ membranaceæ subtus nonnihil rubescentis utraque pagina cystolithis punctiformibus crebre obsita, illis paginæ superioris sub lente modo papillarum eminentibus, petiolo subcentimetro pubescente. Stipulæ basi subamplexicaules, villosæ. Cymæ solitariae, laxæ ramosæ, glomerulis discretis, masc. ex axillis superioribus ortæ, pedunculo pubescente, glomerulis paucis, floribus inter bracteas rotundatas scariosas sessiles, alabastris 1 millim. longis, segmentis perigonii breviter mucronatis; femineæ pedunculo glabro suffultæ (an semper?), perigonii fructiferi segmento intermedio cucullato lateralibus paulo longiore achænioque ovato millimetro lævi vix superato. — Stirps P. ciliari habitu affinis, sed indole cymarum masc. et cæt. charact. optime distincta. (v. s.)

ß, *Cubana*, foliis supra fere omnino glabris, nervis subtus margineque et petiolo pilosulis, cymis masc. folio brevioribus e glomerulis pisiformibus discretis vel approximatis confectis. — In ins. Cubæ regione orientali, prope Monte Verde (Wright. n. 1449). — P. corymbosa Griseb. Pl. Wr. p. 174, non Bl. (l.) teste Griseb. P. ciliaris Griseb. Cat. pl. Cub. p. 60, non Wedd. Folia quam in typo majora, cystolithis laminæ exsiccatæ parum conspicuis, illis inferioris paginæ sparsis linearibusque vel oblongis. Stipulæ oblongæ, 3-5 millim. longæ. Cymæ masc. pedunculis petiolo longioribus glaberrimis suffultæ, glomerulis 1-4, bracteis membranaceis 2-3 millim. longis, inferioribus nudis glomerulum involucentibus. (v. s.)

29. P. STOLONIFERA (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 226; Monogr. p. 192), dioica, foliis caulinis ellipticis vel elliptico-lanceolatis acutis basi obtusatis et subpeltatis acutatisve integris triplinerviis supra glabris nervis subtus villosis, stipulis rotundatis, cymis laxis longe pedunculatis folia plerumque superantibus. ? Ad scopulos et secus flumina ins. Cubæ (Swartz). — *Urtica stolonifera* Swartz, in Act. Holm. (1797), p. 62; Fl. Ind. occid. p. 296; Poir. Encycl. suppl. 4. p. 219; Smith, l. c. 29. — Caulis herbacei, alii elongati filiformes prostrati radicanes et steriles, alii breviores erectiusculi, floriferi breviter villosi. Folia stolonum minima, obovata, atque in nodis extremis sæpe omnino deficientia, illa caulis fertilis 1½-4 centim. longa, 7-16 millim. lata, in ipso apice ejusdem sæpissime quaterna, deorsum magis minusve cuneata, lamina exsiccatâ rigida margine interdum sinuato et primo aspectu serrato, pagina superiore (præsertim ad marginem) cystolithis minimis fusiformibus crebris obsita, petiolis gracilibus dimidio limbo æquilongis villosis. Stipulæ majusculæ, in planta sicca ferrugineæ, in stolonibus (foliis deficientibus) et ad basim inflorescentiarum insignes. Cymæ pedunculis 3-4-centimetralibus

(1) Species Blumcana mihi sequente diagnosi nimis brevi solum cognita : P. *corymbosa* (Bl. Mus. bot. lugd. bat. 2. p. 50), herbacea, foliis longe petiolatis oblique ovatis (13 centim. longis, 8 centim. latis) acutis integerrimis scabris, corymbis longe pedunculatis axillaribus. ? In ins. Guadeloupe. — Wedd. Monogr. p. 186. *Urtica corymbosa* Poir. Encycl. 4. 640.



glabrisque suffultæ, masc. parvæ subcapitatæ, floribus pedicellatis, segmentis perigonii ante apicem longiuscule mucronatis, alabastro 1 millim. circiter longo; fem. ovatæ, glomerulis parvis subdiscretis confluentibusve, perigonii fructiferi segmento intermedio cucullato lateralibus longiore achæniisque late ovato et punctulato dimidio brevior. (v. s. typ. in hb. Holm. et Mus. brit.)

§ 2. — *Heterophyllæ, sive foliis in eodem jugo diversiformibus aut valde inæquimagnis.*

1° Gerontogæ (1).

30. P. ANISOPHYLLA (Wedd. Monogr. p. 193), dioica, foliis subintegerrimis at cæterum valde disparibus, cujusque jugi altero oblongo-vel ovato-lanceolato longiuscule acuminato basi subcordato petiolatoque, altero multo minore plerumque ovato breviter acuminato basique profunde hastato-cordato et amplexicauli, cymis pedunculatis, pedunculo quam petiolus folii majoris plerumque longiore. ½? In Himalaya tropica centrali et orientali: Nepalia (Wallich) Sikkim (J. D. Hooker). *Urtica anisophylla* Wall. Cat. n. 4594. — Herba, caule glaberrimo vel superne (ad nodos præsertim) villosa, simplice aut parce ramoso, angulato, modice crasso, internodiis elongatis. Folia trinervia, glabra vel nervis subtus villosis, majora 6-10 centim. longa, 1½-5 centim. lata, acumine lineari limboque integerrimis vel denticulo hic illic instructis, petiolo 1-2-centimetrali glabro aut magis minusve villosa, minora 1½-4 centim. longa, 10-12 millim. lata, sinu aperto cordata, nervis basilaribus in omnibus usque ad apicem limbi productis, secundariis tenuissimis æqualibus approximatis transverseque parallelis, laminæ exsiccata membranaceæ utraque pagina cystolithis linearibus inspersa, illis superioris minutissimis vix conspicuis et interdum prorsus deficientibus, inferioris sparsis et multo majoribus. Stipulæ parvæ, triangulares. Cymæ masc. parce breviterque ramosæ, pedunculo 2-3-centimetrali, glomerulis parvis subunilateralibus, floribus sessilibus, perigonii lobis ante apicem obtusiuscule mucronatis glaberrimis. Cymæ fem. laxè patulæque ramosæ, glomerulis discretis, floribus breviter pedicellatis sessilibusve, perigonii fructiferi segmento intermedio breviter cucullato, lateralibus multo longiore achæniisque ovato vix millimetrali læviusculo triente vel dimidio brevior. (v. s.)

31. P. MANNIANA, dioica, glaberrima, foliis oblongo-ellipticis acuminatis subintegris sæpiusve in dimidia parte superiore repando-serrulatis cæterum disparibus, altero nempe majore longeque petiolato basi rotundato, altero sæpius omnino sessili basi cordato et amplexicauli, cymis breviter pedunculatis, femineis petiolum folii majoris haud superantibus. ½? In ins. S. Thomas Africa occidentalis (G. Mann). — Suffrutex scandens, caule elongato crassitie calami scriptorii, lævi vel obsolete striato, internodiis 6-10-centimetralibus, siccitate fusco. Folia triplinervia, nervis basilaribus arcuatis usque ad basim acuminis angusti protractis, serraturis paucis distantibus; majora 8-10 centim. longa, 3½-5½ centim. lata, basi nonnunquam submarginata; minora triente vel dimidio minora, lamina exsiccata submembranacea sive rigidiuscula, supra fusco-nigricante, subtus glauca plexuque nervorum et venularum eximie fusco-reticulata, utrinque cystolithis minutissimis oculoque nudo prorsum inconspicuis crebre obsita, petiolo folii majoris tertiam vel dimidiam partem limbi æquante. Stipulæ majusculæ (circiter centimetrales) oblongo-ovatæ, vix persistentes. Cymæ in singulis axillis vulgo geminæ, ovatæ, 1½-4 centim. longæ pedunculo incluso, masculæ majores quam femineæ, floribus breviter pedicellatis; perigonii masculi segmentis juxta apicem brevissime obtusiusculeque mucronatis alabastro orbiculari millimetrum metiens; feminei segmento intermedio oblongo breviter et subacutè cucullato quam lateralia ovata dimidio longiore achæniisque late ovatum submillimetrale læviusculum haud æquante. (v. s. in hb. Kew. et Mus. par.)

(1) Vid. etiam P. rupendiam inter *Dentalas*.

32. *P. INSOLENS*, dioica, glaberrima, foliis ovatis subintegris sæpiusve in parte superiore repando-serrulatis admodum disparibus, altero nempe cujusve jugi (saltem versus apicem ramorum) multo majore longe et anguste acuminato basi rotundato vel subcordato longe petiolato et insigniter peltato, altero parvo vix acuminato basi profunde cordato subsessili et amplexicauli, cymis fem. longiuscule pedunculatis, pedunculo petiolum folii majoris vulgo superante. 4? In declivibus umbrosis prov. Mishmi Bengalæ orient. (Griffith) Caulis herbaceus vel lignescens, gracilis, forsan scandens, parce ramosus, ramis plus minus elongatis flexuosis striato-sulcatis, internodiis elongatis, siccis virentibus. Folia inferiora rami (in specimen unico viso) superioribus minora, basi cordata, haud peltata, serraturis numerosioribus. Superiorum majora 6-8 centim. longa, 3-4 centim. lata, acumine uni-sesqui-centimetrali parce serrulato, nervis basilaribus a petioli insertione arcuatis et usque in acumen protractis, minora 1-2-centimetralia, lamina exsiccata membranacea, pagina superiore fusca cystolithis minutissimis oculo nudo prorsus inconspicuis punctiformibus linearibusque obsita, inferiore fusco-virente, cystolithis majoribus linearibus conspersa, petiolo folii majoris dimidium limbum æquante. Stipulæ 2-4 millim. longæ, oblongo-ovatae, subpersistentes. Cymæ fem. (solum visæ) solitariae, læviuscule patuleque ramosæ, pyramidatae, glomerulis discretis aut nonnullis confluentibus; floribus subsessilibus. Perigonii fructiferi segmentum intermedium apice gibbo dorsali oblongo auctum, achænium ovatum 1½ millim. longum haud æquans lateribusque rotundatis dimidio longius. (v. s. in hb. Kew.)

#### 2° Neogææ (1).

33. *P. GLOMERATA* (Griseb. Cat. pl. cub. p. 69), dioica, suffruticosa, foliis (in eodem jugo valde inæquimagnis) lanceolatis sensim acuminatis acutis basi obtusis rotundatisve et sæpissime ad insertionem petioli oblique emarginatis subintegris integrisque trinerviis sparsissime ciliatis aut fere glabris, stipulis brevibus triangularibus, floribus ex omnibus axillis ortis in glomerulos pisiformes ad basim petiolorum sessiles congestis, achænio ovato lævi. 3 Prope villam Monte Verde, in reg. orient. ins. Cubæ (Wright, n. 529 et n. 2240). — *P. Parietaria* Griseb. Pl. Wr. in Mem. Acad. am. sc. et art. nov. ser. 8 (1860), p. 173 et Cat. pl. cub. p. 60, non Blume. — Caulis 3-6 decim. altus, basi sæpe denudatus, superne præsertim ramosus, ramis gracilibus lævissimis glaberrimisque siccitate sicut caulis ipse fuscis. Folia 1-6 centim. longa, 3-15 millim. lata, in singulis jugis altero quam alterum duplo vel triplo longiore longiusque petiolato, plerumque plus minus inæquilatera, circiter a medio sensim attenuato-acuminata, margine sæpissime vage et inæqualiter suberoso-sinuata nec unquam regulariter denticulata, nervis basilaribus supra impressis circiter ad quartam partem superiorem limbi vel ultra protractis, lamina exsiccata obscure fusco-viridi membranacea, supra cystolithis punctiformibus et solum oculo armato conspicuis obsita, subtus iisdem corpusculis linearibus minimisque crebre conspersa, petiolis gracilibus illo folii majoris 1-15 millim. longo, glabro vel præsertim apice sparsim ciliato. Stipulæ vix millimetales. Glomeruli axillarum oppositarum sæpissime confluentes, quam petiolus longior multo breviores, masc. nonnihil majores, floribus horumce breviter pedicellatis, segmentis perigonii sub apice breviter mucronatis, alabastro 1 millim. longo; floribus fem. sessilibus, perigonii fructiferi segmentis late ovatis subæquiolongis achænio semimillimetrali dimidio brevioribus, intermedio crasse carinato apice sæpe trilobo. (v. s.)

34. *P. HETERONEMA* (Griseb. Pl. Wr. in Mem. Acad. Am. sc. et art. n. ser. 8 (1860) p. 173, Cat. pl. cub. p. 59), dioica, herbacea vel suffrutescens, ramississima, glabra, foliis parvis disparibus rhombæo-ellipticis rotundatisve integerrimis triplinerviis, cujusque jugi altero longiuscule petiolato, altero dimidio minore fere sessili, stipulis inconspicuis, cymulis minimis sessilibus vel brevissime pedunculatis ex omnibus fere axillis nascentibus. 4 In regione

(1) Vid. quoque *P. sessilifloram* inter *Dentatas*.

orientali ins. Cubæ, prope villam Monte Verde (Wright, n. 537 et 1456 pro parte). — Caulis gracilis, basi lignescens, ramis alternis subdistichisque valde foliosis, exsiccatis striatis et cinerascens. Folia distiche patentia, 4-10 millim. longa, vulgo utrinque obtusa basiue sæpe obliqua et subemarginata; majora elliptico-vel obovato-rhombea inferne cuneata, petiolo limbum subæquante vel paulo brevior, minora rotundato-rhombea, petiolo vix millimetrali, nervis basilariibus bene distinctis arcuatis ad quartam partem apicalem limbi protractis, lamina exsiccata membranacea subtus pallida utraque pagina cystolithos lineares exhibente, illis superioris creberrimis, inferioris sparsis. Glomeruli axillares, pauciflori, petiolo folii majoris multo breviores, floribus masc. 3-5 pedicellatis, pedicello perigonium subæquante, segmentis hujusce vix millimetralibus sub apice mucrone brevi conico instructis, floribus fem. quam masc. paulo numerosioribus sessilibus vel brevissime pedicellatis, perigonii fructiferi segmentis rotundatis, intermedio longiore leviter cucullato achæni late ovati scabriusculi dimidium superante. (v. s.)

35. *P. LUCIDA* (Bl. Mus. lugd. bat. 2. p. 48), monoica, fruticulosa, glaberrima, foliis parvis dimorphis, in singulis jugis altero majore pinnatifido-lobato (lobis utrinque 2-3), altero obovato integerrimo, aut interdum (teste Swartzio) aliis in jugis distinctis lobatis aliisque integris, cymis minimis paucifloris, masc. longe pedunculatis, fem. sessilibus.  $\frac{5}{2}$  In Jamaica frigidioris rupibus et ruderalis (Swartz) et Martinica (Sieber n. 51). — Wedd. Monogr. p. 200; Griseb. Fl. Br. W. Ind. p. 156, excl. var. *Urtica lucida* Swartz, in Act. Holm. (1785) p. 31; ejusd. Fl. ind. occid. p. 315; Smith, l. c. n. 48. — Caulis 1-2-decimetralis, diffuse ramosus, inferne denudatus, ramis gracilibus foliosis siccitate rubentibus. Folia 5-10 millim. longa,  $1\frac{1}{2}$ -2 $\frac{1}{2}$  millim. lata, breviter petiolata, majora fere 5-lobata, lobis obovatis subæqualibus obtusis, sinubus inter lobos obtusis, lamina exsiccata membranacea læte viridi supra cystolithis linearibus majusculis sparsim ornata subtus iisdem raro omnino destituta. Stipulæ late ovatæ, petiolis circiter æquilongæ. Cymæ masc. e glomerulo unico paucifloro constantes, pedunculo filiformi folio longiore, floribus sessilibus, perigonii lobis sub apice longiuscule mucronatis, mucronibus in alabastro submillimetrali conniventibus. (v. s. typ. in hb. Holm. et Mus. brit.)

36. *P. CUNEIFOLIA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 212; Monogr. p. 195), monoica, herbacea, glaberrima, foliis disparibus, cujusque jugi altero parvo cuneato-obovato integroque, altero sextuplo majore cuneiformi prope apicem utrinque crenato-dentato penninervio, cymis parvulis pedunculatis quam folia majora brevioribus.  $\frac{2}{3}$  In ins. Jamaica, inter muscos rupium calcarearum (Swartz). — Bl. l. c. p. 50. *P. lucida*  $\beta$  *cuneifolia* Griseb. l. c. *Urtica cuneifolia* Swartz, in Act. Holm. (1787) p. 70; ejusd. Fl. Ind. occ. p. 319; Smith, l. c. n. 50; Poirét, Encycl. suppl. 4. p. 222. — Caulis erectus vel ascendens, 1-3-decimetralis, angulatus, ramosus, ramis fere simplicibus. Folia majora 10-18 millim. longa, 3-6 millim. lata, in petiolum brevem attenuata, lamina exsiccata rigida, supra cystolithis linearibus (in margine numerosioribus) oblique lineolata, subtus albicante cystolithis multo minoribus et parum conspicuis. Stipulæ petiolo etiam breviores, obtusissimæ. Cymæ e glomerulo unico sæpe constantes, pedunculo filiformi folium minus superante, floribus subsessilibus. (v. s. typ. in hb. Holm. et Mus. brit.)

37. *P. NUTANS* (Wedd. Monogr. p. 196. t. 7. f. 11-13), dioica, glaberrima, foliis dimorphis, cujusque jugi altero elliptico vel elliptico-lanceolato utrinque acutiusculo vel obtuso (rarius ovato) petiolato tri-triplici nervio, altero multo minore rotundato vel rotundato-cuneato obtusissimo sessilibus, ultra medium vel trientem inferiorem serratis, cymis minimis sessilibus vel breviter pedunculatis.  $\frac{2}{3}$  In Peruvia subandina, prope Huayvari, etc. (Dombey; Poeppig, n. 1565); in Novæ Granatæ prov. Antioquia (Triana). — *Urtica nutans* Poeppig, mscr. — Herba, caule diffuso v. prostrato et radicante, 2-5 decim. longo, flexuoso, ramoso, inferne longe denudato, ramulis gracilibus foliosis. Folia majora 1-4 centim. longa,  $\frac{1}{2}$ -1 $\frac{1}{2}$  centim. lata, petiolo gracili 2-4 millim. longo suffulta, minora 3-6-millimetralia, serraturis parvis utrinque 6-8 acutiusculis obtusisve, lamina exsiccata submembranacea fere concolore utraque pagina

cystolithis linearibus obsita, illis superioris autem minoribus crebrioribusque. Stipulæ brevissimæ, late triangulares, parum conspicuæ. Cymæ masc. valde paucifloræ, floribus pedicellatis perigonii lobis obtuse mucronatis, alabastris millimetralibus, femineæ quam folium minus breviores, segmento intermedio perigonii fructiferi lateralibus majore achæni oblique rotundati dimidium superante, stigmate fere laterali. (v. s.)

38. *P. CLANDESTINA*, monoica, glaberrima, foliis admodum disparibus, cujusque jugi altero majore oblongo-lanceolato utrinque acuto subsessili in dimidia parte superiore acute serrato triplinervio, altero oblique lanceolato integerrimo sessili, cymis parvis pedunculatis petiolum superantibus.  $\frac{1}{2}$  In ins. Jamaica (Wulfschlägel, n. 1787). — Habitus *P. nutantis*. Caulis 2-3-decimetralis, nudus, ramis foliosis. Folia majora 2 centim. longa, 8-10 millim. lata, serraturis utrinque 5, dente terminali cæteris æquali, nervis basilariibus ultra medium protractis, lamina exsiccata membranacea cystolithis paginæ superioris linearibus elongatis intertextis, inferioris crebrioribus et nonnihil minoribus. Cymæ fem. e glomerulo unico constantes, floribus subfasciculatis longiuscule pedicellatis, perigonii fructiferi segmento intermedio quam achænium lanceolatum dimidio brevior. (v. s. in hb. Monac.)

39. *P. TRICHOSANTHES*, dioica, floribus exceptis fere glabra, foliis dimorphis, in singulis paribus altero majore oblongo acutiusculo vel obtuso basi cuneatim attenuato petiolatoque in dimidia parte superiore grossiuscule serrato penninervio, altero magnitudine varia rotundato vel obovato vel oblique elliptico sessili vel breviter petiolato integerrimo vel parce dentato, cymis paucifloris petiolos æquantibus aut vix superantibus, floribus masc. (solum visis) longe hirsutis.  $\frac{1}{2}$  In sylvis, inter Guaranda et Bodegas, ditione Æquatoris (J. Remy). — Herba, caule elongato gracili ramosissimo prostrato et radicante, ramis adscendentibus ramulisque glaberrimis apice præsertim foliosis. Folia sparsim ciliata, cæterum glabra, majora 8-12 millim. longa, 3-6 millim. lata, petiolo 3-8-millimetrali, serraturis acutiusculis vel obtusis in utroque latere limbi 3-4; folia minora interdum vix bimillimetralia, rotundata et omnino sessilia interdumque oblique obovata aut elliptica atque dimidium limbi folii majoris superantibus serraturis 2-3 aut subnullis, lamina exsiccata membranacea subtus pallidior supra cystolithis linearibus discretis conspicue ornata, subtus iisdem fere destituta. Stipulæ millimetrum longæ, rotundatæ, amplexicaules. Cymæ masc. 3-5-floræ, breviter pedunculatæ vel subsessiles, floribus majusculis congestis, perigonii segmentis circiter bimillimetralibus, acumine seu mucrone subapicali dimidium segmenti totius æquante pilis patulis subfasciculatis dorso munitis. — Habitus præcedentium, sed forma foliorum indoleque nervorum et cymis optime distincta. (v. s. in hb. Mus. par.)

40. *P. FALLAX*, dioica, apice ramorum ramulisque hirsutis, foliis cujusque paris valde inæqualibus, altero ovato vel ovato-lanceolato longiuscule petiolato basique obtuso vel oblique cuneato, altero multo minore ovato breviter petiolato et basi obtuso vel oblique rotundato, ambobus grossiuscule serratis penninerviis supra sparsim pilosis subtus in nervis præsertim hirtis, stipulis cordato-ovatis amplexicaulibus, cymis masc. umbelliformibus pedunculo petiolum superante.  $\frac{1}{2}$  vel  $\frac{3}{5}$  In Andibus regionis æquatorialis (Spruce, n. 6106). — Suffrutescens, caule ascendente basi radicante et omnino glabrato, ramis angulatis siccitate obscure fuscis, internodiis 4-6-centimetralibus, ramulis gracilibus vix elongatis erecto-patulis foliosis. Folia majora 2-4 centim. longa, 12-20 millim. lata, circiter a medio apicem versus sensim attenuata vel subacuminata, serraturis (juxta basim minoribus vel subnullis) acutis extus convexis ciliatis, dente apicali paulo majore ovato acutiusculo, nervis subtus parum prominulis jugo inferiore paulo supra insertionem petioli nascente quam sequens brevior; minora 6-12 millim. longa, 3-8 millim. lata, lamina exsiccata membranacea subtus pallidior cystolithis minimis linearibus ægre perspicuis utrinque crebre conspersa illis paginæ inferioris aliis multo majoribus mixtis, petiolo folii majoris 1-1 $\frac{1}{2}$  centim. longo. Stipulæ (siccæ scariosæ) 3-5 millim. longæ, inæqualiter denticulatæ et ciliatæ, patulæ. Cymæ masc. (solum visæ) e floribus 8-12 pedicellatis majusculis constantes pedunculoque 1-2-centimetrali pubescente suffultæ, pedicello florem æquante perigonioque

hirtulis, segmentis hujusce in secco inferne pallide ferrugineis sub apice longiuscule mucronatis, mucronibus in alabastro circiter trimillimetrali discretis.

β, *glabra*, foliis glabris, cæteris ut in typo. — Crescit in iisdem locis (Spruce, n. 6107). (v. s. cum typo in h. Kew., DC. et Boiss.)

41. *P. RUFÆ* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 220; Monogr. p. 196), monoica, ramis hirsutis. foliis disparibus, in singulis jugis altero elliptico apice attenuato vel subacuminato utrinque acuto petiolatoque, altero multo minore subsessili late ovato vel obovato et utrinque obtuso, supra basim acute serratis pagina utraque (sed præsertim in nervis inferioris) pilis rufescentibus hirsuta, cymis androgynis fere racemiformibus, pedunculis quam petioli longioribus. ¼ - ½ In locis saxosis montium Jamaicae australis (Swartz). — Griseb. Fl. Br. W. Ind. p. 157. *Urtica rufa* Swartz, in Act. Holm. (1787) p. 159; Fl. Ind. occid. 1. p. 292; Poiret, l. c. p. 219; Smith, l. c. n. 15. — Herbacea vel suffruticosa, caule a basi ramoso crasso sinuoso magis minus nodoso 3 decim. circiter alto, ramis ascendentibus teretiusculis ferrugineo-hirsutis, internodiis brevibus. Folia majora 2½-4 centim. longa, 1-1½ centim. lata, a medio utrinque attenuata, nervis basilaribus trientem superiorem limbi fere attingentibus, serraturis triangularibus margine exteriori convexiusculis, lamina exsiccata rigidula subtus glaucescente cystolithisque fusiformibus brevissimis et creberrimis punctulata, pagina superiore inter pilos elongatos cystolithis majusculis lineari-fusiformibus ornata, petiolo 1 centim. longo ut costa ipsa pilis longis rufis hirsuto. Folia minora subcentimetralia, a medio utrinque rotundato-attenuata, cæterum cæteris similia. Stipulæ rotundatæ, semicentimetrales, glabræ, siccitate ferrugineæ. Cymæ vix multifloræ, glomerulis approximatis in rhachi fere simplice subsessilibus, florum masc. perigonii lobis ante apicem breviter mucronatis, mucronibus in alabastro depresso discretis, perigonii fem. segmento intermedio breviter cucullato quam lateralia circiter dimidio longiore achænioque multo breviora. (v. s. typ. in hb. Holm. Monac. Mus. brit.)
42. *P. DIVERSIFOLIA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 212; Monogr. p. 197), dioica, glaberrima, foliis parvis disparibus, cujusque jugi altero lanceolato obtusiusculo basi cuneato attenuato petiolato grosse et obtuse serrato penninervioque, altero oblique rotundato sessili subintegro vel apice crenulato tripplinervio, cymis fem. breviter pedunculatis. ¼ In Peruvia (Pavon). — Herba, caule elato ramosissimo, ramis subfastigiatis, internodiis interdum subfusiformi-inflatis, ramulis gracilibus patulis admodum foliosis. Folia majora sive petiolata 10-12 millim. longa, 2-4 millim. lata, minora 3-6 millim. longa et lata, basi sæpissime oblique cordata, laminæ exsiccatae membranaceæ pagina inferiore cystolithis elongatis dispersa, superiore iisdem corpusculis numerosissimis eleganterque intertextis ornata. Stipulæ minimæ. Cymulæ fem. paucifloræ, perigonii segmento intermedio incurvato segmentis laterali-bus duplo longiore. (v. s. in hb. Webb. et Boiss.)
43. *P. RHIZOBOLA* (Miq. in Mart. Fl. bras. fasc. 12. p. 202), monoica, glaberrima, foliis alternis (scilicet cujusque jugi folio altero minimo s. abortivo) lanceolatis acuminatis basi acutatis excepta basi serratis tri-tripli-nerviis, stipulis triangulari-lanceolatis persistentibus, cymis sessilibus paucifloris petiolo brevioribus v. subæquilongis. ¼ In sylvis Brasiliæ, prope Ilheos (Maxim. Neu-wied. Martius). — Wedd. Monogr., p. 198. *Boehmeria radicans* Nees et Mart., in Nov. Act. Soc. nat. Cur., XI. 35. t. 70. — Herba, caulibus sesquidecime-tralibus, a basi radicante erectis, crassiusculis, sulcatis, foliosis, internodiis inferioribus bi-tri-centimetralibus. Folia majora 4-10 centim. longa, 1½-2½ centim. lata, nervis basilaribus usque ad apicem limbi protensis, petiolo 2-15 millim. longo. Folia abortiva ovata, 5-8-millimetralia, acutiuscula, sessilia v. subsessilia; cystolithis in foliis omnibus crebris punctiformibus-que. Stipulæ basi connatæ vaginamque brevem efformantes. Flores masc. lobis perigonii sub apice mucrone lato instructis, alabastro subtetragono. (v. s. in hb. Mart. et Kew.)
44. *P. TRIANÆANA*, dioica, caule glabriusculo vel pubescente, foliis maxime inæquimagnis sed fere conformibus oblongo-vel ovato-lanceolatis acumina-

tis basi acutis petiolatis triplinerviis supra basim integram crenato-serratis, majoribus amplis, stipulis oblongo-ovatis, cymis fem. laxe multifloris longe pedunculatis. ♀ In sylvis provinciæ Choco, ad Dagua (Triana). — Caulis radicans, modice crassus, in speciminibus obviis apice foliatus floriferque. Folia majora 8-16 centim. longa,  $3\frac{1}{2}$ - $5\frac{1}{2}$  centim. lata, subæqualiter crenato-vel subsinuato-serrata, acumine angusto etiam mucronato, supra glabra, subtus plus minus pubescentia, petiolo 1-3-centimetrali puberulo, lamina exsiccata membranacea cystolithis linearibus utrinque sed præsertim in pagina superiore conspersa; minora  $1\frac{1}{2}$ -4 centim. longa, 6-8 millim. longa. Stipulæ petiolum folii minoris circiter æquantes, persistentes. Cymæ fem. (solum visæ) patule ramosæ, glomerulis parvis subdiscretis, pedunculo petiolum folii majoris superante. Perigonii fructiferi segmentum intermedium oblongum, summo dorso gibbo oblongo auctum et achænium ellipticum sesquimillimetre paulo superans; segmenta intermedia dimidio breviora ovata. Stamina rudimentaria inflexa, quam segmenta ipsa longiora. (v. s. in hb. Mus. par.)

45. P. DENDROPHILA (Miq. in Mart. Fl. bras. fasc. 12. p. 202), dioica, glabra, foliis in singulis jugis admodum dissimilibus, altero nempe elliptico-vel oblongo-lanceolato aut obovato obtuso basi oblique attenuato et superne utrinque grosse bi-quadri-crenato vel serrato, altero obovato-rotundo obtuso obliquissime cuneato vel cordato integerrimoque, cymis sessilibus petiolos non aut vix superantibus. ♀ Ad truncos et scopulos in sylvis Guyanæ (Melinon, Leprieur, etc.), Brasiliæ, juxta flumen Solimoes (Martius), Peruviae orientalis (Spruce, n. 4434) et Novæ Granatæ (Triana). — Wedd. Monogr. p. 185. P. imparifolia Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 212; Monogr. p. 194. t. VII. f. 1-3; Miq. in Mart. Fl. bras. fasc. 12. p. 199. — Herba, caulibus prostratis et plerumque longe repentibus 3-6-decimetralibus subsimplicibus ramosisve, ramis gracilibus inferne magis minusve denudatis. Folia omnia triplinervia; majora 2-5 centim. longa, 8-30 centim. lata, in petiolum gracilem semicentimetralem desinentia; minora 1-2 centim. longa, subsessilia; lamina exsiccata membranacea utrinque viridi cystolithis delicatulis rectis arcuatisve (in pagina superiore majoribus) ornata. Stipulæ parvæ, late amplexicaules, obtusæ. Cymæ masc. sæpius paucifloræ, floribus breviter pedicellatis, lobis perigonii sub apice crasse mucronatis, alabastro 1-1½ millim. longo. Cymæ fem. minimæ, floribus brevissime pedicellatis, perigonii fructiferi segmento intermedio cucullato lateralibus subæquilongis achænioque elliptico ovato compresso et margine scabriusculo fere dimidio brevior. (v. s.)

β, *subintegra*, foliis majoribus late ellipticis obovatisve vel orbicularibus utrinque obtusis vel basi obtusis integerrimis vel rarius prope apicem serratis. — In Brasilia. (v. s. in hb. Mart.)

γ, *canescens*, foliis exsiccatis subtus canis, cystolithis vix conspicuis. — In Guyana gallica. (v. s. in hb. Mus. par.)

δ, *major*, quam varietates præcedentes multo vegetior, caule crasso villis radiculisque papilliformibus undique vestito, foliis majoribus (8-10 centim. longis) in dimidia parte superiore s. apicali crenato-serratis, minoribus ellipticis basi oblique subcordatis, subtus pubescentibus, inflorescentiis. . . . In Novæ Granatæ provinciis S. Buenaventura, Choco y Barbacoas, altit. 300 met. (Triana.)

46. P. CRUGERIANA, dioica, glaberrima, foliis valde disparibus, cufusque paris altero majore lanceolato anguste acuminato basi subinaequali acuto petiolo, altero duplo v. triplo breviori oblique obovato-oblongo breviter acuminato basi rotundato, utroque triplinervio et supra basim integram serrato v. serrulato, cymis sessilibus petiolo folii majoris breviori. ♀ In Venezuela (Cruger). Caulis elongatus, semimetralis et ultra, carnosus, prostratus, ramis alternis adscendentibus, internodiis 1-3-centimetralibus. Folia majora 6-9 centim. longa,  $1\frac{1}{2}$ -2 centim. lata, acumine serrato, nervis basilaribus usque ad basim hujusce protensis, lamina exsiccata membranacea supra obscure subtus glauco-viridi, pagina superiore præsertim cystolithis linearibus substellatisque minimis densissime obsita. Stipulæ breves, triangulares. Cymæ masc. (solum visæ) parvæ sed sat multifloræ, floribus breviter pedicellatis, perigonii seg-

mentis sub apice crasse mucronatis, alabastris in specimine viso vix semimilimetralibus. (v. s. in hb. Kew.)

47. *P. CENTRADENIOIDES* (Seemann, Bot. Herald, p. 194), dioica, caule pubescenti-hirsuto, foliis distichis cujusque paris valde inæqualibus lanceolatis acuminatis grosse serratis basi inæquilateraliter cuneatis triplinerviis, majoribus petiolatis, minoribus sessilibus, stipulis cordato-oblongis, cymis masc. e glomerulo unico pedunculato confectis, perigonii segmentis hirtis et sub apice longe appendiculatis.  $\frac{1}{2}$  In sylvis opacis promontorii Corrientes, in districtu Darien Novæ Granatæ (Seemann), necnon prope Tamana prov. Choco (Triana). — Herba aut (teste Seemann) suffrutex, caule ramisque diffusis radicantibusque. Folia majora 4-8 centim. longa, 8 millim. -  $2\frac{1}{2}$  centim. lata, basi integra, cæterum usque ad apicem acuminis serrata, serraturis acutis extus convexis, intermediis majoribus, supra sparsim pilosa demumque glabrata, nervis subtus pubescentibus; lamina exsiccata membranacea utrinque cystolithis linearibus parum conspicuis inspersa, nervis in pagina superiore sæpe albicantibus; petiolo circiter centimetrali pubescenti-hirto. Folia minora vix centimetrum longa, 2-4 millim. lata, vix acuminata, nonnunquam subintegra. Stipulæ 4-10 millim. longæ, diu persistentes, obtusæ, fuscæ. Cymulæ masc. (solum visæ) 4-5 millim. latæ, pedunculo piloso petiolum folii majoris superante, floribus sessilibus, segmentis perigonii (inclusa appendicula) circiter 3 millim. longis. — Habitus hujusce Pileæ et sequentis illum *Elatostematum* quorundam optime refert. Nondum est exacta hæc cum *Centradenia* similitudo, quoniam *Melatosmaceæ* omnes foliis integris gaudeant. (v. s. in hb. Mus. par.)

48. *P. VARIEGATA* (Seemann Bot. Herald, p. 194?, absque descript., exclus. syn.), dioica, caule hirtulo, foliis distichis ejusdem paris admodum dissimilibus, altero multo majore lanceolato acuminato basi inæquilateraliter cuneato serrato triplinervio petiolato, altero parvulo ovato basi subcordato in dimidia parte superiore serrato sessili, stipulis anguste oblongis, cymis masc..., fem. parvis e glomerulis paucissimis constantibus pedunculatis.  $\frac{1}{2}$  In sylvis calidis prov. Choco et Barbacoas Novogranatensium (Triana). — Habitus et aliquatenus facies præcedentis. Folia in singulis nodis etiam dispariora; majora 6-10 centim. longa, 2-3 centim. lata, supra basim integram subæqualiter serrata, serraturis acutis extus rectiusculis, supra glabra et in vivo longitudinaliter albo-fasciata, nervis subtus strigosis, margine hic et illic ciliolato, laminæ exsiccatae membranaceæ pagina superiore in medio sive juxta nervum medium fuscata et cystolithis linearibus undique pulchre conspersa, pagina inferiore iisdem corpusculis multo minoribus inconspicue notata, petiolo semicentimetrali hirtulo. Folia minora centimetrum longa, 6-8 millim. lata, subabrupte acuminata, serraturis obtusiusculis. Stipulæ 6-8-millimetales, obtusæ, mox deciduæ. Cymulæ fem. (solum visæ) ad glomerulos 2 vel 3 ut plurimum redactæ, pedunculo petiolum folii majoris dimidia longitudine excedente, floribus sessilibus; perigonii fructiferi 2 millim. longi segmenta omnia summo dorso gibbo oblongo et subacuto aucta; intermedium longius, achænium ovatum læviusculum nonnihil superans. (v. s. in hb. Mus. par.)

### § 3. — *Dentatæ, sive foliis nec integris nec in eodem jugo dimorphis aut valde inæquimagnis.*

#### 1° Gerontogæ.

\* Glabræ, sive caule folisque pills omnino destitutis.

A. — *Longi pedunculatæ, sive pedunculis (saltem cymarum fructiferarum) quam petioli plerumque longioribus* (1).

a. Stipulæ breves triangulares persistentes.

49. *P. RUPIPENDIA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186; Monogr. p. 201), monoica aut interdum dioica?, glaberrima, foliis (nonnunquam in eodem

---

(1) Vid. quoque *P. trinerviam* et *P. Celebicam*, inter *Brevipedunculatas*.

jugo valde dissimilibus) ovatis vel rotundatis vel ellipticis vel lanceolatis aut cujusque jugi altero ovato-vel rhomboideo-lanceolato, altero minore late ovato vel rotundato, obtusis aut acutiusculis sæpe acuminatis basi vulgo cuneatis triplinerviis in dimidia parte superiore serratis, stipulis breviter triangularibus, cymis unisexualibus, fem. paucifloris ut plurimum pedunculatis (pedunculo petiolum superante) raro subsessilibus.  $\frac{1}{2}$  In ins. Mascarenis (Commerson, Gaudichaud, etc.); in Africa orientali (Martin). — *P. pendula* et *P. urticæfolia* Bl. Mus. lugd. bat. 2. p. 47. *Parietaria urticæfolia* Linn.? Suppl. p. 434. *Urtica rupipendia* Poir. Encycl. 4. p. 640. *U. pendula* Willd. Sp. pl. 4. p. 358. *Boehmeria urticæfolia* Spreng.? Syst. veg. 3. p. 844. *Dubruella rupipendia* Gaudich. Uran. p. 495. — Herba nonnunquam suffrutescens, caule ramosissimo procumbente, 2-6 decim. longo teretiusculo 4-lineato, exsiccato rubente cystolithisque sparsim punctulato, merithallis inferioribus elongatis, ramulis foliosis. Folia 1-3 centim. longa, 5-12 millim. lata, serraturis utrinque 2-4, dente terminali cæteris æquilongis v. duplo longiore, nervis basilariibus brevibus dimidium limbum vix superantibus; lamina exsiccata membranacea, pagina superiore (nervis exceptis) cystolithis linearibus crebriusculis inspersa, subtus autem hisce fere destituta; petiolo quartam limbi partem longitudine vix æquante. Stipulæ parum conspicuæ. Cymæ in axillis pluribus ramulorum glomeratæ, floribus interdum subverticillatis breviter pedicellatis, perigonii masc. lobis sub apice anguste acutiuscule mucronatis, alabastro bimillimetrali siccitate cystolithis punctulato, perigonii fem. segmento intermedio breviter cucullato quam lateralibus paulo longiore achænioque ovato unisesquimillimetrali circiter æquilongis. (v. s.)

$\beta$ , *homophylla*, foliis omnibus (ut in typo oppositis) æquimagnis et homomorphis plerisque cuneato-obovatis, pedunculis quam petioli haud raro brevioribus. — In ins. Mascarenis. (v. s.)

$\gamma$ , *subverticillata*, foliis nonnullis quaternatim verticillatis basi plerumque rotundatis obtusisve quam in typo sæpe majoribus, cymis folia vulgo superantibus. (v. s.)

50. *P. TERNIFOLIA* (Wedd. Monogr. p. 202), dioica, glaberrima, foliis elongatis sæpius lineari-lanceolatis ternatim verticillatis longe attenuato-acuminatis basi obtusis rotundatisve penninerviis in parte dimidia superiore vel inferiori inæqualiter laxèque dentato-serratis, dentibus subacuminatis, stipulis brevissimis inconspicuis, cymis pauci-multi-floris solitariis vel geminis, pedunculis quam petioli longioribus.  $\frac{1}{2}$  In Indiæ orientalis reg. temp. Himalayæ, prov. Sikkim, alt. 2100-2600 metr. (J. D. Hooker). — Herba, rhizomate repente carnoso sæpe moniliformi-incrassato, caulibus erectis gracilibus statura quam maxime variabili (nempe  $\frac{1}{4}$ -4-decimetralibus) simplicibus inferne aphyllis, merithallis inferioribus elongatis. Folia omnia ternatim verticillata, rarius opposita, 2-10 centim. longa,  $1\frac{1}{2}$ -10 millim. lata, aut nonnihil latiora, acuminate acutissimo v. obtusiusculo, serraturis sæpius paucis ac plerumque argutis, sinubus acutis aut obtusiusculis, nervis basilariibus tenuissimis obliquis, lamina exsiccata tenuiter membranacea supra albo-punctata, cystolithis linearibus subtus solum et in foliis quibusdam conspicuis, petiolis plerumque in eodem jugo subæqualibus brevissimis (1-3-millim.) sed nonnunquam paulo longioribus et tunc inæqualibus. Stipulæ oculo armato vix aut non distinctæ. Cymæ masc. in specim. exam. admodum paucifloræ, et glomerulo nempe unico constantes, pedunculo gracillimo petiolum duplo triplove longitudine superante, floribus breviter pedicellatis, perigonii lobis sub apice mucrone conspicuo instructis, alabastro sesquimillimetrali. Cymæ fem. pauci-multi-floræ, pedunculo 1-2 centim. longo, glomerulis parvis nunc fere solitariis nunc numerosis inflorescentia simul simpliciter vel subcompositè laxèque ramosa, floribus subsessilibus, perigonii fructiferi segmento intermedio oblongo vix cucullato lateralibus multo longiore achænioque millimetro ovato lævi triente vel dimidio brevioris. (v. s. in hb. Kew. et Mus. par.)

51. *P. THOUARSIANA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186; Monogr. p. 203), dioica, glaberrima, foliis late ovatis suborbicularibus obtusis basi rotundatis vel obsolete cordatis excepta basi obtuse serratis tri- vel subquinque-nerviis, stipulis breviter triangularibus, cymis masc. et glomerulis 2-4 inter-



rupte spicatis constitutus, pedunculis quam petioli multo longioribus.  $\frac{1}{2}$  In ins. mascarenis ? (Du Petit Thouars). — Herba, caulibus adscendentibus basi lignescens, radicantibusque 3-5 decim. altis penna corvina vix crassioribus teretiusculis inferne longe nudis, internodiis elongatis, exsiccatis cystolithis minutis albicantibus. Folia  $1\frac{1}{2}$ -3 centim. longa,  $1\frac{1}{2}$ -2 $\frac{1}{2}$  centim. lata, serraturis triangularibus obtusiusculis, dente terminali cæteris multoties majore, nervis basilaribus valde arcuatis fere usque ad apicem limbi productis, lamina exsiccata firmula cystolithis linearibus quidem utrinque sed in pagina superiore (et præsertim in marginibus ejusdem) crebriusculis in pagina inferiore contra raris ornata, petiolis gracilibus 7-15 millim. longis. Stipulæ parum conspicuæ. Cymæ masc. (fem. ignotæ) folia paulo superantes, pedunculo filiformi, glomerulis parvis discretis, floribus sessilibus, perigonii lobis sub apice mucrone semimillimetrali acuto munitis, alabastro sesquimillimetrum circiter longo. (v. s. in hb. Mus. par.)

52. *P. sessilifolia* (Wedd. Monogr. p. 204), dioica, raro monoica, glaberrima, foliis ellipticis rariusve ovatis, superioribus quaternatim verticillatis, utrinque attenuatis acutisque vel subacuminatis et basi rotundatis in margine toto argute serratis aut basi integris subtriplinerviis, stipulis breviter triangularibus persistentibus, cymis unisexualibus e glomerulo unico vulgo constantibus, pedunculis quam folia longioribus.  $\frac{1}{2}$  In ins. mascarenis (Commerson, Gaudichaud). *Urtica sessilifolia* Poir. Encycl. 4. p. 642. *Dubruella sessilifolia* Gaudich. voy. Uran. p. 495. — Herba, caulibus interdum basi lignescens longiuscule radicantibus dein adscendentibus 1-3 decim. altis gracilibus vix ramosis foliosis, exsiccatis cystolithis punctato-striolatis. Folia omnium nodorum interdum verticillata, vulgo  $1\frac{1}{2}$ -3 centim. longa, 6-10 millim. lata, in ramis sterilibus vero sæpe majora ac nonnunquam semidecimetralia, ejusdem jugi magis minus inæqualia, serraturis plerumque acuminatis, nervis lateraliibus usque ad quintam partem superiorem limbi productis, lamina exsiccata membranacea supra læte viridi cystolithis linearibus conspicue ornata subtus sæpe glauco-rufescente cystolithisque tenerrimis conspersa, petiolo subnullo vel sæpius bi-tri-millimetrali. Stipulæ parum conspicuæ. Cymæ in axillis superioribus pedunculis bi-tri-centimetralibus suffultæ, pisi vix magnitudine vel paulo majores; floribus masc. breviter pedicellatis, perigonii lobis sub apice mucrone conico instructis, alabastro bimillimetrali; floribus femineis subsessilibus, perigonii segmento intermedio cucullato lateralibus subdimidio longiore achænioque millimetrali ovato lævi paulo brevior. (v. s. in hb. Mus. par. et Juss.)

53. *P. repanda* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186; Monogr. p. 205), monoica ?, glaberrima, foliis amplis ovato-v. oblongo-lanceolatis acuminatis basi subcordatis fere sessilibus basi excepta repando-serrato-crenatis trinerviis, stipulis triangularibus subpersistentibus, cymis androgynis? laxifloris, pedunculis quam petioli multoties longioribus.  $\frac{3}{5}$  In ins. Mauritio (Commerson ?) — Suffrutescens, elata, ramis simplicibus teretiusculis vix crassis, internodiis semidecimetralibus, exsiccatis cystolithis crebre punctatis. Folia 8-13 centim. longa, 3-5 centim. lata, serraturis magis minusve distantibus obtusisque, acumine vario, nervis basilaribus arcuatis ad acumen usque productis subtus vix prominentibus, lamina exsiccata membranacea cystolithis linearibus inæqualibus aliisque punctiformibus crebre obsita, illis paginæ inferioris nisi in nervis plerumque minoribus, petiolo millimetrali. Stipulæ acutæ, parum conspicuæ. Cymæ in axillis superioribus solitariae, foliis breviores, pedunculis filiformibus 4-5-centimetralibus, ramulis numerosis divaricato-patentibus, glomerulis parvis discretis, floribus masc. .... pedicellis longiusculis in dichotomiis ramulorum alaribus; floribus fem. subsessilibus, perigonii fructiferi segmentis subæquilongis, intermedio breviter cucullato quam achænium ovatum læve millimetrale brevior. (v. s. in hb. Mus. par.)

54. *P. wightii* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186; Monogr. p. 205), monoica, glaberrima, radicans, foliis ovatis vel rarius lanceolato-ovatis acuminatis basi rotundatis subcordatisve in toto margine argute serratis trinerviis, stipulis brevibus triangularibus persistentibus, cymis parvis androgynis (floribus masc. et fem. vel mixtim in eodem vel in distinctis glomerulis) aut

unisexualibus, pedunculis petiolum sæpissime superantibus. 4 In regione temperata montium Nilghiri Indiæ orientalis (Wight, cæt.) et in Zeylanica. — P. Hugelii Bl. Mus. bot. lugd. bat. 2. p. 53†. *Urtica radicans* Wight, Icon. pl. Ind. t. 1974, non Swartz. — Herba, caule elongato succulento crassitie circiter pennæ corvinæ vel aliquantulum crassiore parce ramoso basi longe radicante demum ascendente, exsiccat cystolithis obsoletis punctulato, merithallis sæpe 3-4-centimetralibus. Folia 2-4 centim. longa, 1½-2½ centim. lata, raro fere orbiculata, sensim vel subabrupte acuminata, nervis basilariibus eleganter arcuatis fere usque ad basim acuminis sive ad partem limbi quartam superiorem protensis, secundariis transversis tenuibus, lamina exsiccata membranacea in pagina superiore cystolithis minimis (vix conspicuis) creberrimisque in inferiore vero multo majoribus et rarioribus ornata, petioliis in singulis jugis ut in genere solet sæpe inæquilongis, majoribus centimetrum raro metientibus. Stipulæ 1-2-millimetrales. Cymæ in axillis pluribus aut solum superioribus solitariæ, pedunculo gracili 2-4-centimetrali, glomerulis paucis (nempe plerumque 2-6) approximatis vel discretis et sæpe subspicatis, floribus sessilibus aut breviter pedicellatis; perigonii masc. lobis sub apice conspicue acutiusculeque mucronatis, alabastro 1½ millim. fere lato; perigonii fem. segmento intermedio obtuse cucullato quam lateralia duplo longiore achænioque submillimetrali ovato læviusculo fere triente brevior. (v. s. typ.)

b. *Stipulæ majusculæ, ovatæ vel lanceolatæ.*

55. *P. oxyodon* (Wedd. Monogr. p. 222), dioica, glaberrima, radicans, foliis ovatis vel lanceolato-ovatis acuminatis supra basim obtusam vel rotundatam serrulatis (serraturis argute acuminatis) trinerviis, stipulis ovatis persistentibus, cymis laxè simpliciterque distiche ramosis, pedunculis quam petioli multo longioribus. 4 In India orientali, regione temperata Himalayæ prov. Sikkim. (J. D. Hooker). — Caulis basi longe radicans, 3-5 decim. longus, angulatus, nodis inferioribus denudatis, crassitie pennæ. Folia 2½-7 centim. longa, 1½-3 centim. lata, acumine quoad longitudinem vario ut limbus reliquus serrato, dentibus parum exsertis, acumine submillimetrali, nervis basilariibus usque ad apicem folii fere productis, laminæ exsiccatæ membranacæ pagina superiore cystolithis minimis punctiformibus inferiore (pallidior) iisdem breviter fusiformibus obsita, petioliis gracilibus inæquilongis dimidio limbo brevioribus. Stipulæ 5-6 millim. longæ, scarioso-membranacæ, obtusæ. Cymæ multifloræ ovatæ, pedunculis gracilibus 5-6-centimetralibus, glomerulis parvis approximatis bracteis minimis suffultis in ramulis primariis subspicatis; floribus masc. brevissime pedicellatis, perigonii lobis sub apice breviter mucronatis; achænio ovato ½ millim. longo, in utraque facie linea scabra circinata notato perigonii segmentum majus fere dimidia longitudine excedente. — P. Wightii refert sed stipulis et cæt. optime distincta est. (v. s.)

56. *P. UMBELLATA*, dioica, glaberrima, elata, foliis amplis ellipticis obovatisve superioribus plerumque quaternis acutis vel breviter acuminatis basi acutatis obtusatisve rarius subcordatis ultra basim vel trientem aut etiam dimidium argute serratis tri-tripli-nerviis, stipulis lanceolatis deciduis, cymis densis, masc. capituliformibus pedunculis quam folia persæpe longioribus. 5 In ins. Borbonia (Commerson, Gaudichaud, etc.). *P. Commersoniana* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186, Monogr. p. 221. P. insignis Bl. Mus. lugd. bat. 2. p. 56. — *Urtica umbellata* Bory, Voy. Iles Afr. 3. p. 173. — Suffrutescens, elata, caulibus simplicibus v. a basi ramosis radicanibusque robustis, internodiis inferioribus prælongis, exsiccat cystolithis punctulatis. Folia in nodis duobus superioribus ut plurimum quaternatim verticillata, 5-10 centim. longa, 3-6 centim. lata, serraturis sæpe acuminatis subæqualibus, nervis basilariibus arcuatis ad acumen usque productis subtus prominulis, lamina exsiccata rigidula colore sæpe obscure cyaneo-viridi utrinque cystolithis linearibus subæqualibus (in pagina inferiore, nervis exceptis, paulo minoribus) crebre obsita, petiolo robusto 2-10-millimetrali. Stipulæ subcentimetrales, acutæ. Cymæ in axillis superioribus solitariæ; masculæ congestifloræ, nucem avellanam magnitudine æmulantes, floribus pedicellatis, lobis perigonii sub apice mucrone semimillimetrali obtuso instructis, alabastro

2½ millim. circiter longo; femineæ glomerulis minus congestis subpaniculatis, floribus subsessilibus; perigonii fructiferi segmentis subæqualibus achænio late ovato millimetrali lævi subæquilongis. (v. s.)

B. — *Brevipedunculatæ, sive pedunculis (saltem inflorescentiarum fructiferarum) quam petioli sæpius brevioribus* (1).

a. *Stipulæ breves, triangulares.*

57. *P. SCRIPTA* (Wedd. Monogr. p. 222), dioica, glaberrima, foliis elliptico-lanceolatis lanceolatisve longiuscule acuminatis basi acutis aut obtusatis in triente inferiore integris cæterum acute vel obtusiuscule serrulatis trinerviis, venulis interrupte incrassatis, stipulis brevibus triangularibus, cymis femineis laxis floribundis, pedunculis quam petioli brevioribus. ¼ In regionibus temperatis et subtropicis Himalayæ, Nepalia (Wallich), prov. Sikkim (J. D. Hooker) et Jamu (Thomson) necnon in montibus Khasiæ (J. D. Hooker) et Nilghiri? — *P. scripta* et *P. Goglado* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 187 et 188. *P. scripta* Bl. Mus. lugd. bat. 2. p. 53 et 57. *Urtica scripta* Hamilt. in Don Prodr. Fl. Nepal. p. 59. *U. Goglado* Wall. Cat. n. 4593 et 4593 C. *U. Hamiltoniana* ejusd. l. c. n. 4590. *U. triplinervis* Hamilt. mscr. — Caulis herbacei, submetrales, parce ramosi, exsiccati cystolithis minutissimis punctulati, ramis elongatis foliosis, internodiis 3-4 centim. longis. Folia 7-15 centim. longa, 3-5½ centim. lata, ad petioli insertionem subemarginata, acumine subintegro, nervis basilaribus usque ad apicem limbi productis, secundariis transversis et parallelis, venulis tenuibus subinterruptisque sæpe mire conspicuis et concinne anastomosantibus; lamina exsiccata membranacea subtus pallidior utrinque cystolithis linearibus tenuissimis vix conspicuis inspersa, venulis in pagina superiore plerumque interrupte prominulis, petiolis gracilibus 3-5-centimetralibus in eodem jugo valde inæqualibus. Stipulæ 2-3-millimetrales, a basi anguste acuminatæ, persistentes. Cymæ masc. foliis breviores, ramulis numerosis patentibus, glomerulis approximatis vel discretis, floribus vix non sessilibus, perigonii lobis sub apice brevissime mucronatis, mucronibus in alabastro submillimetrali discretis. Cymæ fem. in axillis pluribus solitariae, pyramidatae, foliis plerumque breviores, ramulis patentibus, glomerulis subdiscretis, floribus ut in cyma masc. subsessilibus sessilibusque; perigonii fructiferi segmento intermedio breviter cucullato lateralibus longiore achænioque ovato millimetrali albedo faciebis conspicue granulosis sed juxta marginem lævibus fere dimidio brevioribus. — A sequente, cui charact. cæt. nimis affinis, plerumque facile distinguenda venulis foliorum irregulariter incrassatis scriptaque hieroglyphica quodammodo referentibus. (v. s.)

β, *stipularis*, stipulis in acumen elongatum productis ¼ in montibus Khasiæ (Griffith). (v. s.)

58. *P. TRINERVIA* (Wight, Icon. pl. Ind. t. 1973), dioica, ut plurimum glaberrima, foliis amplis ellipticis ovatisve longiuscule acuminatis basi subcordatis vel rotundatis rarius acutis in margine toto argute serrulatis vel inferne integris trinerviis, rete venularum æquali, nervis foliorum juniorum interdum villosis, stipulis brevissime triangularibus, cymis majusculis multifloris, pedunculis masculinarum quam petioli multo longioribus, feminearum contra quam petioli brevioribus, achænio sæpissime granulato. ¼ In regionibus tropicis et subtropicis montium peninsulae Indiæ orientalis, Zeylaniae et Javæ frequens. — Wedd. Monogr. p. 224. *P. melastomoides* ejusd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186. *P. pellucida*, *P. melastomoides* et *P. oreophila* Bl. Mus. lugd. bat. 2. p. 54. *P. elongata* Miq. in Zolling. Syst. Verzeichn. p. 102, non Blume. *P. oreophila* ejusd. Fl. Ind. bot. 1 (2). p. 235. t. XVII. *Urtica melastomoides* Vahl mscr. in hb. Juss.; Poir. Encycl. Suppl. 4. p. 223. *U. peduncularis* Sm. in Rees' Cyclop. 37. *Urtica* n. 6. *U. trinervia* Roxb. Fl. Ind. 3.

(1) Vid. quoque *P. rupiendiam* inter *Longipedunculatas*, necnon *P. peploitem* et *P. smilacifoliam* inter *Integrifolias*.

p. 582. *U. elongata* Bl. Bijdr. p. 494. *U. cataractæ* Steud. Nomencl. ed. 2. p. 734; Hassk. Pl. jav. p. 204. *U. oreophila* Miq. Pl. Jungh. p. 27. *Dubruella* melastomoides Gaudich. Uran. p. 495. — *Habitus* præcedentis at plerumque paulo robustior, caulibus herbaceis aut basi lignescentibus, sæpe e basi ramosis, internodiis 3-8-centimetalibus. Folia 6-20 centim. longa, 2½-10 centim. lata, deorsum haud raro subcuneata, necnon ima basi magis minusve emarginata, acumine longitudine variabili acuto sæpissimeque hic et illic serrulato, nervis basilaribus secundariisque in folio adulto semper glaberrimis, venulis haud interruptis sed rete æquale fingentibus; lamina exsiccata membranacea utrinque cystolithis minimis linearibus vel rarius subpunctiformibus crebre aut sparsim obsita, petiolis in eodem jugo inæquilongis limbum dimidium sæpe æquantibus superne aliquando villosis. Stipulæ a basi amplexicauli abruptissime attenuatæ, acutæ, parum conspicuæ. Cymæ masc. in axillis pluribus solitariæ rariusve geminæ (altera multo majore), fere pyramidatae et nonnunquam (juniores præsertim) complanatae, folia superantes iisve (etiam incluso pedunculo interdum plus quam decimetrali) breviores, ramis elongatis patulis, glomerulis approximatis, floribus breviter pedicellatis, perigonii lobis sub apice brevissime mucronatis, mucronibus in alabastrum sesquimillimetrali conniventibus. Cymæ fem. in axillis pluribus plerumque geminæ, quam folia multo breviores, ramulis sæpe numerosissimis divaricatis, glomerulis parvis subsessilibus discretis; perigonii fructiferi segmento intermedio vix cucullato lateralibus paulo longiore achæniumque ¼ millim. longum obtusissimum dimidia fere longitudine superante. (v. s.)

β, *minor*, caule foliisque et inflorescentiis multo minoribus. ¾ In iisdem locis ac typus. (v. s.)

59. *P. PACHYCARPA* (Wedd. Monogr. p. 225), dioica, glaberrima, foliis ellipticis acuminatis basi obtusis glabrisque cæterum inæqualiter acuteque serratis trinerviis, stipulis brevibus triangularibus persistentibus, cymis fem. parvis quam petioli brevioribus subspiciformibus, achænio marmorato nec granulato. ¾ In provincia Assam, Indiæ orientalis (Griffith). — Caulis parce ramosus, angulatus, crassitie pennæ anserinæ vel gracillior, exsiccatus cystolithis minutissimis creberrime punctulatus. Folia a medio ad apicem sensim attenuata demumque acuminata, 5-8 centim. longa, 3-4 centim. lata, serraturis margine exteriori convexiusculis acutis apiculatisve, nervis basilaribus arcuatis summum limbum fere attingentibus, secundariis parallelis vix non transversis, lamina exsiccata membranacea cystolithis minimis linearibus ægre perspicuis obsita, petiolis inæqualibus limbo dimidia parte brevioribus. Stipulæ a basi (petiolo æquilata) subabrupte attenuatæ, 1-2 millim. latæ. Cymæ fem. in axillis pluribus geminæ, petiolos paulo superantes iisve breviores, breviter ramosæ indeque fere spiciformes, glomerulis approximatis, perigonii fructiferi segmento intermedio lanceolato inconspicueque cucullato lateralibus ovata acuta circiter dimidia longitudine superante. Achænium ovatum, 1¼-1½ millim. longum, pericarpio nucamentaceo læviusculo sub lente crebre punctato. — Præcedentem indole refert, inflorescentia autem et achænio distinctissima est. (v. s. in hb. Kew. et Mus. par.)

60. *P. CELEBICA* (Miq. in Zolling. System. Verzeichn. p. 105; Fl. Ind. bat. 1 (2). p. 236), dioica?, carnososucculenta, glabra, foliis ovatis subacuminatis basi rotundatis supraque petiolum leviter emarginatis serratis trinerviis, stipulis parvis, cymulis raro longiuscule sæpius breviter pedunculatis vel subsessilibus quam folia brevioribus, floribus (fem.) confertis. ¾ In ins. Celebes, prope Maros, ad rupes (Zollinger, n. 1065). — Wedd. Monogr. p. 226. — Herba habitu *Peperomia*, caulibus crassis, ramis tetragono-angulatis sulcatis. Folia 1½-7 centim. longa, longiuscule petiolata, petiolis in eodem jugo inæqualibus gracilibus 4-centimetalibus, lamina exsiccata membranacea venosa utrinque cystolithis striolata. Inflorescentia tenuis, folio brevior, dichotome ramulosa, ramulis ultimis spicatum floriferis. (Descript. ex Miq. l. c.)

61. *P. HOOKERIANA* (Wedd. Monogr. p. 226), dioica, elata, glabriuscula, foliis magnis ellipticis abrupte vel subabrupte acuminatis præter basim fere fere rotundatam vel subcuneatam repando- et interdum subobsolete crenato-serratis triplinerviis, stipulis brevissimis, cymis masc. solitariis quam petioli multo

brevioribus e glomerulis paucis pisiformibus discretis constantibus.  $\frac{1}{2}$  In regione subtropica Himalayæ orientalis, prov. Sikkim, altit. 1500 metr. (J. D. Hooker). — Caulis herbaceus, metrum excedens, crassus, succulentus, angulatus, parce ramosus, glaber vel ad nodos subinconspicue villosus, exsiccatus cystolithis oblongis creberrime punctulatus, internodiis elongatis (superioribus nempe etiam 2-5-centimetralibus). Folia quam in omnibus generis speciebus (excepta *P. grandis*) fortassis majora, 1-2 decim. longa, 6-13 centim. lata, acumine subcentimetro obtusiusculo, nervis basilaribus arcuatis circiter ad partem quintam superiorem limbi productis, lateralibus tenuibus oblique transversis, lamina exsiccata submembranacea, cystolithis linearibus æqualibus oculo nudo etiam conspicuis utrinque obsita, petioli in eodem jugo solito more inæquilongis merithallum proximum superantibus interdumque limbum ipsum fere æquantibus, glabris vel juxta laminam utroque latere villosulis. Stipulæ vix sesquimillimetrales, persistentes. Cymæ masc. pedunculo 1-1  $\frac{1}{2}$  centim. longo suffultæ, simplices vel parvis ramosæ, glomerulis in singulis inflorescentiis 3-10 sessilibus vel pedunculatis majusculis fere æquidistantibus, floribus subsessilibus, perigonii lobis sub apice brevissime gibboso-mucronatis siccitate cystolithis striatulis, alabastro subsesquimillimetrali. (v. s. in hb. Kew. et Mus. par.)

62. *P. BOIVINIANA* (Wedd. Monogr. p. 227), dioica, glaberrima, foliis elliptico-vel oblongo-lanceolatis longiuscule acuminatis basi rotundatis et in insertione petioli emarginatis supra basim integram et in acumine ipso laxè serrulatis tri-tripli-nerviis, stipulis breviter triangularibus persistentibus, cymis parvis densifloris subsessilibus pedunculatisve solitariis aut geminis quam petioli plerumque multo brevioribus, achænio acuminato subtiliter circinatim granuloso.  $\frac{1}{2}$  In sylvis, secus rivulos convallium insulæ Nossi-Bé, prope Madagascar (Boivin). — Caulis herbaceus, adscendens, 4-6 decim. longus, basi aphyllus et radicans, crassitie circiter pennæ anserinæ, exsiccatus cystolithis minutissime punctulatus, merithallis superioribus sesqui-bi-centimetralibus. Folia 6-12 centim. longa, 2-4 centim. lata, basi nonnunquam obliqua, acumine sesquicentimetrali ut limbus reliquus serrulato, nervis basilaribus arcuatis usque ad apicem limbi productis margini duplo quam costæ propinquioribus, secundariis plurimis tenuissimis et oblique transversis reticulatisque, lamina exsiccata membranacea cystolithis tenuissimis linearibus utrinque creberrimis obsita, petioli gracilibus inæqualibus dimidio limbo brevioribus. Stipulæ millimetrales. Cymæ masc. non visæ. Cymæ fem. vix centimetrum latæ, pedunculo nullo vel 1-2-centimetrali, glomerulis paucis sæpius confluentibus, floribus sessilibus, perigonii fructiferi segmento intermedio oblongo vix cucullato, lateralibus multo longiore achænioque lenticulari-ovato millimetrali acutangulo tertia vel dimidia parte brevior. (v. s. in hb. Mus. par.)

63. *P. TRILOBATA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186; Monogr. p. 228), glaberrima, foliis parvis plerisque quaternatim verticillatis oblongis apice et basi obtusis paulo infra medium dente obtuso utrinque vel uno latere solum instructis rariis integerrimis penninerviis triplinerviisve, stipulis breviter triangularibus persistentibus, cymis....  $\frac{1}{2}$ ? In insula Mauritio (Commerson). — *Urtica trilobata* Poir. Encycl. p. 639. — Herbacea, caule procumbente, 2-3-decimetrali ramoso, ramis adscendentibus teretiussculis crassitie pennæ corvinæ vel crassioribus foliosis, internodiis inferioribus 3-5 centim. longis, exsiccatis cystolithis crebre punctulatis albicantibusque. Folia 12-20 millim. longa, 3-5 millim. lata, obtusissima, ad insertionem petioli subemarginata, nervis basilaribus dimidium limbi haud attingentibus, lamina exsiccata membranacea pallide virente in pagina superiore cystolithis linearibus oblique transversis creberrimis in inferiore rarioribus et costæ marginique subparallelis interspersa, petiolo gracili quam limbus dimidius plerumque brevior. Stipulæ vix semimillimetrales, obtusæ. (v. s. in hb. Mus. par.)

64. *P. LÆVICAPILLA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186; Monogr. p. 228), dioica, glaberrima, foliis parvis, interdum quaternatim verticillatis ovatis subacuminatis obtusis basi rotundatis subcordatisve trinerviis grosse et obtuse crenato-serratis, stipulis breviter triangularibus persistentibus, cymis masc. minimis racemiformibus petioli subæquilongis, fem....  $\frac{1}{2}$  In insula Mauri-

tio, ad rupes humidæ montis Pouce aliorumque (Belvis). — Cespitosa, caulibus herbaceis 1-3 decim. longis adscendentibus basi longe nudis teretibusculisque crassitie pennæ anatinae (exsiccati albicantibus) simplicibus vel superne ramosis, ramis angulatis foliosis, internodiis inferioribus elongatis, superioribus brevissimis: Folia 1-2 centim. longa, 4-12 millim. lata, nemulla integerrima albis dentatis intermixta, dentibus utrinque 1-3, terminali lateralibus multo majore, nervis basilaribus ad trientem superiorem limbi productis, lamina exsiccata rigidula utraque cystolithifera, cystolithis linearibus in pagina superiore tenuibus creberrimisque, in inferioris intervallis majoribus et rarioribus; in costa punctiformibus; petiolo quam limbus brevior vel eidem æquilongo. Stipulæ parum conspicuæ: Cymæ masc. in speciminibus examinatis rarioræ, petiolo haud superantes, floribus brevissime pedicellatis, perigonii lobis sub apice breviter mucronatis; alabastro vix millimetrali. — Species subdiaba, præcedenti necnon formæ cuidam subdepuperatæ æquantis cognata. (v. s. in hb. Mus. par.):

65. *P. lucens* (Wedd. in Ann. sa. nat. 4<sup>e</sup> sér. 1. p. 186; Monogr. p. 229), dioica, glaberrima, foliis amplis ovatis vulgo longe acuminatis basi rotundatis. subcordatisve ima basi acuminæque exesptis crenato-serratis, trinerviis, stipulis breviter triangularibus persistentibus, cymis floribundis laxis pedunculis quam petioli plerumque brevioribus, achænio subtilissime granuloso vel lævisculo. ①-? In montibus obumbratis ins. mascarenarum (Commerçon, Gaudichaud, etc.; Sieber, n. 511); in ins. Sumatra et Timor (Riedel). — Mtg. Fl. Ind. bat. 1 (2). p. 234. *P. lucens*, *P. subcrenata*, *P. triplinervia* et *P. Riedlei* Bl. Mus. bat. fugd. bat. 2. p. 48. 53 et 55. *Urtica lucens* Poir. Encycl. 4. p. 641. *U. subcordata* Commers. mscr. *U. triplinervia* Rb. L. c. Suppl. 4. 224. *U. cuspidata* Willd. Sp. pl. 4. 349; Smith, l. c. n. 5. *U. triplinervis* Vahl in hb. Juss.; Pers. Synops. 2. p. 552. *Dubreuilia lucens* Gaudich. Uran. 495. *D. Riedlei* Decne Herb. Timor. p. 161. — Herba annua (teste Commerçon) aut subcæspitosa, ramis primariis sive caulibus elongatis teretiusculis foliosis crassitie variabil, exsiccati cystolithis minutis creberrime punctulatis albicantibusque, internodiis 2-6-centimetralibus. Folia 6-15 centim. longa, 2-8 centim. lata, a medio ad apicem sensim attenuata, dentibus ut plurimum obtusis, nervis basilaribus eleganter arcuatis ultra basim acuminis productis subtus parum prominentibus, lamina exsiccata submembranacea nitida utrinque cystolithis crebris linearibus fusiformibusve et inæqualibus obsita, petiolo gracilibus in eodem jugo valde inæquilongis æsqui-quinq. centimetralibus. Stipulæ a basi lata abruptissima attenuatæ, 1-2 millim. vix longæ. Cymæ in axillis pluribus plerumque geminæ quam folia multo breviores, ramulis sæpe numerosissimis divaricatis, glomerulis discretis; floribus masc. pedicellatis, perigonii lobis sub apice breviter mucronatis, alabastro 1-1½ millim. longo; femineis in glomerulis minimis subsessilibus, perigonii fructiferi segmento intermedio vix cucullato lateralibus longiore achænioque late ovato ½ millim. longo et obtusissimo fere dimidia longitudine superato. (v. s.)

β, *minor*, foliis multo minoribus quam in typo. (v. s.).

66. *P. sublucens*, dioica, glaberrima, foliis ovatis acuminatis basi obtusis rotundatisve basi excepta serratis (serraturis subincurvis apiculatisque) tri-triplinerviis, stipulis brevissime triangularibus persistentibus, cymis fem. laxis petiolo brevioribus, achænio ovato lævi. ? In monte Clarence, insulæ Fernando-Po (Mann). — *P. lucens* affinis, sed habitu graciliore. Caulis forsan annuus, 2-4 decim. altus, internodiis elongatis, basi denudatus. Folia 2-5 centim. longa, 1½-3 centim. lata, sensim acuminata, acumine serrato, serraturis omnibus subæqualibus acuminatis dorsoque convexis, limbo ad insertionem petioli subemarginato, lamina exsiccata membranacea cystolithis minimis linearibus sub lente undique dispersa, petiolo gracili dimidio limbo longiore. Stipulæ vix perspicuæ. Cymæ fem. in singulis axillis plerumque geminæ, pedunculatæ, parce ramosæ; glomerulis parvis sessilibus pedunculatisve. Perigonii fructiferi segmentum intermedium lateralibus multo longius, sub apice breviter gibbosum, achæmium ovatum millimetrale æquans. (v. s. in h. Kew.)

b. *Stipulæ*, majusculæ, ovatæ vel oblongæ (1).

67. *P. ANGULATA* (Blume, Mus. bot. lugd. bat. 2. p. 55), dioica, sæpius glaberrima, foliis amplis ellipticis vel elliptico-ovatis acuminatis basi rotundatis obtusisve et integris cæterum grosse serratis (serraturis acutis obtusiusculisve trinerviis), stipulis majusculis ellipticis foliaceis, cymis fem. confertifloris petiolo brevioribus, achænio lævi.  $\frac{2}{4}$  In Indiæ orientalis montibus Bhasia et Sikkim, altit. 1500-2700 metr. (J. D. Hooker et Thomson); in Zeylante (Thwaites, n. 2184), Java (Blume, Junghuhn, Zollinger, n. 869) et ins. Loe-Cheo (Wright, n. 298). — *P. stipulosa* Miq. in Zoll. Syst. Verzeichn. p. 102; Fl. Ind. bat. p. 1 (2). n. 236; Wedd. Monogr. p. 230. *P. petiolaris* ejusd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186. *Urtica angulata* Blume Bijdr. p. 494.  $\frac{1}{4}$  *stipulosa* Miq. Pl. Jungh. p. 28. — Herba elata, caule basi radicante simplicis angulate, sicco cystolithis creberrimis punctato-striolato. Folia 6-10 centim. longa, 4-6 centim. lata, acumine subcentimetroli acutiusculo integro; nervis basilaribus in parte superiore limbi cum secundariis arcuatim anastomosantibus, rarissime pilis quibusdam hic et illic conspersa; lamina exsiccata membranacea supra obscure subtus late virente interdumque fere pellucida utrinque cystolithis crebris linearibus minimis subæqualibusque obsita, petiolo in eodem jugo (ut limbi ipsi) inæquilongis et interdum longissimis. *Stipulæ* deciduæ vel diu persistentes, in speciminibus visis plus quam centimetroli, foliaceæ, obtusissimæ, basi attenuatæ. Cymæ fem. in singulis axillis geminæ, sessiles vel breviter pedunculatæ, vix amplæ, ramulis brevibus, glomerulis subconfluentibus, floribus sessilibus, perigonii fructiferi segmentis late ovatis ac fere æquilongis achænio  $1\frac{1}{2}$  millim. longo obscure fusciscento multo brevioribus. — Stipulis foliaceis insignis. Cl. Blumius præter typum sub nomine *P. angulatæ* in Mus. lugd. bat. inscriptum quatuor enumerat varietates: priores (scilicet  $\beta$  semipellucidam,  $\gamma$  stipulosam et  $\delta$  obscuram) foliis amplis pellucidisque præditas, posteriorem ( $\epsilon$  minorem) foliis multo minoribus opacis et nonnihil cordatis. (v. s.)
68. *P. GRIFFITHII* (Bl. Mus. lugd. bat. 2. p. 55), herbacea vel suffruticosa, foliis longe petiolatis ellipticis vel elliptico-oblongis obtuse acuminatis basi rotundatis obtusis vel subcordatis trinerviis (nervis basilaribus fere ad apicem folii excurrentibus) acute serratis utrinque glabris, stipulis deciduis, cymis pedunculatis subgeminis folio brevioribus.  $\frac{2}{4}$  In Malacca (Griffith). — Wedd. Monogr. p. 231. — Folia 6-9 centim. longa, 3-4 $\frac{1}{2}$  centim. lata, petiolo  $1\frac{1}{2}$ -3 $\frac{1}{2}$  centimetroli, lamina exsiccata præsertim subtus cystolithis inspersa (Descript. ex Blume). — Species a præcedenti dignoscenda, ait cl. auctor, nervis basilaribus foliorum usque ad apicem limbi productis nec non stipulis fugacibus.
69. *P. PELTATA* (F. Hance, in Ann. sc. nat. 5<sup>e</sup> sér. 5. p. 242), dioica, glaberrima, foliis orbiculari-ovatis peltatis basi rotundatis apice acutis in triente inferiore integris cæterum crenato-serratis trinerviis imparibus, stipulis deciduis, cymis fem. laxis multifloris pedunculis petiolo brevioribus, achænio oblique ovato lævi.  $\frac{2}{4}$ ? In cavernulis, ad Kai-kun-shek prov. Canton sinensium (Sampson). Folia  $1\frac{1}{2}$ -3 centim. longa et lata, petiolo lamina longiore breviorive suffulta, lamina exsiccata cystolithis densis linearibus subtus præsertim conspicuis obsita. *Stipulæ* non visæ. Perigonium achænio dimidio brevius. — Species foliis peltatis insignis atque (teste cl. auctore) aspectu exemplaribus parvis R. bracteosæ non absimilis.
70. *P. PETIOLARIS* (Blume, Mus. lugd. bat. 2. p. 53. fig. 18), monoica, glaberrima, foliis amplis longe petiolatis elliptico-vel ovato-oblongis acuminatis basi rotundatis vel obtusis trinerviis grosse dentato-serratis, stipulis lanceolatis foliaceis deciduis, cymis unisexualibus divaricatis subsessilibus quam petioli multo brevioribus, perigonio fem. cupulari-tridentato, achænio scabriusculo.  $\frac{2}{4}$ ? In Japonia (Siebold, Gering; Zollinger, n. 1826). — Wedd. Monogr. p. 232. *Urtica petiolaris* Siebold et Zuccar. in Münch. Abhandl. der math. physik. Kl. der bai. Akad. 4 (3). p. 214; Walp. Ann. bot. 1. p. 643. — Caulis erectus, 3-4 decim. altus, succulentus, obtuse angulatus, merithallis inferiori

(1) Vid. quoque *B. quadrifoliam* inter Rubescentes.

ribus elongatis, superne tantum foliatus. Folia 7-16 centim. longa, 3-8 centim. lata, acumine lanceolato, ima basi integra, serraturis reliqui limbi obtusis, lamina exsiccata tenuiter membranacea læte viridi cystolithis linearibus delicatulis utrinque aequaliter dispersa, petiolis 3-8-centimetalibus. Cymæ masc. in axillis inferioribus enatis (in planta fructifera deficientibus) divaricataramosis, glomerulis subdiscretis, perigonii segmentis sub apice breviter mucronatis. Cymæ fem. in axillis superioribus nascentes, masculis minores, congestifloræ, perigonio fructifero cupuliformi brevissime tridentato quam achænium ovatum sesquimillimetrale ventricosum et acute marginatum plusquam dimidio brevior. — Folia a Zuccarinio alterna dicuntur sed verisimiliter propter lapsum fortuitum nonnullorum horumce. (v. s. in hb. Franqueville.)

1. *P. BOEHMERIOIDES* (Wedd. Monogr. p. 232), dioica, glaberrima, foliis amplis haud verticillatis ovatis vel ellipticis acuminatis basi obtusatis rotundatisve basi excepta sinuato-denticulatis serrulatisve triplinerviis, stipulis oblongis deciduis, cymis sessilibus quam petioli multo brevioribus, masculis capituliformibus, femineis laxiusculis, achæniis..... 4? In ins. mascarenis? (Du Petit Thouars). — Pileæ lucenti habitu valde affinis. Folia foliis ejusdem æquimagna sed basi minus rotundata, acumine acutiusculo plerumque laxè serrulato, denticulis limbi obtusiusculis, nervis basilaribus ad acuminis basim tantum protensis, lamina exsiccata membranacea subtus pallida cystolithis fusiformibus valde inæqualibus in pagina inferiore quam in superiore crebrioribus et in illius nervis majoribus obsita, petiolis gracilibus 2-6 centim. longis. Stipulæ 5-8-millimetrales, jugi communis basi connatæ, obtusæ. Cymæ masc. congestifloræ Boehmeriarum quarundam cymas mentientes, in axillis plurimis solitariae, illis ejusdem nodi subconfluentibus, floribus pedicellatis sessilibusve, perigonii lobis sub apice breviter mucronatis, mucronibus acutis in alabastro bimillimetrali discretis. Cymæ fem. geminæ, vix multifloræ, ramulis brevibus, glomerulis approximatis. (v. s. in hb. Mus. par.)
72. *P. CERATOMERA*, dioica, glaberrima, foliis late ovatis acuminatis basi obtusatis rotundatisve supra basim crenato-serratis tri-tripli-nerviis, stipulis majusculis ellipticis deciduis, cymis fem. quam petioli multo brevioribus sessilibus capituliformibus, segmentis perigonii sub apice longiuscule cornuto-appendiculatis, achænium ovatum læve superantibus. 4 In monte Clarence, ins. Fernando-Po, Africæ occidentalis, altit. 1200 metr. (Mann, n. 626). Habitus nonnihil præcedentis. Caulis erectus, 3-6-decimetralis, simplex, basi nudus, internodiis elongatis. Folia 4-6 cent. longa, 3-5 centim. lata, acumine haud centimetro integrali integro, serraturis obtusis, nervis basilaribus supra medium limbum in margine desinentibus nec unquam basim acuminis attingentibus, lamina exsiccata membranacea concolore cystolithis fusiformibus parum conspicuis utrinque inspersa, petiolis gracilibus 2-4 centim. longis altero cujusque jugi longiore. Stipulæ centimetrum longæ, apice rotundatæ, in sicco fusæ. Cymæ fem. (solum visæ) in nodis singulis confluentes, subsphæricæ, floribus breviter pedicellatis et valde congestis, segmentis perigonii ellipticis, intermedio nonnihil longiore, omnibus sub apice appendice erecto submillimetrali instructis. Achænium fere 2 millim. longum, pallidum, margine acuto, segmentis perigonii fere omnino obtectum. — Stirps admodum insignis foliolis perigonii fem. appendiculatis (v. s. in hb. Kew.)
73. *P. VERBASCIFOLIA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186; Monogr. p. 233), monoica vel dioica?, glaberrima, foliis magnis obovatis acuminatis (acumine acutiusculo) basi longiuscule attenuatis obtusatisque supra trientem inferiorem limbi obtuse inæqualiterque serratis triplinerviis, stipulis oblongis deciduis, cymis unisexualibus androgynise subsessilibus ramosissimis floribundis petiolos nonnihil superantibus, achænio sub lente minutissime punctulato. 4 In sylvis obscuris, ad ripas amnium insulæ Mauritiæ (Commerçon, Bojer, Boivin etc.). Parietaria Huetia Commers. mscr. Urtica verbascifolia Poir. Encycl. 4. p. 641. U. longifolia Willd. Sp. pl. 4. p. 348; Smith in Rees' Cyclop. 37. Urtica n. 4. — Suffrutescens, caule erecto subsimplice fere sexangulo crasso, exsiccato cystolithis punctato-striolato, internodiis elongatis. Folia pauca, 8-20 centim. longa, 5-8 centim. lata (acumine brevi triangulari), deorsum subabrupte angustata, nervis basilaribus a petioli conjunc-



tione cum lamina remotiusculis et usque ad basim acuminis productis subtus prominentibus, lamina exsiccata membranacea utrinque cystolithis punctiformibus linearibusque creberrimis (in pagina inferiore majoribus sed in in ervenio hic et illic deficientibus) obsita, petiolo valido raro centimetrum longo et interdum subnullo. Stipulae 1-1½ centim. longae, obtusae, interdum subspathulatae. Cymae in axillis pluribus ortae, saepe unisexuales, ad originem nervorum basilarium folii haud pertingentes, subcorymbosae, ramulis patulo-divaricatis, glomerulis approximatis; floribus breviter pedicellatis, masculorum perigonii lobis (in vivo rubris) sub apice longe acuteque mucronatis, alabastro 3-millimetrali; femineis breviter pedicellatis perigonii fructiferi segmentis aequimagnis, intermedio cucullato quam achenium millimetrale paulo brevior. (v. s. in hb. Mus. par.)

74. *P. CUNEIFORMIS* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 186; Monogr. p. 234), monoica, glaberrima, foliis obovato-cuneatis ellipticisve obtusis basi acutis in dimidio superiore obtuse serratis subtriplinerviis, stipulis ovatis vix persistentibus, cymis unisexualibus paucifloris sessilibus petiolos vix superantibus. ½ In locis umbrosis montuosisque Mauritii (Commerson, Sieber, n. 129, Bojer, etc.). — Bl. Mus. bot. lugd. bat. 2. p. 47. *Urtica cuneiformis* Poir. Encycl. 4. p. 640. — Suffrutescens, caule basi ramoso radicante, ramis subsimplicibus adscendentibus 1-2-decimetralibus teretiusculis foliosis basi denudatis siccis cystolithiferis, internodiis brevibus. Folia 1-4 centim. longa, 3-16 millim. lata, nervis basilaribus rectiusculis fere usque ad apicem limbi productis, laminae exsiccatae membranaceae pagina superiore cystolithis ut in ipso caule linearibus vel mixtis sparsim suffusa, inferiore isdem tantum in margine et secus nervos ornata. petiolo 1-4-millimetrali. Stipulae bimillimetrales, obtusissimae. Cymae masc. in axillis pluribus ramorum umbelluliformes, floribus longiuscule pedicellatis, perigonii lobis sub apice mucrone conico instructis, alabastro bimillimetrali siccis cystolithis apice striolatis. Cymae fem. glomeratae, floribus breviter pedicellatis, achenio ovato laevi perigonii segmentum intermedium nonnihil superante. (v. s. in hb. Mus. par.)

\*\* Pubescentes, sive caule foliisque puberulis pubescentibus vel pilosis.

- A. — *Longipedunculatae, sive pedunculis (saltem cymarum fructiferarum) quam petioli plerumque longioribus.*

75. *P. UMBROSA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 187; Monogr. p. 243), monoica vel dioica, caule foliisque et ramis cymarum plus minus pubescentibus interdumque pilis mollibus dense hirsutis, foliis vulgo amplis late ellipticis ovatisve saepe inaequaliter subabrupte et anguste acuminatis basi acutis vel obtusatis rotundatisve aut subcoarctatis grosse serratis (serraturis acutis vel obtusis) trinerviis, stipulis oblongis ovatisve persistentibus, cymis compositis floribundis; pedunculis quam petioli longioribus, achenio parvo laeviusculo. ¼ In regionibus temperatis subtropicisque Himalayae, prov. Sikkim et Marri frequentissima, altit. 1200-2700 metr. (Wallich, etc.), necnon in montibus Khasiae altit. 900-1800 metr. (J. D. Hooker et Thomson). — *Urtica umbrosa* Wallich, Cat. n. 4598. A. — Herba, caulibus 2-6-decimetralibus erectis adscendentibusve basi interdum lignescens radicanibusque parce ramosis crassitie pennae anserinae aut gracilioribus angulatis, merithallis elongatis, nodis inferioribus saepe aphyllis, epidermide exsiccata ut plurimum cystolithos parvaucos exhibente. Folia 7-20 centim. longa, 4-13 centim. lata, nonnunquam minora, acumine lineari vel lanceolato integro aut serrato, serraturis limbi margine exteriori convexis, prope basim folii multo minoribus juxtaque petiolum saepe deficientibus, nervis basilaribus arcuatis fere usque ad acumen saepe perductis sed interdum juxta trientem superiorem limbi cum secundariis arcuatis anastomosantibus, lamina exsiccata membranacea cystolithis tenuissimis (haud raro inconspicuis) in utraque pagina conspersa, petiolis in eodem jugo solito more inaequalibus modo glabratis modo pubescentibus vel densiuscule patentimque pilosis. Stipulae longitudine varia at centimetrum raro excedentes, acutae, siccitate

scarioso-ferrugineæ. Cymæ diversa amplitudine, complanato-pyramidatae-ramulis patentibus et pedunculis (petiolum nonnunquam vix æquantibus pubescentibus v. glabratis, glomerulis in ramis secundariis subspicatis confluentibusque; floribus masc. breviter pedicellatis, perigonii lobis (interdum pubescentibus) sub apice brevissime mucronatis, mucronibus concis in alabastro uni-sesqui-millimetrali conniventibus; perigonii fructiferi segmento intermedio cucullato lateralibus multo longiore achænioque albicante ( $\frac{1}{2}$  millim. circiter longo) fere tertia parte brevior. (v. s.)

$\beta$ , *obesa*, undique glabriuscula.  $\frac{1}{2}$  In Indiæ orientalis provincia Nepalia, etc. — P. Wallichii Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 166. *Urtica obesa* Wall. Cat. n. 4587. (v. s.)

76. *P. LEUCOPHÆA* (Blume, Mus. bot. lugd. bat. 2. p. 53), suffruticosa, scandens, foliis longe petiolatis ovato-vel elliptico-oblongis acuminatis basi oblique attenuatis rarissime obtusis trinerviis grosse dentato-serratis ciliatis utrinque atrigoso-puberulis subtus cinerascens, cymis longe pedunculatis.  $\frac{1}{2}$  In ins. Java. — Wedd. Monogr. p. 245. *Urtica leucophæa* Blume, Bijdr. p. 493. — Folia (in eodem jugo inæquimagna) 4-15 centim. longa, 2-6 centim. lata, petiolo 1½-7 centim. longo. (Descript. ex Blume, l. c.)

77. *P. BRACTEOSA* (Wedd. Monogr. p. 245), dioica, pubescens vel glabriuscula, foliis suboblique ovatis lanceolatisve anguste et interdum abrupte acuminatis basi obtusatis aut subcordatis et integris cæterum serratis (serraturis parvis vel grossiusculis) acutis acuminatisve trinerviis supra glabris subtus in nervis præsertim subtiliter pubescentibus, stipulis majusculis ovatis lanceolatisve scariosis, cymis multifloris, pedunculis quam petioli longioribus, ramis muscularum basi ample bracteatis, achænio parce granuloso.  $\frac{1}{2}$  In Indiæ orientalis montibus Khasiæ, reg. subtrop. altit. 900-1860 metr. et in prov. Sikkim altit. 1500-1800 metr. (Hooker et Thomson). — Habitu valde propinqua varietati glabrescenti *P. umbrosæ*. Folia plerumque minus lata interdumque lanceolata ac sæpe valde inæquilateralia, in speciminibus examinatis 3-12 centim. longa, 1-6 centim. lata, acumine lineari utrinque serraturis 1-3 instructo, dentibus limbi margine exteriori convexi rectiusculisve, nervis basilariibus ut plurimum usque ad basim acuminis protensis, secundariis transverse subparallelis, pagina superiore ut videtur fere semper glaberrima, inferiore contra persæpe magis minusve pubescente, lamina exsiccata membranacea utrinque cystolithis minimis breviter linearibus punctiformibusve obsita, petiolis inæqualibus 1-5 centim. longis glabris villosisve. Stipulæ subcentimetales, siccitate ferrugineæ. Cymæ in axillis singulis solitariae, distiche et subsimpliciter ramosæ, glomerulis nempe in ramulis primariis plerumque sessilibus approximatis et subspicatis; masculæ ad basim ramorum glomerulorumque bracteis ovatis 1-3-millimetralibus siccitate ferrugineis postice munitæ, densifloræ, floribus breviter pedicellatis, perigonii glabri lobis sub apice conspicue mucronatis, alabastro subsesquimillimetrali. Cymæ fem. inconspicue bracteate, floribus, subsessilibus, perigonii fructiferi segmento intermedio oblongo breviter cucullato s. gibboso. Achæmium millimetrale, utrinque minute granuloso et juxta marginem lævem acutumque linea elevata insuper notatum (v. s. in hb. Kew. et Mus. par.)

78. *P. SYMMERIA* (Wedd. Monogr. p. 246), dioica, foliis ovato-lanceolatis lanceolatisve interdum inæquilateralibus anguste acuminatis basi emarginatis integrisque cæterum acute serratis vel serrulatis trinerviis supra sparsim breviterque pilosis subtus glabris, stipulis brevibus triangulari-ovatis subpersistentibus, cymis multifloris pedunculis quam petioli longioribus, perigonio fructifero fere cupuliformi. — In regione temperata montium Sikkim Indiæ orientalis, altit. 1800-2700 metr. (J. D. Hooker). Caulis adscendens, angulatus, crassitie circiter pennæ corvinæ, glaberrimus, internodiis inferioribus elongatis, nodis fere omnibus foliatis. Folia in speciminibus quæ examinare contigit 5-10 centim. longa, 1½-4 centim. lata, acumine lineari, serraturis margine exteriori convexi subapiculatis, nervis basilariibus ad basim acuminis productis, secundariis pluribus subtransversalibus et fere parallelis, lamina exsiccata membranacea cystolithis brevissimis ac parum conspicuis utrinque inspersa, petiolis in eodem jugo parum inæqualibus 1-

2-centimetalibus. Stipulae 1-3 millim. longae. Cymae erectiusculae, cum pedunculis foliis subaequilongae, breviter ramulosae, floribus masc. laxe glomeratis (glomerulis approximatis) bracteolis minimis stipatis, perigonii lobis sub apice brevissime mucronatis, alabastro sesqui-bimillimetrali; floribus fem. in glomerulos parvos subdiscretos digestis, perigonii segmentis in cupulam brevem carnosulam achenium ovatum sesquimillimetrale laeviusculum suffulcientem coalitis, intermedio longiore. — Stirps aliquatenus ambigua P. trinerviam, P. bracteosam et P. umbrosam simul referens, ab illis autem forma perigonii fem. optime distincta. A P. petiolarum ad quam hocce characterē precipue accedit, ceterum longe differt. (v. s. in hb. Kew. et Mus. par.)

B. — *Brevipedunculata*, sive *pedunculis* (saltem *gynaeorum fructiferarum*) *quam petioli plerumque brevioribus*. (1).

79. P. *MONGOLICA*, monoica v. dioica, foliis ovatis acuminatis basi late cuneata integris ceterum obtuse serratis utrinque sparsim breviterque pilosis v. subtus glabris, stipulis brevissimis inconspicuis, cymis et glomerulis paucissimis constantibus quam petioli multoties brevioribus (1) 2. ad rivatos, prope Gehol, in Mongolia orientali (Rev. P. David, n. 2043). — Caulis erectus, 2-4-decimetralis, crassitie pennae anserinae, angulatus, glaber, in vivo fere translucentis, exsiccatus virens, internodiis inferioribus valde elongatis. Folia superiora 8-4 centim. longa, 2-3 centim. lata, acumine oblongo obtuso, basi obtusa, inferiora minora haud acuminata basi rotundata paucicrenata; nervis basilariibus tenuibus dimidium limbum superantibus, secundariis e costa adscendentibus; lamina exsiccata tenuiter membranacea viridi cystolithis linearibus sparsis inpersa; petioli parum inaequalibus 1-3 centim. longis gracilibus ad insertionem limbi pilosis. Stipulae absque lente fere omnino inconspicuae. Cymae unisexuales aut androgynae, sessiles, conglomeratae aut breviter patuleque pauci-ramulosae, glabrae, floribus breviter pedicellatis sessilibusve; perigonii masc. segmentis sub apice brevissime mucronatis. Achenium ovatum, 1-1½ millim. longum, laeve, perigonii segmenta lineari-oblonga subaequalia (v. intermedio nonnihil longiore) sub apice breviter cucullata haud dimidia longitudine superans. (v. s. in hb. Mus. par.)

80. P. *HYGROPHILA* (Blume, Mus. bot. Lugd. bat. 2. p. 56), dioica, foliis ellipticis ovatisve acuminatis ultra basin attenuatam vel rotundatam acute aut obtuse serratis trinerviis supra pilis crassis raris inpersis subtus glabriusculis, stipulis majusculis ovato-lanceolatis persistentibus, cymis geminis contractis quam petioli brevioribus, achenio laevi. 2. In Indiae orientalis montibus Nilghiri (Wight) et Khasia, altit. 1500-1800 metr. (J. D. Hooker et Thomson), in Zeylaniam (Gardner) et Java (Junghuhn). — Wedd. Monogr. p. 164; Miq. Fl. Ind. bat. 1 (2). p. 237. Urtica hygrophila Miq. Pl. Jungh. p. 28. Caulis ascendens, basi saepe longiuscule denudatus et non raro radicans, 2-5-decimetralis, angulatus, crassitie pennae anserinae vel gracillior, in specimenibus omnibus visis glaberrimus, internodiis in planta robustiore elongatis. Folia 3½-8 centim. longa, 2½-4 centim. lata, breviter acuminata, basi saepius integra, serraturis margine exteriori convexiusculis, nervis basilariibus usque ad tridentem superiorem limbi circiter productis subtus interdum puberulis, secundariis tenuibus parallele subtransversis, lamina rigida supra nigricante utrinque cystolithis minimis obsita, petioli in eodem jugo valde inaequalibus 1-4 centim. longis gracilibus glabris. Stipulae 3-5 millim. longae, emaridiae, ferrugineae. Cymae breviter saepeque brevissime pedunculatae, modo corymbosae, modo fere racemiformes vel congestae et fasciculiformes, alternatim (nec distiche) ramosae, ut plurimum glaberrimae; floribus masc. brevissime pedicellatis, perigonii segmentis sub apice brevissime mucronatis; floribus fem. saepius confertis, perigonii fructiferi segmentis ovatis subaequimagnis vel intermedio longiore inconspicueque aut distincte

(1) Vid. quoque P. trinerviam et P. Hookerianam inter Glabras.

cucullato. Achænium ovatum, vix millimetrale, sordide fuscescens, perigonio paulo longius. (v. s. in hb. h. b. lugd.-bat. Wight, Kew et Mus. par.)

81. *P. PRODUCTA* (Blume l. c.), monoica vel dioica, foliis ovatis vel ovato-oblongis acuminatis vel cuspidatis basi in petiolum longum productis trinerviis grosse dentato-serratis utrinque pubescentibus et in petiolis nervisque paginæ inferioris subtomentosis, cymis breviter pedunculatis divaricatis. 2/ In montibus Himalayæ, Indiæ orientalis. — Wedd. Monogr. p. 265. Folia 4-18 centim. longa, 2-7 centim. lata, petiolo  $1\frac{1}{2}$  -  $3\frac{1}{2}$  c. ntim. longo. (Descript. ex Blume). — A *P. umbrosa* foliis in petiolum attenuatis et cymis brevius pedunculatis distinguenda sec. cl. auctorem.
82. *P. RIVULARIS* (Wedd. Monogr. p. 266), dioica, foliis late ovatis sensim acuminatis acutisque præter basim rotundatam subcordatamve grosse serratis (serraturis acutis vel obtusis) trinerviis in pagina superiore sparsissime pilosis, stipulis majusculis rotundato-ovatis amplexicaulis persistentibus, floribus masc. fasciculatis, fasciculis paucifloris quam petioli multo brevioribus stipularumque ope involucretis, segmentis perigonii sub apice longe mucronatis. 2/ In sylvis secus rivulos montium insulæ Anjouan, prope Madagascariam (Boivin). — Herba, caule elongato parce ramoso, ramis basi prostratis radicanibusque demum adscendentibus teretiusculis, exsiccatis cystolithis fusiformibus creberrime punctulatis albicantibusque, internodiis 2-5 centim. longis. Folia 2-5 centim. longa,  $1\frac{1}{2}$  -  $3\frac{1}{2}$  lata, serraturis margine exteriori convexis, dente ultimo lanceolato 5-7 millim. longo, nervis basilaribus medium limbum paulo superantibus cum secundariis pariter tenuissimis arcuatum anastomosantibus, lamina exsiccata tenuiter membranacea pallide viridi cystolithis linearibus utrinque sparsim ornata, petiolis gracilibus in eodem jugo inæquilongis glabris sæpiusve ad limbum parvis pilosis. Stipulæ 5-7-millimetralis, obtusissimæ, fere hyalinæ, altera alteram plus minus involvente. Fasciculi masc. densiusculi, plerumque stipulis breviores hisque velati, floribus breviter pedicellatis, perigonii mucronibus segmenta ipsa longitudine fere æquantibus, alabastro elliptico mucrone incluso circiter 3-millimetrali. Inflorescentia feminea ignota. — A sequente forsitan non specificè distincta. Discrepat præsertim foliis et habitu. (v. s. in h. Mus. par.)
83. *TETRAPHYLLA* (Bl. Mus. bot. lugd. bat. 2. p. 50), dioica, humilis, foliis nunc (in planta fem.) paucis ac plerisque in apice caulis subverticillatis, nunc (in pl. masc.) numerosis jugis discretis, late ovatis obtusis acutisve basi vulgo cuneatis crenato-serratis trinerviis, duobus inferioribus sæpe minoribus intergerrimis, omnibus sparsim pilosis aut rarius glabris, stipulis conspicuis rotundatis, cymis utriusque sexus congestifloris brevissime pedunculatis sessilibusve, segmentis perigonii masc. longe aristatis. ① In Abyssiniæ montis Serraba, locis humidis, altit. 1950 metr. (Schimper, n. 436), necnon in montibus Cameroen Africæ occidentalis, altit. 2130 metr. (Mann.) — *P. quadrifolia* A. Rich. Tent. fl. Abyss. 2. p. 263; Wedd. Monogr. p. 199. t. 7. f. 4-10; J. D. Hooker. in Journ. Linn. Soc. 7. p. 216. *Urtica tetraphylla* Hochstett. in Bot. Zeit. 33. p. 260. — Caulis filiformis, 4-20 centim. altus, erectiusculus v. basi radicans, simplex v. parce ramosus, glaber, hyalinus. Folia 1-2 centim. longa, 8-15 millim. lata, basi cuneata v. obtusa, serraturis utrinque 5-8, nervis basilaribus medium limbum vix attingentibus, lamina exsiccata tenuiter membranacea subconcolore utraque pagina cystolithis tenuissimis parumque conspicuis inspersa, petiolo vix centimetrali. Stipulæ 2 millim. longæ et latæ, scariosæ. Cymæ masc. in axillis pluribus dense glomeratæ, floribus majusculis 3-5-meris (teste J. D. Hooker), segmentis perigonii 1. v. pluribus longe aristatis. Cymæ fem. in apice caulis simplicis inter folia congestæ, floribus ut plurimum brevissime pedicellatis, perigonio 4-partito, fructiferi segmento uno cucullato cæteris subæqualibus multoties longiore achæniisque vix semimillimetrali subgranulato subæquilongio. (v. s. in h. Mus. par. et Kew.).

## 2° Neogææ.

\* Glabræ, sive caule foliisque pilis omnino destitutis.

A. — *Longipedunculatæ, sive pedunculis (saltem cymarum fructiferarum) quam petioli plerumque longioribus* (1).

a. Stipulæ breves triangulares (2).

84. P. HILARIANA (Wedd. Monogr. p. 210), dioica, glaberrima, foliis ovatis acuminatis basi obtusis supra trientem inferiorem serratis (serraturis acuminatis) subtriplinerviis, stipulis triangularibus brevibus persistentibus, cymis parvis et glomerulis 1-4 constantibus, pedunculis petiolos superantibus, achænio quam perigonium multo longiore.  $\frac{1}{2}$  In Brasilia meridionali (A. St.-Hilaire). — Herba, caulibus 2-3 decim. longis basi longe radicantibus demum adscendentibus gracilibus parce ramosis foliosis. Folia 1 - 1  $\frac{1}{2}$  centim. longa, 7-10 millim. lata, a medio ad apicem sensim acuminata, serraturis utrinque 4-6, nervis basilaribus ad trientem superiorem limbi protractis haud prominentibus, laminæ exsiccatae membranaceæ rufescentis pagina superiore cystolithis linearibus delicatulis sparsim ornata, inferiore lisdem ut videtur destituta, petiolo gracili 3-8 millim. longo. Stipulæ parum conspicuæ, obtusiusculæ. Cymæ fem. foliis subæquilongis iisve breviores, glomerulis approximatis, floribus subsessilibus, perigonii fructiferi segmento intermedio quam lateralia paulo longiore achænioque ovato millimetrali lævi dimidio brevior. Affinis P. rupipendix. (v. s. in hb. Mus. par.)

85. P. ANOMALA (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 217; Monogr. p. 207), monoica vel dioica, glabra, foliis oblongo-lanceolatis acuminatis (acumine acuto) basi acutis rotundatisve supra partem quartam inferiorem ad apicem usque argute serratis trinerviis, stipulis minimis triangularibus, cymis laxis unisexualibus, pedunculis petiolo longioribus.  $\frac{1}{2}$  In locis humidis umbrosisque sylvarum Boliviz, prov. Enquisivi et Larecaja (Wedd.; Mandon n. 1104.) — Herba, caule 4-8 decim. alto descendente basi que radicante angulato crassitie pennæ anserinæ fistuloso inferne longo nudo, exsiccato cystolithis punctiformibus inconspicue punctulato obscure fusco. Folia 4-12 centim. longa, 2-4 centim. lata, in medio vel in triente inferiore attenuata aut rotundata et ante insertionem petioli subemarginata, acumine angusto serrato; nervis basilaribus sæpe ad acumen usque productis subtus parum prominentibus, laminæ exsiccatae firmulæ subtus pallidioris utraque pagina cystolithis inæqualibus subtilissimis punctiformibus v. fusiformibus (oculo nudo prorsus inconspicuis) conspersa, petiolo gracili 1  $\frac{1}{2}$ -3 centim. longo dessiccatione cystolithis punctulato. Stipulæ a basi lata amplexicauli abruptissime attenuatæ. Cymæ masc. pyramidatae, foliis longiores, laxifloræ, ramulis patulis, glomerulis subdiscretis, floribus brevissimis pedicellatis, alabastris 1  $\frac{1}{2}$  millim. longis, muticis. Cymæ fem. masculis multo minores, e glomerulis paucis approximatis vel subdiscretis constantes, foliis breviores, pedunculo breviter ramoso, perigonii fructiferi segmento intermedio haud cucullato laterales paulo excedente achænioque ovato sesquimillimetrali nitidulo dimidia parte vel triente brevior (v. v.)

β, dentibus foliorum numerosioribus obtusis muticis apiculatisve. (v. s. in hb. Mus. par.)

86. P. SUBSERRATA (Wedd. Monogr. p. 207), dioica, glaberrima, foliis elliptico-ovatis acuminatis (acumine acuto) basi rotundatis emarginatisque in acumine tantum vel prope apicem argute serratis trinerviis, stipulis brevibus triangularibus, cymis masc. laxis, pedunculis quam petioli vix longioribus, feminis. ...  $\frac{1}{2}$  In Nova Granata (Triana, n. 344). — Caulis herbaceus, 3-4 decim. altus, subsimplex, modice crassus, angulatus, inferne denudatus, merithallis

(1) Vid. quoque P. Mutisianam et P. subcrenatam inter Brevipedunculatas.

(2) Vid. quoque P. dauciodoram et P. serrulatam inter species stipulis vulgo majusculis.

longæ, pedunculis parum ramosis, glomerulis discretis, floribus subsessilibus, alabastris 1½ millim. longis, mucronibus subapicalibus distinctis conicis. Cymæ fem. in axillis inferioribus sive segregatim sive cum masc. simul oræ, solitariæ aut geminæ, plerumque foliis etiamque petiolo ipso breviores, glomerulis subsessilibus approximatis, perigonii segmentis dentatis fere æquilongis, intermedio cucullato. Achænium angustissime marginatum, læve. (v. s. in hb. Mus. par. et Kew.)

91. *P. ELLIPTICA* (Hook. fil. Fl. antarct. p. 344), monoica, glabra, foliis ellipticis v. late ovatis utrinque attenuatis obtusisque rarissime subacuminatis in toto margine crenato-serratis (dentibus apiculatis) trinerviis, stipulis late ovatis persistentibus, cymis unisexualibus e glomerulo unico constantibus, pedunculis quam petioli vulgo longioribus iisque interdum æqualibus brevioribusve, achænio orbiculari. ♀ In humidis prov. chilensium Valdivia, Chiloe et Concepcion (Dombey, Gay, Hooker, fil. etc.). — Gay, Fl. chil. 5. p. 364; Wedd. Monogr. p. 211. *P. chilensis* ejusd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 221. *P. pedunculata* Blume Mus. lugd. bat. 2. p. 46. — Herba, caulibus elongatis repentibus demum adscendentibus parce ramosis subsimplicibusve basi sæpe nudis, internodiis inferioribus elongatis, exsiccatis obscure rubentibus cystolithisque parvis oblongis sparsim obsitis. Folia apud typum 8-25 millim. longa, 5-20 millim. lata, in var. ♂ multo majora (nempe 6-8 centim. longa), nervis basilariibus longitudine varia trientem medium limbi plerumque superantibus, petiolis gracilibus in eodem jugo maxime disparibus limbum æquantibus aut multo brevioribus. Cymulæ solitariæ vel geminæ, masculæ superiores; floribus masc. pedicellatis, perigonii lobis sub apice brevissime mucronatis, alabastro bimillimetrali: floribus fem. subsessilibus; perigonii fructiferi segmento intermedio lateralibus dimidio longiore achænioque subsæquimillimetrali et sulco superficiali longitudinali notato paulo brevior. (v. s.)

♂, *Gayana*, foliis majoribus quam in typo, oblongo-vel ovato-lanceolatis subacuminatis, acumine obtusiusculo. ♀ In locis humidis provinciæ chilensis Valdivia (Gay, etc.) *P. elegans* Gay, l. c. non Wedd. (v. s. in hb. Mus. par.)

92. *P. SERRULATA* (Wedd. in Ann. sc. nat. l. c. p. 213; Monogr. p. 212), monoica vel dioica, glabra, foliis parvis oblongo-vel obovato-lanceolatis acutis subacuminatisve basi longe cuneato-attenuatis ultra medium vel trientem inferiorem serrulatis penninerviis, stipulis foliaceis majusculis cordato-reniformibus persistentibus, cymis minimis paucifloris in planta monoica androgynis, pedunculis petiolos superantibus. ♀ In ins. Jamaica (Swartz). — *P. lucida* γ serrulata Griseb. Fl. Br. W. Ind. p. 156. *Urtica serrulata* Swartz, in Act. Holm. ann. 1787. p. 65; ejusd. Fl. Ind. occid. p. 313; Smith, l. c. n. 47. Fruticulosa, caule erecto ramoso circiter bidecimetrali angulato, ramis virgatis inferne denudatis, internodiis brevissimis. Folia 1½-2 centim. longa, 3-5 millim. lata, dentibus obtusis acutisve, lamina exsiccata supra cystolithis linearibus intricatis ornata subtus iisdem corpusculis multo brevioribus conspersa, petiolo brevissimo. Stipulæ conspicuæ, caule multo latiores, subsistentes. Cymæ e glomerulo unico constantes, pedunculo filiformi 6-8 millim. longo, floribus subsessilibus, perigonii masc. segmentis sub apice breviter mucronatis, mucronibus obtusis discretis. (v. s. typ. in hb. Holm. Mus. brit. et Soc. Linn.)

♂, *Cubensis*, dioica, caule superne puberulo, foliis majoribus quam in typo lineari-vel oblongo-vel obovato-lanceolatis ultra medium inæqualiter serrato-vel sinuato-dentatis, stipulis late triangularibus acuminatis, cymis ramosis multifloris, femineis laxis quam masculæ densioribus. ♀ In ins. Cubæ monte Libano, prope Santiago (Linden, n. 1839) et in vicinis villæ Monte Verde (Wright n. 1448). *P. Cubensis* Wedd. in Ann. sc. nat. l. c. p. 214. *Urtica Lindeniana* A. Rich. in Ram. de la Sagra, Hist. ins. Cub. Bot. 2. p. 225. *P. lucida* var. *cuneifolia* Griseb. Cat. pl. cub. p. 59. — Caulis elongatus, procumbens vel diffusus, exsiccatus cystolithis punctiformibus creberrime obsitis. Folia 1½-4 centim. longa, exsiccata firmula, nervis parum conspicuis, pagina superiore cystolithis

elongatis inæqualibus valdeque intricatis eximie reticulata, illis paginæ inferioris nisi in margine et costa paucioribus. Stipulæ basi late amplexicaules, abrupte acuminatæ. Cymæ masc. in axillis superioribus solitariae, glomerulis paucis pisiformibus; femineæ patule ramosæ, glomerulis parvis discretis, perigonii segmento intermedio cucullato ovario segmentisque lateralibus subæquilongis. (v. s.)

γ, *major*, foliis duplo vel triplo majoribus quam in varietate præcedente fere in toto margine grossiuscule sinuato-dentatis, nervis in lamina exsiccata sat conspicuis, cystolithis paginæ inferioris brevioribus crebrioribusque quam in superiore, stipulis ovatis denticulatis laciniatisve, cat. ut in var. præced. 2/ In ins. Cuba (Wright, n. 2236). — P. uninervis Griseb. Cat. pl. cub. p. 59.

93. P. *RHEIXIODES* (Lieb: l. c. p. 304), dioica, glaberrima, foliis elliptico-subfalcatis acuminatis acumine obtusata basi cuneatis vel leviter emarginatis remote et obtuse serratis trinerviis, stipulis ovatis deciduis, cymis masc. paucifloris, pedunculis petiolo subæquilongis, femineis. ... 2/ In imperii mexicani prov. Puebla, altit. 900 metr. (Liebmann). — Herba, caule 7-14 centim. longo a basi decumbente ascendente tetragono articulado. Folia 2 1/2 - 3 1/2 centim. longa, 18 - 23 millim. lata, supra saturate viridia, punctato-aspera, juxta marginem cystolithis linearibus transversis obsita, subtus albicantis, nervis prominulis fusco-viridibus, petiolo semicentimetrali. Stipulæ minutæ, membranaceæ. Cymæ masc. solitariae, pedunculo gracili. (Descript. ex Liebmann).

94. P. *ELEGANS* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 215; Monogr. p. 215. t. 6. f. 11, non Gay), dioica vel monoica, glabriuscula, foliis lanceolatis vel ovato-lanceolatis longe acuminatis (acumine acuto) basi acutis obtusatisve circiter a medio ad apicem serratis (serraturis subacuminatis et interdum subtiliter ciliolatis) triplinerviis, stipulis oblongis deciduis, cymis laxifloris, glomerulis masc. et fem. in planta monoica intermixtis, pedunculis folia æquantibus. 2/ In ins. Antillanis. (Griseb. Fl. Br. W. Ind. p. 157). P. *decora* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 284. P. *elongata* Bl. Mus. lugd. bot. 2. p. 55, non Mig. — Caulis herbaceus aut suffrutescens, erectus, elatus, ramosus vel subsimplex, ramis herbaceis teretiusculis aut obsolete sulcatis, exsiccatis cystolithis minutissime punctulatis, glabris vel superne puberulis, nodis plerisque foliiferis. Folia (superiora plerumque majora) 3-10 centim. longa, 1-4 centim. lata, nervis basilaribus ultra medium limbum protractis subtus prominulis, laminæ exsiccatae firmulæ pagina superiore cystolithis fusiformibus vel linearibus paucioribus numerosioribusve conspersa, inferiore nuda aut illis vix non destituta, petiolo 5-15 millim. longo. Stipulæ 10-12-millimetrales, obtusæ. Cymæ amplæ, ramulis elongatis patulis, glomerulis discretis; floribus masc. longiuscule pedicellatis, alabastris 1 1/2 millim. longis, mucronibus conniventibus obtusis. Flores fem. breviter pedicellati; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achæniumque ovatum 1/2 millim. longum læve non omnino æquante. (v. s.)

β, *latifolia*, caule robusto, foliis circiter dimidio latioribus toto margine subduplicato-serratis, serraturis acuminatis. (v. s.)

γ, *minor*, caule decimetrali, foliis anguste lanceolatis nervis basilaribus fere omnino marginalibus petiolis apice puberulis, stipulis subpersistentibus, cymis parvis densiusculis. — In ins. Portorico (Blauner) (v. s. in hb. Boiss.).

95. P. *SPATHULATA* (Griseb. Cat. pl. cub. p. 59), monoica? glabriuscula, foliis spathulatis acutiusculis in petiolum longe attenuatis margine subintegerrimis v. superne remote serratis uninerviis glabris v. ciliatis, stipulis majusculis laceris, cymis pedunculis folia adæquantibus suffultis, masc. contractis fasciculiformibus, fem. dichotomo-expansis. 2/ In ins. Cubæ reg. orientali (Wright, n. 2235); — Caulis simpliciusculus, internodiis sursum decrescentibus. Folia (incluso petiolo) 3-6 centim. longa, superne 1-1 1/2 centim. lata. — Sequenti nonnihil affinis.

96. P. *SEMIDENTATA* (Wedd. in Ann. sc. nat. l. c. p. 215; Monogr. p. 216), dioica, glaberrima, foliis lanceolatis acuminatis (acumine acuto) inferne vulgo

longe attenuatis in triente v. dimidia parte inferiore integris reliqua aequaliter sinuato v. serrato-dentatis triplinerviis, stipulis magnis ovatis persistentibus, cymis masc. longissime pedunculatis, floribus in glomerulos capituliformes congestis, femineis laxifloris et brevius pedunculatis. — Portorico, S. Thomas et Antigua (Riedlé, etc.) et in Martinica (Sieber — Griseb. Fl. Br. W. Ind. p. 157. *Urtica semidentata* Poir., Encycl. p. 222. — Herba, caule 3-4 decim. alto vel elatiore subsimplice angulato, exsiccato cystolithis crebris albidis punctulato, internodiis ribus elongatis. Folia 3-10 centim. longa,  $1\frac{1}{2}$ - $3\frac{1}{2}$  centum. lata, dentibus acuminatis integrisque vel margine exteriora denticulatis, nervis basi interdum procul a basi limbi ortis et fere usque ad apicem productis prominulis, laminae exsiccatae rigidiusculae pagina inferiore præter axem marginem et nervos ut plurimum cystolithis destituta, superiore composita corpusculis linearibus creberrime inspersa, petiolo 1-2 centim. longo. Folia centimetrum et quod excedit longæ, obtusissimæ, siccae scariosæ ferè. Cymæ in axillis superioribus solitariae; masculae pedunculo decem fultæ, ramulis patulis, glomerulos paucos multifloros distantes et pini magnitudine gerentibus, floribus pedicellatis, perigonii lobis sub apice ter mucronatis, alabastro sesquimillimetrali. Cymæ fem. folia supra ovata, pedunculis petiolo longioribus, glomerulis numerosis parvis paucis discretis, floribus breviter pedicellatis, perigonii segmento intermedio lato lacinias laterales longe excedente achæniunq. submillimetrale ælæve vix non æquante. — Habitu præcedentem refert, ab ea tamen distinguitur stipulis persistentibus et imprimis indole cymæ masculæ.

β, *major*, robustior, foliis ovato-lanceolatis, quam typi amplioribus ipsa basi serratis.

97. *P. VULCANICA* (Liebmann, l. c. p. 313), monoica, glabra, foliis ellipticis minatis basi emarginatis grosse falcato-crenatis (crenis obtusis v. apice trinerviis, stipulis deciduis, cymis petiolos superantibus. ¼ In monte vomo Orizaba imperii Mexicani, altit. 3200 metr. (Liebmann). — Monogr. p. 217. — Herba, caule  $1\frac{1}{2}$ -2 decim. longo, simplice v. rarius tetragono, articulo. Folia  $3\frac{1}{2}$ -6 centim. longa,  $1\frac{1}{2}$ -2½ centim. lata, sub supra obscure viridia, subtus glauco-olivacea, petiolo 12-15. millim. Cymæ solitariae, pedunculo 2½ centim. longo, achænio ovato margine vescente subtilissime punctulato. (Descript. ex Liebmann, l. c.)

†† Nervi basilares usque ad apicem limbi pro lucti.

98. *P. PÆPPIGIANA* (Wedd. in Ann. sc. nat. l. c. p. 225; Monogr. p. dioica, glaberrima, foliis ovato- vel oblongo-lanceolatis acuminatis (acutis) basi acutis in margine toto serrulatis (serraturis apiculatis vel petiolum integris) triplinerviis, stipulis late ovatis mox truncatis, masc. ...., femineis laxiusculis, pedunculo quam petiolum longiore. Peruvia subandina (Poeppig). Herba, caule 2-4 decim. longo, pennæ corvinæ, basi longe radicante, parce folioso, exsiccato cystolithis lato, internodiis 2-3 centim. longis. Folia  $3\frac{1}{2}$ -6½ centim. longa, 2-3 centim. lata, nervis basilaribus fere usque ad acuminis basim excurrentibus, venio venulis quibusdam transversis subparallelis distantibusque nervis latiusculis, lamina exsiccata membranacea et utrinque cystolithis in mibus delicatulis (in pagina inferiore vero minoribus crebrioribusque) in petiolo gracili dimidio limbo subæquilongo. Stipulae 1-2 millim. longæ. Femineæ (solum visæ) in axillis superioribus solitariae, ramulis patulis, glomerulis approximatis, floribus subsessilibus, perigonii fructu merito intermedio cucullato lacinias laterales dimidio longiore achænio ovato millimetrali læviusculo subæquilongo. (v. s. in hb. Mus. par.)

99. *P. GRANDIS* (Wedd. in Ann. sc. nat. l. c. p. 216; Monogr. p. 218), dioica, glabra, foliis ellipticis vel ovato-ellipticis acutis acuminatis (acumine acutiusculo) basi obtusis vel rotundatis subcordatisve basi grosse crenato-serratis trinerviis, stipulis majusculis ovatis deciduis, late multifloris, pedunculis quam petioli longioribus rariusve brevibus



ca et S. Thomæ sylvis umbratis. — Griseb. Kl. Br. W. Ind. p. 157. difolia Blume, Mus. bot. lugd.-bat. 2. p. 52. *Urtica grandifolia?* Jamb. p. 336; Linn. spec. p. 1396; Poir. Encycl. 4. p. 637; Smith, U. diversifolia Plum. Icon. secund. Cl. Rich. — Herba elata, caescentibus robustis obtuse angulatis plerumque simplicibus imae (Cl. Richard) surculiferis neonon in tuberculum elongatum squamula intra terram obliquatum saepe incrassatis, merithalis 3-4 centim. prope bisulcatis (sulcis interdum pubescentibus). Folia vulgo 1-2 longa, 5-10 centim. lata, circiter a medio utrinque attenuata, basi angustata, acumine anguste grosseque serrato, dentibus limbi pleobtusis, nervis basilaribus saepe usque ad basim acuminis productis, proximie reticulato, lamina exsiccata membranacea subius paulo et obscurius punctata utrinque cystolithifera, cystolithis in pagina valde inaequalibus (aliis nempe: fusiformibus aliis punctiformibus), cre contra subaequalibus et fusiformibus sed minoribus crebrioribus, gracili quam internodium proximum plerumque longiore. Stipulae imetrals aut majores, obtusae, latiuscule amplexicaules. Cymae axillis superioribus ortae, foliis breviores, ovatae, pedunculum suum longum subaequant, ramulis patulis, glomerulis approximatis sed floribus pedicellatis, perigonii lobis sub apice longiuscule mucronatis glabris in alabastris 1-1½ millim. longo conniventibus. Cymae scalis ut videtur minores, glomerulis parvis, floribus breviter pedicellatis; perigonii fructiferi segmento intermedio breviter cucullaribus duplo longiore achenioque lato ovato aut fere rotundato imetrals et punctulato paulo brevior (v. s.)

*Urtica*, paulo minor, caulibus apice petiolisque et costa foliorum minus villosis, nervis basilaribus supra basim limbi nascentibus, aturibus obtusius acutiusculisve, cymis minoribus. ♀ In Novæ Granatæ, Barbacoas et Choco, altit. 200 mestr. (Triana, ann. 1868, n. 891).

*CRASSIFOLIA* (Blume l. c. p. 52), dioica, glaberrima, foliis ovatis vel ovatis lanceolatisve sensim acuminatis acumine acutiusculo basi is et in insertione petioli emarginatis interdum subpeltatis supra integram vel in toto margine serratis, stipulis majusculis oblongatis deciduis, cymis masc. densifloris, glomerulis capituliformibus, lis quam petioli multo longioribus, femineis subaxillaribus. ♀ In America (Alexander) et cult. in hort. bot. — *P. densiflora* Kunth, Ind. reg. ber. (1846) p. 12; Wedd. Monogr. p. 219. *Urtica crassifolia* sp. pl. 4. p. 349. *U. reclinata* h. bot. par. — Herba, caulibus inter lignescentibus, ramis herbaceis angulatis nodosis saepe longe nudis, cystolithis punctiformibus crebre obsitis; internodiis brevibus. Folia sim. longa, 1½-3 centim. lata, carnosula, serraturis grossiusculis, nervis basilaribus fere ad apicem limbi pertingentibus, venulis transversis flexuosis admodum conspicuis; lamina exsiccata rigida, cystolithis fusiformibus inaequalibus (in pagina superiore nervisque majoribus) rectiusculis arcuatisve obsita, petiolo 6-10 millim. longo. 6-10-millimetrals, mox decidua. Cymae solitariae, folia excedentes, lis 2-3 centim. longis, ramulis brevibus; masculae glomerulis multis magnitudine sessilibus approximatis, floribus pedicellatis, lobis obtuse mucronatis, alabastris vix millimetralibus. Flores feminei illos in glomerulos minimos discretos vel subconfluentes digesti, pericarpio segmento intermedio cucullato, quam achenium ovatum laeve brevior. (v. s.)

*STRELLATA* (Wedd. Monogr. p. 220), dioica, glaberrima, foliis oblongis atque basi-acutis obtusatisve et haec excepta acute serratis trinerviis, basilaribus apicem limbi attingentibus, stipulis majusculis cordato-stipularis, cymis masc. e glomerulo unico capituliformi compositis, pediculis filiformibus quam petioli longioribus, femineis... ♀ In sylvis humidis prov. Larecaja, prope Tipuani (Wedd.). — Herba, caudato 4-6 decim. alto, crassitie pennam anserinam emulante, sub angulato, emiccate cystolithis minutissimis punctulato, internodiis la. Folia 8-14 centim. longa, 2½-4 centim. lata; acumine subsemit-

metrali ut limbus reliquus serrato, serraturis quam in specie antecedente minoribus et multo numerosioribus, nervis basilaribus usque ad acuminis basin productis, venulis lateralibus rectiusculis obliquis, lamina exsiccata membranacea utrinque cystolithis brevissimis supra rariusculis subtus creberrimis obsita, petiolo 6-10 millim. longo. Cymæ masc. in axillis pluribus solitariae, densifloræ, spheroidæ, 5-8 millim. latæ, pedunculis 1-2 centim. longis, floribus albis brevissime pedicellatis, lobis perigonii brevissime mucronatis, alabastris bimillimetralibus. (v. v.)

B. — *Brevipedunculatæ, sive pedunculis (saltem cymarum fructiferarum) quam petioli brevioribus* (1).

102. P. PTEROPODON, dioica, glaberrima, foliis amplis ovato-oblongis ovatisve acuminatis (acumine obtusiusculo) basi abrupte longissimeque attenuatis (petiolo inde alato) supra quartam partem inferiorem crenato-serratis triplinerviis, nervis basilaribus ante apicem limbi desinentibus, stipulis majusculis oblongis, cymis fem. corymbiformibus pedunculatis petiolo superatis.  $\frac{1}{2}$  In provincia Tuquerres Novæ Granatæ, via de Tuquerres, altit. 80 metr. (Triana). — Caulis sat robustus, in specimine obvio haud elongatus, sulcato-tetragonus. Folia 15-20 centim. longa, 6-10 centim. lata, sensim acuminata, acumene sicut limbus serraturis obtusis notato, lamina exsiccata submembranacea utrinque cystolithis minimis punctiformibus dispersa, petiolo 2-5 centim. longo decurrentiæ limbi ope latiuscule alato. Stipulæ  $1\frac{1}{2}$ -2 centim. longæ, apice rotundatæ, persistentes. Cymæ fem. (solum visæ) floribundæ, fructiferæ corymbiformes, pedunculo erecto bi-sesqui-centimetro, perigonii segmento intermedio oblongo lateralibus ovatis dimidio longiore achæniūque rotundato-ovatum plus quam millimetrale circinatim scabrum fere æquante. (v. s. in h. Mus. par.)

103. P. FLEXUOSA (Wedd. in Ann. sc. nat. l. c. p. 218; Monogr. p. 234), dioica, glaberrima, foliis late ovatis rarius suborbicularibus acuminatis (acumine acutiusculo) a basi rotundata vel subcordata laxiuscule serratis tri-triplinerviis, nervis basilaribus apicem limbi haud attingentibus, stipulis cordato-rotundatis amplexicaulibus, cymis masc. et fem. e glomerulo uno alterove constantibus, pedunculis quam petioli brevioribus rariusve longioribus.  $\frac{1}{2}$  In montibus Quindiuensibus, Novæ Granatæ (Goudot) et in sylvis montis Pichincha, ad truncos arborum (Jameson). — Caules herbacei, simplices v. ramosi, debiles, internodiis 2-4-centimetralibus. Folia 2-7 centim. longa, 2-6 centim. lata, patula, serraturis acutis, nervis basilaribus usque ad trientem superiorem limbi productis vix prominentibus, lamina exsiccata membranacea, subtus pallide virente, cystolithis crebris linearibus oculos allicientibus utrinque ornata, petiolo gracili dimidii limbi circiter longitudine. Stipulæ 4-5-millimetrales, patulæ, insignes, illæ ejusdem nodi interdum in unam connatæ. Cymæ solitariae vel geminae, pedunculis filiformibus 1-4-centimetralibus suffultæ, floribus breviter pedicellatis fere capitatis, perigonii masc. laciniis sub apice longe et acutiuscule mucronatis, mucronibus in alabastro  $2\frac{1}{2}$  millim. longo discretis; perigonio fructifero cystolithis lineolato, segmentis ovatis, intermedio sub apice breviter mucronato-cupullato segmenta lateralia paulo excedente achænioque rotundato vel late ovato bimillimetrali lævi subæquilongo. (v. s. in hb. Mus. par. et Kew.)

104. P. SERRATIFOLIA (Wedd. Monogr. p. 235), monoica, glaberrima, foliis ovatis subacuminatis basi rotundatis vel subcordatis fere in toto margine grossiuscule acutiusculeque serratis (dente apicali obtuso) trinerviis, nervis basilaribus apicem limbi haud attingentibus, stipulis majusculis late ovatis, cymis minimis paucifloris unisexualibus quam petioli brevioribus.  $\frac{1}{2}$  ? In America æquatori li, sylvis reg. occid. montis Pichincha (Jameson) ann. 1850. n. 834). — Herba, caule ascendente vel erecto ramoso succulento angulato folioso crassitie pennæ corvinæ, exsiccato cystolithos vix ullos exhibente, internodiis

(1) Vid. quoque P. grandem et P. flexuosam inter Longipedunculatas, P. argutam inter Pubescentes et P. dendrophilam inter Heterophyllas.

1-3-centimetralibus. Folia 1-3 cm. longa, 1-2½ centim. lata, serraturis marginæ exteriori convexiusculis, dente apicali cæteris minore. nervis basilariibus ad partem quartam superiorem limbi pertingentibus et in ipso margine evanidis; lamina exsiccata rigidula subtus glaucescente, nervis supra impressis cystolithis linearibus utrinque (in pagina superiore vero multo crebrioribus minoribusque) notata, petiolis ejusdem jugi inæquilongis, majore vix centimetrali. Stipulæ 4-6 millim. longæ, amplexicaules et ejusdem nodi sæpe basi connatæ, laxæ patulæ. Cymæ e glomerulo unico ut plurimum constantes, floribus subsessilibus; masculæ plerumque ex axillis superioribus ortæ, stipulis (siccitate ferrugineis) involucratæ, perigonii lobis mucronatis, mucronibus in alabastro subdiscretis; perigonii fructiferi segmento intermedio breviter cucullato segmenta lateralalia longe superante achæniumque millimetrale acute marginatum non omnino æquante. (v. s. in hb. Kew., Webb. et Boiss.)

105. *P. ASTROGRAMMA* (Miq. in Mart. Fl. bras. fasc. 12. p. 201), glaberrima, foliis lanceolatis vel elliptico-lanceolatis longiuscule attenuato-acuminatis basi acutis cuneatis integrisque cæterum serratis vel crenato-serratis triplinerviis, stipulis exilibus triangulari-ovatis, cymis petiolo multo brevioribus. ♀? In Brasiliæ sebastianopolitanæ depressis sylvaticis passim (Martens). — Wedd. Monogr. p. 236. — Herba succulenta, caule 3-6 decim. alto, simplice vel parce ramoso, e basi radicante adscendente. Folia in eodem jugo subæqualia, 6-10 centim. longa, 8-20 millim. lata, in ipso acumine etiam serrata, exsiccata tenuiter membranacea subtus pallida et utrinque cystolithis minutis stellatis (vulgo 3-radiatis) crebre dispersa, petiolo 5 millim. -2½ centim. longo, exsiccato ut caulis ipse cystolithis simplicibus striolato. Stipulæ oculo nudo indistinctæ, petiolo utrinque tanquam parvæ auriculæ adnatæ. Cætera ignota. (v. s. in hb. Monac.)

106. *P. RETICULATA* (Wedd. in Ann. sc. nat. l. c. p. 215; Monogr. p. 236), monoica vel dioica, glabra, foliis suboblique ellipticis oblongisve acuminatis in dimidio vel triente superiore laxè serratis tri-tripli-nerviis, nervis basilariibus apicem limbi haud attingentibus, stipulis triangulari-ovatis persistentibus, cymis unisexualibus androgynisve, pedunculis quam petioli brevioribus. ♀ In petrosis montium Jamaicæ interioris (Swartz). — Griseb. Fl. Br. W. Ind. p. 156. *Urtica reticulata* Swartz, in Act. Holm. (1787), p. 30; ejusd. Fl. Ind. occ. p. 286; Poir. et Encycl. Suppl. 4. p. 49; Smith, l. c. n. 11. — Caulis herbacei vel suffrutescentes, 2-5 decim. alti, simplices, inferne longe nudi, internodiis elongatis. Folia 6-9 centim. longa, 2-3½ centim. lata, sensim breviterque acuminata, prope basim paulo angustiora et juxta petioli insertionem haud raro subcordata, serraturis acutiusculis vel obtusis interdumque fere obsolete, nervis basilariibus ultra medium vel etiam fere usque ad apicem limbi productis, intervenio subtus reticulato; lamina exsiccata rigida, cystolithis in utraque pagina linearibus crebris inæqualibusque in inferiore secus venulas reticulatim ordinatis; petiolo sesquicentimetrali. Stipulæ magnitudine varia. Cymæ petiolum paulo superantes, breviter ramosæ, glomerulis approximatis, pedunculo petiolum raro æquante. (v. s. typ. in hb. Holm., Mus. brit., Soc. Linn. et Kew.)

107. *P. MULTIFLORA* (Wedd. in Ann. sc. nat. l. c. p. 218, addit. spec.), monoica vel dioica, glaberrima, foliis ellipticis elliptico-lanceolatis oblongisve acuminatis (acumine acuto) basi acutatis v. obtusis in dimidio v. triente superiore limbi aut in toto margine argute serrulatis (serraturis sæpe apiculatis) rariusve (nisi in ipso apice folii) integerrimis trinerviis v. subtriplinerviis, nervis basilariibus apicem limbi attingentibus, stipulis brevissimis persistentibus, cymis masc. laxis petiolos longe superantibus, femineis conglomeratis pedunculis quam petioli plerumque brevioribus. ♀ In provinciis Quindiu, Bogota et Popayan Novæ Granatæ, altit. 1850-2600 metr. (Humb. et Bonpl., Goudot, Triana, etc.) et in Peruvia (Jos. de Jussieu). *P. Mutisiana* Wedd. Monogr. p. 237. *P. Mutisiana* et *P. multiflora* ejusd. in Ann. sc. nat. l. c. *Urtica multiflora* Poir. Encycl. suppl. 4. p. 223. *U. melastomoides* et *U. floribunda* Kunth in H. B. Nov. gen. et sp. 2. p. 38. *U. Mutisiana* Spreng. Syst. veg. 3. p. 840. *Dubruelia multiflora* Gaudich. Uran. p. 495. Herba suberecta, caule nonnunquam basi lignescente simplice 3-5 decim. alto angulato-sulcato,

exsiccatō obsolete punctulato, internodiis elongatis, nodis plerisque foliiferis. Folia 6-12 centim. longa, 2-5 centim. lata, inferne attenuata, ad insertionem petioli plerumque submarginata, serraturis laxis et interdum minimis, acumine ipso (nonnunquam centimetrali) etiam denticulato, nervis basilaribus conspicuis eleganter arcuatis margini parallelis usque ad apicem limbi productis subtus prominulis, venulis insigniter parallelis alternatim crassioribus in margine et intervenio concinne anastomosantibus, utraque pagina laminæ exsiccatæ at præsertim superiore cystolithis fusiformibus valde inæqualibus et interdum fere omnibus punctiformibus magis minusve obsita, petiolo plerumque brevi rarius  $1\frac{1}{2}$ -2-centimetrali quandoque longiore. Stipulæ a basi late amplexicauli abrupte attenuatæ, parum conspicuæ. Cymæ in axillis fere omnibus sæpe geminæ; masculæ pedunculo 2-3 centim. longo suffultæ, floribundæ, ramulis patulis, glomerulis approximatis, floribus brevissime pedicellatis, segmentis perigonii sub apice breviter et obtuse mucronatis, alabastro bimillimetrali. Cymæ fem. vix multifloræ, pedunculo petiolum vix æquante aut illo breviorē, ramulis brevibus patulo-recurvis, floribus subsessilibus, perigonii fructiferi segmento intermedio cucullato lateralia longitudine paulo excedente quam achænium autem ovatum sesquimillimetrale læve breviorē. (v. s. in hb. Mus. par.)

108. *P. MARGINATA* (Wedd. Monogr. p. 238), dioica, glaberrima, foliis amplis oblongo-ellipticis acuminatis (acumine angusto acutiusculoque) basi acutis obtusiusve, trinerviis vel subtriplinerviis (nervis basilaribus ad apicem limbi productis), in triente superiore repando-serrulatis, stipulis deciduis, cymis masc. sessilibus vel brevissime pedunculatis quam petioli brevioribus, divaricato-ramosis.  $\frac{4}{5}$  In Peruvia subandina (Poeppig, n. 3045). — *Urtica marginata* Poepp. mscr. — Caulis herbaceus, scandens?, flexuosus, obsolete sulcatus, internodiis longitudine varia. Folia patula, 10-16 centim. longa, 4-6 centim. lata, nervis basilaribus arcuatis ad ipsam basim acuminis productis subtus vix prominentibus, venulis transversis subdistantibus immersis, marginali autem distincta, lamina exsiccatæ subcoriacea subtus pallidiorē et obscure punctata utrinque cystolithis punctiformibus parum conspicuis sed creberrimis obsita marginalibus majoribus linearibusque, petiolo robusto 1-2 centim. longo. Stipulæ.... Cymæ masc. in axillis pluribus solitariae, vix multifloræ, ramulis brevibus, glomerulis discretis approximatisve, floribus subsessilibus, lobis perigonii brevissime mucronatis, alabastro vix millimetrali. — Affinis præcedenti etiamque varietati foliis serratis *P. smilacifoliæ* a qua facillime tamen distinguitur ordinatione cystolithorum. (v. s.)

109. *P. MEXICANA* (Wedd. in Ann. sc. nat. l. c. p. 214; Monogr. p. 239), monoica vel dioica, glaberrima, foliis anguste ellipticis lanceolatisve longe acuminatis (acumine acutiusculo) inferne longe attenuatis in triente superiore acutiuscule serratis trinerviis, nervis basilaribus apicem limbi attingentibus, stipulis brevissimis persistentibus, cymis masc. laxiusculis femineisque breviter pedunculatis petiolos vix æquantibus.  $\frac{4}{5}$  In locis humidis imperii Mexicani (Linden, n. 651; Botteri, n. 541). — Habitus *P. Musitianæ*, caule simili. Folia 6-12 centim. longa, 12-15 centim. lata, in petiolum sensim attenuata et ad insertionem hujusce interdum submarginata, acumine remote serrato, nervis basilaribus usque ad basim acuminis protensis margini propinquis, venulis plerisque in intervenio reticulatim anastomosantibus; laminæ exsiccatæ submembranaceæ pagina inferiore cystolithis fere punctiformibus subtilissimis creberrimisque granulosa, superiore nonnullis punctiformibus aut linearibus fusiformibusve insperso, margine ipso iisdem corpusculis multo majoribus insigniter ornato; petiolo  $\frac{1}{2}$ -1 $\frac{1}{2}$  centim. longo. Stipulæ parum conspicuæ, a basi late amplexicauli abrupte attenuatæ. Cymæ masc. plerumque geminæ, paucifloræ, ramulis brevibus divaricato-recurvis, glomerulis approximatis, floribus vix non sessilibus, segmentis perigonii sub apice brevissime mucronatis, mucronibus in alabastro sesquicentimetrali conniventibus. Cymæ fem. in planta monoica ex axillis superioribus ortæ quam masculæ minoribus, glomerulis discretis aut confluentibus, perigonii fructiferi segmento intermedio breviter cucullato lateralibus longe superantibus achæniisque ovato  $\frac{3}{4}$  millim. longo læviusculo paulo breviorē. — Forsan a sequente non specificè distincta. (v. s.)

β, *vegetior*, foliis integris. ♀ In imperio mexicano (Botteri). — Folia 20 centim. longa, 2-3 centim. lata, exsiccata margine cystolithorum prominentia sinuata. (v. s.)

110. *P. FALCATA* (Liebmänn, l. c. p. 304), monoica, glaberrima, foliis falcato-ellipticis utrinque attenuatis acuminatis basi leviter emarginatis in dimidio superiore remote serratis trinerviis, stipulis deciduis, cymis unisexualibus, femineis in axillis superioribus petiolo brevioribus, achænio margine tuberculato. ♀ In provincia mexicana Oaxaca (Liebmänn). — Wedd. Monogr. p. 240. — Herba, caule sesquidecimetrali tetragono articulo hyalino plerumque indiviso. Folia  $3\frac{1}{2}$ -4 $\frac{1}{2}$  centim. longa,  $1\frac{1}{2}$ -2 $\frac{1}{2}$  centim. lata, pellucida; supra saturate viridia, cystolithis fusiformibus minutissimis dense obsita, subtus glauca, petiolo subbicentimetrali pellucido cystolithis striolato. Cymæ masc. in parte caulis inferiore geminæ, 2 $\frac{1}{2}$  centim. longæ, patulæ, flexuosæ, alterne ramosæ, floribus sessilibus. Cymæ fem. geminæ, reflexæ, achæniis fuscis.
111. *P. CITRIODORA* (Wedd. in Ann. sc. nat. l. c. p. 216; Monogr. p. 240), dioica, glaberrima, foliis cordato-ovatis acuminatis excepta basi et acumine dentato-serratis trinerviis, nervis basilaribus apicem limbi haud attingentibus, stipulis triangulari-ovatis persistentibus, cymis masc. numerosis laxifloris quam folia brevioribus, pedunculis petiolos non æquantibus. ♀ ? In Peruvia (Pavon). — Herba, caule erecto 2-4 decim. alto simplice robusto, nodis inferioribus nudis. Folia 6-10 centim. longa, 2 $\frac{1}{2}$ -4 centim. lata, lamina exsiccata crassiuscula subtus punctata et utrinque cystolithis minimis linearibus creberrime conspersa, petiolo brevi. Stipulæ a basi lata abrupte attenuatæ, acutæ. Cymæ masc. (solum visæ) ramulis patentibus, glomerulis fere discretis, segmentis perigonii sub apice brevissime obtuse mucronatis. — Habitu *P. Goudotianam* refert, cæterum vero ab illa valde discrepat. (v. s. in hb. Webb. et Boiss.)
112. *P. PUNCTATA* (Wedd. in Ann. sc. nat. l. c. p. 222; Monogr. p. 241), monoica, glaberrima, foliis ellipticis vel ovato-ellipticis longe acuminatis (acumine obtuso) deorsum cuneatis excepta basi grossiuscule crenato-serratis (crenis obtusissimis) triplinerviis, nervis basilaribus apicem limbi non attingentibus, stipulis minimis a basi abruptissime attenuatis, cymis androgynis laxiusculis vix multifloris, cum pedunculo petiolum vix æquantibus. ♀ In Peruvia, prov. Jaen (Humb. et Bonpl.). *Urtica punctata* Kunth in H. B. Nov. gen. et spec. 2. p. 38. Caulis herbaceus, 2-4 decim. altus, erectus vel adscendens, simplex, crassitie circiter pennæ corvinæ, exsiccatus angulatus cystolithis striolatus, internodiis 2-3-centimetralibus. Folia 3-8 centim. longa,  $1\frac{1}{2}$ -3 centim. lata, acumine liguliformi laxè serrato, a medio ad basim sensim attenuato-cuneata atque in parte quarta vel quinta inferiore integerrima, nervis basilaribus ad trientem superiorem limbi productis, intervenio late reticulato, lamina exsiccata membranacea et cystolithis fusiformibus delicatulis in utraque pagina (in inferiore autem minoribus numerosioribusque) conspersa, petiolo gracili 10-15 millim. longo. Stipulæ inconspicuæ. Cymæ in axillis omnibus vel tantum in superioribus solitariæ vel geminæ, breviter ramosæ; floribus masc. breviter pedicellatis, perigonii lobis mucrone conico sub apice minutis, alabastro vix millimetrali; perigonii fem. segmento intermedio cucullato lateralibus longiore ovarium autem ovatum non æquante. (v. s. in hb. Mus. par.)
113. *P. SESSILIFLORA* (Wedd. Monogr. p. 242), dioica?, glaberrima, foliis ellipticis vel elliptico-lanceolatis ejusdem paris sæpe valde inæquimagnis utrinque attenuatis acutis in dimidio vel tantum in triente superiore argute serrulatis rarius omnino integris triplinerviis, nervis basilaribus apicem limbi non attingentibus, stipulis brevibus triangularibus, cymis parvis petiolo multo brevioribus. ♀ In sylvis humidis ins. Jamaicæ (Swartz) et promontorii Corrientes in prov. Darien Novæ Granatæ (Seemann). — Griseb. Fl. Br. W. Ind. p. 157. *Urtica sessiliflora* Swartz, Fl. Ind. occ. p. 321; Smith, l. c. n. 61. U. variegata Spreng. Syst. 3. p. 839. — Caulis erectus aut inferne repens, simplex vel basi ramosus, 2-3 decim. altus, siccitate albicans, nodis inferioribus aphyllis, internodiis 2-5-centimetralibus. Folia 3-10 centim. longa,  $\frac{1}{2}$ -5 centim.

lata, nervis basilaribus ad trientem apicalem limbi vel etiam altius productis, lamina exsiccata submembranacea, cystolithis minimis punctiformibus et fusiformibus (in pagina inferiore vix perspicuis) conspersa, venulis nigricantibus, petiolo gracili quam limbus dimidius ut plurimum brevior. Cymæ in speciminibus visis imperfecte evolutæ, densæ, in singulis axillis solitariae. — Habitu præcedenti nonnihil similis. (v. s. typ. in hb. Holm.)

114. *P. RADICANS* (Wedd. in Ann. sc. nat. l. c. p. 223), monoica, glabra, foliis ovatis vel rhombeo-ellipticis obtusis basi cuneatis acutisque in dimidia parte superiore grossiuscule crenatis (crenis utrinque 3-4) triplinerviis, nervis basilaribus apicem limbi non attingentibus, stipulis minimis, cymis parvis androgynis subsessilibus. ½ In sylvis umbrosis Jamaicæ septentrionalis interiorisque (Swartz). — Griseb. l. c. p. 157. *Urtica radicans* Swartz, in Act. Holm. ann. 1785. p. 29; ejusd. Fl. Ind. occ. p. 229; Smith, l. c. n. 38. Herba, caulibus elongatis (4-5 decim.) gracilibus ramosis flexuosis ad truncos repentibus radiculisque brevibus et crassiuscule undique instructis. Folia 1-2 centim. longa, 7-12 millim. lata, nervis basilaribus juxta trientem superiorem limbi in margine desinentibus parum prominulis, lamina exsiccata membranacea supra cystolithis linearibus fusiformibusque inspersa, subtus iisdem fere omnino destituta. Cætera desiderantur. (v. s. in hb. Mus. brit. et Soc. Linn.)
115. *P. LAMIOIDES* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 213; Monogr. p. 198), monoica, glabra, foliis in apice caulis senis subverticillatisque superioribus late ovatis obtusis grosse crenato-dentatis, duobus inferioribus integerrimis paulo minoribus basi subcordatis, stipulis inconspicuis, cymis brevissimis pedunculatis sessilibusve quam petioli brevioribus androgynis. ④ In Peruvia (Dombay, Pavon). — Herba, habitu Lamii purpurei, caule simplice 5-10 centim. alto apice tantum foliigero, internodiis illic brevissimis. Folia 3-24 millim. longa, 6-20 millim. lata, trinervia, serraturis superiorum in utraque margine 5-6, nervis basilaribus ultra medium limbum longiuscule productis, laminae exsiccatae tenuiter membranaceæ concoloris pagina superiore cystolithis linearibus tenuissimis obsita, inferiore iisdem destituta, petiolo gracili 5-8 millim. longo. Stipulae basi late amplexicaules, ovatae, obtusæ. Cymæ densifloræ, conglomeratæ; floribus masc. breviter pedicellatis, perigonii lobis crasse et obtuse mucronatis, floribus fem. subsessilibus, perigonii fructiferi segmento intermedio cucullato, segmentis lateralibus multoties longiore achænioque semimillimetrale vix non æquante. (v. s. in hb. Mus. par.)
116. *P. SUBCRENATA* (non Blume), monoica, glaberrima, foliis parvis ovatis obtusis basi fere rotundatis succulentis in toto margine vel basi excepta subcrenatis enerviis, stipulis brevibus triangularibus, cymis parvis androgynis sessilibus pedunculatisve. ½ In petrosis apricis Andium regionis æquinoctialis (Jameson, ann. 1856. n. 481). — Cæspitosa, caulibus herbaceis adscendentibus 6-8 centim. altis parce ramosis inferne denudatis, merithallis brevibus. Folia 4-5 millim. longa, 2-4 millim. lata, crenis in parte inferiore limbi haud raro deficientibus, nervis omnino immersis, lamina exsiccata coriacea supra planiuscula cystolithisque fusiformibus intricatis creberrime obsita, infra his corpusculis ut videtur destituta et magis minusve lacunosorugosa. Glomeruli e floribus femineis pluribus et masculo unico sæpius constantes, pedunculo subnullo aut petiolum 1-2-millimetrale superante. Flores masc. cito decidui, segmentis perigonii sub apice vix gibbosulis, alabastro sesquicentimetrali; feminei subsessiles, perigonii fructiferi segmento intermedio cucullato semimillimetrali lateralibus lanceolata et achænio oblongo-ovato æquante. — Habitus *P. globosæ*. (v. s.)

\*\* Pubescentes, sive caule foliisve puberulis pubescentibus vel pilosis.

A. — *Longipedunculatæ*, sive pedunculis (saltem cymarum fructiferarum) quam petioli longioribus (1).

117. *P. CERATOCALYX*, monoica, foliis oblongis vel elliptico-ovatis acuminatis

(1) Vid. quoque *P. Chamædryoidem* inter *Brevipedunculatas*; *P. dauciodo-*

in petiolum attenuatus supra trientem inferiorem limbi usque ad apicem acutiuscule serratis triplinerviis utrinque pilosis, stipulis majusculis rotundatis persistentibus, cymis unisexualibus, masculis e glomerulo unico constantibus, floribus longe mucronatis, femineis multo majoribus pedunculo petiolum superante. ♀ In Peruvia? (Poeppig! n. 3046). — Herba, caule basi repente 1-3 decim. longo simplice inferne pubescente et fere aphylo apice hirsuto, nodis superioribus foliiferis approximatisque. Folia 3-7 centim. longa, 1-3 centim. lata, nervis basilaribus supra petioli insertionem ortis, lamina exsiccata membranacea cystolithis minimis fusiformibus punctiformibusque creberrime ornata, petiolo costaque et nervis primariis densiuscule pubescenti-pilosis. Stipulæ in planta sicca subscariosæ, circiter 6 millim. longæ, obtusissimæ, ciliatæ. Cymæ masc. paucifloræ, pedunculo hirtio petiolum æquante, floribus pedicellatis, perigonii hirti segmentis mucrone lineari fere bimillimetrali sub apice instructis. Cymæ fem. folio multo breviores, patule ramosæ, pedunculo ramisque glabris, glomerulis parvis subdiscretis, perigonii fructiferi segmento intermedio breviter cucullato, lateralibus duplo longiore achæniumque semimillimetrale fere æquante. (v. s. in hb. DC.)

118. *P. LATIFOLIA* (Wedd. Monogr. p. 249), dioica, foliis amplis late elliptico-obovatis, breviter acuminatis (acumine sæpius obtuso) basi obtusis subcordatisve in triente inferiore integris cæterum inæqualiter et obtusiuscule serratis tripli- aut subquintupli-nerviis supra glaberrimis subtus (præsertim in nervis) pubescenti-pilosulis, stipulis majusculis ovatis deciduis, cymis masc. densifloris, pedunculis quam petioli longioribus, femineis.... ♀ In sylvis prov. Ocaña Novæ Granatæ (Schlim, Linden, n. 701). — Frutex (teste Schlim), ramis herbaceis lignescentibusve apice pubescentibus, exsiccatis crassitie circiter pennæ anserinæ angulato-sulcatis cystolithisque prope destitutis, internodiis elongatis. Folia 8-12 centim. longa, 6-8 centim. lata, nervis basilaribus subtus parum prominentibus in margine circiter ad quartam superiorem partem limbi desinentibus, lamina exsiccata membranacea subtus pallida sparsimque punctata et utrinque (nisi in punctis paginæ inferioris) cystolithis linearibus creberrimis obsita, petiolo 2-5 centim. longo pubescente. Stipulæ sesquicentimetrales, obtusæ. Cymæ masc. (solum visæ) binæ, abortiente gemma suprema fere terminales, foliis multo breviores, pedunculo crassiusculo pubescente, ramulis brevibus, glomerulis confertis, floribus pedicellatis, perigonii lobis glabris sub apice mucrone angusto longiusculo sæpiusque glabrato munitis, alabastro 2½-3 millim. longo. (v. s.)

119. *P. FASCIATA*, dioica, caule undique dense tomentoso-hirsuto, foliis ovatis acuminatis basi rotundatis in toto margine crebre acuteque dentato-serratis triplinerviis supra asperato-rugosulis pilosulisque subtus pubescenti-hirsutis, stipulis majusculis ovatis persistentibus, cymis unisexualibus subcorymbosis, pedunculis hirtulis petiolos superantibus, achænio ovato læviusculo. ♀ In sylvis prov. Choco et Novita Novo-granatensium, altit. 170 metr. (Triana). — Caulis erectus, robustus, hirsutiae densa et molli vestitus, internodiis superioribus 3-5 centim. longis. Folia 8-10 centim. longa, 5-6 centim. lata, anguste acuminata, acumine acuto sicut limbus dentibus parvis acutis mucronatisve ornato, nervis basilaribus paulo supra petioli insertionem ortis versus quartam partem superiorem limbi evanidis, pagina superiore in planta viva secus nervos albo-fasciata et undique inter venulas verruculis conoideis asperisque obsita, hisce in folio exsiccato cystolithis linearibus radiantibusque ornatis, pagina inferiore lacunosa sive sub verrucis alveolata, nervis densiuscule hirsutis, petiolo 1½-3 centim. longo, hirsutiae ut in caule. Stipulæ uni-sesquicentimetrales, extus sparsim pilosæ. Cymæ fem. (solum visæ) ex axillis superioribus solitarie nascentes, pedunculo quam folium brevius, ramulis patulis, glomerulis subdiscretis. Perigonii fructiferi segmentum intermedium oblongum, sub apice gibbosum, lateralibus ovatis dimidio longius achæniumque sesquimillimetrale fere æquans. (v. s. in h. Mus. par.)

120. *P. PAVONII* (Wedd. in Ann. se. nat., l. c. p. 219; Monogr. p. 257), monoica,

ram, *P. serrulatam*, *P. elegantem*, *P. spathulatam* et *P. grandem* inter *Glabras*; necnon *P. rufam* inter *Heterophyllas*.

superantibus, lamina exsiccata tenuiter membranacea cystolithis linearibus utrinque conspersa, petiolo gracili limbo dimidio æquilongo aut breviori villosa. Stipulæ vix millimetrales, puberulæ. Cymæ in axillis superioribus solitariae, cum pedunculo filiformi glaberrimoque folio subæquilongæ, ramulis divaricatis, glomerulis approximatis; floribus subsessilibus, masculis paucissimis, perigonii lobis sub apice mucronatis, alabastro submillimetrali; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achæniique submillimetrali subtiliter annulatum scabriusculo brevior. — A P. pubescente differt foliis utrinque glabris, stipulis brevibus et achæniis orbicularibus. (v. s.)

127. P. CHRYSOSPLENIODES (Wedd. in Ann. sc. nat. l. c. p. 231; Monogr. p. 270), dioica?, foliis ovato-vel obovato-rotundatis utrinque obtusissimis aut basi subcordatis in dimidio superiore crenatis ciliatis limbo cæterum glabro aut sparsim piloso trinerviis, stipulis late ovatis persistentibus, cymis unisexualibus femineis laxiusculis, pedunculis quam petioli paulo brevioribus. ♀ In Nova Granata, prope Fusaguasuga (Goudot); prope Colonia Tovar (Fendler). — Herba, caulibus radicanibus in specimine unico examinato 1-1½ decim. longis demum adscendentibus pubescenti-pilosis apice rosulato-foliosis, internodiis inferioribus elongatis. Folia ½-5 centim. longa, 4-25 millim. lata, crenis obtusissimis, nervis basilaribus fere ad apicem limbi pertingentibus, lamina exsiccata membranacea, supra obscure viridi cystolithisque nisi in nervis et margine omnino destituta, subtus pallidior et cystolithis undique lineolata, petiolo pubescente quam limbus dimidius plerumque brevior. Stipulæ late amplexicaules, 1½ millim. longæ, cystolithis striatæ. Flores masc. ad basim cymarum fem. subsessiles (an semper?), perigonio hirtello. Cymæ fem. in axillis superioribus solitariae, foliis breviores, ramulis brevibus patulis, glomerulis approximatis; perigonii fructiferi segmento intermedio cucullato laterales superante achæniique ovati semimillimetralis circinatim scabriusculi dimidiam longitudinem circiter æquante. — Habitu P. pubescenti valde proxima. (v. s.)

128. P. MAXIMILIANI (Miq. in Mart. Fl. bras. fasc. 12. p. 200), monoica?, caule superne pubescente, foliis ovatis vel elliptico-ovatis acutiusculis basi rotundatis obtusive serratis tri-vel subquinque-nerviis nervis subtus pubescentibus, stipulis ovato-oblongis serius deciduis, cymis densiusculis multifloris, pedunculis quam petioli sæpius longioribus. ♀ In Brasilia, prope Ilheos (Maxim. Neuwied). — Wedd. Monogr. p. 271. Urtica iners Nees et Mart. in Nov. act. Acad. L. C. Nat. Cur. 11. p. 33. excl. syn. — Herba, caule 3-4½ decim. alto erecto aut basi nonnihil radicante angulato. Folia accrescentia, 3½-10 centim. longa, 2½-6 centim. lata, nervis basilaribus ultra medium limbum productis, lamina exsiccata membranacea cystolithis minutis fusiformibus punctiformibusque conspersa subtusque sub lente hic illic maculis fusciscentibus notata; petiolo 1-3½ centim. longo pubescente. Stipulæ inflorescentiarum basim fere amplexantes. Cymæ fem. glabræ, floribus fasciculatim confertis, perigonii fructiferi segmento intermedio carinato lateralibus multo majore, achænio ovato læviusculo. (v. s. in hb. Mart.)

129. P. HIRTELLA (Miq. in Mart. Fl. bras. fasc. 12. p. 202), foliis ellipticis vel ovato-ellipticis acutiusculis obtusive excepta basi crenato-serratis trinerviis supra sparsim pilosis subtus in nervis hirtis, stipulis late ovatis persistentibus cymis densis longiuscule pedunculatis. ♀ In sylvis Brasiliæ prope Ilheos (Maxim. Neuwied). — Wedd. Monogr. p. 260. Boehmeria repens Nees et Mart. in Nov. Act. Soc. nat. Cur. 11. p. 36. Herba tenera, repens, ramulis petiolisque hirtellis. Folia infima 5 millim. superiora 2-3 centim. longa, lamina exsiccata membranacea, subtus pallida, cystolithis linearibus utrinque dense obsita, petiolo 2-8-millimetrali. Cymarum pedunculi bractea membranacea subtruncata basi vaginati. — Habitus sequentis.

130. P. PUBESCENS (Liebmann, in Vidensk. Selsk. Skr. 5 Række, naturv. og math. Afd. 2. p. 302. in adnot.), monoica vel dioica basi radicans, foliis late ovatis vel elliptico-obovatis rotundatisve utrinque obtusis vel basi subcordatis supra trientem inferiorem limbi grosse crenatis vel crenato-serratis trinerviis utrinque sparsim pilosis margineque ciliatis, stipulis late ovatis rotunda-



tisve persistentibus. cymis vulgo androgynis laxis multifloris, pedunculis quam petioli longioribus. ½ In sylvis humidis Americæ tropicalis sat frequens præsertim in Brasilia. — Wedd. Monogr. p. 258; Miq. in Mart. Fl. bras. fasc. 12. p. 202; Griseb. Fl. Br. W. Ind. p. 158. P. Gaudichaudiana Wedd. in Ann. sc. nat. l. c. p. 226. P. grossecrenata Miq. l. c. p. 199. t. 68. — Herba, caulibus sæpius longe repentibus 2-6 decim. longis demum ramisque adscendentibus pubescentibus, exsiccatis cystolithis parvis fusiformibus undique inspersis, internodiis elongatis (inferioribus interdum decimetralibus). Folia 1½-8 centim. longa, tantumdemque fere lata, superiora cæteris sæpe majora, dentibus obtusissimis acutiusculisve, nervis basilaribus trientem superiorem limbi attingentibus subtus prominulis longiusculeque pilosis, lamina exsiccata membranacea utrinque cystolithis fusiformibus tenuissimis (in inferiore pagina autem minoribus crebrioribusque) inspersa, petiolo longitudine varia pubescente. Stipulæ 2-5 millim. longæ, laxæ, ciliolatæ. Cymæ ex axillis præsertim superioribus ortæ, folia persæpe superantes, interdum unisexuales (masculæ tunc densiores), pedunculo filiformi 2-4 centim. longo, ramulis laxè patulis, glomerulis ut plurimum discretis; floribus masc. in cymis androgynis paucis, breviter pedicellatis, lobis perigonii sub apice longiuscule mucronatis, mucronibus pilosulis in alabastro 2 millim. longo discretis; floribus fem. subsessilibus; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achænioque vix semimillimetrali fere æquilongio. (v. s.)

β, *involutrata*, foliis quam in typo brevius petiolatis dentibus que minoribus instructis superioribus valde approximatis cymas breves involutratibus. — In ins. S. Vincent (Macrae). — *Urtica involutrata* Sims Bot. Mag. 51. t. 2481.

γ, *montana*, foliis late ovatis subacuminatis obtusiusculis supra glaberrimis. — In ins. Antillis et Venezuela (Fendler, n. 1243). P. *montana* Wedd. l. c. n. 228. (v. s.)

δ, *fuscata*, dioica, foliis late ovatis ellipticisve apice attenuatis obtusiusculis que rariusve subacuminatis supra pilosis subtus in nervis venulisque villosis, dentibus obtusis acutisve. — In imperio mexicano, prov. Oaxaca et Vera Cruz (Galeotti, n. 313; Linden, n. 41 et 42). P. *fuscata* Liebm. l. c. p. 300; Wedd. Monogr. p. 255. P. *xalapensis* ejusd. Monogr. p. 255. (v. s.)

ε, *rupicola*, foliis angustius ovatis subacuminatis obtusiusculis subtriplicinerviis utrinque molliter pubescentibus, dentibus obtusis. — In imperio mexicano, prope Tabasco, ad scopulos (Linden, n. 71) et in monte Orizaba (Botteri). *Pilea rupicola* Wedd. in Ann. sc. nat. l. c. p. 224. (v. s.)

ζ, *Guyanensis*, foliis ovatis ellipticisve obtusiusculis supra glabris nervis subtus sparsim pilosis, pedunculis quam petioli brevioribus. — In nemo ribus Guyanæ gallicæ, ad scopulos (Melinon, n. 55; Schomburgk, n. 357). P. *guyanensis* Wedd. in Ann. sc. nat. l. c. p. 221; Monogr. p. 266. (v. s.)

Nonnullæ specierum sequentium huic stirpi tanquam varietates verisimiliter etiam adscribendæ.

131. P. *obtusata* (Liebm. l. c. p. 300, in adnot.), dioica, caule piloso, foliis late ovatis utrinque obtusis grosse et obtuse crenatis subquinque-nerviis supra glabris subtus ad nervos pilosis, stipulis late ovatis, cymis diffuse ramosis longe pedunculatis, glomerulis discretis. ½? In ins. Monserrat (Ryan). — Wedd. Monogr. p. 261. *Urtica obtusata* hb. Vahl. — Herba caule sesquidecimetrali tetragono articulo superne piloso strigulosoque. Folia 5-6 centim. longa, 3½-4 centim. lata, exsiccata utrinque cystolithis minutissimis fusiformibus obsita, supra obscure viridia subtus glaucescentia, petiolo 2½-3½ centim. longo pilosulo. Stipulæ obtusæ. Cymæ solitariae, pedunculis 5 centim. longis, achænio ovato marginato subtilissime granulato. (Descript. ex Liebm. l. c.)

superantibus, lamina exsiccata tenuiter membranacea cystolithis linearibus utrinque conspersa, petiolo gracili limbo dimidio æquilongo aut breviori villosa. Stipulæ vix millimetricæ, puberulæ. Cymæ in axillis superioribus solitariae, cum pedunculo filiformi glaberrimæque folio subæquilongæ, ramulis divaricatis, glomerulis approximatis; floribus subsessilibus, masculis paucissimis, perigonii lobis sub apice mucronatis, alabastro submillimetrali; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achæniæque submillimetrali subtiliter annulatis scabriusculo brevioribus. — A P. pubescente differt foliis utrinque glabris, stipulis brevibus et achæniis orbicularibus. (v. s.)

127. P. CHRYSOSPLENIOIDES (Wedd. in Ann. sc. nat. l. c. p. 231; Monogr. p. 270), dioica?, foliis ovato-vel obovato-rotundatis utrinque obtusissimis aut basi subcordatis in dimidio superiore crenatis ciliatis limbo cæterum glabro aut sparsim piloso trinerviis, stipulis late ovatis persistentibus, cymis unisexualibus femineis laxiusculis, pedunculis quam petioli paulo brevioribus. ♀ In Nova Granata, prope Fusaguasuga (Goudot); prope Colonia Tovar (Fendler). — Herba, caulibus radicanibus in specimine unico examinato 1-1½ decim. longis demum ascendentibus pubescenti-pilosis apice rosulato-foliosis, internodiis inferioribus elongatis. Folia ½-5 centim. longa, 4-25 millim. lata, crenis obtusissimis, nervis basilaribus fere ad apicem limbi pertingentibus, lamina exsiccata membranacea, supra obscure viridi cystolithisque nisi in nervis et margine omnino destituta, subtus pallidiore et cystolithis undique lineolata, petiolo pubescente quam limbus dimidius plerumque brevior. Stipulæ late amplexicaules, 1½ millim. longæ, cystolithis striatæ. Flores masc. ad basim cymarum fem. subsessiles (an semper?), perigonio hirtello. Cymæ fem. in axillis superioribus solitariae, foliis breviores, ramulis brevibus patulis, glomerulis approximatis; perigonii fructiferi segmento intermedio cucullato laterales superante achæniæque ovati semimillimetralis circinatim scabriusculi dimidiam longitudinem circiter æquante. — Habitu P. pubescenti valde proxima. (v. s.)
128. P. MAXIMILIANI (Miq. in Mart. Fl. bras. fasc. 12. p. 200), monoica?, caule superne pubescente, foliis ovatis vel elliptico-ovatis acutiusculis basi rotundatis obtusisve grosse serratis vel subquinque-nerviis nervis subtus pubescentibus, stipulis ovato-oblongis serius deciduis, cymis densiusculis multifloris, pedunculis quam petioli sæpius longioribus. ♀ In Brasiliâ, prope Ilheos (Maxim. Neuwied). — Wedd. Monogr. p. 271. Urtica iners Nees et Mart. in Nov. act. Acad. L. C. Nat. Cur. 11. p. 33. excl. syn. — Herba, caule 3-4½ decim. alto erecto aut basi nonnihil radicante angulato. Folia accrescentia, 3½-10 centim. longa, 2½-6 centim. lata, nervis basilaribus ultra medium limbum productis, lamina exsiccata membranacea cystolithis minutis fusiformibus punctiformibusque conspersa subtusque sub lente hic illic maculis fusciscentibus notata; petiolo 1-3½ centim. longo pubescente. Stipulæ inflorescentiarum basim fere amplectentes. Cymæ fem. glabræ, floribus fasciculatim confertis, perigonii fructiferi segmento intermedio carinato lateralibus multo majore, achænio ovato lævisculo. (v. s. in hb. Mart.)
129. P. HIRTELLA (Miq. in Mart. Fl. bras. fasc. 12. p. 202), foliis ellipticis vel ovato-ellipticis acutiusculis obtusisve excepta basi crenato-serratis trinerviis supra sparsim pilosis subtus in nervis hirtis, stipulis late ovatis persistentibus cymis densis longiuscule pedunculatis. ♀ In sylvis Brasiliæ prope Ilheos (Maxim. Neuwied). — Wedd. Monogr. p. 260. Boehmeria repens Nees et Mart. in Nov. Act. Soc. nat. Cur. 11. p. 36. Herba tenera, repens, ramulis petiolisque hirtellis. Folia infima 5 millim. superiora 2-3 centim. longa, lamina exsiccata membranacea, subtus pallida, cystolithis linearibus utrinque dense obsita, petiolo 2-8-millimetrali. Cymarum pedunculi bractea membranacea subtruncata basi vaginati. — Habitus sequentis.
130. P. PUBESCENS (Liebmann, in Vidensk. Selsk. Skr. 5 Række, naturv. og math. Afd. 2. p. 302. in adnot.), monoica vel dioica basi radicans, foliis late ovatis vel elliptico-obovatis rotundatisve utrinque obtusis vel basi subcordatis supra trientem inferiorem limbi grosse crenatis vel crenato-serratis trinerviis utrinque sparsim pilosis margineque ciliatis, stipulis late ovatis rotunda-

tive persistentibus. cymis vulgo androgynis laxis multifloris, pedunculis quam petioli longioribus. ½ In sylvis humidis Americæ tropicalis sat frequens præsertim in Brasilia. — Wedd. Monogr. p. 258; Miq. in Mart. Fl. bras. fasc. 12. p. 202; Griseb. Fl. Br. W. Ind. p. 158. P. Gaudichaudiana Wedd. in Ann. sc. nat. l. c. p. 226. P. grossecrenata Miq. l. c. p. 199. t. 68. — Herba, caulibus sæpius longe repentibus 2-6 decim. longis demum ramisque adscendentibus pubescentibus, exsiccatis cystolithis parvis fusiformibus undique inspersis, internodiis elongatis (inferioribus interdum decimetralibus). Folia 1½-8 centim. longa, tantumdemque fere lata, superiora cæteris sæpe majora, dentibus obtusissimis acutiusculisve, nervis basilaribus trientem superiorem limbi attingentibus subtus prominulis longiusculeque pilosis, lamina exsiccata membranacea utrinque cystolithis fusiformibus tenuissimis (in inferiore pagina autem minoribus crebrioribusque) inspersa, petiolo longitudine varia pubescente. Stipulæ 2-5 millim. longæ, laxæ, ciliolatæ. Cymæ ex axillis præsertim superioribus ortæ, folia persæpe superantes, interdum unisexuales (masculæ tunc densiores), pedunculo filiformi 2-4 centim. longo, ramulis laxè patulis, glomerulis ut plurimum discretis; floribus masc. in cymis androgynis paucis, breviter pedicellatis, lobis perigonii sub apice longiuscule mucronatis, mucronibus pilosis in alabastro 2 millim. longo discretis; floribus fem. subsessilibus; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achænioque vix semimillimetrali fere æquilongò. (v. s.)

β, *involutrata*, foliis quam in typo brevius petiolatis dentibus que minoribus instructis superioribus valde approximatis cymas breves involucrentibus. — In ins. S. Vincent (Macrae). — *Urtica involutrata* Sims Bot. Mag. 51. t. 2481.

γ, *montana*, foliis late ovatis subacuminatis obtusiusculis supra glaberrimis. — In ins. Antillis et Venezuela (Fendler, n. 1243). P. montana Wedd. l. c. n. 228. (v. s.)

δ, *fuscata*, dioica, foliis late ovatis ellipticisve apice attenuatis obtusiusculisque rariusve subacuminatis supra pilosis subtus in nervis venulisque villosis, dentibus obtusis acutisve. — In imperio mexicano, prov. Oaxaca et Vera Cruz (Galeotti, n. 313; Linden, n. 41 et 42). P. fuscata Liebm. l. c. p. 300; Wedd. Monogr. p. 255. P. xalapensis ejusd. Monogr. p. 255. (v. s.)

ε, *rupicola*, foliis angustius ovatis subacuminatis obtusiusculis subtriplicinerviis utrinque molliter pubescentibus, dentibus obtusis. — In imperio mexicano, prope Tabasco, ad scopulos (Linden, n. 71) et in monte Orizaba (Botteri). Pilea rupicola Wedd. in Ann. sc. nat. l. c. p. 224. (v. s.)

ζ, *Guyanensis*, foliis ovatis ellipticisve obtusiusculis supra glabris nervis subtus sparsim pilosis, pedunculis quam petioli brevioribus. — In nemo ribus Guyanæ gallicæ, ad scopulos (Melinon, n. 55; Schomburgk, n. 357). P. guyanensis Wedd. in Ann. sc. nat. l. c. p. 221; Monogr. p. 266. (v. s.)

Nonnullæ specierum sequentium huic stirpi tanquam varietates verisimiliter etiam adscribendæ.

131. P. OBTUSATA (Liebm. l. c. p. 300, in adnot.), dioica, caule piloso, foliis late ovatis utrinque obtusis grosse et obtuse crenatis subquinque-nerviis supra glabris subtus ad nervos pilosis, stipulis late ovatis, cymis diffuse ramosis longe pedunculatis, glomerulis discretis. ½? In ins. Monserrat (Ryan). — Wedd. Monogr. p. 261. *Urtica obtusata* hb. Vahl. — Herba caule sesquidecimetrali tetragono articulo superne piloso strigulosoque. Folia 5-6 centim. longa, 3½-4 centim. lata, exsiccata utrinque cystolithis minutissimis fusiformibus obsita, supra obscure viridia subtus glaucescentia, petiolo 2½-3½ centim. longo piloso. Stipulæ obtusæ. Cymæ solitariae, pedunculis 5 centim. longis, achænio ovato marginato subtilissime granulato. (Descript. ex Liebm. l. c.)

lata, nervis basilaribus ad trientem apicalem limbi vel etiam altius productis, lamina exsiccata submembranacea, cystolithis minimis punctiformibus et fusiformibus (in pagina inferiore vix perspicuis) conspersa, venulis nigricantibus, petiolo gracili quam limbus dimidius ut plurimum brevior. Cymæ in speciminibus visis imperfecte evolutæ, densæ, in singulis axillis solitariæ. — Habitu præcedenti nonnihil similis. (v. s. typ. in hb. Holm.)

114. *P. RADICANS* (Wedd. in Ann. sc. nat. l. c. p. 223), monoica, glabra, foliis ovatis vel rhombeo-ellipticis obtusis basi cuneatis acutisque in dimidia parte superiore grossiuscule crenatis (crenis utrinque 3-4) triplinerviis, nervis basilaribus apicem limbi non attingentibus, stipulis minimis, cymis parvis androgynis subsessilibus. ♀ In sylvis umbrosis Jamaicæ septentrionalis interiorisque (Swartz). — Griseb. l. c. p. 157. *Urtica radicans* Swartz, in Act. Holm. ann. 1785. p. 29; ejusd. Fl. Ind. occ. p. 229; Smith, l. c. n. 38. Herba, caulibus elongatis (4-5 decim.) gracilibus ramosis flexuosis ad truncos repentibus radiculisque brevibus et crassiuscule undique instructis. Folia 1-2 centim. longa, 7-12 millim. lata, nervis basilaribus juxta trientem superiorem limbi in margine desinentibus parum prominulis, lamina exsiccata membranacea supra cystolithis linearibus fusiformibusque inersa, subtus iisdem fere omnino destituta. Cætera desiderantur. (v. s. in hb. Mus. brit. et Soc. Linn.)
115. *P. LAMIOIDES* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 213; Monogr. p. 198), monoica, glabra, foliis in apice caulis senis subverticillatisque superioribus late ovatis obtusis grosse crenato-dentatis, duobus inferioribus integerrimis paulo minoribus basi subcordatis, stipulis inconspicuis, cymis brevissime pedunculatis sessilibusve quam petioli brevioribus androgynis. ④ In Peruvia (Dombay, Pavon). — Herba, habitu Lamii purpurei, caule simplice 5-10 centim. alto apice tantum foliigero, internodiis illic brevissimis. Folia 3-24 millim. longa, 6-20 millim. lata, trinervia, serraturis superiorum in utraque margine 5-6, nervis basilaribus ultra medium limbum longiuscule productis, lamina exsiccata tenuiter membranacea concoloris pagina superiore cystolithis linearibus tenuissimis obsita, inferiore iisdem destituta, petiolo gracili 5-8 millim. longo. Stipulæ basi late amplexicaules, ovatae, obtusæ. Cymæ densifloræ, conglomeratæ; floribus masc. breviter pedicellatis, perigonii lobis crasse et obtuse mucronatis, floribus fem. subsessilibus, perigonii fructiferi segmento intermedio cucullato, segmentis lateralibus multoties longiore achæniumque semimillimetrale vix non æquante. (v. s. in hb. Mus. par.)
116. *P. SUBCRENATA* (non Blume), monoica, glaberrima, foliis parvis ovatis obtusis basi fere rotundatis succulentis in toto margine vel basi excepta subcrenatis enerviis, stipulis brevibus triangularibus, cymis parvis androgynis sessilibus pedunculatisve. ♀ In petrosis apricis Andium regionis æquinoctialis (Jameson, ann. 1856. n. 481). — Cæspitosa, caulibus herbaceis adscendentibus 6-8 centim. altis parce ramosis inferne denudatis, merithallis brevibus. Folia 4-5 millim. longa, 2-4 millim. lata, crenis in parte inferiore limbi haud raro deficientibus, nervis omnino immersis, lamina exsiccata coriacea supra planiuscula cystolithisque fusiformibus intricatis creberrime obsita, infra his corpusculis ut videtur destituta et magis minusve lacunosorugosa. Glomeruli e floribus femineis pluribus et masculo unico sæpius constantes, pedunculo subnullo aut petiolum 1-2-millimetrale superante. Flores masc. cito decidui, segmentis perigonii sub apice vix gibbosulis, alabastro sesquicentimetrali; feminei subsessiles, perigonii fructiferi segmento intermedio cucullato semimillimetrali lateralibus lanceolata et achænium oblongo-ovatum æquante. — Habitus *P. globosæ*. (v. s.)

\*\* Pubescentes, sive caule foliisve puberulis pubescentibus vel pilosis.

A. — *Longipedunculatæ*, sive *pedunculis* (saltem *cymarum fructiferarum*) quam *petioli longioribus* (1).

117. *P. CERATOCALYX*, monoica, foliis oblongis vel elliptico-ovatis acuminatis

(1) Vid. quoque *P. Chamædryoidem* inter *Brevipedunculatas*; *P. dauciodo-*

in petiolum attenuatis supra trientem inferiorem limbi usque ad apicem acutiuscule serratis triplinerviis utrinque pilosis, stipulis majusculis rotundatis persistentibus, cymis unisexualibus, masculis e glomerulo unico constantibus, floribus longe mucronatis, femineis multo majoribus pedunculo petiolum superante. ¼ In Peruvia? (Poeppig! n. 3046). — Herba, caule basi repente 1-3 decim. longo simplice inferne pubescente et fere aphylo apice hirsuto, nodis superioribus foliiferis approximatisque. Folia 3-7 centim. longa, 1-3 centim. lata, nervis basilaribus supra petioli insertionem ortis, lamina exsiccata membranacea cystolithis minimis fusiformibus punctiformibusque creberrime ornata, petiolo costaque et nervis primariis densiuscule pubescenti-pilosis. Stipulae in planta sicca subscariosae, circiter 6 millim. longae, obtusissimae, ciliatae. Cymae masc. pauciflorae, pedunculo hirsuto petiolum æquante, floribus pedicellatis, perigonii hirti segmentis mucrone lineari fere bimillimetrali sub apice instructis. Cymae fem. folio multo breviores, patule-ramosae, pedunculo ramisque glabris, glomerulis parvis subdiscretis, perigonii fructiferi segmento intermedio breviter cucullato, lateralibus duplo longiore achæniumque semimillimetrale fere æquante. (v. s. in hb. DC.)

118. *P. LATIFOLIA* (Wedd. Monogr. p. 249), dioica, foliis amplis late elliptico-obovatis, breviter acuminatis (acumine sæpius obtuso) basi obtusis subcordatisve in triente inferiore integris cæterum inæqualiter et obtusiuscule serratis tripli- aut subquintupli-nerviis supra glaberrimis subtus (præsertim in nervis) pubescenti-pilosulis, stipulis majusculis ovatis deciduis, cymis masc. densifloris, pedunculis quam petioli longioribus, femineis.... ½ In sylvis prov. Ocaña Novæ Granatæ (Schlim, Linden, n. 701). — Frutex (teste Schlim), ramis herbaceis lignescentibusve apice pubescentibus, exsiccatis crassitie circiter pennæ anserinæ angulato-sulcatis cystolithisque prope destitutis, internodiis elongatis. Folia 8-12 centim. longa, 6-8 centim. lata, nervis basilaribus subtus parum prominentibus in margine circiter ad quartam superiore partem limbi desinentibus, lamina exsiccata membranacea subtus pallida sparsimque punctata et utrinque (nisi in punctis paginæ inferioris) cystolithis linearibus creberrimis obsita, petiolo 2-5 centim. longo pubescente. Stipulae sesquicentimetrales, obtusæ. Cymae masc. (solum visæ) binæ, abortiente gemma suprema fere terminales, foliis multo breviores, pedunculo crassiusculo pubescente, ramulis brevibus, glomerulis confertis, floribus pedicellatis, perigonii lobis glabris sub apice mucrone angusto longiusculo sæpiusque glabrato munitis, alabastro 2½-3 millim. longo. (v. s.)

119. *P. FASCIATA*, dioica, caule undique dense tomentoso-hirsuto, foliis ovatis acuminatis basi rotundatis in toto margine crebre acuteque dentato-serratis triplinerviis supra asperato-rugosulis pilosulisque subtus pubescenti-hirsutis, stipulis majusculis ovatis persistentibus, cymis unisexualibus subcorymbosis, pedunculis hirtulis petiolos superantibus, achænio ovato læviusculo. ¾ In sylvis prov. Choco et Novita Novo-granatensis, altit. 170 metr. (Triana). — Caulis erectus, robustus, hirsutiae densa et molli vestitus, internodiis superioribus 3-5 centim. longis. Folia 8-10 centim. longa, 5-6 centim. lata, anguste acuminata, acumine acuto sicut limbus dentibus parvis acutis mucronatisve ornato, nervis basilaribus paulo supra petioli insertionem ortis versus quartam partem superiorem limbi evanidis, pagina superiore in planta viva secus nervos albo-fasciata et undique inter venulas verruculis conoideis asperisque obsita, hisce in folio exsiccato cystolithis linearibus radiantibusque ornatis, pagina inferiore lacunosa sive sub verrucis alveolata, nervis densiuscule hirsutis, petiolo 1½-3 centim. longo, hirsutiae ut in caule. Stipulae uni-sesquicentimetrales, extus sparsim pilosae. Cymae fem. (solum visæ) ex axillis superioribus solitarie nascentes, pedunculo quam folium breviori, ramulis patulis, glomerulis subdiscretis. Perigonii fructiferi segmentum intermedium oblongum, sub apice gibbosum, lateralibus ovatis dimidio longius achæniumque sesquimillimetrale fere æquans. (v. s. in h. Mus. par.)

120. *P. PAVonii* (Wedd. in Ann. se. nat., l. c. p. 219; Monogr. p. 257), monoica,

ram, *P. serrulatam*, *P. elegantem*, *P. spathulatam* et *P. grandem* inter *Glabras*; necnon *P. rufam* inter *Heterophyllas*.

superantibus, lamina exsiccata tenuiter membranacea cystolithis linearibus utrinque conspersa, petiolo gracili limbo dimidio æquilongus aut brevior villosus. Stipulæ vix millimetrales, puberulæ. Cymæ in axillis superioribus solitariae, cum pedunculo filiformi glaberrimoque folio subæquilongæ, ramulis divaricatis, glomerulis approximatis; floribus subsessilibus, masculis paucissimis, perigonii lobis sub apice mucronatis, alabastro submillimetrali; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achæniisque submillimetrali subtiliter annulatis scabriusculo brevioribus. — A P. pubescente differt foliis utrinque glabris, stipulis brevibus et achæniis orbicularibus. (v. s.)

127. P. CHRYSOSPLENIOIDES (Wedd. in Ann. sc. nat. l. c. p. 231; Monogr. p. 270), dioica?, foliis ovato-vel obovato-rotundatis utrinque obtusissimis aut basi subcordatis in dimidio superiore crenatis ciliatis limbo cæterum glabro aut sparsim piloso trinerviis, stipulis late ovatis persistentibus, cymis unisexualibus femineis laxiusculis, pedunculis quam petioli paulo brevioribus. ♀ In Nova Granata, prope Fusaguasuga (Goudot); prope Colonia Tovar (Fendler). — Herba, caulibus radicanibus in specimine unico examinato 1-1½ decim. longis demum adscendentibus pubescenti-pilosis apice rosulato-foliosis, internodiis inferioribus elongatis. Folia ½-5 centim. longa, 4-25 millim. lata, crenis obtusissimis, nervis basilaribus fere ad apicem limbi pertingentibus, lamina exsiccata membranacea, supra obscure viridi cystolithisque nisi in nervis et margine omnino destituta, subtus pallidior et cystolithis undique lineolata, petiolo pubescente quam limbus dimidius plerumque brevior. Stipulæ late amplexicaules, 1½ millim. longæ, cystolithis striatæ. Flores masc. ad basim cymarum fem. subsessiles (an semper?), perigonio hirtello. Cymæ fem. in axillis superioribus solitariae, foliis breviores, ramulis brevibus patulis, glomerulis approximatis; perigonii fructiferi segmento intermedio cucullato laterales superante achæniique ovati semimillimetralis circinatim scabriusculi dimidiam longitudinem circiter æquante. — Habitu P. pubescenti valde proxima. (v. s.)

128. P. MAXIMILIANI (Miq. in Mart. Fl. bras. fasc. 12. p. 200), monoica?, caule superne pubescente, foliis ovatis vel elliptico-ovatis acutiusculis basi rotundatis obtusisve grosse serratis tri-vel subquinque-nerviis nervis subtus pubescentibus, stipulis ovato-oblongis serius deciduis, cymis densiusculis multifloris, pedunculis quam petioli sæpius longioribus. ♀ In Brasiliâ, prope Ilheos (Maxim. Neuwied). — Wedd. Monogr. p. 271. Urtica iners Nees et Mart. in Nov. act. Acad. L. C. Nat. Cur. 11. p. 33. excl. syn. — Herba, caule 3-4½ decim. alto erecto aut basi nonnihil radicante angulato. Folia accrescentia, 3½-10 centim. longa, 2½-6 centim. lata, nervis basilaribus ultra medium limbum productis, lamina exsiccata membranacea cystolithis minutis fusiformibus punctiformibusque conspersa subtusque sub lente hic illic maculis fusciscentibus notata; petiolo 1-3½ centim. longo pubescente. Stipulæ inflorescentiarum basim fere amplectentes. Cymæ fem. glabræ, foliis fasciculatim confertis, perigonii fructiferi segmento intermedio carinato lateralibus multo majore, achænio ovato lævisculo. (v. s. in hb. Mart.)

129. P. HIRTELLA (Miq. in Mart. Fl. bras. fasc. 12. p. 202), foliis ellipticis vel ovato-ellipticis acutiusculis obtusisve excepta basi crenato-serratis trinerviis supra sparsim pilosis subtus in nervis hirtis, stipulis late ovatis persistentibus cymis densis longiuscule pedunculatis. ♀ In sylvis Brasiliæ prope Ilheos (Maxim. Neuwied). — Wedd. Monogr. p. 260. Boehmeria repens Nees et Mart. in Nov. Act. Soc. nat. Cur. 11. p. 36. Herba tenera, repens, ramulis petiolisque hirtellis. Folia infima 5 millim. superiora 2-3 centim. longa, lamina exsiccata membranacea, subtus pallida, cystolithis linearibus utrinque dense obsita, petiolo 2-8-millimetrali. Cymarum pedunculi bractea membranacea subtruncata basi vaginati. — Habitus sequentis.

130. P. PUBESCENS (Liebmann, in Vidensk. Selsk. Skr. 5 Række, naturv. og math. Afd. 2. p. 302. in adnot.), monoica vel dioica basi radicans, foliis late ovatis vel elliptico-obovatis rotundatisve utrinque obtusis vel basi subcordatis supra trientem inferiorem limbi grosse crenatis vel crenato-serratis trinerviis utrinque sparsim pilosis margineque ciliatis, stipulis late ovatis rotunda-

tisve persistentibus. cymis vulgo androgynis laxis multifloris, pedunculis quam petioli longioribus. ♀ In sylvis humidis Americæ tropicalis sat frequens præsertim in Brasilia. — Wedd. Monogr. p. 258; Miq. in Mart. Fl. bras. fasc. 12. p. 202; Griseb. Fl. Br. W. Ind. p. 158. P. Gaudichaudiana Wedd. in Ann. sc. nat. l. c. p. 226. P. grossecrenata Miq. l. c. p. 199. t. 68. — Herba, caulibus sæpius longe repentibus 2-6 decim. longis demum ramisque adscendentibus pubescentibus, exsiccatis cystolithis parvis fusiformibus undique inspersis, internodiis elongatis (inferioribus interdum decimetralibus). Folia 1½-8 centim. longa, tantumdemque fere lata, superiora cæteris sæpe majora, dentibus obtusissimis acutiusculisve, nervis basilaribus trientem superiorem limbi attingentibus subtus prominulis longiusculeque pilosis, lamina exsiccata membranacea utrinque cystolithis fusiformibus tenuissimis (in inferiore pagina autem minoribus crebrioribusque) inspersa, petiolo longitudine varia pubescente. Stipulæ 2-5 millim. longæ, laxæ, ciliolatæ. Cymæ ex axillis præsertim superioribus ortæ, folia persæpe superantes, interdum unisexuales (masculæ tunc densiores), pedunculo filiformi 2-4 centim. longo, ramulis laxè patulis, glomerulis ut plurimum discretis; floribus masc. in cymis androgynis paucis, breviter pedicellatis, lobis perigonii sub apice longiuscule mucronatis, mucronibus pilosulis in alabastro 2 millim. longo discretis; floribus fem. subsessilibus; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achænioque vix semimillimetræli fere æquilongæ. (v. s.)

β, *involutrata*, foliis quam in typo brevius petiolatis dentibus que minoribus instructis superioribus valde approximatis cymas breves involutrantibus. — In ins. S. Vincent (Macrae). — *Urtica involutrata* Sims Bot. Mag. 51. t. 2481.

γ, *montana*, foliis late ovatis subacuminatis obtusiusculis supra glaberrimis. — In ins. Antillis et Venezuela (Fendler, n. 1243). P. *montana* Wedd. l. c. n. 228. (v. s.)

δ, *fuscata*, dioica, foliis late ovatis ellipticisve apice attenuatis obtusiusculisque rariisve subacuminatis supra pilosis subtus in nervis venulisque villosis, dentibus obtusis acutisve. — In imperio mexicano, prov. Oaxaca et Vera Cruz (Galeotti, n. 313; Linden, n. 41 et 42). P. *fuscata* Liebm. l. c. p. 300; Wedd. Monogr. p. 255. P. *xalapensis* ejusd. Monogr. p. 255. (v. s.)

ε, *rupicola*, foliis angustius ovatis subacuminatis obtusiusculis subtriplicinerviis utrinque molliter pubescentibus, dentibus obtusis. — In imperio mexicano, prope Tabasco, ad scopulos (Linden, n. 71) et in monte Orizaba (Botteri). Pilea *rupicola* Wedd. in Ann. sc. nat. l. c. p. 224. (v. s.)

ζ, *Guyanensis*, foliis ovatis ellipticisve obtusiusculis supra glabris nervis subtus sparsim pilosis, pedunculis quam petioli brevioribus. — In nemo ribus Guyanæ gallicæ, ad scopulos (Melinon, n. 55; Schomburgk, n. 357). P. *guyanensis* Wedd. in Ann. sc. nat. l. c. p. 221; Monogr. p. 266. (v. s.)

Nonnullæ specierum sequentium huic stirpi tanquam varietates verisimiliter etiam adscribendæ.

131. P. *OBTUSATA* (Liebm. l. c. p. 300, in adnot.), dioica, caule piloso, foliis late ovatis utrinque obtusis grosse et obtuse crenatis subquinque-nerviis supra glabris subtus ad nervos pilosis, stipulis late ovatis, cymis diffuse ramosis longe pedunculatis, glomerulis discretis. ♀ ? In ins. Monserrat (Ryan). — Wedd. Monogr. p. 261. *Urtica obtusata* hb. Vahl. — Herba caule sesquidecimetrali tetragono articulato superne piloso strigulosoque. Folia 5-6 centim. longa, 3½-4 centim. lata, exsiccata utrinque cystolithis minutissimis fusiformibus obsita, supra obscure viridia subtus glaucescentia, petiolo 2½-3½ centim. longo pilosulo. Stipulæ obtusæ. Cymæ solitariae, pedunculis 5 centim. longis, achænio ovato marginato subtilissime granulato. (Descript. ex Liebm. l. c.)

vix prominentibus, lamina exsiccata membranacea cystolithis fusiformibus linearibusve utrinque sed supra præsertim conspersa subtus pallida, petiolo pubescente quam limbus dimidiis sæpe brevior. Stipulæ 1-2 millim. longæ. Cymæ vix pisi minoris magnitudine, pedunculis nunc petiolum vix æquantibus nunc multo longioribus, glomerulis paucis subdiscretis confluentibusve, floribus masc. perigonio glabriusculo segmentis sub apice mucrone conico longiusculo instructis, alabastro sesquimillimetrali; perigonii fructiferi segmento intermedio cucullato achænio ovato lævi vel scabriusculo brevior. .

β, *pusilla*, caulibus foliisque multo minoribus quam in typo.

γ, *strigosa*, foliis utrinque at præcipue in nervis paginæ inferioris sicut rami strigoso-pilosis, cymis masc. e glomerulo unico interdum ad basim cymarum fem. sessili constantibus, perigonio pilosulo. ♀? In Peruvia (Mathews, n. 2031). — *P. strigosa* Wedd. in Ann. sc. nat. l. c. p. 225; Monogr. p. 262. (v. s. in hb. Mus. par.)

139. *P. ROTUNDATA* (Griseb. Fl. Br. W. Ind. p. 157; Cat. pl. cub. p. 60), monoica, repens, foliis parvis rotundatis obovatisve in dimidia parte superiore obtuse pauciserratis sparsim pilosis trinerviis, stipulis rotundatis, cymis unisexualibus e glomerulo unico constantibus, pedunculis petiolum superantibus. ♀? In ins. Jamaica. — Caulis 3-10 centim. longus, demum adscendens, glaber, nodis inferioribus aphyllis radicanibusque. Folia 6-20 millim. lata, basi subtruncata, pilis paucis diaphanis conspersa aut glabra, exsiccata membranacea, cystolithis linearibus in pagina superiore præsertim conspicuis ad marginem crebrioribus. Cymæ ex axillis superioribus nascentes, 5 millim. latæ, pedunculo folium interdum æquante. — Habitus *Peperomiæ* minoris.

β, *subintegrifolia* (Griseb. Pl. Wr. in Mem. Am. Acad. sc. et art. n. ser. 8 1860, p. 174), foliis obovato-rotundatis apice vage crenatis supra pilosis, petioliis superne villosis. ♀? Prope Monte Verde, in Cuba orientali (Wright, n. 1457). Caules filiformes, repentes, intricati. Folia 3-6 millim. longa, omnino integra vel apice subcrenata, basi obtusa, nervis basilariibus in lamina exsiccata bene conspicuis dimidium limbi parum superantibus, cystolithis subtus nullis, petiolo gracili 1-3-millimetrali. Cymæ minimæ, pedunculo filiformi folium interdum superante suffultæ, floribus masc. et fem. breviter pedicellatis; perigonii masc. glabri segmentis sub apice longiuscule mucronatis alabastro sesquimillimetrali; perigonii fructiferi segmento intermedio vix cucullato lateralia longe superante et quam achænium ovatum sesquimillimetrale brevior. (v. s.)

140. *P. TRICHOTOMA* (Liebm. l. c. p. 303), monoica, caule basi radicante pilosulo, foliis ovatis basi obtusis leviterque emarginatis obtuse crenatis trinerviis ciliatis supra molliter pilosis subtus puberulis, stipulis ovato-rotundatis, cymis androgynis iterum atque iterum trichotomis, pedunculo petiolum superante. ♀? Cult. in hort. bot. Hafn. ex Mexico (Liebmann). — Wedd. Monogr. p. 262. — Herba, caule 1½-3 decim. adscendente terete. Folia 3½ centim. longa, 2-2½ centim. lata, supra saturate viridia, subtus pallidiora, petiolo subcentimetrali puberulo. Stipulæ cucullatæ, pellucidæ, ciliatæ. Cymæ solitariae vel geminae, pedunculo 1-1½ centim. longo glabro; floribus masc. ad basim inflorescentiæ rariusculis quam feminei majoribus pedicellatis carneis. Achænium ovatum. — Stirps affinis *P. densifloræ* (teste auctore), sed distincta foliis pilosis et cæt.
141. *P. AURICULATA* (Liebm. l. c. p. 299), dioica, foliis rotundato-rhombeis, basi cuneatis grosse crenatis trinerviis ciliatis supra sparsim pilosis subtus glabriusculis, stipulis ovatis persistentibus, cymulis pedunculatis petiolos paulo superantibus. ♀? In Costarica, prope Cartago (Ersted). — Wedd. Monogr. p. 253. — Herba, caule a basi ramoso 6-11 centim. longo tetragono glabro nodoso. Folia 8-14 millim. longa, 5-12 millim. lata, supra obscure viridia, cystolithis minutissimis fusiformibus obsita, subtus glauco-viridia et strigulosa, petiolo laminæ subæquilongo glabro. Stipulæ scariosæ, obtusæ, revolutæ, glabræ. Cymulæ paucifloræ; perigonii fructiferi segmentum intermedium lateraliibus majus crassum, auriculatum, achænio majusculo ovato lævi marginato multo brevius. (Descript. ex Liebmann).



142. *P. URTICELLA*, monoica, caulibus debilibus repentibusque apice petiolisque pubescentibus, foliis parvis ovatis obtusiusculis basi rotundata excepta grossiuscule subinciso-serratis tri-tripli-nerviis supra sparsim pilosis subtus in nervis hirtulis, stipulis rotundatis scariosis persistentibus, cymulis e glomerulo unico constantibus androgynis, pedunculo folium superantibus. (1) In Venezuela, inter Valencia et Campanero, prope Colonia Tovar, altit. 1200 metr. (Fendler, n. 2429). — Herba tenera, caule simplice 10-15 centim. longo in dimidia parte inferiore decumbente et radicante dein adscendente, internodiis 1-3-centimetralibus. Folia apicalia multo majora quam inferiora et magis approximata, 1-1½ centim. longa, 5-10 millim. lata, serraturis obtusiusculis, dente apicali oblongo cæteris nonnihil minore, triente vel quarta parte inferiore limbi integro, nervis basilaribus nonnunquam paulo supra basim e costa nascentibus medietatem limbi vix superantibus; lamina exsiccata tenuiter membranacea, nervis subtus bene conspicuis, utrinque cystolithis linearibus sparsim notata; petiolo gracili 5-8 millim. longo. Cymulæ paucifloræ, globulosæ, in axillis superioribus solitariæ, pedunculo tenui 15-20 millim. longo glabro suffultæ; floribus subsessilibus pedicellatisve, perigonii masc. segmentis sub apice conspicue mucronatis, alabastro mucronibus inclusis circiter bimillimetrali, perigonii fem. segmento intermedio longiuscule cucullato lateralibus longe superantibus achænioque ovato millimetrali lævi parum brevioris. — Specimina minora *Urticæ* urentis bene refert. (v. s.)

143. *P. PULEGIFOLIA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 213; Monogr. p. 235), dioica, ramosissima, ramis ramulisque villosis, foliis parvis ovatis obtusis deorsum cuneatis crenato-serratis (crenis in utroque margine 2-5) trinerviis glabris, stipulis conspicuis ovatis persistentibus, cymis masc. e glomerulo unico constantibus, pedunculis quam petioli paulo longioribus. (2) In Peruvia (Dombey). — *Urtica pulegifolia* Poir. Encycl. suppl. 4. p. 224. — Herba, caulibus elongatis prostratis? haud radicantibus basi interdum lignescentibus gracilibus angulato-sulcatis, internodiis 1-3-centimetralibus, ramis patentibus subdistichis valde foliosis. Folia in eodem jugo inaequimagna, 3-9 millim. longa, 2-6 millim. lata, nervis basilaribus subtus haud prominentibus dimidio limbo vix longioribus, lamina exsiccata firmula subtus glaucescente cystolithis linearibus in pagina superiore majusculis plerisque transversalibus et in inferiore minoribus potiusque longitiorum dispositis ornata, petiolo 1-3 millim. longo villosulo. Stipulæ sæpe petiolis subæquilongæ, caulem parum amplectentes. Cymulæ masc. (femineis non visis) in axillis ramulorum superiorum solitariæ, foliis breviores, 2-5-floræ, pedunculo puberulo, floribus breviter vel longiuscule pedicellatis, lobis perigonii sub apice mucrone conico brevi instructis, alabastris 1½-2 millim. longis (v. s. in hb. Mus. par. et Juss.)

B. — *Brevipedunculatæ, sive pedunculis (saltem cymarum fructiferarum) quam petioli longioribus.*

144. *P. HIRUTA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 220. exclus. var. β; Monogr. p. 251), dioica, caule elato magis minusve villosulo, foliis majusculis ovatis ellipticisve acuminatis basi cuneatis obtusatis rotundatisve fere in toto margine grossiuscule crenato-serratis insigniter triplinerviis supra mox glabratiss nervis primariis subtus petioloque ferrugineo-villosis, stipulis majusculis ovatis deciduis, cymis masc. et fem. floribundis laxè patuleque ramosis ex axillis pluribus ortis subsessilibus. 3 In Nova Granata (Pavon). *Urtica hirsuta* Pav. mscr. — Caulis erectus, robustus, angulato-sulcatus, internodiis elongatis, apice plerumque hirsutis densa (in sicco rufa) indutus, inferne in sulcis solum villosus vel omnino glabratus. Folia 8-15 centim. longa, 3-7 centim. lata, acumine angusto grosse serrato, limbo juxta petiolum interdum subabrupte angustato integroque, serraturis plerumque acutis extus haud raro denticulo instructis, nervis basilaribus in limbo inferne cuneato altius nascentibus quam in folio basi rotundato, lamina exsiccata utrinque fere concolore rigide membranacea cystolithis parvis fusiformibus subæqualibus undique crebre obsita, petiolo 1-5 centim. longo dense ferrugineo-hirsuto aut demum fere glabrato. Stipulæ subcentimetrales, vix amplexicaules, mox

superantibus, lamina exsiccata tenuiter membranacea cystolithis linearibus utrinque conspersa, petiolo gracili limbo dimidio æquilongo aut brevior villosa. Stipulæ vix millimetrales, puberulæ. Cymæ in axillis superioribus solitariae, cum pedunculo filiformi glaberrimoque folio subæquilongæ, ramulis divaricatis, glomerulis approximatis; floribus subsessilibus, masculis paucissimis, perigonii lobis sub apice mucronatis, alabastro submillimetrali; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achænioque submillimetrali subtiliter annulatum scabriusculo brevior. — A P. pubescente differt foliis utrinque glabris, stipulis brevibus et achæniis orbicularibus. (v. s.)

127. P. CHRYSOSPLENIODES (Wedd. in Ann. sc. nat. l. c. p. 231; Monogr. p. 270), dioica?, foliis ovato-vel obovato-rotundatis utrinque obtusissimis aut basi subcordatis in dimidio superiore crenatis ciliatis limbo cæterum glabro aut sparsim piloso trinerviis, stipulis late ovatis persistentibus, cymis unisexualibus femineis laxiusculis, pedunculis quam petioli paulo brevioribus. ♀ In Nova Granata, prope Fusaguasuga (Goudot); prope Colonia Tovar (Fendler). — Herba, caulibus radicanibus in specimine unico examinato 1-1½ decim. longis demum adscendentibus pubescenti-pilosis apice rosulato-foliosis, internodiis inferioribus elongatis. Folia ½-5 centim. longa, 4-25 millim. lata, crenis obtusissimis, nervis basilaribus fere ad apicem limbi pertingentibus, lamina exsiccata membranacea, supra obscure viridi cystolithisque nisi in nervis et margine omnino destituta, subtus pallidior et cystolithis undique lineolata, petiolo pubescente quam limbus dimidius plerumque brevior. Stipulæ late amplexicaules, 1½ millim. longæ, cystolithis striatæ. Flores masc. ad basim cymarum fem. subsessiles (an semper?), perigonio hirtello. Cymæ fem. in axillis superioribus solitariae, foliis breviores, ramulis brevibus patulis, glomerulis approximatis; perigonii fructiferi segmento intermedio cucullato laterales superante achænioque ovati semimillimetralis circinatim scabriusculi dimidiam longitudinem circiter æquante. — Habitu P. pubescenti valde proxima. (v. s.)
128. P. MAXIMILIANI (Miq. in Mart. Fl. bras. fasc. 12. p. 200), monoica?, caule superne pubescente, foliis ovatis vel elliptico-ovatis acutiusculis basi rotundatis obtusive grosse serratis tri-vel subquinque-nerviis nervis subtus pubescentibus, stipulis ovato-oblongis serius deciduis, cymis densiusculis multifloris, pedunculis quam petioli sæpius longioribus. ♀ In Brasilia, prope Ilheos (Maxim. Neuwied). — Wedd. Monogr. p. 271. Urtica iners Nees et Mart. in Nov. act. Acad. L. C. Nat. Cur. 11. p. 33. excl. syn. — Herba, caule 3-4½ decim. alto erecto aut basi nonnihil radicante angulato. Folia accrescentia, 3½-10 centim. longa, 2½-6 centim. lata, nervis basilaribus ultra medium limbum productis, lamina exsiccata membranacea cystolithis minutis fusiformibus punctiformibusque conspersa subtusque sub lente hic illic maculis fusciscentibus notata; petiolo 1-3½ centim. longo pubescente. Stipulæ inflorescentiarum basim fere amplectentes. Cymæ fem. glabræ, floribus fasciculatim confertis, perigonii fructiferi segmento intermedio carinato lateralibus multo majore, achænio ovato læviusculo. (v. s. in hb. Mart.)
129. P. HIRTELLA (Miq. in Mart. Fl. bras. fasc. 12. p. 202), foliis ellipticis vel ovato-ellipticis acutiusculis obtusive excepta basi crenato-serratis trinerviis supra sparsim pilosis subtus in nervis hirtis, stipulis late ovatis persistentibus cymis densis longiuscule pedunculatis. ♀ In sylvis Brasiliæ prope Ilheos (Maxim. Neuwied). — Wedd. Monogr. p. 260. Boehmeria repens Nees et Mart. in Nov. Act. Soc. nat. Cur. 11. p. 36. Herba tenera, repens, ramulis petiolisque hirtellis. Folia infima 5 millim. superiora 2-3 centim. longa, lamina exsiccata membranacea, subtus pallida, cystolithis linearibus utrinque dense obsita, petiolo 2-8-millimetrali. Cymarum pedunculi bractea membranacea subtruncata basi vaginati. — Habitus sequentis.
130. P. PUBESCENS (Liehmann, in Vidensk. Selsk. Skr. 5 Række, naturv. og math. Afd. 2. p. 302. in adnot.), monoica vel dioica basi radicans, foliis late ovatis vel elliptico-obovatis rotundatisve utrinque obtusis vel basi subcordatis supra trientem inferiorem limbi grosse crenatis vel crenato-serratis trinerviis utrinque sparsim pilosis margineque ciliatis, stipulis late ovatis rotunda-

tisve persistentibus, cymis vulgo androgynis laxis multifloris, pedunculis quam petioli longioribus. ♀ In sylvis humidis Americæ tropicalis sat frequens præsertim in Brasilia. — Wedd. Monogr. p. 258; Miq. in Mart. Fl. bras. fasc. 12. p. 202; Griseb. Fl. Br. W. Ind. p. 158. P. Gaudichaudiana Wedd. in Ann. sc. nat. l. c. p. 226. P. grossecrenata Miq. l. c. p. 199. t. 68. — Herba, caulibus sæpius longe repentibus 2-6 decim. longis demum ramisque adscendentibus pubescentibus, exsiccatis cystolithis parvis fusiformibus undique inspersis, internodiis elongatis (inferioribus interdum decimetralibus). Folia 1½-8 centim. longa, tantumdemque fere lata, superiora cæteris sæpe majora, dentibus obtusissimis acutiusculisve, nervis basilaribus trientem superiorem limbi attingentibus subtus prominulis longiusculeque pilosis, lamina exsiccata membranacea utrinque cystolithis fusiformibus tenuissimis (in inferiore pagina autem minoribus crebrioribusque) inspersa, petiolo longitudine varia pubescente. Stipulæ 2-5 millim. longæ, laxæ, ciliolatæ. Cymæ ex axillis præsertim superioribus ortæ, folia persæpe superantes, interdum unisexuales (masculæ tunc densiores), pedunculo filiformi 2-4 centim. longo, ramulis laxè patulis, glomerulis ut plurimum discretis; floribus masc. in cymis androgynis paucis, breviter pedicellatis, lobis perigonii sub apice longiuscule mucronatis, mucronibus pilosis in alabastro 2 millim. longo discretis; floribus fem. subsessilibus; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achænioque vix semimillimetrali fere æquilongio. (v. s.)

β, *involutrata*, foliis quam in typo brevius petiolatis dentibus que minoribus instructis superioribus valde approximatis cymas breves involutratibus. — In ins. S. Vincent (Macrae). — *Urtica involutrata* Sims Bot. Mag. 51. t. 2481.

γ, *montana*, foliis late ovatis subacuminatis obtusiusculis supra glaberrimis. — In ins. Antillis et Venezuela (Fendler, n. 1243). P. *montana* Wedd. l. c. n. 228. (v. s.)

δ, *fuscata*, dioica, foliis late ovatis ellipticisve apice attenuatis obtusiusculisque rariusve subacuminatis supra pilosis subtus in nervis venulisque villosis, dentibus obtusis acutisve. — In imperio mexicano, prov. Oaxaca et Vera Cruz (Galeottii, n. 313; Linden, n. 41 et 42). P. *fuscata* Liebm. l. c. p. 300; Wedd. Monogr. p. 255. P. *xalapensis* ejusd. Monogr. p. 255. (v. s.)

ε, *rupicola*, foliis angustius ovatis subacuminatis obtusiusculis subtriplicinerviis utrinque molliter pubescentibus, dentibus obtusis. — In imperio mexicano, prope Tabasco, ad scopulos (Linden, n. 71) et in monte Orizaba (Botteri). Pilea *rupicola* Wedd. in Ann. sc. nat. l. c. p. 224. (v. s.)

ζ, *Guyanensis*, foliis ovatis ellipticisve obtusiusculis supra glabris nervis subtus sparsim pilosis, pedunculis quam petioli brevioribus. — In nemo ribus Guyanæ gallicæ, ad scopulos (Melinon, n. 55; Schomburgk, n. 357). P. *guyanensis* Wedd. in Ann. sc. nat. l. c. p. 221; Monogr. p. 266. (v. s.)

Nonnullæ specierum sequentium huic stirpi tanquam varietates verisimiliter etiam adscribendæ.

131. P. *OBTUSATA* (Liebm. l. c. p. 300, in adnot.), dioica, caule piloso, foliis late ovatis utrinque obtusis grosse et obtuse crenatis subquinque-nerviis supra glabris subtus ad nervos pilosis, stipulis late ovatis, cymis diffuse ramosis longe pedunculatis, glomerulis discretis. ♀? In ins. Monserrat (Ryan). — Wedd. Monogr. p. 261. *Urtica obtusata* hb. Vahl. — Herba caule sesquidecimetrali tetragono articulo superne piloso strigulosoque. Folia 5-6 centim. longa, 3½-4 centim. lata, exsiccata utrinque cystolithis minutissimis fusiformibus obsita, supra obscure viridia subtus glaucescentia, petiolo 2½-3½ centim. longo pilosulo. Stipulæ obtusæ. Cymæ solitariae, pedunculis 5 centim. longis, achænio ovato marginato subtilissime granulato. (Descript. ex Liebm. l. c.)

132. *P. SANCTÆ CRUCIS* (Liebm. l. c. p. 301, in adnot.), dioica, foliis late ovatis acutis basi obtusis et leviter emarginatis grosse et obtuse crenatis (crenatis obsolete denticulatis v. repandis) 5-nerviis (nervis 2 lateralibus tenuissimis) ciliatis cæterum glabriusculis, stipulis cordato-rotundatis, cymis fem. longe pedunculatis quam folia longioribus diffuse ramosis, glomerulis discretis, achænio lenticulari-ovato marginato subtilissime granulato quam perigonium duplo longiore. ♀? In insula Sanctæ Crucis Antillarum (Cæsted). — Wedd. Monogr. p. 261. — Herba, caule e basi decumbente et radicante adscendente, sesquidecimetrali et quod excedit, tetragono, exsiccato superne cystolithis minutis insito. Folia 7-9 centim. longa,  $4\frac{1}{2}$ -7 centim. lata, supra obscure viridia, petiolo  $3\frac{1}{2}$  centim. longo angulato adnate striguloso. Stipulæ semiamplexicaules, scariosæ, fuscæ, glabræ. Cymæ fem. (solum visæ) solitariae, pedunculo 5-6 centim. longo angulato glabro, floribus minimis pedicellatis, achænio lenticulari-ovato fusco-olivaceo emarginato subtilissime granulato quam perigonium duplo longiore. (Descript. ex Liebmann.)
133. *P. LONGIPES* (Liebm. l. c. p. 301), dioica, foliis late ovatis acuminatis basi subcordatis rotundatisve grosse serratis (serraturis acutis) quinquenerviis (nervis 2 lateralibus tenuioribus) ciliatis supra sparsim setosis ac pilosis subtus sparsim pilosis stipulis strigulosisque, stipulis late ovatis, cymis fem. longis longe pedunculatis. ♀? In Mexici prov. Oaxaca, locis humidis umbrosisque, altit. 1600 metr. (Liebmann). — Wedd. Monogr. p. 250. — Herba, caule  $1\frac{1}{2}$ -3-decimetrali ramoso tetragono inferne nudo superne piloso et fusco-pubescente. Folia inferiora minora, superiora gradatim magnitudinæ increscentia, 7-9 centim. longa,  $4\frac{1}{2}$ -7 centim. lata, supra obscure viridia, subtus glaucescentia, exsiccata utrinque cystolithis minimis fusiformibus conspersa, petiolo  $3\frac{1}{2}$ -5 centim. longo pubescente. Stipulæ membranaceæ, fuscæ. Cymæ fem. solitariae aut geminae, 7-12 centim. longæ, pedunculo 5-6-centimetrali patentissime ramosæ, floribus breviter pedicellatis, achænio marginato fusco subtilissime granuloso. (v. s.)
134. *P. ACUMINATA* (Liebm. l. c. p. 302), dioica, caule adscendente glabro, foliis ovato-ellipticis longe acuminatis basi cordatis grosse acuteque crenatodentatis trinerviis ciliatis supra sparsim pilosis nervis subtus fusco-pilosis, stipulis lanceolatis repandis, cymis laxis longe pedunculatis. ♀? In Mexici locis humidis umbrosisque prope Mirador (Liebmann). — Wedd. Monogr. p. 248. — Herba, caule  $1-2\frac{1}{2}$  decim. longo, basi prostrato et radicante, simplice vel bifurco. Folia  $4\frac{1}{2}$ -6 centim. longa, 15-22 millim. lata, supra obscure viridia, subtus glauca, lamina exsiccata utrinque cystolithis inpersa, petiolo 1-2 centim. longo fusco-piloso. Stipulæ 4 millim. longæ. Cymæ masc. diffuse ramosæ, pedunculo  $3\frac{1}{2}$ -4 centim. longo glabro, glomerulis discretis, floribus breviter pedicellatis; femineæ solitariae, masculis conformes, achænio ovato. (v. s.)
- β, *Cruzensis*, caule apice villosa. ♀? In imperii mexicanæ prov. Vera Cruz (Linden, n. 42). — *P. cruzensis* Wedd. in Ann. sc. nat. l. c. p. 228. Cymæ laxissimæ; perigonii masc. segmenta sub apice longiuscule mucronata. (v. s.)
135. *P. DIFFUSA* (Wedd. in Ann. sc. nat. l. c. p. 222; Monogr. p. 249), monoica, prostrata, foliis rhomboideo-vel lanceolato-ovatis magis minus acuminate basi obtusis ultra trientem vel dimidium inferiorem acute grossiusculaque serratis trinerviis supra glabris nervis subtus pubescentibus aut utrinque sparsim pilosis, stipulis ovato-oblongis persistentibus, cymis androgynis laxis, pedunculo petiolum longe superante. ♀? In ins. Jamaica (Swartz). — Griseb. Fl. Br. W. Ind. p. 158. *Urtica diffusa* Swartz, in Act. Holm. ann. 1785. p. 30; ejusd. Fl. Ind. occ. p. 290; Poir. Encycl. Suppl. 4. p. 218; Smith, l. c. n. 13. — Herba, caule prostrato radicante et interdum basi lignescente, glabro vel superne puberulo, ramis diffusis elongatis etiam radicantibus at demum adscendentibus, internodiis elongatis nodis plerisque aphyllis. Folia 2-6 centim. longa,  $1\frac{1}{2}$ -3 centim. lata, sensim attenuata, serraturis fere æqualibus, nervis basilaribus ad trientem superiorem limbi protractis; lamina exsiccata membranacea supra obscure viridi subtus pallida, cystolithis in pagina superiore oculo armato solum conspicuis linearibus, in inferiore nullis; petiolo fere

centimetrali gracili puberulo. Stipulæ 4-6 millim. longæ, obtusissimæ, margine interdum revolutæ. Cymæ parce et laxæ ramosæ, ramulis brevibus, glomerulis perisque discretis, pedunculo gracili folium haud raro longitudine subæquante. Achænium ovatum, læve. (v. s. typ. in hb. Swartz nunc Holm., Mus. brit. et Monac.)

136. P. LAXA (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 228), dioica, prostrata, foliis ovatis vel elliptico-ovatis longiuscule acuminatis basi acutis rotundatisve aut subcordatis fere in toto margine dentato-serratis (dentibus sæpe inæqualibus) trinerviis supra glabris subtus in nervis pubescentibus hirtulisve aut demum glabratis, stipulis ovatis persistentibus, cymis longiuscule pedunculatis, masculis capituliformibus vel e glomerulis paucissimis constante, femineis laxifloris. ½ In fruticetis, secus rivulos ins. S. Domingo (Swartz). — Griseb. Fl. Br. W. Ind. p. 158. Urtica laxa Swartz, in Act. Holm. (1787) p. 58; ejusd. Fl. Ind. occid. p. 288; Smith, l. c. n. 12. — Herba, caulibus gracilibus ramosis prostratis nonnunquam plus quam metralibus, ramis diffuse patentibus striatis glabris vel haud raro superne pubescentibus. Folia 2½-8 centim. longa, 1½-4½ centim. lata, nervis basilaribus circiter ad trientem superiorem limbi productis, dentibus sive serraturis juxta petiolum et in acumine parvis, intermediis contra grossiusculis obtusiusculisque; laminæ exsiccatæ membranaceæ pagina inferiore pallide virente cystolithisque nunc destituta et nunc punctiformibus notata, superiore iisdem corpusculis aliis brevissimis aliisque elongatis obsita, petiolo gracili pubescente vel glabrato quam dimidius limbus vulgo brevior. Stipulæ 3-5 millim. longæ, membranaceæ. Cymæ masc. densæ, pisiformes, pedunculo filiformi quam petiolus longiore, floribus pedicellatis, perigonii glabri laciniis sub apice mucrone conico instructis, mucronibus in alabastro semimillimetrali discretis. Cætera ignota. (v. s. in hb. Holm., Mus. brit. et Soc. linn.)
137. P. HEDERACEA (Wedd. in Ann. sc. nat. l. c. p. 222; Monogr. p. 263), dioica, repens, foliis rotundatis vel late ovatis obtusis basi cordatis subcordatisve grosse crenato-dentatis trinerviis ciliatis, pagina superiore pilis paucis conspersa vel omnino glabra, venis subtus patentibus pubescentibus, stipulis suborbicularibus persistentibus, cymis densis subsphæricis lobatisve longe pedunculatis. ½ In Saxosis umbrosisque ins. S. Domingo et Guadelupe (Swartz, Poiteau, Cl. Richard, etc.). — P. betulæfolia Wedd. in Ann. l. c. p. 227. Urtica betulæfolia Swartz, in Act. Holm. ann. 1787. p. 59; Fl. Ind. occ. p. 291; Smith, l. c. n. 14. U. hederacea Lam. dict. 4. p. 642. — Herba, habitu Glechomæ hederaceæ, caulibus elongatis prostratis ramosis ad nodos radicantibus villosis glabratissive, exsiccatis striatis cystolithisque punctulatis, ramis adscendentibus villosis. Folia 1-6 centim. longa et lata, nervis basilaribus trientem superiorem limbi attingentibus, lamina exsiccatæ membranaceæ cystolithis tenuibus linearibus punctiformibusque utrinque conspersa, petiolo pubescente 5-10 millim. longo. Stipulæ 2-4 millimetrales, laxæ, scariosæ. Cymæ masc. in axillis intermediis ramorum natæ, paucæ, pisi circiter magnitudine v. majores, pedunculis filiformibus quam folia longioribus, floribus pedicellatis, perigonii lobis glabris, sub apice mucronatis, mucronibus in alabastro vix millimetrali discretis. Cymæ fem. pedunculis plerumque folio brevioribus suffultæ, perigonii fructiferi segmento intermedio lateralibus multoties longiore achæniumque ovatum vix semimillimetrale læve non omnino æquante. (v. s. in hb. Mus. par., Holm. et Mus. brit.)
138. P. REPENS (Wedd. in Ann. sc. nat. l. c. p. 220; Monogr. p. 269), monica, foliis late ovatis vel elliptico-rotundatis obovatisve obtusis basi late cuneatis rotundatisve in dimidia parte superiore crenatis aut crenato-serratis supra glabris subtus in nervis pubescentibus trinerviis, stipulis late ovatis persistentibus, cymis parvis densifloris androgynis, pedunculo filiformi petiolo sæpius longiore. ④? In ins. Antillis S. Domingo (Swartz), Cuba (Wright, n. 539, 1454 et 1455), Portorico, etc. — Griseb. Fl. Br. W. Ind. p. 157; Cat. pl. cub. p. 60. Urtica repens Swartz in Act. Holm. ann. 1787. p. 61. t. 1. f. 1; Fl. Ind. occ. p. 294; Smith, l. c. n. 28. — Herba, caulibus elongatis pubescentibus ad nodos radicantibus simplicibus aut parce ramosis demum ramosisque adscendentibus apice præsertim foliosis. Folia 1-3 centim. longa, 1-2½ centim. lata, nervis basilaribus ad trientem superiorem limbi productis subtus

vix prominentibus, lamina exsiccata membranacea cystolithis fusiformibus linearibusve utrinque sed supra præsertim conspersa subtus pallida, petiolo pubescente quam limbus dimidiis sæpe brevior. Stipulæ 1-2 millim. longæ. Cymæ vix pisi minoris magnitudine, pedunculis nunc petiolum vix æquantibus nunc multo longioribus, glomerulis paucis subdiscretis confluentibusve, floribus masc. perigonio glabriusculo segmentis sub apice mucrone conico longiusculo instructis, alabastro sesquimillimetrali; perigonii fructiferi segmento intermedio cucullato achænio ovato lævi vel scabriusculo breviora.

β, *pusilla*, caulibus foliisque multo minoribus quam in typo.

γ, *strigosa*, foliis utrinque at præcipue in nervis paginæ inferioris sicut rami strigoso-pilosis, cymis masc. e glomerulo unico interdum ad basim cymarum fem. sessili constantibus, perigonio pilosulo. ♀? In Peruvia (Mathews, n. 2031). — *P. strigosa* Wedd. in Ann. sc. nat. l. c. p. 225; Monogr. p. 262. (v. s. in hb. Mus. par.)

139. *P. ROTUNDATA* (Griseb. Fl. Br. W. Ind. p. 157; Cat. pl. cub. p. 60), monoica, repens, foliis parvis rotundatis obovatisve in dimidia parte superiore obtuse pauciserratis sparsim pilosis trinerviis, stipulis rotundatis, cymis unisexualibus e glomerulo unico constantibus, pedunculis petiolum superantibus. ♀? In ins. Jamaica. — Caulis 3-10 centim. longus, demum adscendens, glaber, nodis inferioribus aphyllis radicanibusque. Folia 6-20 millim. lata, basi subtruncata, pilis paucis diaphanis conspersa aut glabra, exsiccata membranacea, cystolithis linearibus in pagina superiore præsertim conspicuis ad marginem crebrioribus. Cymæ ex axillis superioribus nascentes, 5 millim. latæ, pedunculo folium interdum æquante. — Habitus *Peperomiæ* minoris.

β, *subintegrifolia* (Griseb. Pl. Wr. in Mem. Am. Acad. sc. et art. n. ser. 8 1860, p. 174), foliis obovato-rotundatis apice vage crenatis supra pilosis, petiolis superne villosis. ♀? Prope Monte Verde, in Cuba orientali (Wright, n. 1457). Caules filiformes, repentes, intricati. Folia 3-6 millim. longa, omnino integra vel apice subcrenata, basi obtusa, nervis basilariibus in lamina exsiccata bene conspicuis dimidium limbi parum superantibus, cystolithis subtus nullis, petiolo gracili 1-3-millimetrali. Cymæ minimæ, pedunculo filiformi folium interdum superante suffultæ, floribus masc. et fem. breviter pedicellatis; perigonii masc. glabri segmentis sub apice longiuscule mucronatis alabastro sesquimillimetrali; perigonii fructiferi segmento intermedio vix cucullato lateralia longe superante et quam achæmium ovatum sesquimillimetrale breviora. (v. s.)

140. *P. TRICHOTOMA* (Liebm. l. c. p. 303), monoica, caule basi radicante pilosulo, foliis ovatis basi obtusis leviterque emarginatis obtuse crenatis trinerviis ciliatis supra molliter pilosis subtus puberulis, stipulis ovato-rotundatis, cymis androgynis iterum atque iterum trichotomis, pedunculo petiolum superante. ♀? Cult. in hort. bot. Hafn. ex Mexico (Liebmann). — Wedd. Monogr. p. 262. — Herba, caule 1½-3 decim. adscendente terete. Folia 3½ centim. longa, 2-2½ centim. lata, supra saturate viridia, subtus pallidiora, petiolo subcentimetrali puberulo. Stipulæ cucullatæ, pellucidæ, ciliatæ. Cymæ solitariae vel geminae, pedunculo 1-1½ centim. longo glabro; floribus masc. ad basim inflorescentiæ rariusculis quam feminei majoribus pedicellatis carnis. Achæmium ovatum. — Stirps affinis *P. densifloræ* (teste auctore), sed distincta foliis pilosis et cæt.

141. *P. AURICULATA* (Liebm. l. c. p. 299), dioica, foliis rotundato-rhombeis, basi cuneatis grosse crenatis trinerviis ciliatis supra sparsim pilosis subtus glabriusculis, stipulis ovatis persistentibus, cymulis pedunculatis petiolos paulo superantibus. ♀? In Costarica, prope Cartago (Ersted). — Wedd. Monogr. p. 253. — Herba, caule a basi ramoso 6-11 centim. longo tetragono glabro nodoso. Folia 8-14 millim. longa, 5-12 millim. lata, supra obscure viridia, cystolithis minutissimis fusiformibus obsita, subtus glauco-viridia et strigulosa, petiolo laminæ subæquilongo glabro. Stipulæ scariosæ, obtusæ, revolutæ, glabræ. Cymulæ paucifloræ; perigonii fructiferi segmentum intermedium lateraliibus majus crassum, auriculatum, achænio majusculo ovato lævi marginato multo brevius. (Descript. ex Liebmann).

142. *P. URTICELLA*, monoica, caulibus debilibus repentibusque apice petiolisque pubescentibus, foliis parvis ovatis obtusiusculis basi rotundata excepta grossiuscule subinciso-serratis tri-tripli-nerviis supra sparsim pilosis subtus in nervis hirtulis, stipulis rotundatis scariosis persistentibus, cymulis e glomerulo unico constantibus androgynis, pedunculo folium superantibus. (1) In Venezuela, inter Valencia et Campanero, prope Colonia Tovar, altit. 1200 metr. (Fendler, n. 2429). — Herba tenera, caule simplice 10-15 centim. longo in dimidia parte inferiore decumbente et radicante dein ascendente, internodiis 1-3-centimetralibus. Folia apicalia multo majora quam inferiora et magis approximata, 1-1½ centim. longa, 5-10 millim. lata, serraturis obtusiusculis, dente apicali oblongo cæteris nonnihil minore, triente vel quarta parte inferiore limbi integro, nervis basilaribus nonnunquam paulo supra basim e costa nascentibus medietatem limbi vix superantibus; lamina exsiccata tenuiter membranacea, nervis subtus bene conspicuis, utrinque cystolithis linearibus sparsim notata; petiolo gracili 5-8 millim. longo. Cymulæ paucifloræ, globulosæ, in axillis superioribus solitariæ, pedunculo tenui 15-20 millim. longo glabro suffultæ; floribus subsessilibus pedicellatisve, perigonii masc. segmentis sub apice conspicue mucronatis, alabastro mucronis inclusis circiter bimillimetrali, perigonii fem. segmento intermedio longiuscule cucullato lateralibus longe superantibus achænioque ovato millimetrali lævi parum brevior. — Specimina minima Urticæ urentis bene refert. (v. s.)

143. *P. PULEGIFOLIA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 213; Monogr. p. 235), dioica, ramosissima, ramis ramulisque villosis, foliis parvis ovatis obtusis deorsum cuneatis crenato-serratis (crenis in utroque margine 2-5) trinerviis glabris, stipulis conspicuis ovatis persistentibus, cymis masc. e glomerulo unico constantibus, pedunculis quam petioli paulo longioribus. (1?) In Peruvia (Dombey). — Urtica pulegifolia Poir. Encycl. suppl. 4. p. 224. — Herba, caulibus elongatis prostratis? haud radicantibus basi interdum lignescentibus gracilibus angulato-sulcatis, internodiis 1-3-centimetralibus, ramis patentibus subdistichis valde foliosis. Folia in eodem jugo inæquimagna, 3-9 millim. longa, 2-6 millim. lata, nervis basilaribus subtus haud prominentibus dimidio limbo vix longioribus, lamina exsiccata firmula subtus glaucescente cystolithis linearibus in pagina superiore majusculis plerisque transversalibus et in inferiore minoribus potiusque longitrorsum dispositis ornata, petiolo 1-3 millim. longo villosulo. Stipulæ sæpe petiolis subæquilongæ, caulem parum amplectentes. Cymulæ masc. (femineis non visis) in axillis ramulorum superiorum solitariæ, foliis breviores, 2-5-floræ, pedunculo puberulo, floribus breviter vel longiuscule pedicellatis, lobis perigonii sub apice mucrone conico brevi instructis, alabastris 1½-2 millim. longis (v. s. in hb. Mus. par. et Juss.)

B. — *Brevipedunculatæ, sive pedunculis (saltem cymarum fructiferarum) quam petioli longioribus.*

144. *P. HIRUTA* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 220. exclus. var. β; Monogr. p. 251), dioica, caule elato magis minusve villosulo, foliis majusculis ovatis ellipticisve acuminatis basi cuneatis obtusatis rotundatisve fere in toto margine grossiuscule crenato-serratis insigniter triplinerviis supra mox glabratissimis nervis primariis subtus petioloque ferrugineo-villosis, stipulis majusculis ovatis deciduis, cymis masc. et fem. floribundis laxè patuleque ramosis ex axillis pluribus ortis subsessilibus. 2/ In Nova Granata (Pavon). Urtica hirsuta Pav. mscr. — Caulis erectus, robustus, angulato-sulcatus, internodiis elongatis, apice plerumque hirsutie densa (in sicco rufa) indutus, inferne in sulcis solum villosus vel omnino glabratus. Folia 8-15 centim. longa, 3-7 centim. lata, acumine angusto grosse serrato, limbo juxta petiolum interdum subabrupte angustato integroque, serraturis plerumque acutis extus haud raro denticulo instructis, nervis basilaribus in limbo inferne cuneato altius nascentibus quam in folio basi rotundato, lamina exsiccata utrinque fere concolore rigide membranacea cystolithis parvis fusiformibus subæqualibus undique crebre obsita, petiolo 1-5 centim. longo dense ferrugineo-hirsuto aut demum fere glabrato. Stipulæ subcentimetales, vix amplexicaules, mox

superantibus, lamina exsiccata tenuiter membranacea cystolithis linearibus utrinque conspersa, petiolo gracili limbo dimidio æquilongo aut brevior villoso. Stipulæ vix millimetrales, puberulæ. Cymæ in axillis superioribus solitariae, cum pedunculo filiformi glaberrimoque folio subæquilongæ, ramulis divaricatis, glomerulis approximatis; floribus subsessilibus, masculis paucissimis, perigonii lobis sub apice mucronatis, alabastro submillimetrali; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achæniisque submillimetrali subtiliter annulatis scabriusculo brevioribus. — A *P. pubescente* differt foliis utrinque glabris, stipulis brevibus et achæniis orbicularibus. (v. s.)

127. *P. CHRYSOSPLENIOIDES* (Wedd. in Ann. sc. nat. 1. c. p. 231; Monogr. p. 270), dioica?, foliis ovato-vel obovato-rotundatis utrinque obtusissimis aut basi subcordatis in dimidio superiore crenatis ciliatis limbo cæterum glabro aut sparsim piloso trinerviis, stipulis late ovatis persistentibus, cymis unisexualibus femineis laxiusculis, pedunculis quam petioli paulo brevioribus. ♀ In Nova Granata, prope Fusaguasuga (Goudot); prope Colonia Tovar (Fendler). — Herba, caulibus radicanibus in specimine unico examinato 1-1½ decim. longis demum adscendentibus pubescenti-pilosis apice rosulato-foliosis, internodiis inferioribus elongatis. Folia ½-5 centim. longa, 4-25 millim. lata, crenis obtusissimis, nervis basilaribus fere ad apicem limbi pertingentibus, lamina exsiccata membranacea, supra obscure viridi cystolithisque nisi in nervis et margine omnino destituta, subtus pallidiore et cystolithis undique lineolata, petiolo pubescente quam limbus dimidius plerumque brevior. Stipulæ late amplexicaules, 1½ millim. longæ, cystolithis striatæ. Flores masc. ad basim cymarum fem. subsessiles (an semper?), perigonio hirtello. Cymæ fem. in axillis superioribus solitariae, foliis breviores, ramulis brevibus patulis, glomerulis approximatis; perigonii fructiferi segmento intermedio cucullato laterales superante achæniisque ovati semimillimetralis circinatim scabriusculi dimidiam longitudinem circiter æquante. — Habitu *P. pubescenti* valde proxima. (v. s.)

128. *P. MAXIMILIANI* (Miq. in Mart. Fl. bras. fasc. 12. p. 200), monoica?, caule superne pubescente, foliis ovatis vel elliptico-ovatis acutiusculis basi rotundatis obtusisve grosse serratis tri-vel subquinque-nerviis nervis subtus pubescentibus, stipulis ovato-oblongis serius deciduis, cymis densiuscule multifloris, pedunculis quam petioli sæpius longioribus. ♀ In Brasilia, prope Ilheos (Maxim. Neuwied). — Wedd. Monogr. p. 271. *Urtica iners* Nees et Mart. in Nov. act. Acad. L. C. Nat. Cur. 11. p. 33. excl. syn. — Herba, caule 3-4½ decim. alto erecto aut basi nonnihil radicante angulato. Folia accrescentia, 3½-10 centim. longa, 2½-6 centim. lata, nervis basilaribus ultra medium limbum productis, lamina exsiccata membranacea cystolithis minutis fusiformibus punctiformibusque conspersa subtusque sub lente hic illic maculis fusciscentibus notata; petiolo 1-3½ centim. longo pubescente. Stipulæ inflorescentiarum basim fere amplectentes. Cymæ fem. glabræ, floribus fasciculatim confertis, perigonii fructiferi segmento intermedio carinato lateralibus multo majore, achænio ovato læviusculo. (v. s. in hb. Mart.)

129. *P. HIRTELLA* (Miq. in Mart. Fl. bras. fasc. 12. p. 202), foliis ellipticis vel ovato-ellipticis acutiusculis obtusisve excepta basi crenato-serratis trinerviis supra sparsim pilosis subtus in nervis hirtis, stipulis late ovatis persistentibus cymis densis longiuscule pedunculatis. ♀ In sylvis Brasiliæ prope Ilheos (Maxim. Neuwied). — Wedd. Monogr. p. 260. *Boehmeria repens* Nees et Mart. in Nov. Act. Soc. nat. Cur. 11. p. 36. Herba tenera, repens, ramulis petiolisque hirtellis. Folia infima 5 millim. superiora 2-3 centim. longa, lamina exsiccata membranacea, subtus pallida, cystolithis linearibus utrinque dense obsita, petiolo 2-8-millimetrali. Cymarum pedunculi bractea membranacea subtruncata basi vaginati. — Habitus sequentis.

130. *P. PUBESCENS* (Liebmann, in Vidensk. Selsk. Skr. 5 Række, naturv. og math. Afd. 2. p. 302. in adnot.), monoica vel dioica basi radicans, foliis late ovatis vel elliptico-obovatis rotundatisve utrinque obtusis vel basi subcordatis supra trientem inferiorem limbi grosse crenatis vel crenato-serratis trinerviis utrinque sparsim pilosis margineque ciliatis, stipulis late ovatis rotunda-



tive persistentibus. cymis vulgo androgynis laxis multifloris, pedunculis quam petioli longioribus.  $\frac{1}{2}$  In sylvis humidis Americæ tropicalis sat frequens præsertim in Brasilia. — Wedd. Monogr. p. 258; Miq. in Mart. Fl. bras. fasc. 12. p. 202; Griseb. Fl. Br. W. Ind. p. 158. P. Gaudichaudiana Wedd. in Ann. sc. nat. l. c. p. 226. P. grossecrenata Miq. l. c. p. 199. t. 68. — Herba, caulibus sæpius longe repentibus 2-6 decim. longis demum ramisque adscendentibus pubescentibus, exsiccatis cystolithis parvis fusiformibus undique inspersis, internodiis elongatis (inferioribus interdum decimetralibus). Folia  $1\frac{1}{2}$ -8 centim. longa, tantumdemque fere lata, superiora cæteris sæpe majora, dentibus obtusissimis acutiusculisve, nervis basilaribus trientem superiorem limbi attingentibus subtus prominulis longiusculeque pilosis, lamina exsiccata membranacea utrinque cystolithis fusiformibus tenuissimis (in inferiore pagina autem minoribus crebrioribusque) inspersa, petiolo longitudine varia pubescente. Stipulæ 2-5 millim. longæ, laxæ, ciliolatæ. Cymæ ex axillis præsertim superioribus ortæ, folia persæpe superantes, interdum unisexuales (masculæ tunc densiores), pedunculo filiformi 2-4 centim. longo, ramulis laxè patulis, glomerulis ut plurimum discretis; floribus masc. in cymis androgynis paucis, breviter pedicellatis, lobis perigonii sub apice longiuscule mucronatis, mucronibus pilosis in alabastro 2 millim. longo discretis; floribus fem. subsessilibus; perigonii fructiferi segmento intermedio cucullato lateralibus duplo longiore achænioque vix semimillimetrali fere æquilong. (v. s.)

$\beta$ , *involutrata*, foliis quam in typo brevius petiolatis dentibus que minoribus instructis superioribus valde approximatis cymas breves involutratibus. — In ins. S. Vincent (Macrae). — *Urtica involutrata* Sims Bot. Mag. 51. t. 2481.

$\gamma$ , *montana*, foliis late ovatis subacuminatis obtusiusculis supra glaberrimis. — In ins. Antillis et Venezuela (Fendler, n. 1243). P. montana Wedd. l. c. n. 228. (v. s.)

$\delta$ , *fuscata*, dioica, foliis late ovatis ellipticisve apice attenuatis obtusiusculisque rariusve subacuminatis supra pilosis subtus in nervis venulisque villosis, dentibus obtusis acutisve. — In imperio mexicano, prov. Oaxaca et Vera Cruz (Galeotti, n. 313; Linden, n. 41 et 42). P. fuscata Liebm. l. c. p. 300; Wedd. Monogr. p. 255. P. xalapensis ejusd. Monogr. p. 255. (v. s.)

$\epsilon$ , *rupicola*, foliis angustius ovatis subacuminatis obtusiusculis subtriplici nervi utrinque molliter pubescentibus, dentibus obtusis. — In imperio mexicano, prope Tabasco, ad scopulos (Linden, n. 71) et in monte Orizaba (Botteri). Pilea rupicola Wedd. in Ann. sc. nat. l. c. p. 224. (v. s.)

$\zeta$ , *Guyanensis*, foliis ovatis ellipticisve obtusiusculis supra glabris nervis subtus sparsim pilosis, pedunculis quam petioli brevioribus. — In nemo ribus Guyanæ gallicæ, ad scopulos (Melinon, n. 55; Schomburgk, n. 357). P. guyanensis Wedd. in Ann. sc. nat. l. c. p. 221; Monogr. p. 266. (v. s.)

Nonnullæ specierum sequentium huic stirpi tanquam varietates verisimiliter etiam adscribendæ.

131. P. obtusata (Liebm. l. c. p. 300, in adnot.), dioica, caule piloso, foliis late ovatis utrinque obtusis grosse et obtuse crenatis subquinque-nerviis supra glabris subtus ad nervos pilosis, stipulis late ovatis, cymis diffuse ramosis longe pedunculatis, glomerulis discretis.  $\frac{1}{2}$  ? In ins. Monserrat (Ryan). — Wedd. Monogr. p. 261. *Urtica obtusata* hb. Vahl. — Herba caule sesquidecimtrali tetragono articulo superne piloso strigulosoque. Folia 5-6 centim. longa,  $3\frac{1}{2}$ -4 centim. lata, exsiccata utrinque cystolithis minutissimis fusiformibus obsita, supra obscure viridia subtus glaucescentia, petiolo  $2\frac{1}{2}$ -3 $\frac{1}{2}$  centim. longo pilosulo. Stipulæ obtusæ. Cymæ solitariae, pedunculis 5 centim. longis, achænio ovato marginato subtilissime granulato. (Descript. ex Liebm. l. c.)

staminodii post anthesin sæpe valde accrescentibus laciniasque perigonii demum superantibus fructu haud raro simul abortiente; achænium ovatum, sesquimillimétrale, marmoratum, quam perigonium dimidio longiore. (v. s.)

151. *P. HYALINA* (Fenzl, Denkschrift. d. Wien. kaiserl. Akad. d. Wissensch. 1. p. 4), monoica, foliis rhombeis ellipticis ovatisve acutis basi sæpius cuneatis acutisque aut obtusatis in triente inferiore limbi integris cæterum acute et plerumque grosse serratis trinerviis subtus glabris glabriusculisve supra pilis diaphanis conspersis, stipulis minimis, cymis sessilibus quam petioli brevioribus ramosissimis, floribus masc. (perigonio bipartito) et fem. intermixtis. ① In locis humidis umbrosisque Brasilæ præsertim orientalis frequens, necnon in imperio mexicano (Botteri, n. 283 et 286), in Venezuela (Fendler, n. 1826) et Peruvia (Matthews, n. 2032, Poeppig, n. 1539, etc.). — Wedd. in Ann. sc. nat. l. c. p. 227; Monogr. p. 271, t. 6, f. 8; Miq. in Mart. Fl. bras. fasc. 12. p. 200, t. 69. *Urtica arvensis* Poepp. mscr. in sched. l. c. *U. longipes* Mart. hb. *Annua*, caule herbaceo erecto pellucide succulento 1-5 decim. alto subsimpliciter pyramidato-ramoso glabro, internodiis plerisque elongatis, inferioribus mox denudatis. Folia patula, 1-5 centim. longa, 9 millim. - 3½ centim. lata, serraturis subæqualibus utrinque 5-9 raro in margine exteriori denticulo dorsali munitis, nervis basilaribus parum prominulis trientem superiorem limbi attingentibus, lamina exsiccata tenuiter membranacea cystolithis linearibus subtilibusque (non semper distinctis) utrinque obsita, petiolis in eodem jugo inæquilongis gracilibus limbo brevioribus vel sæpissime longioribus sub illius insertione pilosiusculis. Stipulæ vix conspicuæ. Cymæ in axillis omnibus nascentes, magis minusve ramosæ, ramulis ex ipsa axilla ortis spurieque verticillatis patentibus, glomerulis approximatis vel discretis; floribus masc. paucissimis cito deciduis interdumque in cymulis multis deficientibus, perigonio 2-3- sed nunquam 4-partito, alabastro vix semimillimétrali; floribus fem. subsessilibus; perigonii fructiferi segmento intermedio lateralibus multo longiore achæniique late ovati vix semimillimétralis scabriusculi fuscescentis dimidiam partem vix attingente. (v. v. et s.)
152. *P. LUNDII* (Liebmann, l. c. p. 299), monoica, foliis ovatis acutis basi late cuneatis crenatis trinerviis supra sparsim pilosis sed demum subtusque glabris, stipulis deciduis, cymis sessilibus densifloris petiolo multo brevioribus. ①? In Mexici regione calida orientali (Liebmann); in Costa-Rica (Ersted) et in Brasilia (Lund). — Herba, caule 5-10 centim. longo erecto et indiviso vel prostrato et opposite ramoso gracili tetragono pellucido glabro. Folia 1½-2 centim. longa, 11-13 millim. lata, supra læte viridia, subtus pallidiora, petiolo 10-15-millimétrali. Cymæ dense ramulosæ, floribus minutissimis, achænio oblique obovato subtilissime granulato fusco. (Descript. ex Liebmann). — A *P. hyalina* non videtur discrepare nisi stipulis deciduis, sed hæc ob exiguitatem suam a cl. auctore forsân prætermisissæ fuerunt.
153. *P. SCROBICULATA* (Liebmann, l. c. p. 300), dioica, foliis elliptico-ovatis utrinque acutis acute serratis serraturis 1-2-denticulatis supra setulosus subtus glabris scrobiculatis, cymis sessilibus petiolo brevioribus. ①? In prov. mexicana Oaxaca (Liebmann). — Wedd. Monogr. p. 272. — Herba, caule erecto 1½ decim. alto indiviso tetragono glabro. Folia subcentimétralia, 12-14 millim. lata, serraturis patulis, nervis non excurrentibus, supra fusco-viridia, petiolo 14-18 millim. longo glabro. Cymæ patulæ vel suberectæ. (Descript. ex Liebmann). — Affinis præcedenti, secundum cl. Liebmann.
154. *P. SWARTZII* (Wedd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 228; Monogr. p. 274), monoica, repens, foliis ellipticis vel elliptico-ovatis utrinque acutiusculis sæpiusve apice obtusis vix acuminatis excepto triente vel dimidio inferiore limbi crenato-serratis serratisve margine et nervis utriusque paginæ ciliatis, stipulis parvis triangularibus persistentibus, cymis parvis plerumque unisexualibus sessilibus, perigonii masc. segmentis longe mucronatis. ②? In locis saxosis nemorum Jamaicae interioris (Swartz) et Cubæ (Wright, n. 533). — Griseb. Fl. Br. W. Ind. p. 157. *P. ciliata* Bl. Mus. lugd. bat. 2. p. 46. *Urtica ciliata* Swartz, in Act. Holm. ann. 1787. p. 32; Fl. Ind. occ. p. 298; Smith, l. c. n. 37. — Herba, caule gracili 1-3-decimétrali repente diffuse ramoso, merithallis 2-3-centim. longis, ramis adscendentibus subsimplicibus glabris,

nodis hirsutis. Folia  $2\frac{1}{2}$ – $3\frac{1}{2}$  centim. longa, 10–13 millim. lata, deorsum attenuata, dentibus parvis acutiusculis, dente apicali obtuso, lamina exsiccata rigida cystolithis punctiformibus parumque conspicuis adspersa, petiolo brevi glabrato. Cymæ paucifloræ, nempe e glomerulo unico constantes, floribus masc. pedicellatis in axillis superioribus subfasciculatis, mucronibus perigonii segmenta ipsa æquantibus hirtulis, alabastro trimillimetrali. Achænium ovatum, læve, perigonio longius. (v. s. typ. in hb. Holm., Mus. brit., Soc. linn. et DC.)

155. P. CHAMÆDRYS (Wedd. in Ann. sc. nat. l. c. p. 229; Monogr. p. 253), monoica, foliis ovato-vel rhomboideo-oblongis basi cuneata integris cæterum grosse crenato-serratis (serraturis integris crenulatisve) triplinerviis ciliolatis supra glabris nervis venisque paginæ inferioris subtilissime pilosulis, stipulis brevibus persistentibus subtruncatis, cymis minimis androgynis e glomerulo unico vel duobus constantibus subsessilibus aut pedunculo petiolum superante suffultis. ♀ In ins. Portorico (Bertero). — *Urtica collina* Bertero mscr. ?; Spreng. Syst. pl. 3. p. 839? — Herba, caulibus repentibus ramosis demum adscendentibus 1–2 decim. altis apice foliosis puberulisque, internodiis inferioribus elongatis. Folia 1–2 centim. longa, 7–13 millim. lata, in apice ramorum approximata, serraturis sive crenis in utroque margine 4–5 subæqualibus, dente apicali cæteris æquimagno aut minore obtuso, nervis basilaribus ultra dimidium limbi productis, lamina exsiccata membranacea utrinque sed præsertim ad marginem paginæ superioris cystolithis fusiformibus inpersa. Stipulæ millimetrum haud metientes, abruptissime attenuatæ. Cymæ paucifloræ, in axillis superioribus solitariae, pedunculo quum exstat filiformi petiolique nonnunquam duplæ longitudinis glabro; floribus utriusque sexus subsessilibus, perigonii masc. segmentis sub apice longiuscule mucronatis. (v. s. in hb. Webb. et DC.)

156. P. SPRUCKANA, dioica, stolonifera, foliis oblongis ellipticisve acutis aut obtusis basi cuneatis obtusatisve ultra trientem inferiorem acutiuscule serratis trinerviis ciliatis supra glabris subtus petioloque villosis, stipulis late ovatis ciliatis persistentibus, cymis fem. subsessilibus e glomerulis paucis confectis petiolum non aut vix superantibus. ♀ In Peruvia orientali, prope Tarapoto (Spruce, n. 4376). — Caulis herbaceus, ascendens, in specimine unico suppetente decimetralis, crassitie pennæ corvinæ, parce villosus, ex axilla inferiore stolonem gracilem horizontalem sesquidecimetralem emittens, internodiis 3–1 centim. longis. Folia in singulis nodis conspicue inæquimagna, 4–8 centim. longa,  $2\frac{1}{2}$ – $3\frac{1}{2}$  lata, nonnulla apice rotundata, serraturis æqualibus inæqualibusve, petiolo  $1\frac{1}{2}$ – $2\frac{1}{2}$  centim. longo robusto plus minus villosio, lamina exsiccata rigida sub lente cystolithis punctiformibus subconfluentibusque utrinque creberrime obsita. Stipulæ bi-tri-millimetrales. Cymæ fem. (solum visæ) ex axillis superioribus ortæ, parce ramosæ, glomerulis minimis subdisscatis; perigonii fructiferi segmento intermedio cucullato quam achænium subrotundatum semimillimetrale circinatim scabrum paulo brevior. — Folia stolonis suppetentis distantia, rotundata, vix semicentimetrum lata. (v. s. in hb. Kew.)

157. P. ARGUTA (Wedd. in Ann. sc. nat. l. c. p. 218; Monogr. p. 275), dioica, foliis lanceolatis v. oblongo-aut ovato-lanceolatis longe acuminatis (acumine acuto) basi acutis rariusve rotundatis ultra basim limbi et ad apicem usque argute serratis tripli-v. quintupli-nerviis initio puberulis sed mox glabratiss, stipulis lineari-lanceolatis persistentibus, cymis unisexualibus parvis fere sessilibus petiolo brevioribus. ♀ In temperatis Andium caracasensium (Humb. et Bonpl.) et Æquatoris, monte Pichincha (Jameson ann. 1850. n. 63 et 838). — *Urtica arguta* Kunth in H. B. Nov. gen. et sp. 2. p. 39. Herba, caule procumbente (anne ita semper?) 3–5 decim. longo obtuse angulato diffuse ramoso, ramis gracilibus foliosis apice puberulis. Folia  $1\frac{1}{2}$ – $3\frac{1}{2}$  centim. longa, 12–25 centim. lata, in ramis sæpissime abortu alterna, a medio vel altius sensim acuminata, serraturis parvis subacuminatis, nervis basilaribus utrinque geminis a margine et costa necnon inter se fere æquidistantibus ad trientem superiorem limbi v. etiam ulterius pertingentibus, lamina exsiccata firmula cystolithis linearibus utrinque ornata, illis paginæ superioris quidem creberrimis mini-

mis lentisque sine ope haud perspicuis, in inferioris intervenio contra (non autem in nervis ipsis) majoribus rarioribus et oculo nudo facile evidentibus, petiolo 5-15 millim. longo subtilissime puberulo. Stipulae vix anaplexicaules, 2-5-millimetrales. Cymæ densae, e glomerulo unico plerumque constantes; floribus masc. breviter pedicellatis, perigonii lobis sub apice inconspicue mucronatis, alabastro obovato v. globoso sesquimillimetrali; floribus fern. subsessilibus; perigonii fructiferi segmentis subaequilongis, intermedio cucullato achænium  $\frac{1}{2}$  millim. longum, ovatum vel ellipticum laeve superante. (v. s.)

*Species imperfecte cognitæ.*

158. *P. KAKURANG* (Blume, Mus. lugd. bat. 2. p. 57), foliis lanceolatis vel rhomboideo-lanceolatis obtusiusculis basi angustatis triplinerviis inæqualiter glanduloso-serratis vel repandis membranaceis glabris. ? In Banda, ubi Kakurang dicitur. — Wedd. Monogr. p. 276. Herba erecta. Folia 4-6 centim. longa, 2-4 centim. lata. (Descript. ex Blume.)
159. *P. VENULOSA* (Blume, l. c.), foliis elliptico-oblongis lanceolatisve acuminatissimis basi angustatis triplinerviis medio ad apicem repando-glanduloso-serrulatis utrinque pube minutissima innata obsitis glabris subtus venulis crebris arcuatis reticulatis. ? In insulis Celebes. — Wedd. l. c. — Herba erecta. Folia 1-2 decim. longa, 3 $\frac{1}{2}$ -7 centim. lata, petiolo 1 $\frac{1}{2}$ -6-centimetrali. (Descript. ex Bl.)

*Index specierum et synonymorum generis Pilea.*

(N. B. Numeri numeros specierum indicant.)

- Pilea acuminata* 134, *alsinifolia* 25, *angulata* 67, *anisophylla* 36, *anomala* 85, *arguta* 157, *articulata* 11, *astrogramma* 185, *auriculata* 141;  
*betulæfolia* 137, *boehmerioides* 71, *Boliviana* 62, *bracteosa* 77;  
*callitrichioides* 3, *capitellata* 101, *carnosula* 37, *Celebica* 66, *centradenioides* 47, *cephalaantha* 21, *ceratocalyx* 117, *ceratomera* 72, *Chamædrys* 135, *Chilensis* 91, *chrysosplenoides* 127, *ciliaris* 124; *ciliaris* 28, *ciliata* 154, *citrinodora* 111, *clandestina* 38, *Commersoniana* 36, *Crugeriana* 46, *Cruzensis* 134, *Cubensis* 92, *cuneifolia* 36, *cuneiformis* 74;  
*dauciflora* 88, *decora* 94, *deltoides* 8, *densiflora* 100, *dendrophila* 45, *depressa* 148, *diffusa* 135, *diversifolia* 42, *Dombeyana* 90;  
*elegans* 94, *elegans* 91, *elliptica* 91, *elongata* 94 et 58;  
*falcata* 110, *fallax* 40, *fasciata* 119, *flexuosa* 103, *Forsythiana* 14, *Franquevilleana* 6, *fuscata* 130;  
*Gaudichaudiana* 130, *glaberrima* 12, *globosa* 4, *glomerata* 33, *gnidioides* 7, *Goglado* 12 et 57, *Goudotiana* 89, *grandifolia* 99, *grandis* 99, *Griffithii* 68, *grossereolata* 130, *Guadalupensis* 125, *Guyanensis* 130;  
*Haitiensis* 145, *heteracea* 137, *herniarifolia* 1, *herniarioides* 1, *heteronoma* 34, *Hilariana* 84, *hirsuta* 144, *hirtella* 129, *Hookeriana* 61, *Hugelii* 54, *hyalina* 161, *hygrophila* 80;  
*imparifolia* 45, *inæqualis* 126, *insignis* 56, *insolens* 32, *integrifolia* 27;  
*Jamesoniana* 89;  
*Kakurang* 158;  
*laevicaulis* 64, *lamioides* 115, *lanceolata* 13, *latifolia* 118, *laxa* 136, *leucophæa* 76, *Lindeniana* 28, *Lindensis* 28, *longipes* 133, *lucens* 65, *lucida* 35, *lucida* 36 et 92, *Lundii* 152;  
*Manniana* 31, *marginata* 108, *Maximiliani* 128, *melastomoides* 58, *Mexicana* 109, *miconiæfolia* 12, *microphylla* 1, 8 et 20, *mollis* 121, *Mongolica* 79, *montana* 130, *multiflora* 107, *muscosa* 1, *Mutisiana* 107;

*nana* 149, *nitida* 23, *nudicaulis* 15, *nudicaulis* 15, *nummularifolia* 147, *nuttans* 37;

*obtusata* 131, *oreophila* 58, *ovalis* 146, *oxyodon* 55;

*pachycarpa* 59, *Parietaria* 24, *Pavonii* 120, *pedunculata* 91, *pellucida* 58, *petata* 69, *pendula* 5, *pendula* 49, *peperomiaefolia* 1, *peplidifolia* 3, *peplodes* 10, *petiolaris* 70, *petiolaris* 67, *Pœppigiana* 98, *Portoricensis* 124, *Portula* 1, *portulacina* 1, *producta* 81, *propinqua* 26, *pteropodon* 102, *pubescens* 180, *pulegiifolia* 143, *pumila* 150, *punctata* 112, *puncticaulis* 24 *pygmaea* 10; *quadrifolia* 83;

*radicans* 114, *repanda* 53, *repens* 133, *reticulata* 106, *rhexioides* 93, *rhizobola* 43, *rhombea* 19, *rhombea* 17, *Riedlei* 65, *Rivoria* 24, *rivularis* 82, *rotundata* 139, *rubrifolia* 22, *rufa* 41, *rugosa* 145, *rupicola* 49, *rupipendia* 49.

*salicifolia* 15, *Sanctæ Crucis* 132, *sarcophylla* 9, *scripta* 57, *scrobiculata* 153, *semidentata* 96, *serpyllacea* 3, *serpyllifolia* 1, *serratifolia* 104, *serrulata* 92, *sessiliflora* 113, *sessilifolia* 52, *smilacifolia* 12, *spathulata* 95, *Spruceana* 156, *stelluligera* 17, *stipulosa* 67, *stolonifera* 29, *strigosa* 138, *subrenata* 116, *subcrenata* 65, *sublucens* 66, *submissa* 123, *subserrata* 86, *succulenta* 122, *succulenta* 1, *Swartzii* 154, *Symmeria* 78;

*tenerrima* 1, *ternifolia* 50, *tetraphylla* 83, *Thouarsiana* 51, *thymifolia* 4, *Trianaeana* 44, *trianthemoides* 2, *trichosanthes* 39, *trichotoma* 140, *trilobata* 63, *trinervia* 58, *triplinervia* 65.

*umbellata* 56, *umbrosa* 76, *umcidens* 88, *uninervis* 62, *urticaefolia* 49, *Urticella* 142;

*variegata* 48, *venulosa* 159, *verbascifolia* 73, *virgata* 16, *vulcanica* 97;

*Wallichii* 75, *Wightii* 54, *Wrightiana* 27, *Wydleri* 24;

*Xalapensis* 130.

### *Species exclusæ.*

*P. JAVANICA* Wedd. = *Pellionia Javanica*.

*P. SUBPUBERA* Miq. = *Achudemia Javanica*.

XII. *ACHUDEMA* Blume, *Mus. bot. lugd. bat.* 2. p. 57; *Wedd. Monogr.* p. 278, *Miq. Fl. Ind. bat.* 1. (2), p. 238. — *Pileæ spec. Miq. et Wedd. olim.*

Flores polygami, masculi feminei et hermaphroditi in eodem glomerulo, glomerulis laxè paniculato-cymosis, cymis in singulis axillis solitariis, pedicellis florum masc. articulatis. *Masc.* Perigonium quinquepartitum, segmentis inæqualibus in præfloratione margine subimbricatis. *Stamina* 5. Pistilli rudimentum minutissimum. *Fem.* Perigonium quinquepartitum, foliolis parum inæqualibus. *Staminum rudimenta* 6, perigonii foliolis opposita, squamiformia, inflexa. *Ovarium liberum, subobliquum.* *Ovulum basilare.* *Stigma parum excentricum, sessile, villosiusculum.* *Hermaphr.* Perigonium et pistillum ut in fl. fem. *Stamina* 5, ut in fl. masc. *Achenium lenticulari-compressum, nonnihil inflatum, perigonii foliolis persistentibus inclusum.* *Semen erectum.* *Embryo in axi albuminis parci rectum, cotyledonibus plano-convexiusculis, radícula brevi.* — *Herba javanica, facie Pileæ. Folia decussatim opposita, æquilateralis, petiolata (alternis minoribus et brevius petiolatis), dentata, trinervia.* *Stipulæ axillares.* *Cymæ longepedunculatæ parce ramosæ, glomerulis subspicatis unilateralibus, floribus fem. subsessilibus basi bracteolatis.* — *Genus a Pileæ distinctum floribus polygamis et perigonio fem. sicut masculum 5-partito nec tripartito.*

A. *JAVANICA* (Blume, l. c. t. 20), foliis ovatis acuminatis excepta basi rotundata vel acutiuscula dentato-serratis, junioribus præsertim utrinque sparsim pilo-

sis, stipulis triangulari-ovatis, pedunculis quam folia multo longioribus, perigonii segmentis sub apice hirtulo brevissime mucronatis.  $\frac{1}{2}$  In ins. Java (Blume; Zollinger, n. 870). — Wedd. Monogr. p. 278. t. 9. A; Miq. Fl. Ind. bat. 1 (2). p. 238. Pilea subpubera Miq. in Zolling. System. Verzeichn. p. 105; Fl. Ind. bat. 1. c. p. 236; Wedd. Monogr. p. 244. — Caulis repens (teste Blumeo), ramis gracilibus adscendentibus superne villosulis demumque glabris, internodiis elongatis. Folia 1½-4 centim. longa, serraturis acutis ciliatis, nervis basilaribus arcuatis dimidio limbo vix longioribus; lamina exsiccata subius pallida utrinque cystolithis linearibus minutis conspersa, iis paginæ inferioris paucioribus, et interdum subnullis, petiolo 1-2 centim. longo gracili fere glabro. Stipulæ deciduæ (sec. cl. Bl.). Cymæ fructiferae (atque a me non visæ) pedunculis 3-7 centim. longis glabris subtrifidis, ramis patentibus glomerulis subdiscretis confluentibusve in singulis ramis 2-5, pedicellis defloratis rarissimis (fl. ♂ ?) hic et illic inter perigonia fructifera sessalia perspicuis. Achænium latiuscule ovatum, vix millimetrale, verruculosum. (v. s. in hb. DC.)

**XIII. LECANTHUS** Wedd. in *Ann. sc. nat.* 4<sup>e</sup> sér. 1. p. 187; *Monogr.* p. 279; *Miq. Fl. Ind. bat.* 1 (2). p. 238. — Procridis vel Elatostematis sp. auct.

Flores monoici v. dioici, receptaculo discoideo peltatoque v. campanulato insidentes, involucri nullo, receptaculis in singulis axillis semper solitariis. *Masc.* Perigonium 4- v. 5-partitum, petaloideum, segmentis parum inæqualibus dorso gibboso-cucullatis, alabastro obconico. Stamina 4 v. 5, lobis antherarum oblongis aut subglobosis. Pistilli rudimentum conicum, glaberrimum. *Fem.* Perigonium 3-phyllum s. profunde 3-partitum, segmentis inæqualibus nunc (in floribus fertilibus) planiusculis denticulatisque, nunc (in floribus sterilibus) magis minusve cucullatis. Ovarium rectiusculum, ovatum, subcompressum, perigonio longius. Ovulum imo loculo funiculo brevi affixum. Stigma sessile, penicillatum, evanidum. Stamina rudimenta squamiformia, inflexa, in floribus sterilibus plerumque majuscula laciniasque ipsi perigonii magnitudine æmulantia. Achænium ovatum, rectum, compressiusculum, striatum, subnudum. Semen pericarpio conforme. Embryo cotyledonibus ellipticis radícula conica longioribus; albumine crassiusculo. — Herba annua, in India orientali et Abyssinia obvia, erecta. Folia opposita, in eodem jugo inæquimagna, æquilateralia, petiolata, ut plurimum serrata, trinervia, patentia, cystolithis linearibus inspersa. Stipulæ axillares. Capitula pedunculata, masculis nonnunquam minimis flores mentientibus; floribus plerumque pedicellatis ebracteatis; masculis albo-fuscis; femineis virentibus, fertilibus sterilibusque (majoribus) inter se commixtis; receptaculo margine denticulato. — Genus inter Pileam et Elatostema quodammodo intermedium, a priori imprimis floribus in receptaculo congregatis a posteriore foliis decussatim oppositis primo aspectu discrepans.

**L. PEDUNCULARIS**, monoica, foliis sæpius ovatis obtuseque serratis rarissime (in planta depauperata) subintegra tri-tripli-nerviis supra pilis raris longiusculisque conspersis subius sparsim pubescentibus, capitulis unisexualibus, masculis minimis turbinatis. ④ In locis humidis umbrosisque montium Indię orientalis temperatæ et subtropicæ (Wallich, Griffith, Wight, etc.), in ins. Java (ex Miquel), in montibus Abyssiniæ, prope Asega (W. Schimper), necnon in ins. Fernando Po altit. 2400 metr. (Mann). — L. Wightii Wedd. in *Ann. sc. nat.* 1. c., *Monogr.* p. 280. t. 9; Miquel Fl. Ind. bat. 1 (2), p. 238. Procris peduncularis Wall. Cat. n. 4634; Royle, *Illustr.* t. 83. P. obtusa Royle, 1. c. Elatostema ovatum Wight, *Icon. pl. Ind.* t. 1985. — Herba, interdum pusilla (nempe vix 2-3-centimetralis), sæpius 1-2 decim. alta, quandoque elatior, caule gracili succulento simplice aut parce ramoso erecto vel adscendente basique

aphyllo et haud raro radicante angulato-sulcato glabriusculo. Folia amplitudine quam maxime variabili, in speciminibus diminutis sæpe vix semicentimetralia, in procerrimis contra 8-10 centim. longa, 4-4½ centim. lata, apicem ver-us attenuata v. acuminata (acumine vario acutiusculo v. obtuso), basi late cuneata aut rotundata et juxta petiolum integra, cæterum grossiuscule serrata, serraturis raro in planta pusilla obsolete subobsoletisve, nervis basilariibus medium limbum superantibus, laminæ exsiccatae tenuiter membranaceæ subtus pallidæ pagina superiore cystolithis crebriusculis linearibus tenuibus pue ornata, petiolo gracili pro magnitudine foliorum 3-15 millim. longo glaberrimo v. sparsissime pubescente. Stipulæ 2-5 millim. longæ, plerumque obtusæ, scariosæ, integræ v. denticulatæ, glabræ aut puberulæ. Capitula magnitudine valde variantia, mascula plerumque minima turbinata breviterque pedunculata; feminea diametro transversali 1-12 millim. metientia v. adhuc latiora, pedunculo filiformi ut plurimum 1-3 centim. longo, receptaculo nunc glabro nunc etiam puberulo. Flores masc. pedicellati, perigonii 4-5-partiti segmentis obtusis glabris, alabastro obtuse angulato superne depresso. Flores fem. fertiles minimi, segmentis quam achænium ⅓-1 millim. longum triplo v. quadruplo brevioribus. (v. s.)

β, *Wallichii*, dioica, foliis acutiuscule serratis, capitulis masc. et fem. coniformibus longe pedunculatis. ① In India orientali. L. Wallichii Wedd. in Ann. sc. nat. l. c. (v. s.)

γ, *major*, monoica, foliis obtusiuscule serratis, floribus utriusque sexus in eodem capitulo intermixtis. ② In India orientali. L. major Wedd. l. c. Elatostema oppositifolium Dalzell, in Hook. Journ. of bot. 3, p. 179. (v. s.)

XIV. PELLIONIA *Gaudich. Bot. Voy. Uran. p. 494; Wedd. Monogr. p. 282; Benth. Fl. Hongkong. p. 330; Seem. Fl. Vit. p. 239.*

Flores dioici vel monoici, vulgo glomerati seu dense (rarius laxiuscule) cymosi, inflorescentiis sessilibus pedunculatisve in axillis singulis sæpius solitariis. *Masc.* Perigonium 4-sæpiusve 5-partitum, segmentis obtusissimis margine membranaceis summo dorso mucronatis in præfloreatione valde imbricatis, alabastro subsphærico mucronibus distinctis. Stamina 5, lobis antherarum oblongo-reniformibus. Pistilli rudimentum conicum, glaberrimum. — *Fem.* Perigonium profunde 4-sæpiusve 5-partitum, segmentis æqualibus inæqualibusve nudis aut infra apicem mucronatis. Ovarium ellipticum, parum compressum, perigonio brevius. Ovulum imo loculo affixum, erectum, funiculo brevi aut longiusculo sustentum. Stigma sessile, penicillatum. Staminum rudimenta squamiformia, tot quot perigonii segmenta iisque opposita, inflexa. Achænium magis minusve compressum aut fere cylindricum, læve et punctulato-pictum sæpiusve tuberculatum, perigonio persistente vestitum. Embryo cotyledonibus ut plurimum rotundatis radiculæ crassa conica duplo longioribus; albumine parcissimo. — Herbæ (raro suffrutescentes) asiaticæ vel oceanicæ, glabræ rariusve villosæ. Folia distiche subopposita (cujusque jugi altero minimo aut fere inconspicuo) aut (hocce omnino abortiente) alterna, inæquilateralia, margine lateris angustioris limbi sursum constanter spectante, integerrima vel serrata, penninervia vel trinervia, cystolithis fusiformibus inspersa. Flores bracteis parvis triangulari-lanceolatis stipati; glomerulis seu fasciculis involucri vero destitutis. — Habitus plerumque Elatostematis, inflorescentia potius Pileæ. Character essentialis generis a numero segmentorum perigonii feminei sumptus est.

A. — *Folia serrata.*

1. P. GRIFFITHIANA, dioica, caule puberulo, foliis abortu alternis oblique oblongis lanceolatisve longiuscule acuminatis basi semicordatis supra basim inte-

gram subæqualiter crenato-serratis breviter petiolatis subtus puberulis, stipulis persistentibus, cymis masc. longe pedunculatis laxè dichotome ramosis.  $\frac{1}{2}$  In montibus Palkaya Bengalæ orientalis (Griffith). — Caulis herbaceus, crassitie pennæ anatinæ, ut videtur erectus, angulatus, basi aphyllus, internodiis superioribus 2-4-centimetralibus. Folia in speciminibus obvisis 7-15 centim. longa, 2-4 $\frac{1}{2}$  centim. lata, acumine lanceolato 2-3-centimetrali, serraturis convexis nonnullisque apiculatis, nervis primariis utrinsecus 5-8 intra marginem arcuatim anastomosatis, jugo inferiore breviorè, lamina exsiccata rigida membranacea fuscata cystolithis in utraque pagina etiam sub lente inconspiculis. Stipulæ subulatæ, 5-8-millimetrales. Inflorescentiæ masc. (solum visæ) in axillis foliatis solitariæ, pedunculo 3-6 centim. longo ramisque patulis pubescentibus, floribus in apice ramulorum subsessilibus singulisque bractea lineari suffultis, perigonii 5-partiti segmentis 3 millim. longis sub apice rotundato mucrone submillimetrali instructis. — Affinis sequenti. (v. s. in hb. Kew.)

2. *P. HETEROLOBA* (Wedd. Monogr. p. 283. t. 5. f. 15), dioica, glabriuscula, foliis alternis (cujusque jugi nempe altero abortiente) oblique ellipticis oblongisve acuminatis basi semicordatis supra partem quartam dimidiamve inferiorem inæqualiter serratis breviter petiolatis, stipulis brevibus deciduis, cymis brevissimè vel longiuscule pedunculatis, perigonii segmentis admodum inæqualibus, aliis sub apice longe aristato-mucronatis, aliis nudis, achænio oblongo v. angustè elliptico ventricosè grossiuscule tuberculato.  $\frac{1}{2}$  In Indiæ orientalis prov. Sikkim, altit. 1520 metr. (J. D. Hooker). — Caulis herbaceus, ut plurimum simplex, ascendens, 1-3 decim. altus, crassitie pennæ corvinæ v. anatinæ, basi teretiusculus aphyllus sæpeque radicans, superne angulatus, glaber, internodiis 1-2-centimetralibus. Folia 3-8 centim. longa, 1 $\frac{1}{2}$ -centim. lata, acumine 5-10-millimetrali, serraturis margine exteriorè convexis, nervis primariis utrinsecus 5-10 ante marginem arcuatim anastomosantibus subtus prominulis, jugo inferiore sequentibus multo breviorè, lamina exsiccata membranacea pallide fusco-virente concolore cystolithis in pagina superiore paucis linearibus et in inferiore nullis, petiolo 1-3-millimetrali. Stipulæ lineares vel subulatæ, vix 2-millimetrales. Inflorescentiæ in singulis nodis solitariæ, ex axillis foliatis solum enascentes, masculæ (in specimine unico visæ et quidem incomplete evolutæ) corymbulosæ, pedunculo 1-2-centimetrali gracili puberulo, bracteis lanceolatis; floribus perigonio 5-partito, segmentis fere rotundatis et in præfloratione solito more imbricatis sub apice longiuscule mucronatis. Capitula fem. pisiformia, pedunculo brevissimo vel haud raro centimetrali tenui erectiusculoque suffulta, floribus numerosissimis confertis breviter pedicellatis; perigonii fructiferi 4-5-partiti laciniis aliis brevissimis inermibusque, aliis sub apice s. in summo dorso mucrone iis æquilongè instructis. Achæmium  $\frac{1}{2}$ - $\frac{3}{4}$  millim. longum, laciniis perigonii majoribus æquilongum sed eorum mucronibus longe superatum, magis minusve tuberculatum, tuberculis inæqualibus. Staminodia vulgo parva, sed in floribus quibusdam sterilibus laciniis minores perigonii interdum excedentia. (v. s. in hb. Kew. et Mus. par.)

3. *P. SCABRA* (Benth. l. c.), dioica, foliis alternis oblique ovatis oblongisve acuminatis basi subsemicordatis in dimidia parte superiore serratis supra magis minusve scabro-hispidis subtus pubescentibus, cymis masc. laxis pedunculatis, femineis densis sessilibusque, perigonii fructiferi segmentis admodum inæqualibus, majore (sub apice interdum mucronato) achænio ellipticum tuberculatum æquante.  $\frac{1}{2}$  In ins. Hongkong (Hance, n. 1095; Wright, n. 447, etc.) et Japonia (Oldham, n. 474). — Herba, caule gracili 2-5-decimetro prostrato simplici vel ramoso basi lignescente et aphylo, internodiis superioribus 1-2-centimetralibus brevioribusve. Folia vulgo (teste Benth.) 3 $\frac{1}{2}$ -7 $\frac{1}{2}$  centim. longa, sed interdum (præsertim ad basim ramorum) multo minora, acumine longiusculo triangulari-lanceolato, serraturis acutis aut obtusis, lamina exsiccata membranacea supra cystolithis paucissimis oblongis linearibusve notata subtus hisce destituta, petiolo circiter millimetrali. Stipulæ lineares, 1-sequi-millimetrales, persistentes. Inflorescentiæ ex axillis foliatis præsertim ortæ; masculæ pedunculis gracilibus 5-10 millim. longis suffultæ. Perigonii vulgo 4-partiti segmentis ovatis bimillimetralibus mucrone viridi



sub apice instructis. Glomeruli feminei 2-3 millim. lati, omnino sessiles sicut flores ipsi; perigonii fructiferi sæpius 4-partiti segmentis tribus subæqualibus lanceolatis acutissimis margine laxè ciliatis, quarto cæteris triente majore oblongo ut plurimum sub apice mucrone cylindrico dimidium segmenti circiter sequente munito. Achænium  $\frac{1}{2}$  millim. vix longum, compressum, inæqualiter tuberculatum, segmentum majus perigonii æquans vel paulo brevius. Staminodia parva. — Affinis præcedenti, præsertim habitu. (v. s. in hb. Mus. par.)

ß, foliis lanceolatis basi obtusis læviusculis. — Cum præcedente.

4. *P. BREVIFOLIA* (Benth. Fl. Hongkong, p. 330), dioica, foliis alternis oblique elliptico-obovatis obtusis basi valde inæquilateralibus crenatis breviter petiolatis pubescentibus, cymis fem. subsessilibus vel breviter pedunculatis, perigonii segmentis valde inæqualibus linear-lanceolatis, achæniq. tuberculato.  $\frac{1}{2}$  In ins. Hongkong (Willford). — Herba, caule prostrato repente, ramis foliiferis 15-20 centim. longis hispids. Folia admodum obliqua, 12-15 millim. longa, petiolo raro bimillimetrali. Cymæ fem. circiter 6 millim. latæ, pedunculo 2-7 millim. longo. Staminodia lanceolata medio inflexa. (Descript. ex Bentham.)
5. *P. UMBELLATA*, dioica, caule glaberrimo, foliis alternis oblique ellipticis rhombelave longe acuminatis basi latiuscule semicordatis a medio limbo ad apicem utrinque dentibus 3-4 grossis et obtusiusculis instructis subsessilibus supra sparsissime hispids subtus glabris, cymis masc. paucifloris umbelluliformibus longe pedunculatis.  $\frac{1}{2}$  ? In Japonia (Siebold). — Procris umbellata Sieb. et Zuccar. in Abhandl. der Math. phys. Klasse der bairisch-Akad. 4 (3). p. 217; Walp. Ann. 1. p. 647. *Elatostema japonicum* Wedd. Monogr. p. 315. *E. umbellatum* Bl. Mus. bot. lugd. bat. 2. t. 19. — Caulis herbaceus, gracilis, in specim. visis circiter 2-decimetralis, basi radicans, apice tantum foliiferus, internodiis brevibus. Folia 5-6 centim. longa, 2 centim. circiter lata, nervis primariis utrinque 5-6 in margine desinentibus vix prominulis jugo inferiore sequenti æquilongis; lamina exsiccata tenuiter membranacea subtus pallida cystolithis linearibus parumque conspicuis suffusa, petiolo subnullo. Stipulæ 1-2-millimetrales. Cymæ masc. parvæ, pedunculo  $1\frac{1}{2}$  centim. longo suffultæ, in singulis nodis solitariæ, pauci- et laxi-floræ, bracteis bracteolisque lanceolatis glabriusculis, perigonii 5-partiti segmentis glaberrimis mucrone subapicali brevissimo munitis. — Flores femineos non vidi, sed inflorescentia mascula absque dubio est Pellionis nec *Elatostematis*.
6. *P. RADICANS*, monoica v. dioica caule ramisque (radicantibus) subtiliter puberulis, foliis alternis oblique ovatis ellipticisve acuminatis basi semicordatis in dimidia parte superiore laxè obtuseque serratis breviter petiolatis glabriusculis vel supra sparsim pilosis, cymis masc. laxifloris longe pedunculatis, femineis densissimis sessilibusque, perigonii fructiferi segmentis admodum inæqualibus, duobus multo majoribus sub apice longe subulato-acuminatis.  $\frac{1}{2}$  In sylvis umbrosis Japoniæ (Siebold, Oldham, n. 473). — Procris radicans Sieb. et Zuccar. l. c.; Walp. l. c. p. 648. *Elatostema radicans* Wedd. Monogr. p. 332. — Caulis herbacei, 4-8 decim. longi, graciles, prostrati, ramosi, undulati, ad nodos radicanes, internodiis 2-4-centimetralibus. Folia  $1\frac{1}{2}$ -4 centim. longa, 1-2 centim. lata, acumine triangulari-ovato acutiusculo vel obtuso, serraturis utrinque 4-5, lobo basali rotundato, nervis primariis utrinque 2-5 ante marginem arcuatim anastomosantibus, lamina exsiccata submembranacea et haud raro nigricante cystolithis paucis linearibus conspersa: petiolo 1-2-millim. longo puberulo. Stipulæ subulatæ. Cymæ masc. ex axillis superioribus ramorum ortæ, folio breviores, laxiusculæ, subdichotomæ, pedunculo sequi-bi-centimetrali puberulo, bracteis numerosis conspicuis subulato-lanceolatis, floribus majusculis pedicelatis, segmentis sub apice longe aristato-mucronatis, alabastris obovatis mucronibus inclusis trimillimetralibus. Glomeruli fem. parvi (vix 3 millim. lati), floribus sessilibus, segmentis duobus majoribus perigonii 2 millim. circiter longis, segmentis minoribus ovarium vix sequantibus oblongis lanceolatisve subintegris denticulatisve. Achænium... (v. s. in hb. Mus. par., Kew. et Monac.)
7. *P. VITRENSIS* (Asa Gray, in litt. absque descript.), dioica, caule ramisque basi

radicantibus puberulis, foliis alternis oblongo-lanceolatis obtusiusculis basi obtusa vix semiauriculatis fere in toto margine laze crenato-serratis brevissime petiolatis glabris, cymis masc. longiuscule pedunculatis e glomerulo unico constantibus, femineis. ...  $\frac{1}{2}$  In ins. Viti s. Fiji (Harvey). — Caulis gracilis, 1-3 decim. longus, nisi apice prostratus denudatusque, parce ramosus. Folia  $1\frac{1}{2}$ -3 centim. longa, 5-10 millim. lata, a medio utrinque sensim attenuata, crenis s. serraturis depressis obtusis utrinque 4-6, nervis primariis in latere longiore folii 4-5 et in breviori 1-2; laminae exsiccatae rigidae pagina superiore cystolithis linearibus parvis crebriusculis inspersa, inferiore iisdem corpusculis secus nervos praesertim ornata; petiolo vix millimetrali puberulo. Stipulae ovatae, 4 millim. longae, anguste acuminatae. Cymae masc. paucae, pedunculo quam folium breviori glabro, floribus paucis, evolutis pedicello brevi strigoso suffultis, bracteis parvis parumque conspicuis, segmentis perigonii bimillimetralibus summo dorso breviter mucronatis, (v. s. in hb. Kew.)

8. *P. FILICOIDES* (Seem. Fl. vit. p. 239), erecta, foliis alternis sessilibus parvis oblongis v. lanceolatis acutis basi 1-auriculatis obtuse serratis 1-nerviis supra pubescentibus demum glabris subtus ramulisque sparse pilosis, inflorescentiis axillaribus longe pedunculatis.  $\frac{3}{4}$  In sylvis ad ripas amnis Navua, in insula Viti Levu (Seemann, n. 421) necnon in ins. Viti (Williams, Harvey). — Praecedenti nonnullis charact. sat affinis, habitus autem ut videtur valde diversus.

9. *P. STENOCARPA* (Wedd. Monogr. p. 284. t. 5. f. 13 et 15), monoica aut interdum dioica, glabra, foliis abortu alternis oblique oblongis vel elliptico-lanceolatis inaequilateralibus acuminatis basi acutis obtusatisve ultra tertiam vel dimidiam partem inferiorem grossiuscule acuteque serratis brevissime petiolatis, cymis masc. longe pedunculatis, femineis sessilibus, perigonii fructiferi segmentis parum inaequalibus infra apicem longiuscule mucronatis mucrone parce setuloso, achenio anguste lanceolato subcylindrico minute verrucoso.  $\frac{1}{2}$  In India orientali, regione tropica Khasiae, altit. 0-1200 metr. (J. D. Hooker et Thomson). — Caulis herbaceus, erectus, in speciminibus suppetentibus vix decimetrum excedens, gracilis, striatus, simplex, basi denudatus, internodiis superioribus 5-10 millim. longis. Folia 2-5 centim. longa, 8-20 millim. lata, acumine s. dente apicali 3-8-millimetrali acutiusculo, serraturis utrinque 5-8 margine exteriori rectiusculo; nervis primariis utrinsecus 5-6 in margine evanescentibus subtus vix prominentibus, jugo inferiore sequentibus breviori, costa supra papillosa; lamina exsiccata membranacea subtus (exceptis nervis fusciscentibus) albicante, intervenio cystolithis crebriusculis oblongis linearibusve reticulato, pagina superiore iisdem parcissime conspersa; petiolo circiter millimetrali. Stipulae in nodis singulis ut solet binae (quarum altera folii abortivi), lineares, sesqui-tri-millimetrales, persistentes. Inflorescentiae ex axillis foliatis tantum ortae, masculae corymbulosae, pedunculo gracili  $1\frac{1}{2}$ -4 centim. longo erecto-patente; floribus pedicellatis cito deciduis, perigonio 5-partito, segmentis sub apice longiuscule mucronatis; femineae capitatae, inflorescentiam Elatostematis referentes, sed bracteolis multo minoribus et quam pedicelli florum brevioribus: perigonii fructiferi segmentis fere aequalibus, mucronibus autem plerumque inaequalibus. Achenium  $\frac{1}{2}$  millim. circiter longum, a medio ad apicem sensim attenuatum perigonioque fere dimidia parte longius, pericarpio tenui aequaliter verruculoso. (v. s.)

10. *P. AMBIGUA* (Wedd. Monogr. p. 285), monoica vel dioica, glabriuscula, foliis suboppositis admodum disparibus, altero cuiusque jugi amplo oblique oblongo-lanceolato vel elliptico acuminato basi obtuso rotundato vel subcordato supra tertiam vel dimidiam partem inferiorem grosse serrato perrimino, altero parvo lineari-lanceolato subintegerrimo, cymis masc. longe pedunculatis, femineis sessilibus, perigonii fem. segmentis subaequalibus sub apice longiuscule mucronatis, mucrone glabro.  $\frac{1}{2}$  In regione tropica montium Khasiae Indiae orientalis (J. D. Hooker et Thomson). — Caulis herbaceus, 2-3-decimetralis, ascendens, simplex, basi radicans, crassitie pennae corinae vel paulo crassior, angulatus, merithallis superioribus centimetralibus, nodis inferioribus subaphyllis. Folia normalia 5-7, superiora ampliora 8-12 centim.

longa 2-3 centim. lata, inferiora multo minora, sæpius sensim acuminata, dente apicali subcentimetro, nervis primariis utrinsecus 5-10 valde inæqualibus, jugo inferiore brevisimo, costa supra villosula s. papillosa, lamina exsiccata subtus nisi in nervis glaucescente, cystolithis ut in *P. stenocarpa* sed minoribus, petiolo 1-2 millimetrali. Folia abortiva centimetro vel paulo minora, sesquimillimetrum lata, acuta, patentia, integerrima aut parvisime denticulata. Stipulæ lanceolatæ, parum conspicuæ. Cætera verisimiliter ut in specie supra memorata, sed inflorescentias omnino evolutas examinare mihi non licuit. (v. s. in hb. Kew. et Mus. par.)

11. *P. FLATOSTEMOIDES* (Gaudich. Bot. Voy. Bon. t. 119), dioica, glaberrima, foliis suboppositis admodum disparibus, altero cujusque jugi amplo suboblique elliptico-vel oblongo-obovato abrupte acuminato basi acuto ultra medium laxè serrato, altero contra minimo ovato integroque, inflorescentiis fem. parvis sessilibus, perigonii fructiferi segmentis subæqualibus summo dorso longe aristato-mucronatis, achænio compresso læviusculo marmorato. ♀ In ins. Moluccis (Gaudichaud), Noukahiva (Hombron), Nova Guinæa (Le Guillou) et Fidgi (Milne). — Wedd. Monogr. p. 286. t. 6. A; Miq. Fl. Ind. bat. 1 (2). p. 239. *Procris* rostrata Reinw. mscr.; Bl. Bijdr. p. 510. — Caulis 3-4-decimetralis, simplex, adscendens, succulentus, superne flexuosus, striatus, internodiis 1-2½ centim. longis, nodis infernis sæpius aphyllis. Folia majora 8-15 centim. longa, 2½-6 centim. lata, acumine subcentimetro angusto et acuto, serraturis limbi inæqualibus obtusiusculis utrinque 6-8 paucioribusve, nervis primariis utrinsecus 7-10 in margine desinentibus subtus vix prominentibus jugo inferiora sequenti subæquilongis, laminæ exsiccata tenuiter membranaceæ subtus pallidæ pagina superiore cystolithis fusiformibus minimis inæqualibusque crebriusculis inspersa, petiolo brevissimo limbo decurrente alato. Stipulæ vix conspicuæ. Fasciculi fem. in singulis nodis sæpe gemini, floribus pedicellatis 1-2-millimetralibus suffultis, perigonio fructifero cum mucronibus 4-5 millim. longo achænium late ovatum et circiter sesquimillimetrale includente. (v. s. in hb. Mus. par., Lessert, Mus. lugd. bat. etc.)
12. *P. AUSTRALIS*, dioica, foliis suboppositis admodumque disparibus, in quoque jugo altero amplo elliptico subabrupte et anguste acuminato basi obtuso vel acuto fere in toto margine inæqualiter serrato triplinervio nervis subtus hirtulis, altero abortivo minimo fere deltoideo seu acute trilobo, cymis masc. densifloris pedunculatis, pedunculo quam petiolus folii majoris longiore. ♀ In ins. Ovalaho archip. Viti s. Fidgi (Vieillard). — Caulis 3-4-decimetralis simplex? crassitie pennæ anserinæ, obtuse angulatus, inferne lignosus, pubescentihirtus aut demum glabratus, exsiccatus griseus, merithallis pler squæ bicentimetalibus. Folia majora (in nodis superioribus ampliora) 6-13 centim. longa, 3-7 centim. lata, acumine 1-2-centimetro serrulato acuto, serraturis limbi obtusiusculis in utroque margine plurimis; nervis primariis utrinsecus 6-8 in margine desinentibus, basilariibus in latere altero limbi fere usque ad basim acuminis productis; lamina exsiccata submembranacea subtus pallida sive glaucescente utrinque cystolithis linearibus inspersa; petiolo 3-8 millim. longo hirtio-pubescente. Folia abortiva (in nodis superioribus speciminis visi solum conspicua) vix 3 millim. lata, longitudine etiam minore. Stipulæ lineari-lanceolatæ, sesquicentimetales, pubescentes, cito deciduæ. Cymæ masc. (solum visæ) ex axillis superioribus ortæ, breviter ramosæ, 1½ centim. circiter latæ, pedunculo puberulo, floribus pedicellatis, perigonii 5-partiti segmentis obovatis sub apice breviter mucronatis. (v. s. in hb. Lenormand.)

B. — *Folia integerrima.*

13. *P. HEYNEANA* (Wedd. Monogr. p. 287. t. 5), monoica?, foliis suboppositis valde imparibus integris trinerviis supra glabris subtus in nervis villosis glabratissive, cujusque jugi altero amplo inæquilateraliter oblongo acuminato basi oblique cordato petiolato, altero minimo subcordato sessili, cymis utriusque sexus pedunculatis, pedunculo quam petiolus folii majoris multo longiore, perigonii fructiferi segmentis subæqualibus sub apice breviter mucronatis mucronibus pilosulis, achænio compresso tuberculato. ♀ - ♂ In montibus tropicis Zeylanicæ (Thwaites, n. 2179) et provinciarum australium Indiæ

lateralia; deorsum nonnunquam cuneata, acumine lineari 1-2-centimetro integrali; nervis primariis seu costulis longitudine et numero admodum variabilibus tenuibus, illis lateris latioris limbi 7-11 inferiore et sequente cæteris sæpe brevioribus, plerisque ante marginem arcuatim anastomosatis, illis lateris angustioris folii paucioribus pariterque inæqualibus; lamina exsiccata membranacea obscure viridi supra cystolithis minimis oblongis obsita; petiolo brevissimo vel 2-4 millim. metiente. Stipulæ 6-10 millim. longæ, obtusæ vel acutiusculæ, membranaceæ, glabræ. Flores monoici, inflorescentiis masc. et fem. in iisdem nodis, prioribus ex axillis foliatis, femineis autem ex axillis foliorum abortivorum plerumque nascentibus. Inflorescentiæ masc. juniores 10-15 millim. longæ, ficum parvulam omnino referentes, receptaculo membranaceo subcarnoso deorsum in pedunculo vulgo 3-6 centim. longum interdumque longissimum attenuato apice contra abrupte breviterque acuminato extus lævi glaberrimoque intus cavo et undique bracteis linearibus glabris floribusque breviter pedicellatis confertissimis obducto, acumine apicali circiter bimillimetro tubuloso ore subintegro; floribus perigonio 5-partito, segmentis sub apice breviter mucronatis. Capitula fem. (in specim. visis incomplete evoluta) 2 millim. vix lata, bracteolis linearibus breviter ciliatis, perigonii laciniis oblongis ovario quarta vel tertia parte brevioribus superne ciliato-denticulatis. — Foliis et habitu stirpem sequentem refert. (v. s. in hb. Kew. et Mus. par.)

§ 2. *Elatostema*. — *Inflorescentia mascula sæpius distincte involuocrata, receptaculo vulgo discoido.*

A. — *Folia abortu alterna.*

\* Folia conspicue sæpissimeque abrupte acuminata seu cuspidata (1).

2. **E. SESSILE** (Forster, l. c. p. 106), caule ramisque adpresse pubescentibus villosive interdumque glabris, foliis alternis oblique oblongis vel obovato-lanceolatis sensim aut subabrupte acuminatis basi auriculato-semicordatis fere in toto margine (incluso acumine) acute serratis triplinerviis multicostulatis nervis basilariibus sæpius dimidium limbum haud attingentibus. In Indiæ orientalis regione temperata et subtropica montium Nilghiri et Himalaya, altit. 600-2280 metr., in ins. Java (Junghuhn, Zollinger), Manilla, ins. Societatis, Taiti, Nova Hollandia, etc. — Brongn. l. c. p. 208; Guillem. in Ann. sc. nat. 2<sup>e</sup> sér. 7. p. 183; Wedd. Monogr. p. 294. t. 9. f. 9. E. serratum Forst. mscr. E. pubescens Pers. Synops. 2. p. 557. Procris sessilis Hook. et Arn. Bot. Voy. Beech. p. 70. P. australis Spreng. Syst. pl. 3. p. 846. — Caulis herbaceus vel nonnunquam basi lignescens, 2-4-decimetralis, simplex aut parce ramosus, ascendens rariusve prostratus, ad nodos inferiores sæpissime radicans, magis minusve angulatus. Folia 5-15 centim. longa, 2½-6 centim. lata, serraturis limbi interdum grossiusculis acutis rariusve obtusis æqualibus inæqualibusve, nervis primariis in utroque latere 7-10; lamina exsiccata ut plurimum membranaceæ ac sæpissime læte virentis pagina superiore cystolithis linearibus crebriusculæ obsita, iisdem in inferiore sæpius etiam perspicuis. Stipulæ fere centimetricæ, obtusæ. Flores monoici aut dioici, capitulis ad singulos nodos solitariis geminisve aut ternis. Capitula masc. depressoglobosa, sæpe centimetrum lata, pedunculo nullo aut brevi, bracteis exterioribus late ovatis plerumque acuminatis sive crasse mucronatis, bracteolis ovatis oblongis spathulatisque ciliatis; perigonii 4-partiti lobis sub apice longiuscule mucronatis, mucrone pilis rigidis instructo. Capitula fem. masculis conformia et sæpius minora, involucri pubescente, bracteolis lineari-spathulatis longeque ciliatis, perigonii laciniis modo brevissimis fere triangularibus glabrisque modo elongatis sublanceolatis et longe ciliatis. Achænium ellipticum, punctulatum, perigonium longe superans eive æquilongum.

(1) Species dignoscendæ sunt præsertim characteribus nervationis foliorum, sed propter varietates difficile est eas in agmina bene distincta contrahere.

- β, *punctatum*, capitulis fem. in singulis nodis pluribus fasciculatis pisi-formibus, cæt. ut in typo. ♀ In India orientali. — E. polycephalum Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 189. Procris punctata Hamilt. in Don Prodr. fl. Nep. p. 61. P. polycephala Wall. Cat. n. 4629, pro parte. — Capitula fem. non sunt in re vera plura quam apud typum, sed profunde lobata, lobis singulis pedunculatis inflorescentiamque distinctam mentientibus.
- γ, *cyrtandræfolium*, serraturis foliorum minoribus subacuminatis. ♀ In ins. Java (Zollinger, n. 1712 a). E. cyrtandræfolium Miq. in Zolling. Syst. Verzeichn. p. 102, non Wedd. Monogr.
- δ, *angustifolium*, foliis minoribus quam in typo lanceolatis anguste acuminatis inferne attenuatis basi que acutis vel subsemicordatis fere in toto margine (incluso acumine) argute serratis serrulatisve, stipulis subulatis. ♀ In ins. Java (Zollinger, 3760); in monte Clarence, districtu Fernando Po Africæ occid. (Mann, n. 632).
- ε, *ulmifolium*, foliis oblique obovato-ellipticis oblongisve anguste longiusculeque cuspidatis (cuspidate integro) basi semirobundatis acutatisve supra basim acute grossiusculeque serratis. ♀ In ins. Java (Junghuhn). — E. ulmifolium Miq. Pl. Jungh. 1. p. 21. — Nervi primarii inferiores in latere angustiore folii ultra dimidium limbum sæpe protracti.
- ζ, *cuspidatum*, foliis oblongo-lanceolatis cuspidato-acuminatis (cuspidate integro) basi obtusis acutatisve in dimidio vel triente apicali tantum serratis crenatisve. ♀ In India orientali (Wight, Hohenack. n. 980, etc.). — E. cuspidatum Wight, Icon. pl. Ind. t. 2091. f. 1 et t. 1983. E. affine Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 188.
- η, *tomentosum*, foliis oblongo-lanceolatis attenuato-acuminatis basi acutis obtusiusculisve supra tertiam partem inferiorem laxè serratis subtus incano-pubescentibus. ♀ In ins. Philippinis. — E. tomentosum Wedd. l. c. p. 189.
- θ, *grande* caule frutescente? foliis illis typi conformibus sed paulo majoribus, capitulis masc. longe pedunculatis. ♂? In ins. Lord Howe (Milne, Mac Gillivray). — E. grande Wedd. mscr. — Forsan ulterius specificè distinguendum.
- ι, *Brongniartianum*, pusillum, foliis multo minoribus quam in cæteris varietatibus oblique obovato-ellipticis utrinque attenuatis vel basi hinc auriculatis breviter acuminatis raro insuper cuspidatis excepta basi serratis. ♀ In Manilla (Cuming, n. 629), in India orientali necnon in Australia (F. Mueller). E. Brongniartianum Wedd. l. c. p. 190. Nervi primarii pauciores quam in antecedentibus.
3. E. MOLLE (Wedd. Monogr. p. 298), herbaceum, caule pubescente, foliis alternis elliptico-vel oblongo-lanceolatis longe acuminatis basi anguste semicordatis supra partem quartam inferiorem limbi æqualiter serratis (serraturis acutiusculis) triplinerviis multicostulatis (nervo inferiore lateris angustioris limbi ad medietatem ejusdem producto) supra glabris nervis subtus molliter et patentem pubescentibus, stipulis lineari-lanceolatis deciduis, capitulis sessilibus. ♀ In insula Pinang (Phillips). — Procris mollis Wall. Cat. n. 4633. — Caulis 2-3 decim. longus, simplex, crassitie circiter pennæ anatinæ, angulato-sulcatus, internodiis superioribus vix centimetralibus. Folia 6-10 centim. longa, 18-25 millim. lata, inæquilateralia, circiter a medio utrinque attenuata, latere inferiore folii (in planta erecta) longiore juxtaque petiolum angustato et auriculato, serraturis parvis margine exteriori convexis integris, acumine fere centimetrali ut limbus reliquus serrato, nervis primariis in latere latiore limbi 8-10 valde inæqualibus, jugo inferiore et sequente minimis, majoribus ante marginem arcuatum conjunctis; lamina exsiccata rigida læte viridi supra cystolithis linearibus creberrimis subtus autem paucis (præsertim in nervis) instructa, petiolo brevissimo 1-2 millim. vix metiente. Stipulæ sesquicentimetales, acutissimæ, adpresse pilosulæ. Flores dioici? Capitula masc. in singulis nodis solitaria, depressa, 8-15 millim. lata, vix lobata, bracteis exterioribus rotundatis obtusis parce pubentibus, bracteolis oblongis apice ciliatis, peri-

gonii 4-partiti lobis acuminato-macronatis, mucrone glabro. Capitula fem. ignota. — A præcedente paucis notis distinctum. (v. s. in herb. Soc. linn. et Kew.)

4. *E. FAGIFOLIUM* (Gaudich. Uran. p. 493), caule ramisque subtiliter pubescenti-tomentosis villosisque, foliis alternis magnis elliptico- v. obovato-lanceolatis anguste acuminatis excepta basi acuta crenato-serratis penninerviis (costulis inferioribus quam sequentes brevioribus) glaberrimis, stipulis fere linearibus prælongis diu persistentibus, capitulis sessilibus vel breviter pedunculatis. ½ In sylvis densis humidisque insularum Mascarenarum ad truncos vetustos (Commerson, etc.). — Brongn. Bot. Voy. Coq. p. 208; Wedd. Monogr. p. 299. *Procris hypocrypta* Commers. mscr. P. *fagifolia* Poir. Encycl. 4. p. 639. *Boehmeria fagifolia* Pers. Synops. 2. p. 556. Caulis herbaceus, simplex v. parce ramosus, crassus, exsiccatus teretiusculus sæpeque rugosus. Folia vulgo 10-25 centim. longa, 3-7 centim. lata, apicem versus sensim acuminata (acumine acuto magis minusve elongato atque ut ipse limbus serrato), basi interdum subæquilateralia, serraturis parvis acutis obtusiusculisve, sinubus plerumque obtusis; nervis primariis 10-15 ante marginem arcuatim reticulativæ anastomosantibus, jugis inferioribus quam sequentes brevioribus; laminæ exsiccata pagina superiore tota nervisque inferioris cystolithis fusiformibus parvis inspersis, intervenio sæpe glanduloso-punctato; petiolo plerumque centimetrali vel paulo longiore aut breviori puberulo vel glabrato. Stipulæ 1-4-centimetrales, internodio longiores, acutissimæ, sparsim pilosulæ vel glabratae. Flores dioici, capitulis in singulis nodis plerumque solitariis. Capitula masc. hemisphærica, centimetralia vel ampliora, involucri bracteis exterioribus ovatis obtusiusculis pubescenti-hispidis glabrativæ, bracteolis linearibus vel obovato-oblongis, perigonii 4-partiti lobis ovatis mucroneque subapicali elongato pilosis. Capitula fem. masculis minorâ, hemisphærica, involuero pubescente, perigonii laciniis brevissimis. — Stipulis prælongis ab affinis facillime distinguendum. (v. s.)

5. *E. MACROPHYLLUM* (Brongn. Bot. Voy. Coq. p. 207. t. 45), caute breviter et subadpresse pubescente vel glabro, foliis alternis magnis oblique oblongis vel obovato-oblongis breviter acuminatis basi cuneatis acutatisque supra basim vel partem tertiam dimidiamve inferiorem limbi serratis vel sinuato-serratis penninerviis nervis inferioribus quam sequentes brevioribus pagina superiore glabra inferiore sparsim subtiliterque pubescente vel glabrata nervis villosis pilosisve, stipulis lanceolatis majusculis, capitulis masc. breviter pedunculatis involucriatis. ½ In ins. Amboyna (Duperré, Barclay), Java (Junghuhn) et Fiji (Harvey) — Wedd. Monogr. p. 300; Miq. Fl. Ind. Bat. 1 (2), p. 241. E. spectabile Miq. Pl. Jungh. 1, 19. Cautis interdum plus quam metralis, succulentus, angulato-sulcatus, exsiccatus haud raro rugosus, internodiis superioribus bi-quadri-centimetralibus. Folia 15-30 centim. longa, 5-12 centim. lata, inæquilateralia, acumine acuto, deorsum subcuneato-attenuata, serraturis acutis obtusisve, nervis primariis utrinsecus 9-16 subtus prominulis ad marginem reticulativæ anastomosantibus, laminæ exsiccatae utraque pagina cystolithis parvis oblongis (in inferiore reticulativæ ordinatis) inspersa, petiolo centimetrali sicut costa densiuscule puberulo. Stipulæ 1½-3 centim. longæ, pubescentes vel ciliolatae. Flores dioici. Capitula masc. in singulis nodis gemina, depressa, subdiscoidea, adulta diametro sesquicentimetrali, receptaculo subdiscoideo, bracteis involucriantibus obtusissimis extus puberulis, bractæolis oblongis vel obovatis ciliatis; perigonii 4-partiti lobis ovatis glaberrimis, mucronibus subapicalibus brevibus conicis. Capitula fem. masculis sæpe majora, depressa, inter-lum sublobata, sessilia, involuero pubescente, bractæolis lineari-spathulatis longe ciliatis, perigonii laciniis brevibus triangulari-lanceolatis æchanio elliptico multo brevioribus. (v. s.)

β, *paludosum*, foliis subobovato-ellipticis basi subdimidiato-auriculatis. ½ In ins. Java (Junghuhn). — *E. paludosum* Miq. Pl. Jungh. 1. p. 19, excl. syn. Bl.; Fl. Ind. bat. l. c. (v. s.)

γ, *pedunculatum*, foliis ut in var. β, capitulis longius pedunculatis. ½ In ins. Sumatra (Junghuhn). — *E. pedunculatum* Miq. Pl. Jungh. 1. p. 21. — Folia glabriuscula, sicca nigrescentia. Pedunculi 3-6-centimetrales,

tenues. Bracteæ duæ exteriores capituli masc. abrupte apiculatæ, ciliolatæ. (v. s.)

6. *E. PLATYPHYLLUM* (Wedd. Monogr. p. 301), caule glabro rariusve patentim piloso, foliis alternis magnis oblique oblongis vel oblongo-obovatis anguste longiusculeque acuminatis (acumine dentato) basi hinc anguste auriculatis s. semi-cordatis illic acutis obtusisque supra basim vel in toto margine acutiuscule serratis triplinerviis nervis basilaribus (præsertim in latere angustiore folii) dimidium limbum superante pagina superiore glaberrima aut sparsim pilosa intervenio inferioris etiam glabro sed nervis haud raro subinconspicue pilosulis margine nonnunquam ciliolato, stipulis majusculis lanceolatis, capitulis subsessilibus vel breviter pedunculatis, involucri nullo.  $\frac{1}{2}$  In regionibus tropicis Himalayæ, a Bhotan (Griffith) ad Kumaon, necnon in montibus Khasia, altit. 900-1520 metr. (Hook. fil. et Thomson) et in provinciis Assam et Chittagong. — Habitus omnino præcedentis. Caulis crassitie variabili, angulato-sulcatus, siccus cystolithis oblongis punctiformibus creberrimisque notatus, internodiis ut plurimum 2-4-centimetalibus. Folia in specim. floriferis vulgo 10-15 centim. longa, 5-7 centim. lata, in aliis vero multo majora, nempe interdum 4-decimetralia, acumine lineari 1-5-centimetroali sæpius crebre serrulato, serraturis reliqui limbi sæpe minimis nunquam magnis, nervis primariis utrinque 7-10 frequenter valde inæqualibus dimidium limbum superante, sequentibus ante marginem arcuatis et inter se anastomosantibus, lamina exsiccata rigida utrinque cystolithis lineari-fusiformibus in inferiore pagina juxta nervulos eximie reticulatim dispositis obsita, petiolo 3-12 millim. longo valido. Stipulæ 2-5 centim. longæ, acuminatæ, binerves, glabræ. Flores dioici, capitulis in singulis nodis plerumque geminis. Capitula masc. depressa, 12-18 millim. lata, receptaculo (Dorsteniæ receptaculum æmulante) discoideo et latitudine capituli ipsius extus glaberrimo, margine sub-4-lobato. involucri ut videtur nullo, bracteolis numerosis obovatis margine villosulis; floribus pedicellatis, perigonii 4-lobati segmentis obtusissimis glabris sub apice brevissime mucronatis. Capitula fem. quam mascula multo minora (saltem in speciminibus examinatis) breviusque pedunculata, receptaculo profundius lobato margine laciniato bracteatoque, bracteolis spathulatis apice incrassatis denseque ciliatis necnon cystolithis majusculis fusiformibus obductis, perigonii segmentis linearibus raro piliferis, achænio elliptico vel ovato rubro-punctulato dimidia parte circiter brevioribus. — Nervatione foliorum a præcedente optime distinctum. (v. s. in herb. Kew. et Mus. par.)

7. *E. NIGRESCENS* (Miq. in Zolling. Syst. Verzeichn. p. 101, ex parte), caule apice puberulo foliis alternis oblique ellipticis vel oblongis anguste acuminatis basi cuneatis obtusiusculisve aut obsolete semicordatis supra limbi partem tertiam inferiorem dimidiamve acute serratis triplinerviis submulticoctulatis (nervis basilaribus dimidium limbum superantibus) utrinque sparsim hispidulis glabrativæ, stipulis majusculis subsistentibus, capitulis masc. pedunculatis, femineis sessilibus.  $\frac{1}{2}$  In paludibus montium occid. Javæ, altit. 1000-1800 metr. (Zollinger, n. 1712; Junghuhn). *E. cyrtandræfolium* Miq. Pl. Jungh. p. 21; Wedd. Monogr. p. 297. *Procris cyrtandræfolia* Zolling. et Moritz? Syst. Verzeichn. p. 74. — Caulis elongatus, gracilis, simplex, basi longe radicans, exsiccatus striatus, internodiis sesqui-bi-centimetalibus. Folia 6-14 centim. longæ, 3-6 centim. lata, acumine fere centimetroali, dentibus triangularibus, æqualibus nervis primariis utrinsecus 6-8 ante marginem arcuatum anastomosantibus subtus parum prominentibus, lamina exsiccata tenuiter membranacea supra cystolithis fusiformibus minimis dispersa, petiolo gracili 2-3 millim. longo. Stipulæ 1-2 centim. longæ, scariosæ, obtusæ v. acutiusculæ. Flores monoici, capitulis in singulis nodis solitariis. Capitula masc. pedunculo filiformi 3-4 centim. longo glabroque suffulta, 1 centim. circiter lata, bracteis exterioribus involucri ovatis acuminatis villosis, perigonii 4-partiti segmentis glaberrimis, mucronibus subapicalibus 1 millim. longis obtusiusculis. Capitula fem. pial circiter magnitudine, involucri villosulo, bracteolis oblongis linearibusve orasse ciliatis, majoribus in dimidia parte superiore cylindricis glabrativæ, floribus brevissime pedicellatis, perigonii segmentis quam stamina abortiva brevioribus, achænio oblongo. — Sub numero citato collectionis Zollingerianæ duæ exstant species, scilicet hæc (cujus vidi specimen

typ. Miquelianum in herb. Zolling.) et *E. cyrtandraefolium* Miq. in Zolling. Syst. Verzeichn. p. 102 (non Pl. Jungh.) quod ad *E. sessile* referendum est. (v. s.)

8. *E. DECEPIENS*, caule glabro vel superne puberulo, foliis alternis oblique oblongis vel oblongo-ellipticis anguste acuminatis basi subsemicordatis supra quartam partem inferiorem acute serratis triplinerviis vix multicostulatis nervis basilaribus dimidium limbum superantibus supra sparsim hispidis demumque glabris nervis subtus puberulis, stipulis oblongis deciduis, capitulis masc. sphaericis breviter pedunculatis bracteis binis exterioribus concavis abrupte acuminatis conniventibusque involutis. 2 In Indiæ orientalis prov. Mishmi (Griffith). — Caulis modice crassus, angulato-sulcatus, simplex aut parce ramosus, internodiis varie elongatis. Folia 10-15 centim. longa, 3½-6 centim. lata, acumine lineari sesquicentimetrali integro, dentibus limbi sæpius inæqualibus nonnullisque subacuminatis, nervis basilaribus vulgo supra petioli insertionem ortis, illo lateris angustioris limbi usque ad basin acuminis protracto, costulis lateris latioris inæqualibus et parum prominentibus; laminæ exsiccatae membranaceæ, pagina superiore cystolithis linearibus tenuissimis ornata, inferiore his corpusculis ut videtur fere destituta; petiolo brevissimo puberulo. Stipulæ sesquicentimetrales, acutæ. Capitula masc. (solum visa) in nodis singulis plerumque bina, quæ adsunt (verisimiliter non plene evoluta) pisiformia apiceque acuminibus bractearum involucrentium instructa; floribus sessilibus pedicellatisque, segmentis perigonii sub apice breviter mucronatis. — Bracteæ exteriores concavæ capituli junioris flores amplectentes receptaculum ficoideum globosum imitantur. (v. s. in hb. Kew.)
9. *E. RETICULATUM* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 188; Monogr. p. 302), glaberrimum, foliis alternis lanceolatis vel oblongo-lanceolatis acuminatis basi anguste semicordatis in toto margine crebriusculè dentato-serratis multipenni-nerviis, costulis inferioribus brevissimis, stipulis lanceolatis deciduis, capitulis masc. longe pedunculatis, femineis sessilibus. 2 In Nova Hollandia, juxta flumen Hunter's river, inter Filices arboreas (Leichard). — Caulis simplex crassiusculus, vix sinuatus, siccus striatus. Folia 10-15 centim. longa, 2½-3½ centim. lata, circiter a medio utrinque attenuata apiceque sensim acuminata, acumine acuto, serraturis parvis acutis, nervis primariis utrinsecus 10-12 ante marginem inter se et cum nervis secundariis eleganter reticulatim connexis, jugo inferiore brevissimo, lamina exsiccata membranacea subtus pallida supra (in folio adulto) cystolithis parvis fusiformibus creberrimis inspersa, petiolo brevissimo. Stipulæ subcentimetrales, acutæ, glabræ. Flores monoici, capitulis in singulis nodis solitariis. Capitula masc. 1 centim. circiter lata, pedunculo glaberrimo 3-5-centimetrali suffulta, bracteis exterioribus ovatis acutiusculis ciliolatis, perigonii 5-partiti segmentis glabris, mucronibus cylindricis obtusiusculis apicem segmentorum paulo superantibus. Capitula fem. in specimine unico viso minima, ex axillis inferioribus orta, involucre hispidio, bracteoilis spatulatis ciliatis, laciniis perigonii brevissimis, acheniis acutis. (v. s.)
10. *E. MACULATUM* (Gaudich. Uran. 493), caule breviter pubescente, foliis alternis amplis oblique elliptico-obovatis acuminatis basi obtusis in dimidio superiore grossiusculè et inæqualiter serratis triplinerviis utrinque sparsim hispidulis nervis subtus subtiliter pubescentibus, stipulis ovatis brevibus mox deciduis, capitulis masc. sessilibus. 2 In ins. Java (Commerson). — Brongn. Bot. Voy. Coq. p. 208; Wedd. Monogr. p. 303; Miq. Fl. Ind. Bat. 1 (2) p. 242. Procris maculata Poir. Encycl. 5. p. 628. Boehmeria maculata Pers. Synops. 2. p. 556. Caulis simplex? crassiusculus, teres. Folia 15 centim. longa, 5½ centim. lata, inæquilateralia, in triente superiore sensim attenuata breviterque acuminata, deorsum attenuato-cuneata, ima basi obtusata paululumve emarginata, serraturis acutiusculis hinc s. in latere ampliore folii 8-10 illinc 5-6, nervis primariis utrinsecus 7-8 intra marginem arcuatim anastomosantibus, inferioribus quam sequentes longioribus, laminæ exsiccatae rigida utraque pagina et præsertim superiore cystolithis fusiformibus minimis inspersa, petiolo brevissimo pubescente. Stipulæ 3-4 millim. vix longæ, puberulæ. Flores dioici? Capitula masc. (solum visa) in singulis nodis ge-



mina, depressa, diametro subcentimetroli, bracteis exterioribus ovatis obtusis pubescentibus, bracteolis oblongo-spathulatis ciliatis, perigonii 5-partiti segmentis ovatis dorso et apice pilosis, mucrone subapicali in segmentis aliis longiusculo pilosoque in aliis subnullo. (v. s. in hb. Juss.)

11. *E. RUGOSUM* (A. Cunningham. in Ann. nat. hist. 1. p. 215. t. 9. f. 5-8), caule glabrato, foliis alternis oblongo-vel obovato-lanceolatis sæpius abrupte acuminatis auriculato-semicordatis ab ima basi usque ad acumen argute serratis (sinubus inter serraturas obtusis) penninerviis utrinque hispidulis supraque asperatis subtus interdum pubescentibus, stipulis lanceolatis mox deciduis, capitulis brevissime pedunculatis.  $\frac{1}{2}$  In sylvis densis Novæ Zelandiæ ins. septentr. (Cunningham, J. D. Hooker, etc.). — Hook. fil. Fl. Nov. Zel. p. 227; Wedd. Monogr. p. 304. — Caulis elongatus, succulentus, vix flexuosus, crassitie pennæ anserinæ v. gracilior; exsiccatus obsolete angulatus, internodiis sesquicentimetralibus. Folia 8-25 centim. longa, 2-6 $\frac{1}{2}$  centim. lata, auricula baseos ut reliquis limbus serrata, acumine lanceolato integro, serraturis magnis minusve acuminatis integris aut nonnunquam in margine exteriori denticulo munitis, nervis primariis utrinsecus 9-12, ante marginem arcuatim connexis subtus vix prominentibus, jugo inferiore quam sequentes breviores, laminæ exsiccatae membranaceæ sæpius læte viridis utraque pagina cystolithis linearibus brevibusque inspersa, petiolo brevissimo. Stipulæ 1-2 centim. longæ, acuminatæ, puberulæ. Capitula in singulis nodis solitaria, depresso hemisphærica, pedunculis 2-3-millimetralibus: mascula majuscula, bracteis exterioribus late ovatis vel rotundatis puberulis glabrativæ, duabus extimis longe et abrupte acuminatis: floribus breviter pedicellatis, perigonio 4-partito glabro, mucronibus subapicalibus lobis ipsis æquilongis. Capitula fem. masculis subæquimagna, involucrio subtiliter pubescente, bracteolis spathulatis breviterque ciliatis, perigonii segmentis brevissimis, achænio obtusiuscule acuminato. (v. s.)
12. *E. NEMOROSUM* (Seemann, Acc. gov. miss. Fij. Isl. app. p. 441; Fl. Vit. p. 240. t. 61), caule pilosulo, foliis alternis interdum maximis obovato-oblongis abrupte acuminatis basi in petiolum decurrentibus et juxta insertionem auriculatis toto margine argute serratis (sinubus inter serraturas acutissimis) penninerviis supra pilosis levibusque subtus molliter pubescentibus, stipulis magnis oblongo-lanceolatis, capitulis subsessilibus.  $\frac{1}{2}$  In ins. Fiji s. Viti (Seemann, n. 422). — *E. umbrosum* A. Gray? in litt. — Folia 2-3 decim. longa, 7-12 centim. lata, acumine lanceolato 1-2-centimetroli serrato, serraturis limbi triangularibus versus apicem folii sensim majoribus et haud raro denticulo dorsali instructis, nervis primariis s. costulis in utroque latere limbi circiter 12, jugo inferiore minore; lamina exsiccata membranacea supra obscure viridi subtus pallida, cystolithis in nervis solum perspicuis minimis oblongis punctiformibusque; petiolo brevissimo. Stipulæ nonnunquam 4-5 centim. longæ, acutæ, medio dorso longitudinaliter pilosæ. Capitula fem. (solum visa) bracteis exterioribus ovatis acutis pubescenti-pilosis, bracteolis densè hirsuto-ciliatis; achænio ovato apiculato, perigonii segmentis subulatis inæqualibusque longiore. — *E. rugosum* habitu refert. (v. s.)
13. *E. RUPESTRÆ* (Wedd. Monogr. p. 304), caule præsertim apice puberulo, foliis alternis elliptico-vel obovato-lanceolatis longe et anguste acuminatis basi acutis supra tertiam partem inferiorem limbi acumineque serratis (serraturis interdum mucronulatis) tripli-vel subpenni-nerviis utrinque glabris nervis subtus pubescentibus, stipulis lanceolatis deciduis persistentibusve, capitulis sessilibus vel breviter pedunculatis.  $\frac{1}{2}$  In Indiæ orientalis prov. Nepalia (Wallich) et in regione tropica montium Khasiæ, altit. 600-900 metr. (Hooker et Thomson). — *Procris rupestris* Don, Fl. Nepal. p. 60. P. punctata Wall. Cat. n. 4627, pro parte, non Hamilt. Caulis herbaceus, e basi radicante demumque lignea ascendens, 3-4 decim. altus, simplex, obtuse angulatus, crassitie circiter pennæ anserinæ, vix flexuosus, teretiusculus, internodiis superioribus brevibus. Folia 10-15 centim. longa, 1-4 $\frac{1}{2}$  centim. lata, inæquilateralia, deorsum cuneata vel longiuscule attenuata, serraturis parvis integris rariusve denticulo in margine exteriori instructis, acumine sesquicentimetroli etiam serrato rariusve integro, nervis primariis utrinsecus 8-10 subtus parum pro-

minentibus ante marginem arcuatim connexis, jugo inferiore illinc s. in latere angustiore folii medium limbum sæpius superante; laminæ exsiccatae rigida pagina superiore cystolithis lineari-oblongis creberrime obsita, inferiore iisdem nisi in nervis prorsus destituta; petiolo 2-3 millim. longo breviter pubescente. Stipulae uni-sesqui-centimetales, siccitate scariosae, acutae, glabrae. Capitula dioica in singulis nodis ut plurimum solitaria, depressa globosa, 10-15 millim. lata, receptaculo haud multum dilatato, bracteis exterioribus rotundatis obtusissimis sub apice brevissime mucronatis puberulis vel pubescenti-tomentosis; mascula bracteis binis involucrentibus majusculis munita, bracteolis oblongis dorso et apice lanato-pilosis, floribus numerosis, perigonii 4-partiti segmentis mucrone subapicali brevissimo longiusculeque piloso instructis; feminea bracteolis elongatis linearibus longe densequè ciliatis, floribus breviter pedicellatis, perigonii fructiferi segmentis lineari-triangularibus apice piliferis achenii ovati dimidia fere parte brevioribus. (v. s.)

β, *salicifolium*, foliis anguste lanceolatis fere æquilateralibus longe attenuato-acuminatis supra basim acutam serrulatis. ½ In ins. Java juxta rivulum Tjasmarea ad Prabakti, altit. 700 metr. (Zollinger, n. 3531). — E. Zollingerianum Miq. in Zolling. Syst. Verzeichn. p. 105. (v. s.)

14. E. *PAIVÆANUM*, caule glabro aut vix puberulo, foliis alternis oblique ellipticis longiuscule acuminatis basi angustatis et subsemiauriculatis ima basi excepta in toto margine (incluso acumine) dentato-serratis serratisve triplicivel subperenni-nerviis, limbo adulto supra omnino glabro nervis subtus villosis, stipulis lineari-lanceolatis mox deciduis, capitulis masc. et fem. subsessilibus. ¾ In monte Clarence Peak dicto, ins. Fernando-Po Africæ occidentalis, altit. 1200 metr. (Mann). — Caulis herbaceus, simplex, gracilis, nonnihil flexuosus, 2-4 decim. altus, basi radicans, in dimidia parte superiore foliatus, internodiis 2-4-centimetalibus. Folia 6-12 centim. longa, 3-5 centim. lata, inæquilateralia, acumine sesqui-bi-centimetro utrinque 1-2-dentato, serraturis limbi extus sæpius convexis et haud raro subacuminatis, costulis lateris lateris limbi 6-8 (infra quam sequens brevior), nervo basilari lateris angustioris medietatem limbi superante; lamina exsiccata membranacea supraque cystolithis linearibus tenuibus inspersa, infra vero hisce destituta; petiolo 2-3-millimetrali villosis. Stipulae centimetrum circiter longæ, apice subulatæ. Flores monoici aut dioici, capitulis in singulis nodis solitariis. Capitula masc. (in planta monoica superiora) 1-1½ centim. lata, depressa, nonnunquam pedunculo brevissimo villosa suffulta, bracteis involucrentibus late ovatis vix acuminatis basi pilosulis margineque ciliatis, floribus pedicellatis, perigonii segmentis 2 millim. longis glabris mucrone lineari 1 millim. longo piloso sub apice instructis. Capitula fem. masculis minora, bracteis bracteolisque lanceolatis aut linearibus ciliatis, acheniis breviter vel longiuscule pedicellatis, segmentis perigonii fere inconspicuis. (v. s. in hb. Kew.)

15. E. *MANNII*, caule apice strigosulo, foliis alternis vix oblique obovato-lanceolatis subabrupte acuminatis basi cuneatis integrisque cæterum grossiuscule mucronato-serratis (acumine serrato) triplinerviis, nervis subtus sparsim hispidis, limbo cæterum glaberrimo, stipulis lanceolatis persistentibus, capitulis masculis pedunculo gracili suffultis, femineis in eodem caule sessilibus. ¾ In districtu Fernando-Po Africæ occidentalis (Mann, n. 1431). — Caulis herbaceus, gracilis, erectus, 2-4 decim. altus, simplex, in dimidia parte inferiore omnino nudus, internodiis superioribus haud centimetalibus. Folia alterna 4-7 centim. longa, 1½-2 centim. lata, pleræque vix inæquilateralia, acumine lanceolato circiter centimetro utrinque sub-3-dentato, serraturis limbi plerumque obtusis, costulis utrinsecus 6-7 vix prominulis arcuatim connexis, nervo basilari lateris superioris (sive interioris) limbi medietatem ejusdem paulo superante, lamina exsiccata submembranacea supra cystolithis linearibus crebre inspersa; pagina inferiore iisdem corpusculis paucioribus brevioribusque ornata, petiolo vix ullo aut 1-3-millimetrali. Stipulae semicentimetrum longæ, apice subulatæ, in internodiis superioribus fere imbricatæ. Flores monoici, inflorescentiis masculis femineisque sæpe in eodem nodo oppositis. Capitula mascula 3-5 millim. lata, pedunculo glabriusculo internodium superante, bracteis involucrentibus ovatis breviter acuminatis puberulis, floribus

pedicellatis, perigonii segmentis (incluso mucrone lineari longiusculo) bimillimetralibus. Capitula feminea sessilia masculis minora, bracteis bracteolisque linearibus ciliatis flores breviter pedicellatos superantibus, perigonii laciniis lanceolatis fimbriato-ciliatis achenium fere æquantibus. (v. s. in hb. Kew.)

16. *E. rostratum* (Miq. in Zolling. Syst. Verzeichn. p. 102, non Hassk.), herbaceum vel suffrutescens, caule glabrato, foliis alternis amplis oblique elliptico-vel oblongo-ovatis longe et anguste cuspidato-acuminatis obtusatis supra trientem inferiorem inæqualiter crenato-serratis tri-nerviis subpaucicostulatis nervo basilari lateris angustioris limbi costam fere æquante subtusque (sicutque nervi cæteri) pilosulo, stipulis ovato-lanceolatis deciduis, capitulis fem. sessilibus.  $\frac{2}{3}$  In ins. Java (Zollinger, n. 1530). — *E. thibaudæfolium* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 188; Miq. Fl. Ind. bat. 1 (2), p. 242. — Caulis elongatus, basi pennæ anserinæ circiter crassitie, flexuosus, exsiccatu angulatus, internodiis inferioribus 4-5 centim. longis, superioribus centimetralibus longioribusve. Folia 10-18 centim. longa, 4-7 centim. lata, acumine bi-tri-centimetrali lineari acutissimo, dentibus obtusis mucronulatisve nonnunquam etiam fere obsolete, nervis primariis hinc s. in latere latiore limbi 5-7 ante marginem arcuatim anastomosantibus inferiore medium limbum vix attingente, illic in unicum basilem limbo æquilongum rectiusculumque reductis, subtus parum prominulis, lamina exsiccata subcoriacea supra cystolithis crebris minimis fusiformibus inspersa, petiolo 3-5 millim. longo valido. Stipulæ 5 millim. circiter longæ, acutissimæ, pubescentes. Capitula dioica (feminea solum visa) multiflora, pisi circiter magnitudine, supra glabriuscula, bracteolis lineari-lanceolatis sparsim ciliatis, perigonii subquinquepartiti segmentis ovatis parce ciliatis achenii elliptici maculati dimidiam partem superantibus, staminodiis post anthesim valde auctis. (v. s.)

$\beta$ , *Manihense*, caule apice pubescente vel glabrato, foliis amplis elliptico-vel oblongo-lanceolatis lanceolatisve longe acuminatis tri-triplinerviis utrinque glabris, stipulis lanceolatis.  $\frac{3}{4}$  In ins. Manilla (Cumming, n. 786; Barthe, etc.). — *E. manillense* Wedd. in Ann. sc. nat. 1. c. p. 189; Monogr. p. 310. — Folia 2-2  $\frac{1}{2}$  decim. longa, 6-7 centim. lata, sensim acuminata, acumine obsolete denticulato sive subserrato, nervis primariis subtus prominulis, jugo inferiore plerumque brevissimo proximo contra longissimo, lamina exsiccata coriacea, petiolo 4-6 millim. longo adpresse pubescente. Stipulæ 15 millim. longæ, acutissimæ, subtiliter pubescentes. Inflorescentiæ masc. in nodis singulis geminæ, 6-8 millim. latæ, bracteis parvis ovatis crassis pubescentibusque stipatæ, bracteolis lanceolatis, perigonii 5-partiti segmentis pubescentibus, mucrone subapicali brevissimo. (v. s.)

17. *E. integrifolium*, fruticosum, caule ramisque glaberrimis aut (junioribus præsertim) pubescentibus, foliis alternis oblique oblongo-vel elliptico-lanceolatis lanceolatisve longe acuminatis basi acutis integerrimis aut dente uno vel duobus infra acumen instructis triplinerviis subpaucicostulatis nervo basilari lateris angustioris limbi sæpe fere usque ad apicem hujusce producto, stipulis lanceolatis mox deciduis, capitulis plerumque sessilibus, masculis glomeruliformibus.  $\frac{3}{4}$  In India orientalis prov. Assam, Khasia, Chittagong, etc. altit. 300-1520 metr. (J. D. Hooker, Thomson, Griffith, etc.), necnon in montibus Java (Junghuhn, Zollinger, n. 426). — *E. sesquifolium* Hassk. Hort. bogor. p. 79; Wedd. Monogr. p. 308; Miq. Fl. Ind. bat. (2). p. 243. *E. cuspidiferum* Miq. Pl. Jungh. 1. p. 22. *E. Miquelianum* et *E. Zollingerianum* Wedd. in Ann. sc. nat. 1. c. p. 188. *Procris integrifolia* Don Prodr. Fl. nepal. p. 61. *P. piperifolia* Wall. Cat. n. 4628. *P. sesquifolia* Reinw. mscr.; Blume Bijdr. p. 54. — Caulis metralis etiamque elatior, suberectus, ramosus, ramis obtuse angulato-sulcatis haud raro flexuosis demum glabris, exsiccatu fuscis cystolithisque minutissimis punctulatis. Folia 5-15 centim. longa, 1  $\frac{1}{2}$ -4 centim. lata, sæpe vix inæqualateralia, a medio ut plurimum utrinque attenuata, acumine lineari 1  $\frac{1}{2}$ -3-centimetrali acutissimoque, nervis primariis utrinusque 4-7 rarius numerosioribus, nervo basilari in latere angustiore folii cæteris multo longiore margini parallelo supraque medium limbum cum sequentibus arcuatim anastomosante vel interdum usque ad acumen ipsum producto, lamina exsiccata rigida olivacea glaucescente aut late viridi utraque pagina

cystolithis linearibus in inferiore autem multo minoribus et plerisque reticulatim ordinatis obsita, petiolo brevi. Stipulæ 6-10 millim. longæ, acuminate, siccitate ferrugineæ, glabræ aut pilosulæ. Capitula monoica vel dioica in singulis nodis vulgo gemina, parva (nempe vix semicentimetralia), bracteis exterioribus rotundatis glabriusculis; mascula bracteolis ovatis oblongisve dorso pilosis, floribus (nonnunquam numerosissimis) longiuscule pedicellatis, perigonii 4-partiti segmentis glabris vel pubescentibus sub apice mucrone conico instructis. Capitula fem. ex axillis superioribus ramorum orta, vulgo masculis conformia, bracteolis linearibus longe ciliatis, perigonii fructiferi segmentis ovatis denticulatis apiceque piliferis achænio ovato fere dimidio brevioribus. (v. s.)

β, *semiserratum*, foliis oblongo-vel elliptico-lanceolatis, in triente vel dimidio superiore grosse crenato-serratis glabris aut cum ramis subtus pubescentibus. γ In peninsulæ Indiæ orientalis prov. Conca (herb. Stock.) et in regione tropica montium Kumaon, altit. 620-1250 metr. (Thomson). (v. s. in hb. Kew. et Mus. par.)

18. *E. HIRSUTUM* (Miq. Fl. Ind. bat. 1 p. 2. p. 243), suffrutescens, foliis alternis? suboblique lanceolatis longe attenuato-acuminatis basi acutis integerrimis vel infra acumen repandulis fugaciter ciliolatis cæterum glabris triplinerviis, nervis lateralibus paucis nervo basilari lateris angustioris limbi fere usque ad acumen producto, capitulis (teste Blumio) dimidiato-verticillatis hirsutissimis. γ In ins. Java. Procris hirsuta Blume Bijdr. p. 509. Folia 15 centim. longa, exsiccata chartacea cystolithis utrinque lineolata. (Descript. ex Miq. l. c.) — A præcedente videtur discrepare præsertim hirsutie capitulorum.
19. *E. PROCRIDIOIDES*, herbaceum, glabrum, foliis alternis oblique vel suboblique elliptico-vel obovato-oblongis oblongisve subabrupte acuminatis basi acutis integris triplinerviis subpaucicostulatis, nervo basilari lateris angustioris limbi medietatem ejusdem superante, stipulis lanceolatis deciduis, capitulis masc. longe pedunculatis. γ In Indiæ prov. Mishmi et Assam et mont. Khasia Bengalæ orientalis necnon in imperio birmanico (Griffith). — Caulis a basi radicante erectus, 3-5 decim. altus, crassitudine calami scriptorii vel gracilior, siccitate obscure fuscus, internodiis superioribus magis minusve elongatis. Folia in parte superiore caulis solum persistentia, erectiuscula, 6-15 centim. longa, 2-6 centim. lata, nonnunquam fere æquilateralia, acumine triangulari-lineari 1-sesquicentimetrali acuto, nervis primariis in utroque latere limbi 5-7, nervo basilari angustioris supra medium cum cæteris arcuatim anastomosato; laminæ exsiccatae membranacæ subtus pallidioris utraque pagina cystolithis linearibus inspersa, petiolo 3-10-millimetrali. Stipulæ 5-10 millim. longæ, apice subulatæ. Capitula dioica; mascula (solum visa) in nodis singulis plerumque solitaria, omnino evoluta sæpe 2-3 centim. lata, depressa, receptaculo peltato bracteis ovatis acuminatis margine instructo, duobus horumce oppositis majoribus longiusque acuminatis, floribus numerosis plerisque pedicellatis bracteolis elongatis cuneiformibus apiceque crasse cucullatis glabris intermixtis, perigonii segmentis sub apice mucrone conico munitis. — Species habitu distinctissima. (v. s. in hb. Kew.)
20. *E. HOOKERIANUM* (Wedd. Monogr. p. 309), herbaceum, glaberrimum, caule subsimplice, foliis alternis obliquissime oblongo-lanceolatis longiuscule acuminatis basi late semiauriculatis in dimidio vel triente superiore (acuminibus) grossiuscule acutæque serratis triplinerviis paucicostulatis nervis basilari-bus medium limbum longe superantibus, stipulis linearibus subpersistentibus, capitulis sessilibus, masculis laxifloris. γ In regione subtropica Indiæ orientalis prov. Sikkim, altit. 1200-2480 metr. (J. D. Hooker). — Caulis ascendens, bi-quadri-decimetralis, basi longe radicans denudatusque, angulatus, crassitie variabili, siccitate fusco-virens cystolithisque ut videtur destitutus, internodiis superioribus subcentimetralibus. Folia, 2-10 centim. longa,  $\frac{1}{2}$ -2  $\frac{1}{2}$  centim. lata, valde inæquilateralia, sensim acuminata, serraturis ipsis interdum subacuminatis superioribusque multo minoribus, nervis primariis (quorum jugum inferius solum bene est conspicuum) paucis, secundariis duobus basilari-bus in latere ampliore folii quam primarii superioris distinctioribus; laminæ exsiccatae membranacæ viridis subtus pallidioris pagina superiore

cystolithis linearibus sæpissime albo-marginata cæterum illis vix punctulata, pagina inferiore lisdem ut videtur omnino destituta; petiolo subnullo. Stipulæ 3-8-millimetrales, acutæ, siccitate subscariosæ. Capitula dioica in singulis nodis solitaria; mascula 5-8 millim. lata, receptaculo angusto, bracteïs involucri ovatis vix amplis subacuminatis, bracteolïs oblongis obovatisve, margine ciliato-laceris dorso carinatis glabris, perigonii 4-partiti segmentis sub apice breviter mucronatis glabris, alabastro fere bi-millimetrali. Capitula fem. in specimine unico visa masculis multo minora, bracteïs involucriantibus bracteolisque exterioribus oblongis longe ciliatis sub apice crasse mucronatis, bracteolïs internis linearibus perigonio brevi. (v. s. in hb. Kew. et Mus. par.)

21. *E. STRIGOSUM* (Hassk. Cat. h. bog. p. 79), herbaceum, caule ramisque breviter griseo-villosis, foliis alternis oblique obovato-ellipticis abrupte acuminatis basi acutiusculis supra medium grosse et inæqualiter crenato-serratis triplinerviis, nervis basilariibus vix ad medium limbum perductis lateralibusque paucis subtus villosis, stipulis parvis lineari-lanceolatis deciduis, capitulis sessilibus dimidiato-verticillatis villosulis. 2/ In ins. Java. — Miq. Fl. Ind. bat. 1 (3) p. 246. *Procris strigosa* Blume, Bijdr. p. 512. — Folia 2½-12 centim. longa, supra glabra, exsiccata cystolithis dense striolata, petiolo villosa. — Ab *E. thibaudiæ* folio cui (teste cl. Miquelio) haud absimile est, differt nervis basilariibus non adeo alte perductis et indumento.
22. *E. ACUMINATUM* (Brongn. Bot. Voy. Coq. p. 211), herbaceum, caule ramisque glaberrimis, foliis alternis elliptico-lanceolatis longe acuminatis basi acutis circiter a medio limbo vel ultra trientem inferiorem laxè obtusiusculeque serratis triplinerviis paucicostulatis jugo inferiore medietatem limbi vix superante, stipulis parvis lanceolatis, capitulis masc. (glomeruliformibus) femineisque sessilibus. 2/ In ins. Java (Commerson). — Wedd. Monogr. p. 311. t. 9. D. f. 1-4; Miq. l. c. p. 244. *Procris acuminata* Poir. Encycl. 4. p. 629. *P. membranacea* Reinw. mscr.; Bl. Bijdr. p. 512. *Boehmeria acuminata* Pers. Synops. 2. p. 556. *Langevelia acuminata* Gaudich. Uran. p. 494. — Caulis gracilis, ramosus, 2-4-decimetralis, exsiccatus striato-sulcatus, ramis parum flexuosis, internodiis longitudine variabili. Folia 3½-6 centim. longa, 1-2 centim. lata, nervis primariis utrinsecus 4-5 ante marginem arcuatim anastomosantibus subtus vix prominentibus, laminæ exsiccatae membranacæ pallide virentis pagina superiore cystolithis minimis punctiformibus inspersa, petiolo 1-2 millim. longo. Stipulæ 1-3 millim. longæ, illis foliorum deficientium quam cæteræ sæpius longioribus et plerumque subulatis. Flores monoici v. dioici, capitulis in singulis nodis subgeminis. Capitula masc. (plantæ monoicæ) in axillis superioribus ramorum plerumque orta, pisi circiter magnitudine, multiflora, bracteïs exterioribus minimis bracteolisque ovatis; perigonio 5-partito, lobis glabris, mucronibus subapicalibus brevibus obtusis. Capitula feminea masculis sæpius minora, involucrio puberulo, bracteolïs lineari-spathulatis ciliatis, perigonii laciniis linearibus apice parce ciliato-dentatis, achænio ovato-elliptico subæquilongis. — Sequenti habitu proximum. Stipula extra-axillaris quam illa folii normalis magis evoluta limbum folii rudimentarii nonnihil refert. (v. s. in hb. Juss., Delessert, et h. lugd. bat.)
23. *E. LINEOLATUM* (Wight, Icon. fl. Ind. t. 1984), fastigiatim ramosum, ramis pubescentibus, foliis alternis elliptico-vel oblongo-lanceolatis longe lineari-acuminatis ante acumen integrum hinc seu in dimidio latiore limbi serraturis 2-3 illinc dente solitario instructis triplinerviis paucicostulatis (costula basilari lateris angustioris limbi ultra medium ejusdem producta) glabris vel solum subtus aut rarius utrinque (præsertim in nervis) adpresse pilosulis pilosisque, stipulis lineari-lanceolatis vix persistentibus, floribus dioicis, capitulis utriusque sexus sessilibus solitariisque, masculis multifloris. 2/ In *Zeylania* et provinciis humidioribus peninsulæ Indiæ orientalis Carnatic, Malabar et Concan frequentissimum, necnon in montibus Khasiæ altit. 1500-1800 metr. et in Himalaya tropica. — Wedd. Monogr. p. 312. — Caulis herbaceus aut suffrutescens, simpliciter vel subdichotome ramulosus, 2-4 decim. altus, ramis gracilibus pennæ corvinæ sæpe vix crassitie breviter vel longiuscule pubentibus demumque glabris obtuse angulatis, exsiccatis oliva-

ceæ vel pallidi-viridibus, cystolithis plus minus punctulatis. Folia 3-4 centim. longa, 1 centim. circiter lata, acumine 10-12-millimetrali, serraturis acuminæque acutis obtusiusve, nervis primariis hinc 3-6 ante marginem arcuatim anastomosantibus, jugo inferiore margini parallelo, laminæ exsiccatæ membranaceæ pagina superiore præsertim cystolithis linearibus oculo nudo valde conspicuis ornata in inferiore autem nunc sparsis (illisque superioris conformibus aut multo minoribus et in nervulis præcipue evidentibus) nunc vix ullis conspersa, petiolo millimetrali aut subnullo. Stipulæ 2-5 millim. longæ, glabriusculæ aut dorso adpresse pilosulæ. Capitula in axillis singulis semper solitaria et sessilia; mascula pisi minoris circiter magnitudinis, bracteis exterioribus rotundatis obtusissimis pubescentibus, bracteolis oblongis in nervo dorsali et apice pilosis, floribus subfasciculatis, perigonii 4-partiti segmentis glabris sub apice obtusissimo breviter mucronatis; femina masculis minoræ, bracteis involucri parvis pubescentibus, bracteolis lineari-spathulatis longeque ciliatis; perigonii fructiferi laciniis minimis denticulatis, æchenio elliptico acuto plus quam dimidia parte brevioribus. (v. s.)

β, *major*, foliis quam in typo majoribus (interdum decimetrum longis) serraturis hinc 4-6 illinc 2-3.

γ, *falcigerum*, foliis minoribus quam in typo (sæpe vix sesquicentimetralibus), acumine lineari subfalcato reliquum limbum longitudine subæquante, nervis basilaribus medium limbum non attingentibus, stipulis fere subulatis persistentibus.

δ, *lineare*, foliis (5-6 centimetralibus) linearibus vel lineari-lanceolatis subæquilateralibus integerrimis aut prope apicem hinc vel inde 1-2-denticulatis.

24. *E. EURHYNCHUM* (Miq. Fl. Ind. bat. 1 p. 2, p. 244), herbaceum, puberum vel glabrescens, foliis alternis oblique ellipticis abrupte longeque acuminatis basi semimarginatis hinc supra basim illinc supra dimidium limbi grossiuscule serratis (acumine integro) tri-triplinerviis, stipulis lineari-lanceolatis, capitulis femineis geminis brevissime pedunculatis exinvolucrat. ¼ In ins. Java (Zollinger, n. 1451). — E. parvum β Miq. in Zolling. Syst. Verzeichn. p. 102. — Caulis ramique tenuiter pubescentes vel glabrescentes. Folia majora 5 centim. circiter longa, nervo basilari lateris angustioris folii infra medium vel in medio limbo evanido, subtus demum lacunoso-reticulata, supra raro pilis rigidis inspersa, exsiccata cystolithis lineolata. Stipulæ tenere membranaceæ. Capitula fem. receptaculo discoideo exinvolucrato (Descript. ex Miq.). — E. lineolato sat affine videtur, præsertim varietati β, sed in planta Miqueliana capitula fem. gemina sunt absque involucrio.

25. *E. SUBINCISUM* (Wedd. Monogr. p. 314), herbaceum, caule simplici vel ramoso glabro, foliis alternis parvis oblique oblongo-lanceolatis acuminatis basi semiauriculatis supra trientem inferiorem inciso-serratis (serraturis obtusiusculis) triplinerviis paucicostulatis nervis basilaribus medium limbi parum superantibus margine hispidulis at cæterum glabris, stipulis lineari-lanceolatis persistentibus, capitulis dioicis masculis latiuscule involucriis brevissimisque pedunculatis multifloris ad nodos singulos solitariis. ¼ In regione temperata Indiæ orientalis, prov. Sikkim, altit. 2130 metr. (J. D. Hooker). — Caulis gracilis, 1-3 decim. longus, simplex vel parce ramosus, basi radicans, striatus, internodiis superioribus brevibus. Folia 1½-3½ centim. longa, 6-12 millim. lata, dente apicali s. acumine cæteris paulo longiore obtusiusculoque, nervis primariis vix prominentibus, in latere ampliore folii 3-5, in angustiore contra tantum 2-3; lamina exsiccata tenuiter membranacea supra obscure subtus pallide viridi cystolithis parvis oblongis sparsisque suffusa, petiolo subnullo. Stipulæ 5-6 millim. longæ, millimetrum circiter latæ. Capitula masc. 3-5 millim. lata, bracteis duabus exterioribus sub apice longiuscule crasseque mucronatis, sequentibus carinatis, bracteolis oblongis apice ciliato-laceris, perigonii 4-partiti segmentis (duobus?) sub apice longiuscule mucronatis glabriusculis. (v. s.)

26. *E. DISSECTUM* (Wedd. Monogr. p. 314), herbaceum, glabrum, caule adscendente, foliis alternis oblique oblongo-lanceolatis acuminatis basi acutis supra

partem quartam vel tertiam inferiorem grossiuscule crenato-serratis serraturis acutiusculis aut apiculato-acuminatis triplinerviis subpaucicostulatis nervis basilaribus medium limbi parum superantibus, stipulis parvis fere subulatis superpersistentibus, capitulis masc. longe pedunculatis solitariis multifloris, bracteis duabus exterioribus sub apice longe mucronatis, receptaculo demum subdichotome ramoso s. dissecto, capitalis fem. sessilibus. ♀ In Indiæ orientalis regione temperata montium prov. Sikkim et Khasia, altit. 1200-2400 metr. (J. D. Hooker et Thomson). — Caulis simplex, 1-4 decim. altus, basi radicatus, crassitie pennæ anserinæ aut graciliore, angulato-sulcatus, succulentus, inferne sæpius denudatus, exsiccatus cystolithis punctulatus. Folia 6-15 centim. longa, 1-5 centim. lata, interdum fere æquilateralia, acumine sublineari aut lanceolato integro, serraturis laxiusculis margine exteriori convexis, nervis primariis in latere latiore folii 5-6, lamina exsiccata tenuiter membranacea pallide virente, cystolithis linearibus (in pagina superiore præsertim manifestis) utrinque obsita, petiolo brevi. Stipulæ 4-7-millimetrales. Capitula monica vel dioica, mascula 1-2 centim. lata, pedunculo 1-4-centimetroali, bracteis binis extimis ovatis mucrone 2-5 millim. longo munitis, sequentibus lanceolatis mucroneque destitutis, receptaculi primum rotundati dein subfastigiatim ramosi sive profunde 4-partiti segmentis magis minusve lobatis bracteoas lanceolato-v. lineari-oblongas glabrasque margine exhibentibus, perigonii 4-partiti glabri lobis sub apice brevissime mucronatis; feminea raro breviter pedunculata, bracteis involucri triangulari-lanceolatis, bracteois oblongo-spathulatis apice plus minus incrassatis obtusive margine longiuscule ciliatis, perigonii fructiferi lacinilis breviter triangularibus denticulatis, achænio obovato in dimidio superiore carnosulo. (v. s. in hb. Kew. et Mus. par.)

27. *E. LEPIDOTULUM* (Miquel, Fl. Ind. bat. suppl. 1. p. 413), ramis glabris, foliis alternis oblongo-sublanceolatis-ellipticisve parum inæquilateralibus abrupte acuminatis supra medium minute repando-denticulatis subpauci-penni-nerviis subtus squamellis orbicularibus membranaceis fuscis adspersis, glomerulis axillaribus solitariis globosis glabris. ♀ In Sumatræ occidentalis prov. Padang (Teysmann). — Rami læves, albid. Folia 6-15 centim. longa, costulis utrinque 6-7 patulis æqualibus, limbo cæterum subevenio, exsiccata chartacea subtus pallida utrinque cystolithis fusiformibus notata. Achænia pallide rubiginoso-punctata. (Descript. ex Miq. l. c.)

\*\* Folia sensim acuminata vel absque acumine (1).

28. *E. CORNUUM* (Wedd. Monogr. p. 316), caule repente glabro, foliis alternis oblique elliptico-oblongis vix acuminatis basi semi-cordatis toto margine vel ultra tertiam partem inferiorem acute grossiusculeque serratis triplinerviis subpaucicostulatis, nervis basilaribus medium limbum superantibus glabriusculis vel supra sparsim pilosulis nervis subtus hispidulis, stipulis lanceolatis subpersistentibus, capitulis masc. longe pedunculatis solitariis multifloris, bracteis sex exterioribus sub apice longissime mucronato-appendiculatis. ♀ In Indiæ orientalis prov. Khasia, altit. 0-1200 metr. (J. D. Hooker et Thomson). — ? *Procris gibbosa* Wall. Cat. n. 7273. — Caulis herbaceus, gracilis, parce ramosus, ramis elongatis apice vix puberulis ex nodis omnibus radices agentibus, internodiis elongatis. Folia  $3\frac{1}{2}$ -8 centim. longa, 2-4 centim. lata, basi hinc auriculata, illinc semirotundata vel acutata, serraturis triangularibus margine exteriori vix convexis, dente apicali lanceolato acuto vix centimetroali, nervis primariis in latere latiore folii 5-8, jugo inferiore in utraque parte medium limbum superante, lamina exsiccata tenuiter membranacea pallide viridi subtus glaucescente cystolithis linearibus tenuibusque inspersa, petiolo 1-8-millimetrali. Stipulæ siccitate scariosæ, acutissimæ, circiter 5 millim. longæ Capitula masc. depressa, centimetrum lata, pedunculo 3-6-centimetroali glabro, bracteis 6 exterioribus glabris rotundatis sub apice obtusissimo mucrone semicentimetroali lineari acutissimoque instructis, bracteois obovatis oblongisve ciliatis, perigonii 4-partiti lobis glabris sub apice brevissime mucronatis. — Proceridem gibbosam ad hanc speciem dubitanter retuli. Habitus

(1) Vid. quoque inter species præcedentes *E. lineolatum* &

plantæ Wallichianæ illum E. Cornuti mentitur, sed caulis hirsutus est foliaque sinuato-crenata. Flores non vidi. (v. s.)

29. *E. LONGIFOLIUM* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 189; Monogr. p. 317), caule apice pubescente, foliis alternis suboblique elongato-lanceolatis acuminatis basi acutis a triente inferiore ad apicem laxè et obtusiuscule v. subacutè serratis subtriplinerviis subpaucicostulatis (nervis basilariibus limbo dimidio brevioribus) supra glaberrimis costa subtus adpresse hispidulo-pilosa, stipulis linearibus aut subulatis diu persistentibus, capitulis masc. multifloris sessilibus, femineis sessilibus.  $\frac{1}{2}$  In ins. Philippinis (Cuming, n. 456). — Miq. l. c. Caulis elongatus, crassitie pennæ corvinæ, demum glabratus, subsimplex, exsiccatus striatus, internodiis elongatis. Folia 6-10 centim. longa, 12-20 millim. lata; nervis primariis utrinsecus 6-7 ante marginem arcuatim anastomosantibus subtus parum prominentibus, jugo inferiore margini propinquo paralleloque quam dimidius limbus brevior; laminæ exsiccatae rigidulæ pagina superiore cystolithis fusiformibus crebriusculis inspersa, inferiore pallidiorè iisdem corporeculis reticulatim ornata; petiolo 3-5 millim. longo gracili puberulo. Stipulæ semicentimetales, acutissimæ, extus puberulæ. Capitula monoica in singulis nodis solitaria; mascula ex axillis inferioribus orta, depresso-hemisphærica, 6-8 millim. lata, multiflora, bracteis exterioribus majusculis rotundatis mucronatis extus breviter pubescentibus, perigonii 4-partiti? lobis mucrone lineari  $\frac{1}{2}$  millim. longo infra apicem instructis; fem. ex axillis superioribus orta, piso circiter æquimagna, involucri pubescente, bracteolis ciliatis. — Foliis sensim acuminatis stipulis longioribus diutiusque persistentibus et capitulis monoicis a var. majore *E. lineolati* (cui tamen habitu sat affine) recedit. (v. s.)

30. *E. URVILLEANUM* (Brongn. Bot. Voy. Coq. p. 210. t. 46. f. A), caule glabro, foliis alternis oblique obovato-lanceolatis acuminatis basi acutis in dimidio inferiore integerrimis cæterum serratis triplinerviis paucicostulatis utrinque glaberrimis, stipulis oblongis lanceolatisve persistentibus, capitulis masc. (multifloris) et fem. breviter pedunculatis.  $\frac{1}{2}$  In ins. Moluccana Ualan (d'Urville). — Wedd. Monogr. p. 318; Miq. Fl. ind. bat. 1 pars 2. p. 245. Caulis parum ramosus, ramis elongatis gracilibus, internodiis 1-1 $\frac{1}{2}$  centim. longis. Folia 5-5 $\frac{1}{2}$  centim. longa, 15-18 millim. lata, circiter a medio ad basim cuneata, nervis primariis utrinsecus 4-6 jugo inferiore medium limbum superante, lamina exsiccata utrinque cystolithis numerosis lineari-fusiformibus obducta, petiolo 2-3 millim. longo. Stipulæ acutæ, 3-5-millimetales. Capitula monoica vel dioica in singulis nodis solitaria, parva, hemisphærica, pedunculo stipulis æquilongo suffulta; mascula ex axillis superioribus orta, bracteis exterioribus ovatis acuminatis glabris, bracteolis obovatis haud ciliatis, perigonii 4-partiti lobis glabris mucrone subapicali brevi conico; feminea brevissime pedunculata, masculis subæquimagna, bracteolis lineari-spathulatis ciliatis, perigonii subbifidi laciniis lineari-oblongis ciliatis ovario fusiformi brevioribus. (v. s. in hb. Brongn.)

$\beta$ , *parvifolium*, foliis minoribus oblique ovatis vel ovato-lanceolatis supra trientem inferiorem serratis, stipulis lineari-lanceolatis, capitulis fem. sessilibus.  $\frac{1}{2}$  In ins. Ualan (d'Urville). *F. parvifolium* Brongn. l. c. f. B; Wedd. Monogr. p. 323; (Miq. l. c.). Folia 1-3 centim. longa, 4-10 millim. lata, floribus fem. perigonio bi-tri-fido, segmentis dimidio ovario brevioribus. (v. s.)

31. *E. LANCIFOLIUM* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 190; Monogr. p. 318), ramis prostratis altero latere strigosis, foliis alternis lanceolatis vel oblongo-lanceolatis attenuato-acuminatis basi rotundato-semicordatis integerrimis vel prope apicem inconspicue crenulato-serrulatis triplinerviis subpaucicostulatis supra glabris subtus in nervis margineque parvis pilosis, stipulis minimis lanceolatis, capitulis monoicis masculis (laxifloris) femineisque subsessilibus.  $\frac{1}{2}$  In ins. Java (Zollinger n. 2314). — Miq. l. c. p. 244. — Caulis herbaceus, longe diffuseque ramosus, ramis gracilibus exsiccatis striatis. Folia 2 $\frac{1}{2}$ -3 centim. longa, 6-10 millim. lata, interdum subfalcata, acuta, nervis primariis utrinsecus 5-7 subimmersis ante marginem arcuatim anastomosantibus, jugo inferiore medium limbum frequentius superante, laminæ exsiccatae



membranaceæ utraque pagina at præsertim superiore cystolithis linearibus inspersa, petiolo brevissimo. Stipulæ ciliolatæ. Capitula mascula et feminea in ramis distinctis collocata, in singulis nodis solitaria; mascula pauci- et laxiflora, bracteis exterioribus ovatis acutis tenuiter membranaceis pubescentibus, floribus longiuscule pedicellatis, perigonii 4-partiti lobis superne pilosis mucrone subapicali brevissimo apice piligero; feminea minima, involuacro piloso, bracteolis lineari-spathulatis longe ciliatis, perigonii lacinis brevissimis, achænio elliptico acutiusculo. (v. s.)

32. *E. Goudotianum* (Wedd. Monogr. p. 319), caule subtiliter pubescente, foliis alternis elliptico- vel oblongo-lanceolatis acuminatis basi obtusiusculis supra trientem inferiorem acute serrulatis triplinerviis subpaucicostulatis supra glabris subtus in nervis adpresse pubescentibus, stipulis lineari-lanceolatis diu persistentibus, capitulis dioicis, masculis multifloris sessilibus. ♀ In ins. Madagascaria, prope Tananarivo (Goudot, n. 29). — Caulis elongatus, crassitie pennæ corvinæ, rectiusculus, obtuse angulatus, internodiis subcentimetricibus. Folia 5-8 centim. longa, 1-1½ centim. lata, fere a medio ad basim subcuneata, serraturis parvis sinibus acutis sejunctis, nervis primariis hinc 6-8 illinc 3-4 ante marginem arcuatim anastomosantibus inferioribus medium limbum superantibus, laminæ exsiccatae pagina supericre cystolithis minimis fusiformibus inspersa, petiolo vix bimillimetrali gracili pubescente. Stipulæ 5-6 millim. longæ, acutissimæ, glabriusculæ. Capitula masc. in singulis nodis solitaria, subhemisphærica, 5-8 millim. lata, multiflora, bracteis exterioribus late ovatis mucronatis sparsim hispida, bracteolis oblongis vel obovato-linearibus ciliatis, perigonii 4-partiti segmentis dorso sparsim pilosis, mucrone subapicali lineari submillimetrali glabro. (v. s. in herb. Delessert.)

33. *E. Madagascariense* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 189; Monogr. p. 320), caule superne dense strigoso, foliis alternis parvis suboblique lanceolatis apice et basi acutis vel basi obtusiusculis utrinque subinciso-serratis fere penninerviis supra glabris nervis subtus strigosis, stipulis lanceolatis persistentibus, capitulis dioicis, femineis sessilibus. ♀ In ins. Marossi, prope Antongil-Bé (Bojer.). — Caulis 1-3-decimetralis, subsimplex, basi radicans, internodiis vix semicentimetricis. Folia 2-2½ centim. longa, 6-7 millim. lata, nervis primariis utrinsecus 6 in margine evanidis parum prominentibus, jugo inferiore sequentibus vix longiore, laminæ exsiccatae rigidulæ supra obscure virentis subtus pallidioris pagina superiore cystolithis fusiformibus (in pagina inferiore etiam conspicuis) ornata, petiolo 2 millim. longo pubescente. Stipulæ 3 millim. longæ, acutæ, glabriusculæ. Capitula fem. (solum visa) parva, involuacro pubescente, bracteolis linearibus longiuscule ciliatis, perigonii lacinis staminodia paulo superantibus sed achænio anguste ovato multoties brevioribus. (v. s. in herb. Mus. par. et Delessert.)

34. *E. rigidum* (Wedd. Monogr. p. 320), caule apice pubescente, foliis alternis suboblique oblongo-lanceolatis sensim acuminatis basi acutis a triente inferiore ad apicem laxè et obtusiuscule serratis trinerviis paucicostulatis (nervo basilari lateris angustioris limbi ad apicem ejusdem producto) supra sparsissime hispido-pilosis subtus in nervis adpresse pilosis, stipulis lanceolatis deciduis, capitulis dioicis, masculis glomeruliformibus sessilibus. ♀ In ins. Samar Philippinarum (Cumming, n. 1674). — Miq. l. c. p. 245. — Caulis elongatus, crassiusculus, rectus, rigidus, exsiccatus obtuse quadrangularis, utrinque anguste sulcatus, simplex vel parce ramosus, internodiis centimetrum vel ultra longis. Folia alterna, 5-10 centim. longa, 1-2 centim. lata; nervis primariis utrinsecus 3-4 ante marginem arcuatim anastomosantibus subtus prominentibus, jugo inferiore utrinque vel altero tantum latere ad apicem limbi pertingente marginique subparallelo; laminæ exsiccatae subcoriaceæ pagina superiore demum glaberrima cystolithis fusiformibus inspersa; petiolo 3-5-millimetrali pubescente. Stipulæ 4-6 millim. longæ, acutæ, extus pubescentes. Capitula masc. in singulis nodis gemina, at in speciminibus visis nondum perfecte evoluta, involuacro minimo, bracteis ovatis pubescentibus, pedicellis filiformibus pilosis quam perigonium pubescens (?) longioribus. (v. s.)

35. *E. incisum* (Wedd. Monogr. p. 321), caule superne dense strigoso, foliis alternis parvis suboblique obovato-oblongis deorsum anguste cuneato-attenuatis

- in dimidio superiore grosse inciso-serratis (dentibus utrinque 2-3) subpenninerviis paucicostulatis supra glabris subtus in nervis strigosulis, stipulis lanceolatis-subulatis persistentibus, capitulis dioicis, masculis paucifloris sessilibus. ♀ In Madagascaria, ad radices arborum, in locis humidis (Boivin, n. 269). — Caulis gracilis, repens, ramis ascendentibus, internodiis 2-3 millimetralibus. Folia 1-2 centim. longa, 5-8 millim. lata, dentibus obtusiusculis (apicali cæteris subæquimagno); nervis primariis utrinsecus 2-4 in apice dentium desinentibus vix prominentibus, jugo inferiore sequentibus æquilongo; laminae exsiccatae membranacæ supra læte viridis subtus palidioris pagina superiore cystolithis linearibus conspicuis in margine et costa crebrioribus ornata, pagina inferiore lisdem ut videtur prorsus carente; petiolo brevissimo glabrisculo. Stipulæ 3-millimetrales. Capitula masc. (solum visa)  $\frac{1}{2}$  centim. lata, pauciflora, bracteis exterioribus ovatis acutis ciliolatis, interioribus linearilanceolatis longiuscule ciliatis, floribus pedicellatis, perigonii 4-partiti lobis ipsi æquilongo instructis. (v. s. in hb. Mus. par.).
36. E. STIPITATUM (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 190; Monogr. p. 322. t. 9. D. f. 11-16), caule ramisque radicanibus strigosis, foliis alternis oblongis vel oblongo-lanceolatis apice attenuatis acutisque basi oblique semicordatis sessilibus ima basi excepta grossiuscule crenato-serratis triplinerviis paucicostulatis utrinque sparsim hispidulis subtusque in nervis subadpresse pilosis, stipulis lanceolatis diu persistentibus, capitulis dioicis, femineis longe pedunculatis. ♀ In Nova Hollandia (Verreaux, n. 145.). — Caulis interdum basi nudus glabratusque, lignescens, ramis gracilibus striatis, internodiis 1-2 $\frac{1}{2}$  centim. longis. Folia 2 $\frac{1}{2}$ -4 centim. longa, 1-1 $\frac{1}{2}$  centim. lata, dentibus acutis obtusive et haud raro acuminato-mucronatis: nervis primariis utrinsecus 2-5 in ipso margine folii desinentibus subtus prominulis, jugo inferiore dimidium limbum plerumque superante; laminae exsiccatae membranacæ subtus pallidæ pagina superiore cystolithis linearibus tenuissimis ornata. Stipulæ 4-5 millim. longæ, pilosulæ. Capitula fem. (solum visa) pisi magnitudine, pedunculo filiformi 8-15 millim. longo glabrisculo dimidiumque internodium longitudine superante suffulta, involucri pubescente, bracteis spatulatis longiuscule ciliatis, acheniis fere ellipticis quam perigonii lacinie multoties longioribus. (v. s. in hb. Mus. par. etc.)
37. E. POROPHYLLUM (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 190; Monogr. p. 323. t. 9. f. 17), ramis breviter strigosis s. strigoso-hispidis, foliis alternis parvis oblique oblongis obtusis acutisque basi semi-cordatis in triente superiore utrinque acute bi-tri-dentatis triplinerviis paucicostulatis, glabris, stipulis parvis lanceolatis, capitulis dioicis, masculis paucifloris sessilibus. ♀ In ins. Manilla (Cuming, n. 789). — Miq. l. c. — Caulis gracilis, fastigiato-ramosus, ramis basi fere nudis (exsiccatis teretiusculis), internodiis semicentimetralibus. Folia 1-1 $\frac{1}{2}$  centim. longa, 4-7 millim. lata, inferiora plerumque minima v. abortiva, dentibus acutiusculis: nervis primariis utrinsecus 2-4 reticulatim connexis subtus vix prominentibus, jugo inferiore quam dimidium limbus longiore; lamina exsiccata rigida subtus pallida, supra cystolithis linearibus valde conspicuis dispersa; petiolo subnullo. Stipulæ millimetrum vix longæ. Capitula masc. (solum visa) in singulis nodis solitaria, pauciflora, bracteis exterioribus ovatis acuminatis bracteolisque linearibus hirsutis, perigonii 4-partiti segmentis hirsutis, mucrone subapicali lineari  $\frac{1}{2}$  millim. circiter longo. (v. s. in hb. Mus. par.)
38. E. ELLIPTICUM, glaberrimum, caulibus longe repentibus, foliis alternis vix oblique ellipticis obtusis basi rotundatis subcordatis integerrimis triplinerviis paucicostulatis, capitulis dioicis, masculis breviter pedunculatis, juniorum bracteis exterioribus arcte conniventibus floresque omnino obtegentibus. ♀ In prov. Mishmi Bengalie orientalis (Griffith). — Caulis graciles, crassitudine pennæ corvinæ etiamque graciliores, ramosi, tota longitudine radican-tes, ramique in parte inferiore aphylli. Folia 1-4 centim. longa, 7-25 millim. lata, fere æquilateralia, basi plus minus obliqua, apice nonnunquam vage acuminata (acumine obtusissimo); nervis primariis utrinque 2-3, basilaribus arcuatis ultra medium extensis; laminae exsiccatae rigida pagina superiore cystolithis linearibus impositæ (illis marginis sepe majoribus), inferiore hinc

fore omnino carente, petiolo 6-8 millim. longo. Stipulæ parvæ, ovato-lanceolatæ, deciduæ. Capitula masc. (juniora solum visa) in singulis nodis solitaria, æte evolutionem florum ovoidea, subacuminata, floribus ut videtur paucis. — Habitus *Ficus* repentis. Capitula masc. examinata vix bimillimetralia. (v. s. in hb. Kew.)

39. *E. obtusum* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 190; Monogr. p. 324), caule retrorsum hispidio, foliis alternis parvis oblique obovatis vel obovato-oblongis obtusissimis basi semicordatis in dimidio inferiore aut altius integris cæterum crenatis 3-5 grossiusculis obtusissimisque munitis triplinerviis paucicostulatis (nervis basilariibus dimidium limbum superantibus) supra glabris hispidulisve nervis subtus margineque sparsim setoso-hispidis, stipulis lineari-lanceolatis v. lanceolatis persistentibus, capitulis dioicis, masculis paucifloris longiuscule pedunculatis. ♀ In Himalaya temperata a Sikkim ad Simla et in montibus Khasiæ (Wallich, etc.) necnon in ins. Manilla (Barthe). — Procris obtusa Wall. Cat. n. 4636. — Caulis gracilis, flexuosus, 1-2-decimetralis, radicans, subsimplex, basi sæpe aphyllus, exsiccatus teretiusculus aut obsolete tetragonus, internodiis  $\frac{1}{2}$ -1 centim. longis. Folia 1-2-centimetralia, ciliolata, nervis primariis hinc 2-3 subtus prominulis; lamina exsiccata rigidula supra obscure viridi subtus pallidiore, adulta cystolithis linearibus supra inspersa subtus lisdem destituta; petiolo subnullo. Stipulæ  $1\frac{1}{2}$ -2 millim. longæ, acutæ. Capitula masc. in singulis nodis solitaria, pedunculo filiformi hispidulo suffulta, laxi-et pauci-flora, bracteis exterioribus ovatis bracteolisque sub apice hispidio-mucronatis; floribus majusculis (nempe 2-3 millim. longis), perigonii 4-partiti segmentis sparsim hispidulis, mucrone subapicali  $\frac{1}{2}$ - $\frac{2}{3}$  millim. longo hispidio. (v. s.)

β, *delicatumum*, caule filiformi atrigoso, foliis oblique ellipticis (subtus glaucescentibus) serraturas 1-2 obtusas utrinque monstrantibus. ♀ In ins. Philippinis (Callery, Barthe). — *E. delicatumum* Wedd. in Ann. sc. nat. 1. c. *E. glaucescens* β *delicatumum* ejusd. Monogr. p. 325. — Caulis valde flexuosus. Folia interdum majora quam in typo, supra sparsim pilosa, nervis subtus setoso-pilosis, exsiccata tenuiter membranacea cystolithis paucis linearibus in pagina inferiore ornata. (v. s. in hb. Mus. par.)

40. *E. glaucescens* (Wedd. Monogr. p. 325, exclus. var.), caule crasse breviterque et subadpresse piloso, foliis alternis parvis oblique ellipticis vel obovato-oblongis utrinque obtusis vel basi subsemicordatis in dimidio superiore serraturis crenisve 1-5 grossiusculis munitis cæterum integris paucicostulatis (nervis primariis inferioribus quam sequentes paulo longioribus) supra sparsim piloso-hispidis subtus in nervis strigoso-pilosis, stipulis parvis lineari-lanceolatis, capitulis femineis sessilibus, masculis paucifloris pedunculatis. ♀ In ins. Luzon (Commerson) et Manilla (Callery, Cuming, n. 629). — Miq. 1. c. — Caulis radicans, teres, 1-2-decimetralis, internodiis 5-10 millim. metientibus. Folia 8-28 millim. longa, 5-18 millim. lata, nervis primariis utrinque 3-4, ante marginem arcuatim reticulatimve connexis, subtus prominentibus; laminae exsiccatae subtus glaucescentis pagina superiore (nervis exceptis) cystolithis ut videtur prorsus carente, inferiore hisce crebriuscule ornata; petiolo brevissimo. Stipulæ 1-2 millim. longæ, adpresse pilosæ. Capitula in singulis nodis solitaria, mascula parva pedunculo 3-5-millimetrali suffulta pauci- et laxiflora, involucri bracteis exterioribus bracteolisque lanceolatis tenuiter membranaceis pilosulis, perigonii 4-partiti lobis setoso-pilosis, mucrone subapicali elongato pilifero. Capitula fem. minima, supra dense hirsuta, involucro pubescente, bracteolis lineari-spathulatis longe densissimeque ciliatis, perigonii laciniis brevissimis. (v. s.)

41. *E. monticolum* (Hook. fil. in Journ. Linn. Soc. 7. p. 216), caule gracili retrorsum strigoso-pubescente, foliis alternis oblique obovato-oblongis basi acutis vel semi-rotundatis grosse serrato-dentatis utrinque aut solum supra sparse setulosis paucicostulatis, stipulis lineari-lanceolatis petiolum superantibus, capitulis dioicis, femineis sessilibus depressis. ♀ ? In montibus Cameroons Africæ occidentalis altit. 2150 metr. (Mann). — Caulis 15-20 centimetralia. Folia 3-4 centim. longa, nervis primariis utrinque 2-3, exsiccata mem-

- *brancea viridia*, pagina superiore cystolithis oblongis linearibusve inspersa, petiolo brevi. *Bracteæ involucri* ♀ *exteriore ovatæ acutæ.* (v. s. in hb. Kew.)
42. *E. OBOVATUM* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 190; Monogr. p. 326), caule ramisque breviter crasseque pilosis, foliis alternis oblique obovatis obtusissimis supra basim latiuscule semicordatam integramque crenato-serratis triplinerviis paucicostulatis (nervis basilaribus medium limbum superantibus) supra glaberrimis subtus papilloso-pilosulis, stipulis lanceolatis cito deciduis, capitulis dioicis, femineis sessilibus. ¼ In ins. Manilla (Callery, Cuming, n. 52 et 628). — Miq. l. c. — Caulis filiformis, 1 decim. et quod excedit longus, teres, parce ramosus v. simplicissimus, radicans, internodiis 5-10-millimetralibus. Folia 1-2 centim. longa, 6-15 millim. lata, serraturis s. crenis parvis 1-2 millim. latis; nervis primariis utrinsecus 3-4 ante marginem arcuatim anastomosantibus subtus prominulis, jugo inferiore medium limbum plerumque superante; lamina exsiccata rigidula supra cystolithis crebris linearibus nervis parallelis oculoque nudo manifestis ornata; petiolo subnullo. Stipulæ 1-2 millim. longæ, pilosulæ. Capitula fem. (solum visa) in singulis nodis solitaria, piso fere æquimagna, involucrio pubescente, bracteis linearibus longe ciliatis, perigonii lacinii brevissimis, achæniis ellipticis subacuminatis. (v. s.)
43. *E. FILICOIDES* (Seemann, in sched. absque descript.), caule distiche ramoso glabro, foliis alternis parvis oblongis obtusis basi oblique subcordatis in toto margine laxè crenato-serratis penninerviis nervo medio piloso, stipulis lineari-lanceolatis, floribus. . . ¼ In ins. Viti (Seemann, n. 421, ex parte). — Caulis ut videtur sat elatus cylindricus, ramis distiche patentibus pilosulis. Folia numerosa, sat approximata, 2-2½ centim. longa, nervis primariis vix conspicuis, lamina exsiccata submembranacea pagina superiore nisi in nervo medio cystolithis destituta inferiore vero iisdem sparsim notata. — Inter specimina Seemanniana sub nomine *E. filicoidis* dispensata duæ exstant plantæ, altera scilicet filicoidea supra descripta, altera *Filix genuina.* (v. s.)
- B. — *Folia opposita sed maxime disparia, altero cujusque jugi minimo interdumque ægre conspicuo.*
44. *E. JUNGHUNIANUM* (Miq. Pl. Jungh. p. 18; Fl. Ind. bat. 1 p. 2. p. 246), frutescens, ramis junioribus pilosulis, foliis oppositis in singulis jugis autem maxime inæqualibus, altero amplo oblique elliptico longe acuminato basi obtuso acutove et integerrimo cæterum grosse inæqualiterque serrato triplinervio (nervo basilari lateris angustioris limbi longissimo) supra glabro subtus in nervis breviter et subadpresse piloso, altero abortivo minimo subrotundato inciso, stipulis lanceolatis deciduis, capitulis masc. multifloris sessilibus in singulis nodis geminis. ⅓ In sylvis montium Sumatræ, altit. 1550-1800 metr. (Junghuhn). — Wedd. Monogr., p. 527. — Frutex 1-1½-metralis (teste Junghuhn), ramis angulatis, adultioribus glabris. Folia 10-16 centim. longa, 4-6 centim. lata, acumine lineari sesquicentimetrali; nervis primariis hinc 4-6, illinc contra seu in latere angustiore folii ad nervum basilarem (limbo ipso æquilongum) reductis; lamina exsiccata rigidula nigricante utrinque cystolithis minimis fusiformibus obsita; petiolo 3-5 millim. longo. Stipulæ circiter 8-millimetrales, dorso adpresse pilosulæ. Capitula dioica ?, mascula in speciminibus visis 5-8 millim. lata, receptaculo involucrioque parum conspicuis, bracteis ovato-lanceolatis carinatis dorso ut ipsi pedicelli pilosis, perigonii 5-partiti segmentis sub apice mucrone conico crasso instructis superne pilosulis. (v. s. in hb. hort. bot. lugd. bat.)
45. *E. PAPILLOSUM* (Wedd. Monogr. p. 327), caule subtiliter pubescente, foliis oppositis sed maxime disparibus, cujusque jugi altero oblique oblongo-vel elliptico-lanceolato anguste acuminato basi acuto supra tertiam aut quartam partem inferiorem crenato vel crenato-serrato triplinervio (nervis basilaribus medium limbum superantibus) supra glabro subtus in nervis villosulo vel puberulo, altero abortivo minimo ovato v. lanceolato, stipulis lanceolatis diu persistentibus, capitulis dioicis in singulis nodis solitariis, masculis multifloris pedunculatis, femineis sessilibus. ¼ In Indiæ orientalis regione tropica prov. Cachar et Khasia, altit. 600-1200 metr. (J. D. Hooker et Thomson) —

*Procris peduncularis* Wall. Cat. n. 4634, B. — Caulis herbaceus, adscendens, 2-3-decimetralis, simplex, inferne denudatus imaque basi radicans, obsolete angulatus, modice crassus, siccitate (sicut tota planta) fuscescens cystolithisque ut videtur parvissime instructus. Folia majora 5-17 centim. longa, 2-5 centim. lata, acumine interdum bicentimetrali integro vel laxo serrulato, serraturis s. crenis sæpe inæqualibus margine exteriori convexis rectiusculisve obtusissimis acutisve vel apiculatis; nervis primariis hinc s. in latere latiore limbi 6-8, jugo inferiore s. majore medium limbum (illinc præsertim) sæpius superante; lamina exsiccata membranacea supra cystolithis minimis oblongis fusiformibus conspersa; petiolo 1-4-millimetrali sicut costa villosa vel subtiliter pubescente. Folia abortiva 2-4 millim. longa, integerrima v. paucidentata. Stipulæ 5-8 millim. longæ, 1 millim. circiter latæ. Capitula masc. (in specimine unico visa) centimetrum circiter lata, receptaculo indiviso, bracteis exterioribus late ovatis rotundatisque, bracteolis oblongis vel ovatis glabris, perigonii 4-partiti segmentis sub apice longiuscule aristato-mucronatis. Capitula fem. masculis subconformia, bracteolis linearibus margine et dorso papillis conicis crassiusculisque præditis, perigonii fructiferi segmentis brevibus triangularibus, achænio ovato vel elliptico conspicue costato. (v. s. in hb. Kew. et Mus. par.)

46. *E. STRACHEYANUM* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 188; Monogr. p. 328 t. 9. fig. 18), caule pubescente, foliis oppositis sed admodum disparibus, in singulis jugis altero oblongo-lanceolato attenuato-acuminato deorsum cuneato v. semi-rotundato excepta basi argute serrato triplinervio subpaucicostulato (nervis basilariibus dimidium limbum superantibus) supra sparsim piloso subtus in nervis hispido-pilosulo, altero abortivo minimo ovato sæpius integro, stipulis ellipticis diu persistentibus, capitulis dioicis sessilibus ad singulos nodos geminis v. solitariis, masculis submultifloris. ♀ In temperatis Himalayæ occid. ad Kumaon (Strachey et Winterbottom, n. 5) et in montibus Khasiæ altit. 1500-1800 metr. (J. D. Hooker et Thomson). — Caulis herbaceus, adscendens, basi radicans, in speciminibus visis 2-3 decim. longus, simplex, gracilis, exsiccatus teretiusculus. Folia 4-6 centim. longa, 1-2 centim. lata, utrinque supra partem quartam inferiorem limbi serratis, serraturis interdum subacuminatis, nervis primariis utrinsecus 5-6 reticulatim arcuatimque anastomosantibus, lamina exsiccata membranacea cystolithis linearibus sparsissime ornata, petiolo 2-5 millim. longo pubescente. Folia abortiva valde evidenter, 2 millim. circiter longa, ciliata, integra vel paucidentata. Stipulæ 4-5 millim. longæ, acutæ, scariosæ, puberulæ. Capitula masc. in singulis nodis gemina, subbiloba, pisi vix magnitudine, pauciflora, bracteis exterioribus rotundatis obtusissimis extus pubescentibus, bracteolis obovato-oblongis, perigonio 4-partito, segmentorum mucrone subapicali brevi conico. Capitula fem. masculis minora, subsolitaria, involucri pubescente, bracteolis obovatis longe ciliatis, floribus subsessilibus, perigonii segmentis quam staminodia brevioribus, achæniis late ellipticis obtusiuscule acuminatis. — Foliis oppositis et charact. capituli masc. distinguendum ab *E. sessili* cui primo aspectu valde simile. (v. s.)

47. *E. DIVERSIFOLIUM*, glabriusculum, foliis oppositis admodum disparibus, cusque jugi altero majore nunc (in nodis superioribus) lanceolato et plerumque attenuato-acuminato basi obtusiusculo vel semirobundato et utrinque supra medium dentibus 2-4 (rarius 5-10) instructo triplinervio, nunc (in jugis inferioribus) rotundato v. ovato et subintegro, altero abortivo fere lineari subintegroque, stipulis minimis, capitulis dioicis, masculis paucifloris pedunculatis, femineis sessilibus. ♀ ? In montibus temperatis Zeylanæ (Thwaites, n. 2181) et Himalayæ a Sikkim ad Kumaon, altit. 1200-2700 metr. et in montibus Khasiæ et Nilghiri (Wight, etc.). — ♂. surculosum Wight, Icon. fl. Ind. t. 2091. f. 4; Wedd. Monogr. p. 329. *Procris diversifolia* Wall. Cat. n. 4631. *P. elegans* ejusd. l. c. n. 4632. — Caulis herbaceus, gracilis, 1-2-decimetralis, fere simplex, basi radicans et stolonibus filiformibus sæpe instructus, glaber vel hispidiusculus. Folia majora alia 1½-7 centim. longa, alia 2-3 millim. vix metientia, glaberrima sæpiusve ciliolata et interdum sparsissime papillosa, serraturis magnitudine variabili interdum fere lanceolatis obtusiusculis, dente apicali triangulari aut lineari; nervis primariis in

foliis superioribus utrinsecus 4-6 subimmersis in margine desinentibus jugo inferiore medium limbum paulo superante; lamina exsiccata tenuiter membranacea ut plurimum late viridi subtus pallidior, pagina superiore cystolithis linearibus sparsim ornata, inferiore eadem ut videtur destituta; petiolo brevissimo. Folia abortiva evidentissima, elliptico-ovata v. lanceolata, 2-5 millim. longa, integerrima. Stipulae ovatae, submillimetrales. Capitula in singulis nodis solitaria, mascula laxe pauciflora, pedunculo 5-20 millim. longo, bracteis exterioribus ovatis ciliolatis glaberrimisve, perigonii 4-partiti lobis breviter lineari-mucronatis (mucrone parce piloso) aut fere muticis. Capitula fem. pisi circiter magnitudine aut multo minora, involucri foliolis ciliatis, bracteis obovato-spathulatis, floribus plerisque pedicellatis, perigonii laciniis brevissimis, achenio maturo subcylindrico dessiccatione (?) sulcato. (v. s.)

β, *latum*, foliis normalibus omnibus obovatis. γ In Nepalla (Wallich). — E. *latum* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 190. Procris *lata* Wall. Cat. n. 463. P. monandra? Hamilt. mscr. (v. s.)

48. E. APPROXIMATUM, caule in altera facie hirtulo vel strigoso, foliis oppositis valde imparibus, cujusque jugi altero majore oblique obovato obtuso acutiusculove deorsum latiuscule cuneato et integro in dimidio superiore grossiuscule crenato tripplinervio supra nervisque subtus sparsim hispido-piloso, altero minime ovato integro, stipulis ovatis persistentibus, capitulis fem. sessilibus ☉ in Indis orientalis montibus Nilghiri (Wight) et prov. Concan (Stocks) necnon in ins. Java (Labillardière). — E. *cuneatum* Wight, Icon. pl. Ind. 6. t. 2091. f. 3; Wedd. Monogr. p. 330; Miq. Fl. Ind. bat. 1 pars 2 p. 246. E. *webbianum* Wedd. in Ann. sc. nat. 1. c. Procris *approximata* Wall. Cat. n. 4630. Herba, caule gracili 1-2-decimetrali basi radicante exsiccato striato simplice v. parum ramoso, ramis filiformibus. Folia majora alia (superiora)  $\frac{1}{2}$ -4 $\frac{1}{2}$  centim. longa, 5-20 millim. lata, alia (inferiora) minima, nonnumquam subintegra, nervis primariis hinc 3-4 ante marginem reticulatim vel arcuatim connexis subtus vix prominentibus, jugo inferiore dimidium limbum superante; lamina exsiccata tenuiter membranacea obscure virente, pagina superiore cystolithis linearibus tenuissimis inspersa; petiolo 1-2 millim. longo. Folia abortiva minima et interdum aegre conspicua, ovata. Stipulae petioli circiter longitudine. Flores dioici? Capitula fem. (solum visa) in singulis nodis subsolitaria, bracteis involucri pubescentibus floresque superantibus, bracteis oblongo-spathulatis breviter ciliatis, perigonii laciniis brevissimis, achenio elliptico obtusiuscule acuminato. (v. s.)

β, *alternifolium*, foliis omnibus (abortivis nempe nullis) oblique obovatis ellipticisve utrinque attenuatis basi oblique subsemicordatis in dimidia parte superiore grossiuscule crenato-serratis orenatisve. ☉ In peninsula Malaica ad Mergui (Griffith). — Caulis in angulis strigosus, cæt. ut in typo. (v. s. in hb. Kew.)

*Species non satis cognita.*

49. E. SINUATUM (Hassk. Cat. hort. Bog. p. 79), herbaceum, foliis (alternis?) elliptico-oblongis acuminatis basi oblique attenuatis supra basin grosse sinuatis exserte serratis glabris membranaceis in costa subtus puberulis, floribus conferte axillaribus. γ In ins. Java. — Miq. Fl. Ind. bat. 1 pars 2 p. 247. Procris *sinuata* Blume, Bijdr. p. 511. (Descript. ex Bl. l. c.) — E. *nigrescenti* affine secundum cl. Miquel.

50. E. GRACILE (Hassk. l. c.), herbaceum, foliis (oppositis sec. cl. Miquel) lanceolatis acutis basi angustatis subsessilibus supra basin serratis glabris, stipulis membranaceis carinatis, capitulis subsessilibus involucriatis. γ? In ins. Java. — Miq. l. c. Procris *gracilis* Bl. l. c. p. 513.

51. E. PARVUM (Bl. macr. Miq. l. c. p. 247), herbaceum, foliis (oppositis sec. cl. Miquel) oblique ovato-oblongis obtusis supra basin acute serratis in venis subtus capitulisque dimidiato-verticillatis hirsutis. γ? In ins. Java. — Procris *parva* Bl. Bijdr. p. 512.

*Species exclusæ.*

- E. FRUTESCENS* Hassk. = *Procris frutescens*.  
*E. JAPONICUM* Wedd. = *Pellionia umbellata*.  
*E. LÆVIGATUM* Hassk. = *Procris lævigata*.  
*E. LUCIDUM* Forst. = *Procris pedunculata*.  
*E. MONTANUM* Endl. = *Procris?* montana.  
*E. OPPOSITIFOLIUM* Dalz. = *Lecanthus peduncularis*.  
*E. OVATUM* Wight = *Lecanthus peduncularis*.  
*E. PEDUNCULATUM* J. R. et G. Forst. = *Procris pedunculata*.  
*E. RADICANS* Wedd. = *Pellionia radicans*.  
*E. ROSTRATUM* Hassk. = *Pellionia elatostemoides*.  
*E. SUCCOSUM* Miq. = *Procris pedunculata*.  
*E. TRINERVE* Hochst. = ?  
*E. UMBELLATUM* Blume = *Pellionia umbellata*.  
*E. WIGHTIANUM* Wedd. = *Procris lævigata*.

XVI. *PROCRIS* Commerson, mscr.; Juss. Gen. p. 403; Wedd. Monogr. p. 333.  
 t. 6. B. Seem. Fl. Vit. p. 141. — *Procridis* spec. quorund. — Boehmeria spec.  
 Pers. synopsis. 2. p. 556. — *Sciophila* Gaudich. Voy. Uran. p. 493.

Flores monoici vel dioici, masculi glomerati, glomerulis (raro capitulisformibus) cymosis, feminei in receptaculum carnosum globosum vel clavatum densissime aggregati, cymis capitulisque (ebracteatis) solitariis. Masc. Perigonium 5-partitum, laciniis obovatis carnosulis muticis. Stamina 5. Pistilli rudimentum globosum v. obovatum. Fem. Perigonium tri-tetraphyllum, segmentis obovatis cucullatis carnosulis. Ovarium ovatum, perigonio brevius. Ovulum imo loculo affixum, funiculo brevifutrum. Stigma sessile, longe penicillatum, mox evanidum. Staminum rudimenta nulla? Achænium ovatum vel ellipticum, subbaccatum, striatopictum, perigonii laciniis carnosis obtectum, capitulo fructifero demum fragariiformi. Embryo turbinatus, cotyledonibus late ellipticis radícula conica longioribus, albumine paucissimo vel deficiente. — Suffrutescentes succulenti frutescensve, in iisdem regionibus ac *Elatostemata* obvia, erectiusculi vel adscendentes, sæpius glabri. Folia disticha, inæquilateralia et valde inæquimagna (folio nempe majore in utroque latere caulis cum folio abortivo seu bracteiformi alternante), integerrima vel sinuato-dentata, penninervia, cystolithis linearibus minimis inspersa. Stipulæ axillares integræ. Capitula feminea sessilia aut pedunculata; floribus bracteolis lineari-spathulatis apice sæpe glandulosis stipatis; cymæ masc. pedunculatæ, pedicellis ebracteatis.

1. *P. PEDUNCULATA*, foliis majoribus oblique lanceolatis vel oblongo-lanceolatis acuminatis basi acutis, integerrimis aut prope apicem obsolete sinuato-serratis, cymis masc. sæpius parvis bracteis parum conspicuis, capitulis fem. solitariis sessilibusque. ♀ Ad rupes et truncos emortuos in convallibus montium ins. Javæ (Junghuhn, Zollinger, n. 912), Timor (Jacquinot, Societatis (Forst.), Taiti (Bertero, Moerenhout, etc.), Mascarenis (Commerson, Gaudichaud, etc.). *P. cephalda*, Commers. mscr., Poir. Encycl. 4. p. 629, Wedd. Monogr. p. 334. t. 6. B., Miq. Fl. Ind. bot. 1 (2) p. 258. *P. longifolia* Bl. Bijdr. p. 508. *P. lucida* Spreng. Syst. veg. 3. p. 846. *B. Torresiana* Steud. Repert. p. 396. *Elatostema pedunculatum* J. R. et G. Forst. Char. gen. pl. ins. austr.

- p. 106. t. 53. E. lucidum eorund. ex Guillemain in Ann. sc. nat. 2. sér. 7. p. 184, Pers. Synops. 2. p. 557. E. succosum Miq. Pl. Jungh. p. 23. Dorstenia lucida G. Forst. Prodr. fl. ins. austr. n. 58. Boehmeria Cephalida Pers. Synops. 2. 556. Sciophila Torresiana Gaudich. l. c. — Caulis succulentus, 2-5-decimetralis, simplex aut parce ramosus; ramis teretiusculis magis minusve sinuatis, internodiis brevibus. Folia majora patentia, 6-20 centim. longa, 1½-4 centim. lata, basi oblique attenuata, acumine sublanceolato v. lineari interdum arcuato sæpe obtusiusculo, nervis primariis utrinque 6-8 supra immersis subtus in lamina exsiccata (rigida) vix prominentibus ante marginem inconspicue arcuatum conjunctis, petiolo vix centimetro pulvino prominente insidente. Folia abortiva fere lanceolata 5-15 millim. longa, subsessilia, axilla plerumque (ut majorum) fertili. Cymæ masc. pedunculis 1-2 centim. longis suffultæ, e glomerulis paucis sæpiusque approximatis compositæ, floribus brevissime pedicellatis sessilibusve, perigonii 5-partiti segmentis inæqualibus obovatis obtusissimis carnosulis glaberrimisque. Capitula fem. caulem sæpe magis minusve amplectentia, initio albida, sub maturitate globosa aut varie difformia, diametro vario at cerasum haud raro magnitudine æmulantia, colore rubente, floribus minutissimis densissimeque confertis bracteolis linearibus ciliolatisque intermixtis, perigonio fructifero e foliolis tribus spatulatis superne incrassatis margineque minutissime glanduloso-ciliolatis plerumque constante achænia ovata pericarpio extus tenuiter carnosio prædita superante. (v. s.)
2. P. FRUTESCENS (Bl. Bijdr. p. 510), foliis majoribus suboblique obovato-oblongis abrupte angustate acuminatis basi acutis in tertia aut quarta parte inferiore integris cæterum laxè sinuato-serrulatis acumine fere integro, cymis masc. laxis, bracteis minutis, capitulis fem. solitariis sessilibusque. 3 Ad arbores scandens, in montibus ins. Javæ altit. 1000-1500 metr. (Junghuhn, Zollinger, n. 510 et 1291 Z). — Wedd. Monogr. p. 335; Miq. Fl. Ind. bat. 1 (2). p. 248. Elatostema frutescens Hassk. Cat. hort. bogor. p. 79; Miq. Pl. Jungh. 1. p. 19. — Caulis ligneus, repens, ramis erectis subsimplicibus crassis glabris siccitate pallidis. Folia majora 12-16 centim. longa, 4-6 centim. lata, acumine lineari uni-sesqui-centimetro, nervis primariis utrinque 8-10 sub lente (in planta exsiccata) squamuloso-puberulis, lamina exsiccata membranacea supra nigricante cystolithis minimis fusiformibus parce conspersa, petiolo 1-sesqui-centimetro. Folia abortiva lanceolata, centimetralia vel paulo longiora. Inflorescentiæ masc. ex axillis inferioribus ramorum ortæ, pedunculatæ, subdichotome ramosæ, ramulis ultimis fasciculato-floriferis; perigonii 4-phylli segmentis membranaceis. Capitula fem. ut in præcedente cui stirps ipsa valde est affinis. (v. s. in hb. hort. lugd. bat. et Franqueville.)
3. P. LÆVIGATA (Bl. Bijdr. p. 508), foliis majoribus suboblique lanceolatis acuminatis basi acutis in dimidia parte superiore laxè serratis, acumine integro, inflorescentia masc. pauciflora racemiformi?, capitulis fem. pedunculatis fasciculatis. 4 Ad truncos arborum pseudoparasitica, in montibus Zeylanæ, Peninsulæ Indiæ orientalis, Khasiæ et Himalayæ, altit. 300-1200 metr. frequens, necnon in sylvis Javæ et in insulis africanis Comore (Boivin) et Fernando Po (Mann). — Miq. Fl. Ind. bat. 1 (2). p. 248. P. Wightiana Wall. Cat. n. 4636; Wedd. Monogr. p. 336. Elatostema lævigatum Hassk. Cat. hort. bogor. p. 79. E. Wightianum Wedd. in Ann. sc. nat. 4. sér. 1. p. 188. — Caulis adscendens, 2-4-decimetralis, succulentus, basi sæpiissime digiti auricularis crassitie, ut tota planta glaberrimus, merithallis brevibus, nodis superioribus s. hornotinis solum foliiferis, epidermide exsiccata rugosa pallida. Folia majora 8-15 centim. longa, 15-25 centim. lata, fere æquilateralia, acumine lanceolato, acutiusculo, serraturis inæqualibus, nervis primariis utrinsecus 6-7, lamina exsiccata tenuiter membranacea cystolithis minimis fusiformibus sparsim inspersa olivaceo-aut fusco-virente, petiolo 3-4-millimetroli Stipulæ ut in speciebus præcedentibus brevissime triangulares, obtusæ. Folia abortiva oblongo-lanceolata, semicentimetralia, obtusiuscula, subsessilia. Capitula fem. quam in cæteris speciebus minor, nempe 3-5 millim. tantum lata, globulosa, sed cum pedunculo crasso 3-9-millimetroli claviformia, in singulis axillis (sæpe aphyllis) gemina terna aut quaterna, fasciculata, perigonii fructiferi sub-4-phylli foliolis rotundatis cucullatis hyalinis bracteolisque (floribus inter-



positis) conformibus quam achænia ovata parum compressa acutiuscula dimidio brevioribus. (v. s.)

β, *cuneata*, foliis e basi inæquali anguste cuneato-oblongis abrupte obtuse breviterque acuminatis præter apicem (in plerisque repandum) integerrimis. 2. In sylvis Javæ (Zollinger n. 912 pro parte). — *P. cuneata*. Bl. l. c. p. 509; Miq. l. c. 249. *Elatostema lævigatum* β subcuneatum ejusd. in Zolling. Syst. Verzeichn. p. 102. (v. s.)

4. *P. GRANDIS* (Wedd. Monogr. p. 337), foliis majoribus amplis ovatis vix inæquilateralibus basi rotundatis subcordatisve acuminatis integerrimis, floribus masc. dense capitatis, capitulis orbicularibus laxè dichotomo-cymosis, ramis primariis cymæ bracteis majusculis subcordato-rotundatis suffultis. 5 In Nova Guinea? — Miq. Fl. Ind. bat. 1 (2). p. 249. — Caulis ramosus, ramis crassis teretibus lævibus siccitate olivaceis. Folia majora 20-30 centim. longa, 10-16 centim. lata, penninervia, acumine subcentimetrali acutiusculo, nervis primariis 10-12, jugo inferiore sequentibus minore, lamina exsiccata rigida petioloque 6-8 centim. longo cystolithis punctulatis. Folia abortiva oblongo-spatulata, 3-4 centim. longa, 6-8 millim. lata, subsessilia, obtusissima, enervia. Cymæ masc. majusculæ, petiolos foliaque minora longe superantes, ter quaterque laxè dichotomæ, pedunculo ramisque 1-4 centim. longis, puberulis glabratissive, bracteis fere rotundatis cochleatis obtusissimis v. mucronatis centimetrum circiter longis, capitulis sæpe majusculis ramulos cymæ summos terminantibus orbicularibus. (v. s. in hb. Kew.)

*Species dubia.*

5. *P. LATIFOLIA* (Bl. Bijdr. p. 509), suffruticosa, dioica?, foliis longe petiolatis elliptico-oblongis obtusis acuminatis basi obtusiusculis integerrimis glabris, capitulis subsessilibus (Blume). 5 In ins. Java. — Wedd. Monogr. p. 337; Miq. l. c.

*Stirps dubiæ sedis.*

6. *P. MONTANA* (Steud. Nomencl. 2. p. 398), foliis magnis oblique oblongis cuspidatis petiolatis serratis scabriusculis, floribus masc. paniculatis, capitulis fem. per ramos sparsis breviter pedunculatis. 5 In insula Norfolk. — Wedd. Monogr. p. 338. *Elatostema montanum* Endlich. Prodr. Fl. Norf. p. 39. — Rami dichotome ramulosi. Ramuli digitum minimum crassi, teretiusculi, gemma conica acuminata 4-centimetrali terminati, internodiis centimetralibus. Folia 3-decimetralia, 4-5 centim. lata, valde inæquilateralia, latere superiore limbi in ramo erecto quadrante angustiore basi que abrupte angustato, inferiore in petiolum subdecurrente necnon ima basi libero et rotundato, remote et obsolete serrulata, nonnulla subintegerrima, utrinque pilis brevissimis crassiusculis adpressiusculisque in pagina superiore rarioribus rigidioribusque in inferiore copiosioribus et mollioribus conspersa; petioli basi dilatata vaginantibus, summo margine discolore libero stipulari caulem annulatim ambiente. Paniculæ masc. subdichotomæ, 15-20 millim. longæ, solitariae binæ vel ternæ, pedunculo pedicellis que crassis, pilis clavatis articulatis crystallinis ad apicem pedicelli assurgentibus floremque cupulatim recipientibus, perigonii 5-partiti segmentis ovatis concavis carinatis, filamentis staminum usque ad medium adnatis. Capitula fem. depressiuscule subglobosa, receptaculo carnosio exalbido hyalino nucis avellanæ mole bullato-verrucoso, ovariis plurimis ovato-subglobosis nudis pilis articulatis densissime stipatis, stigmatibus plurimis piliformibus. Achænia magnitudine et colore seminis cannabini, stigmatibus persistentibus coronata, sinibus receptaculi inclusa. (Descript. ex Endlich.) — Planta characteribus notabilis sed pluribus horumce a genere recedens. Synonymum Cunninghamianum ab Endlicherio laudatum (*Boehmeria castaneæfolia*) ad *Morum pendulinam* pertinere videtur secundum ipsum Cunninghamium.

*Species exclusæ.*

- P. ACUMINATA* Poir. = *Elatostema acuminatum*.  
*P. APPROXIMATA* Wall. = *Elatostema approximatum*.

- P. AUSTRALIS Spr. = *Elatostema sessile*.  
 P. CANTONIENSIS Spr. Syst. 3. p. 846 = alius ordinis.  
 P. CAULESCENS Poir. = *Dorstenia caulescens*, fide Stend.  
 P. CHINENSIS Spr. Syst. 3. p. 846 = alius ordinis.  
 P. COCHINCHINENSIS Syst. 3 p. 846 = alius ordinis.  
 P. CYRTANDRÆFOLIA Zoll. et Moritz = *Elatostema nigrescens*.  
 P. DIVARICATA Steud. = *Pellionia divaricata*.  
 P. DIVERSIFOLIA Wall. = *Elatostema diversifolium*.  
 P. ELATOSTEMOIDES Steud. = *Pellionia elatostemoides*.  
 P. ELEGANS Wall. = *Elatostema diversifolium*.  
 P. EXCELSA Steud. = *Boehmeria excelsa*.  
 P. FAGIFOLIA Poir. = *Elatostema fagifolium*.  
 P. FICOIDEA Wall. = *Elatostema ficoideum*.  
 P. GIBBOSA Wall. = *Elatostema cornutum*?.  
 P. GLABRA Hook. et Arn. = *Urera sandwicensis*.  
 P. GRACILIS Bl. = *Elatostema gracile*.  
 P. HEYNEANA Wall. = *Pellionia Heyneana*.  
 P. HIRSUTA Bl. = *Elatostema hirsutum*.  
 P. HYPOCRYPTA Commers. = *Elatostema fagifolium*.  
 P. INTEGRIFOLIA Don = *Elatostema integrifolium*.  
 P. LÆTA Wall. = *Elatostema diversifolium*.  
 P. LATERIFLORA Poir. = *Boehmeria cylindrica*.  
 P. LITTORALIS Poir. = *Boehmeria cylindrica*.  
 P. LONGIFOLIA Poepp. = *Phenax angustifolius*.  
 P. MACULATA Poir. = *Elatostema maculatum*.  
 P. MELASTOMOIDES Spr. Syst. 3. p. 847 = ?  
 P. MEMBRANACEA Reinw. = *Elatostema acuminatum*.  
 P. MOLLIS Wall. = *Elatostema molle*.  
 P. MONANDRA Hamilt. = *Elatostema diversifolium*.  
 P. NIVEA Gaudich. = *Boehmeria nivea*.  
 P. NUDIFLORA Poir. Suppl. 4. p. 454 = ?  
 P. OBTUSA Wall. = *Elatostema obtusum*.  
 P. PALUDOSA Bl. = *Elatostema macrophyllum*.  
 P. PARVA Bl. = *Elatostema parvum*.  
 P. PEDUNCULARIS Wall. = *Elatostema papillosum*.  
 P. PEDUNCULARIS Wall. = *Lecanthus peduncularis*.  
 P. PIPERIFOLIA Wall. = *Elatostema integrifolium*.  
 P. POLYCEPHALA Wall. = *Elatostema sessile*.  
 P. PUNCTATA Hamilt. = *Elatostema sessile*.  
 P. PUNCTATA Wall. = *Elatostema rupestre*.  
 P. QUINQUENERVIA Spr. Syst. 3. p. 846 = alius ordinis.  
 P. RACEMOSA Royle = *Pilea Wightii*.  
 P. RADICANS Sieb. et Zucc. = *Pellionia radicans*.  
 P. ROSTRATA Reinw. = *Pellionia elatostemoides*.

- P. *RUGOSA* Poir. = *Phenax ballotæfolius*.  
 P. *RUPESTRIS* Don = *Elatostema rupestre*.  
 P. *SESSILIS* Hook. et Arn. = *Elatostema sessile*.  
 P. *SESQUIFOLIA* Reinw. = *Elatostema integrifolium*.  
 P. *SIEBERI* Steud. (Sieb. Martin. n. 105) = ?  
 P. *SINUATA* Bl. = *Elatostema sinuatum*.  
 P. *SONNERATII* Spr. = *Phenax Sonneratii*.  
 P. *SPLENDENS* Lindl. = *Boehmeria australis*.  
 P. *STRIGOSA* Bl. = *Elatostema strigosum*.  
 P. *TRINERVATA* Poir. = *Gyrotænia cephalantha*.  
 P. *URTICÆFOLIA* Poir. = *Phenax urticæfolius*.  
 P. *UMBELLATA* Sieb. et Zucc. = *Pellionia umbellata*.

Tribus III. *BOEHMERIÆ* Gaudich, pro parte; Wedd.  
 Monogr. p. 340.

Frutices v. arbores v. suffrutices aut rarissime herbæ, inermes. Folia alterna v. decussatim opposita, rarissime ternatim verticillata, homomorpha aut symmetrice diversiformia, quandoque inæquilateralia, cystolithis sæpius punctiformibus inspersa. Stipulæ caulinae (axillares aut interpetiolares), rarius petiolares, plerumque liberæ. Inflorescentiæ exinvolucratæ. Flores diclini. *Masc.* : perigonio 4-5-partito rarissime 3-partito, segmentis muticis raro dorso gibbosis aut mucronatis in præfloratione valvatis imbricatisve, pistilli rudimento forma variabili glabro v. lanato, alabastro sæpe acuminato. *Fem.* : perigonio interdum plane deficiente (1) vel brevissimo, sæpius autem tubuloso-ventricosus, nunc libero nunc ovario connato, postea nthesin sicco v. carnosus, ore ut plurimum contracto 2-4-dentato v. edentulo; stigmate vario sed numquam vere penicillato; staminum rudimentis nullis. Fructus siccus v. bacca-tus, glaberrimus v. setosus. Embryo cotyledonibus ellipticis v. oblongis rarissime rotundatis apiceve aut basi emarginatis, radicu-la conica, albumine vario.

Subtrib. I. *EUBOEHRERIÆ* Wedd. Monogr. p. 341. Perigonium fem. liberum aut ovario magis minusve adhærens, fructiferum siccum s. membranaceum læve aut varie costatum vel alatum. Stigma sæpius lineare, modo cum apice ovarii continuum et in fructu persistens, modo basi valde constrictum s. articulatum et mox deciduum.

XVII. *BOEHMERIA* Jacq. *Stirp. amer. hist.* p. 216; *Juss. Gen. pl.* p. 403; *Michx. in Mart. Fl. Bras.* 12. p. 185; *Wedd. Monogr.* p. 343; *Benth. Fl. hongk.* p. 331; *Seem. Fl. Vit.* p. 241. — *Boehmeria*, Duretii et Procridis spec. *Gaudich. Bot. Voy. Uran.* p. 499. — *Boehmeria* spec. *quorumd.* — *Splitgerbera Michx. Comment. phyt.* p. 134. — *Urticæ* spec. *auct. antiq.*

Flores monoici vel dioici, semper glomerulati, bracteis brevibus scarioris stipati, glomerulis (in planta monoica sæpius unisexualibus) nunc

(1). In describendis generibus *Boehmeriearum*, perigonium dixi nullum quem nulla vestigia oris integumentis talis sunt manifesta.

solitariis axillaribusque nunc spicatis aut rarius in cymam paniculiformem dispositis. *Masc.* Perigonium 4-partitum vel 4-lobum, rarissime 3- aut 5-partitum, segmentis ovatis subacuminatis vel infra apicem breviter mucronatis, in præfloratione valvatis. Stamina tot quot perigonii segmenta. Pistilli rudimentum clavatum vel fere globosum, glaberrimum aut basi breviter lanatum pilosulumve. *Fem.* Perigonium tubulosum, compressum vel plus minus ventricosum, ore sæpius contracto conspicueque 2-4-dentato. Ovarium inclusum, a perigonio discretum v. eidem adhærens, sessile vel pedicellatum. Ovulum imo loculo v. juxta fundum loculi funiculo brevi affixum, erectum aut adscendens. Stigma elongato-filiforme, cum summo ovario continuum, v. go altero latere solum pilulosum sæpiusque diu persistens. Achænium ovario subconforme, perigonio marcescente inclusum eique haud raro cohærens, pericarpio crustaceo tenui vel nucamentaceo ut plurimum fusco. Albumen varium, sed nunquam deficiens. Embryo cotyledonibus ellipticis, radícula conica sæpius paulo longioribus. — Frutices vel arbusculæ vel suffrutices, inter tropicos utriusque orbis obvii, pauci extratropici, plerique sylvicolii, sæpius magis minusve pubentes. Folia opposita aut alterna (etiam in eodem ramo), homomorpha et æquilaterialia vel dimorpha et interdum inæquilaterialia, varie dentata, in unica specie biloba, trinervia, lævia aut rugosa, petiolata, cystolithis punctiformibus parum conspicuis. Stipulæ axillares, ut plurimum liberæ vel basi tantum coalitæ, rarissime in unam integram connatæ, vulgo deciduæ. — Habitu solo non distinguendæ sunt Boehmeriæ a multis aliis stirpibus ejusdem tribus; characteribus autem floris feminei: stigmate scilicet lineari cum apice ovarii continuo et persistente, necnon perigonio sub maturitate fructus sicco, nullo modo difficile est eas dignoscere.

De indole perigonii feminei celeberrimus Blumius recte monet sequentia: « Textura, ait, plerumque est tenerrima membranacea sicca, neque ulla parte cum pericarpio cohæret; in aliis autem est magis vegeto-herbaceum et, quamquam primo liberum, tum interdum cum pericarpio maturescente cohæsione plus minus arcta jungitur; in aliis vero speciebus in simili statu perigonii vel in fructu liberum est. » — Exempla insignia hujus singularis casus in *B. nivea* non infrequenter observentur.

Dixi marginem perigonii feminei in nonnullis post anthesin incrassatum vel alatum fieri, atque sane advertit auctor supra laudatus pericarpium etiam (cum perigonio cohærente) in hoc statu sæpe plus minus affici, quod in stirpibus perigonio libero nunquam accidit.

Quod autem ad delimitationem generis attinet a Blumio valde dissensior. Ita, ut alia levioris momenti taceam, clarissimus auctor, fere omnino nescius characteris optimi ex ovarii cum stigmate connexionē desumpti, nullumque videns aliud discrimen quam quod pendet a præsentia costarum vel alarum in perigonio fructifero, immo verisimiliter immemor levis pretii (sua propria sententia sacrati) hujus characteris, species horum generum alias aliis infauste commiscuit. Ulterius dabitur forsân locus redeundi ad animadversiones cl. auctoris de affinitatibus Boehmeriæ cum aliis nonnullis generibus.

### § 1. *Species neogææ.*

#### A. — *Glomeruli axillares.*

\* Folia alterna.

1. *B. RAMIFLORA* (Jacq. Stirp. Amer. hist. p. 246. t. 157), monoica vel dioica, foliis alternis dimorphis, aliis lanceolatis basi acutatis obtusatisve aut ra-

rius subcordatis petiolatis, aliis cum præcedentibus alternantibus minoribus ovato-lanceolatis basi semi-cordatis sessilibusque, omnibus longe angustaque acuminatis serratis (minoribus ab ipsa basi) sæpe rugosis supra sparsim pilosulis v. asperato-hispidis subtus pubescentibus, glomerulis axillaribus unisexualibus, floribus masc. triandris, perigonio fructifero lanceolato apice longiusculo attenuato pubescente, achænio sessili.  $\frac{3}{5}$ - $\frac{5}{5}$  In ruderatis et fruticetis montium ins. Antillarum. — Swartz Prodr. p. 34; ejusd. Fl. Ind. occid. l. p. 283; Bl. Mus. bot. lugd. bat. 2. f. 13; Wedd. Monogr. p. 344; Griseb. Fl. Br. W. Ind. p. 159. *Caturus ramiflorus* Linn. Mantiss. p. 127. *Procris ramiflora* Poir. Encycl. 5. p. 630; Lam. Illustr. gen. t. 76. f. 1. — *Arbuscula bi-metr* lis, ramulis teretiusculis pubescentibus demumque glabratiss, internodiis brevibus. Folia maiora erecto-patula, 8-20 centim. longa,  $2\frac{1}{2}$ - $5\frac{1}{2}$  centim. lata, margine altero convexiore, minora 2-7-centimetralia, priora basi plerumque integra, serraturis reliqui limbi triangularibus acutiusculis, acumine  $1\frac{1}{2}$ -3 centim. longo integerrimo; nervis basilaribus inæquilongis in dimidio latiore folii medium limbum vix attingentibus, in angustiore contra limbo fere æquilongis rectiusculisque; lamina exsiccata membranacea læte virente, petiolo gracili 1-4 centim. longo pilosulo. Stipulæ lanceolatæ, centimetrales, in nervo dorsali adpresse pilosæ, ciliolatæ. Glomeruli pisiformes, valde multiflori, nunc omnino discreti, nunc fere confluentes, masculi et feminei in distinctis ramulis vel in eodem ramulo nascentes, feminei superiores. Flores masc. brevissime pedicellati, bracteis numerosissimis diuque persistentibus intermixti, perigonii 3-lobi lobis extus pubescentibus iuxta apicem gibboso-cucullatis, pistilli rudimento clavato glabro. Flores fem. subsessiles, perigonio lanceolato quam stylus gracillimus triplo quadruplo brevior in dimidio superiore præsertim pilis rectis uncinatisve obsito ore bidentato, fructifero sesquimillimetrali vix compresso margine acutiusculo inferne glabrato achænio subconforme rectiusculum includente. (v. s.)

$\beta$ , *cuspidata*, foliis omnibus petiolatis, minoribus basi acutatis vel rotundatis subcordatisve nec semicordatis,  $\frac{3}{5}$  In iisdem locis ac typus, necnon in monte Orizaba imperii mexicani (Botteri, n. 280). — *B. cuspidata* Wedd. l. c. p. 345. (v. s. in hb. Mus. par.)

$\gamma$ , *subreticulata* (Blume, l. c. p. 199), ramulis petiolisque quam in typo densius pubescenti-hirtulis, foliis (minoribus ovatis, majoribus ovato-oblongis) minus inæquilateralibus et minus falcatis pro rata parte latioribus, serraturis copiosioribus minoribusque, utraque pagina et præsertim inferiore subvelutino-pubescenti. — Cult. in horto Cæs. Schoenbrunensi. — Ad hanc formam verisimiliter adscribendum est specimen Fendlerianum e Venezuela allatum sub n. 1266.

2. *B. FALLAX* (Wedd. Monogr. p. 346), dioica aut monoica, foliis alternis in nodis proximis dimorphis, aliis majoribus oblongo- v. ovato- v. elliptico-lanceolatis acuminatis basi obtusis rotundatisve vel ad insertionem petioli submarginatis toto margine serratis aut ima basi et acumine integris, aliis (cum præcedentibus alternantibus) minimis v. interdum deficientibus rotundatis obtusissimis sessilibus paucidentatis, omnibus plus minus rugosis supraque in intermedio fastigiatim v. substellatim hispido-pilosis et subtus pubescentibus nervis hirtis, glomerulis axillaribus androgynis unisexualibusque, floribus masc. tetrandris, perigonio fructifero elliptico-ovato piloso, achænio sessili.  $\frac{3}{5}$  In Peruvia (Mathews, n. 2036; Poeppig, n. 1623). — Fruticosa, ramulis gracilibus breviter hirtis puberulisve internodiis brevibus. Folia patula, maiora  $2\frac{1}{2}$ -6 centim. longa, 8-20 millim. lata, sæpe nonnihil inæquilateralia, nervis basilaribus ad partem quartam superiorem limbi vel in altero latere altius productis subtus vix prominentibus, intermedio reticulato, lamina exsiccata vix asperula, petiolo 2-3-millimetrali. Folia minora sæpe ægre conspicua etiamque omnino deficientia, axilla horumce nihilominus sicut cætera florifera. Stipulæ lineari-lanceolatæ, 3-5-millimetrales, nervo dorsali pilosæ. Glomeruli vix pisi magnitudine, masculi sæpe confluentes folioque nullo axillati, floribus masc. breviter pedicellatis sessilibusve, perigonii 4-lobi segmentis sub apice acutissimo mucrone conico obtuso munitis sparsim hispidis, pistilli rudimento obovato inferne pilis paucis rectis instructo. (v. s. in hb. Mus. par.)

- β, *cordata*, foliis majoribus ovato- v. oblongo-lanceolatis longiuscule acuminatis basi cordatis ad extremum apicem serratis, minoribus prorsus deficientibus. ♂ In Peruvia (Pavon). — Rami pubescentes. Folia 3-15 centim. longa, apice h. u. d. raro falcata. Glomeruli in speciminibus visis omnes feminei, parvi. Perigonia fructifera vix millimetralia, ventricosa, utrinque attenuata, superne præsertim hirta-pilosa, pilis paucissimis uncinatis cæteris commixtis, ore bidentato dentibus lanceolato-triangularibus. (v. s. in hb. Boissier.)
- γ, *ulmifolia*, foliis majoribus basi subacutis supra undique hispido-pilosis asperatisque, minoribus ovatis. ♂ In sylvis imperii mexicani prope Xalapa, altit. 1500 metr. (Galeotti, n. 311). B. ulmifolia Wedd. Monogr. p. 347; Bl. l. c. p. 226. — Ramuli breviter hirti, internodiis brevibus. Folia majora 5-10 centim. longa, minora semicentimetralia vel paulo majora, acumine in prioribus lanceolato et sæpe integro, serraturis limbi subacuminatis, rugis vix prominulis. Glomeruli parvi, floribus masc. in specimine suppetente inter femineos rarissimis, perigonio hortumce ut in var. β. (v. s. in hb. Mus. par.)
3. B. *CELTIDIFOLIA* (Kunth in H. et B. Nov. gen. et sp. 2. p. 34), dioica, foliis alternis in nodis proximis valde inæquimagnis ovatis ellipticisve aut lanceolatis acuminatis basi obtusis acutiusculisve aut rotundatis argute serratis rugosis supra strigosis asperulisque subtus pubescenti-hirtis vel tomentosis, glomerulis axillaribus, floribus masc. ...., perigonio fructifero lineari- vel obovato-lanceolato basi longe attenuato superne subadpresse hispido, achænio pedicellato. ♂ In locis aridis lapidosisque Andium quitenium, altit. 2900 metr. (Humb. et Bonpl.); in prov. Quindiu et Antioquia Novæ Granatæ (Triana). — Wedd. Monogr. p. 348; Bl. Mus. bot. lugd. bat. 2. p. 200. — Fruticosa, ramulis gracilibus teretiusculis adpresse s. strigoso-pubescentibus, internodiis brevibus. Folia (majora erecto-patentia, 5-10 centim. longa, 1 1/2-3 centim. lata, minora patuli bicentimetralia vel sæpe breviora) ima basi sæpius integra, serraturis reliqui limbi subacuminatis; nervis basilaribus ad 1/2 superiorem limbi productis supra impressis subtus prominulis, intervenio nonnunquam lacunoso-impresso, lamina exsiccata coriacea rugosa rugis obtusis pilis brevibus adpressis crebriuscule hispidis, petiolis 2-8 millim. longis pubescentibus. Stipulæ lanceolæ æ. 4-5 millim. longæ, extus puberulæ vel adpresse pilosæ. Glomeruli fructiferi p. s. minori æquimagni, vix densi, stylo et perigonio fere æquilongis singulis circiter 3-millimetralibus, hocce vix compresso ore bidentato longiuscule hispido achæmium parvum (pedicello haud raro brevius) laxiuscule includente. (v. s. in h. Mus. pav.)
4. B. *ASPERA* (Wedd. Monogr. p. 349. t. 11. f. 24-28), monoica, foliis alternis dimorphis, aliis ovato-lanceolatis lanceolatisve superne attenuatis, aliis multo minoribus ovatis, omnibu basi acutis obtusatisve crenulato-serratis rugosis supra adpresse exasperato-strigosis subtus pubescenti-hirtis, glomerulis axillaribus, perigonio fructifero ovato apice attenuato dense pubescenti-hispido, achænio subsessili. ♂ In Andibus Novo-Granatensium, prope Quindiu (Goudot). — Ramuli densiuscule pubescentes, internodiis brevibus. Folia patentia; majora 6-10 centim. longa, 1 1/2-3 centim. lata, minora vix 3-centimetralia et 1-1 1/2 centim. lata, limbo infimo ad insertionem petioli haud emarginato, nervis basilaribus ad trientem superiorem folii productis supra impressis; lamina exsiccata coriacea eximie bullato-rugosa, rugis paginae superioris plerisque conoideis pilisque crebriusculis valde adpressis radiatisque (cystolithos referentibus) notatis; petiolis 5-10 millim. longis pubescentibus. Stipulæ triangulari-lanceolatæ, semicentimetrales, acutissimæ, nervo dorsali adpresse piloso, persistentes. Flores in glomerulos avillares sessiles pisiformes densissime congesti, glomerulis plerumque unisexualibus superioribus fem. inferioribus masc. Flores masc. pubescentes, perigonii 4-lobi lobis latè ellipticis sub apice breviter et obtuse mucronatis, pistilli rudimento basi pilis quibusdam rigidis stipato, perigonio femineo (ore bidentato) stylo æquilongo, fructifero vix compresso millimetrali densiuscule hispido, achænio elliptico ventricoso utrinque attenuato. (v. s. in hb. Mus. par.)
5. B. *PAVONII* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 202; Monogr. p. 354), mo-

noica, foliis alternis in nodis propinquis inæquimagnis et subheteromorphis lanceolatis longe acuminatis supra basim integram argute serrulatis breviter petiolatis membranaceis lævibus aut vix rugosis supra glabris vel subtiliter et adpresse puberulis subtus pube tenuissima albicante vestitis nervis hirtellis, glomerulis axillaribus sæpius unisexualibus, perigonio fructifero utrinque attenuato subcompresso anguste membranaceo-marginato. ♂ In Peruvia (Pavon). — Arbuscula, ramulis novellis teretiusculis canescenti-puberentibus, internodiis uni-sesqui-centimetralibus. Folia majora 10-20 centim. longa, 3-7 centim. lata, basi obtusa, acumine plus minus elongato, minora 3-5-centimetralia, latitudine varia, nonnunquam ovata basiue rotundata, acumine quam in foliis majoribus sæpius pro rata parte multo breviora, serraturis omnium parvis et ad basim limbi obsoletis, nervis basilaribus interdum fere ad acuminis basim productis subtus parum prominulis sparsim et breviter hirtulis demumque fere glabris; lamina exsiccata discolora, juniore membranacea lævique et supra haud raro nitida, vetustiore sæpe plus minus rugosula; petiolo pro limbi magnitudine variæ longitudinis, etiam in minoribus glomerulum superante. Stipulæ lanceolatæ, deciduæ. Glomeruli pisum minorem crassitudine æmulantes, masculi et feminei in eodem ramo vel in distinctis ramis orti, floribus masc. subsessilibus, perigonii 4-lobi segmentis extus sparsim pubescentibus: perigonio fructifero circiter 2 millim. longo in parte superiore præsertim pilis brevibus rectis et quibusdam uncinatis intermixtis hirtis, ore 2-4-dentato. (v. s. in hb. Boiss.)

β, *diversifolia*, foliis magis heteromorphis, majoribus lanceolatis vel oblongo-lanceolatis longiuscule acuminatis, minoribus ovatis lanceolatisve vel oblongis non aut breviter acuminatis basi rotundatis cordatisve, omnibus crenulato-serratis et plus minus bullato-rugosis subcoriaceis supra glaberrimis vel sparsim puberulis subtus pubescenti-hirtis et sæpe canescentibus. ♂ ? In Peruvia, prope Palca (Dombey; Mathews, n. 2034); prope Tarapoto (Spruce, n. 4029 et 4951), Nova Granata (Linden, n. 479). — B. *diversifolia* Wedd. Monogr. p. 350. B. *strigosa* Bl. l. c. 226. (v. s. in hb. Mus. par., Webb., Kew, etc.)

γ, *anomala*, foliis majoribus tantum evolutis, cæteris omnino nullis loco tamen horumce præsentia glomerulorum designata. ♂ Secus rivulos Peruviae orientalis, prope Tarapoto (Spruce, n. 3962). — Arbor debilis, circiter 5-metralis, patule ramosa, ramulis gracilibus flexuosis subtiliter puberulis demumque glabratis. Folia 10-15 centim. longa, lanceolata vel oblongo-lanceolata, anguste acuminata, basi obtusa vel rotundata, crenato-serrata, lamina exsiccata submembranacea et fere concolore. Glomeruli minores quam in typo (v. s.).

6. B. *LINDENIANA* (Bl. Mus. bot. lugd. bat. 2. p. 226), monoica?, foliis alternis conformibus æquilateralibusque ellipticis vel elliptico-lanceolatis acuminatis basi obtusis vel rotundatis argute serratis coriaceis supra subrugosis et hispidulo-strigosis subtus velutino-tomentosis, glomerulis axillaribus unisexualibus, perigonio fem. subpubescente apice longiuscule attenuato quadridentato. ♂ In Nova Granata (Linden, n. 164). — Ramuli petiolique strigosuli. Folia 5-10 centim. longa, 2-4  $\frac{1}{2}$  centim. lata. Ovarium pedicellatum, perigonio obsolete 4-dentato inclusum.

7. B. *RHYNOPHYLLA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 202; Monogr. p. 354), dioica, foliis alternis sæpius inæquilateralibus elliptico- vel ovato-lanceolatis longissime et angustissime acuminatis basi fere cuneatis integrisque cæterum acute serratis planis supra molliter pilosiusculis subtus pubescentibus, glomerulis axillaribus, femineis petiolo multo brevioribus, perigonio fructifero elliptico-ovato valde compresso alato-marginato longiuscule acuminato. ♂ In Venezuela prov. Caracas (Funck, n. 430). — Ramuli graciles, inferne puberuli vel glabrati, apice incano-pubescentes, internodiis magis minus elongati. Folia late patentia, 1-2 decim. longa, 4  $\frac{1}{2}$ -6 centim. lata, margine altero convexiore, serraturis brevissime acuminatis, acumine ipsius limbi lineari 2-3-centimetrali basi tantum serrato, nervis basilaribus parum prominentibus inæquilongis altero nempe usque ad basim acuminis producto atque altero vix dimidium limbum superante, lamina exsiccata tenuiter membranacea

molliuscula, petiolo gracili 2-6 centim. longo adpresse pubescente. Stipulæ lanceolatæ, in nervo dorsali pilosæ. Glomeruli fem. (solum visi) pisi magnitudine, floribus sessilibus, perigonio superne pilosulo quam stylus longiuscule villosus multo brevior, fructiferi ( $1\frac{1}{4}$ - $1\frac{1}{2}$  millim. longi) facie utraque in medio convexuscula achæmium parvum arcte includente ore 2-dentato. (v. s. in hb. Mus. par.)

8. *B. BREVIROSTRIS* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 201; Monogr. p. 353), dioica, foliis alternis isomorphis ovatis acuminatis subcordatis crenato-serratis planis supra adpresse hispidis subtus subtiliter pubescentibus, glomerulis fem. petiolo brevioribus, perigonio fructifero rotundato parum compresso haud marginato. § In Peruvia (Mathews, n. 2039). — Ramuli novelli obtuse tetragoni, strigosi villosulique, internodiis brevibus. Folia patula, 4-8 centim. longa.  $3-4\frac{1}{2}$  centim. lata, serraturis parvis grossiusculisve acutis vel obtusiusculis, acumine etiam dentato, nervis basilaribus usque ad acuminis basim interdum productis subtus parum prominentibus, venulis conspicuis, lamina exsiccata rigidula, petiolo subcentimetroli strigoso. Stipulæ lanceolatæ, 4-6 millim. longæ, extus pilosulæ. Glomeruli fem. (solum visi) hemi-phærici, magnitudine pisi majoris, ramulum fere amplectentes, inferiores ut plurimum aphylli, floribus subsessilibus, perigonio elliptico omnino vel in dimidia parte superiore pilis rectis hispido quam stylus superne villosus multo brevior, ore plerumque breviter 4-dentato. fructifero  $1\frac{1}{4}$  millim. longo inferne glabrato faciebus convexusculis margine obtuso apice brevissime acuminato achæmiumque conforme arcte involvente, pericarpio nucamentaceo. (v. s.)

9. *B. EXCELSA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 202; Monogr. p. 352), monoica, foliis alternis et fere conformibus elliptico-ovatis apice attenuatis acuminatisve basi acutis aut obtusatis crenatis serratisque planis supra sparsim pilosulis sed mox glabratibus subtus breviter incano-tomentosis atque concinne reticulatis, glomerulis axillaribus sæpe unisexualibus, perigonio fem. ovario subadnato, fructifero ovato apice longiuscule attenuato compresso marginato pubescente. § In sylvis umbrosis montium et secus rivulos insulæ Juan Fernandez ubi dicitur Manzano (Bertero, Cl. Gay). — Bl. Mus. bot. lugd. bat. 2. p. 205. Procris excelsa Beitr. in Bot. Zeit. 33. p. 261; Walp. Ann. 3. p. 414. Splitgerbera denudata Cl. Gay, Fl. chil. 5. p. 362. — Arbor 5-10-metralis vel procerior, diametro trunci fere 3-decimetrali, cortice ejusdem rimoso fuscoque vel ferrugineo, ramulis novellis teretiusculis pubescentibus apice sæpe incanis, internodiis brevibus infimis sæpius aphyllis. Folia 4-10 centim. longa;  $1\frac{1}{2}$ -5 centim. lata, basi integra, dentibus reliqui limbi acutiusculis obtusisve, nervis basilaribus circiter ad trientem superiorem limbi productis costaque et venulis pubescentibus nec incanis, lamina exsiccata rigidula supra sub lente eximie albo-punctulata subtus nisi in venis glaucescenti-incana, petiolo centimetroli vel longiore pubescente. Stipulæ lanceolatæ, extus pubescentes. Glomeruli utriusque sexus vulgo in diversis ramulis siti, sed interdum in eodem commixti, raro androgyni, discreti aut fere confluentes, pisi circiter magnitudine. Flores masc. sessiles, perigonii 4-partiti lobis extus cano-pilosus juxta apicem acute mucronatis, mucronibus in alabastro conniventibus, pistilli rudimento clavato basi lanato. Flores fem. fere sessiles, perigonio ovato sive lageniformi quam stylus villosus brevior, fructifero compresso subincano-pubescente achænio lenticulari arcte cohærente, ore bi-quadridentato. (v. s. in hb. Mus. par., etc.)

\*\* Folia opposita.

10. *B. HIRTA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 202; Monogr. p. 358, non Swartz), dioica, foliis oppositis isomorphis ovatis acuminatis basi rotundatis vel fere cordatis crenato-serratis planis utrinque dense hirsutis, glomerulis axillaribus, perigonio fem. elliptico hirtio libero. § ? In Peruvia? (Mathews, n. 2039). — Ramuli tetragoni, pilis longis patulis undique dense hirsuti, apice siccitate fusciscentes. Folia (in eodem jugo inæquimagna) 6-10 centim. longa, 3-6 centim. lata, dentibus (prope basim limbi minoribus) sæpe subacuminatis, acumine ipsius limbi acutissimo serratoque, nervis basilaribus ad partem quartam superiorem limbi productis costaque dense hirsutis,



lamina exsiccata membranacea hirsutiae paginae superioris rigidula inferioris molli, petiolis foliorum ejusdem nodi inaequalibus 1-3-centimetralibus dense hirsutis. Stipulae anguste lanceolatae, pilosae, deciduae. Glomeruli fem. (solum visi) sphaeroides, 8-10 millim. lati, floribus sessilibus, perigonio utrinque attenuato quam stylus villosus paulo brevior. (v. s.)

11. B. REPENS, monoica, caulibus filiformibus lignescentibus diffusis repentibusque, foliis parvis oppositis subisomorphis sed in eodem pare valde inaequimagnis ovatis acutis supra basim integram pauciserratis utrinque sparsim pilosis, glomerulis axillaribus paucifloris unisexualibus, perigonio fem. oblongo-ovato superne angustato hirsuto ovario adnato. ♂ In Cuba orientali, ad muros, prope villam Monte Verde (Wright, n. 1459) — Pouzolzia repens Griseb. in Mem. Acad. am. sc. et art. n. ser. 8 (1860). p. 173. — Caulis 1-3-decimetralis, ramosi, ad nodos radicales, glabrati, ramulis adscendentibus pubescentibus. Folia 3-12 millim. longa, majora nonnunquam subacuminata, basi obtusa vel rotundata, serraturis utrinque 2-6, petiolo quam limbus dimidiis brevior pubescente. Stipulae ovatae, acuminatae, persistentes. Glomeruli 3-6-flori, floribus masc. pedicellatis, perigonii 4-partiti segmentis breviter acuminatis pubescenti-hirsutis; femineis sessilibus, perigonio fructifero late ovato fere bimillimetrali ore bidentato, pericarpio crasso crustaceo. (v. s.)

β, *major*, robustior, caulibus 3-5 decim. longis minus ramosis basi radicans, internodiis longioribus, foliis majoribus 3-5 centim. longis acuminatis, serraturis numerosis, petiolo 1-2-centimetrali. ♂ In eisdem locis ac typus (Wright, n. 533). — Specimina Wrightiana sub n. 540 dispensata sunt nonnulla typica atque alia inter typum et varietatem intermedia. (v. s.)

B. — *Glomeruli plerique spicati, spicis nonnunquam apice foliiferis; foliis oppositis aut in ramulis alternis.*

12. B. CAUDATA (Swartz, Prodr. p. 34; Fl. Ind. occid. p. 279), dioica, foliis oppositis ovatis vel ellipticis acuminatis basi acutis aut obtusis vel rotundatis serratis supra planis rugulosive sparsimque hispidis subtus pubescentibus, spicis simplicibus aphyllis quam folia longioribus patulis pendulisve, perigonio fructifero late elliptico vel obovato valde compresso membranaceo-marginato puberulo. ♂-♀ In regionibus tropicis et subtropicis Mexici, Antillarum, Novae Granatae, Peruviae Boliviae et Brasiliae frequens. — Miq. in Mart. Fl. bras. XII, 185, t. 44 et 45; Wedd. Monogr. p. 360: Bl. Mus. bot. lugd. bat. 2. p. 215. t. 56. C; Griseb. Fl. Br. W. Ind. p. 159. B. ampullacea et B. petiolaris Kunth in H. et B. Nov. gen. et sp. 2. p. 35 et 36. B. flagelliformis Liebm. in k. danske Vidensk. Selsk. Skr. 5 Røek. nat. og math. Afd. 2. p. 310. Frutex vel arbuscula 2-5-metralis, ramulis teretiussculis aut obtuse quadrangulis praesertim apice pubescentibus rarius pubescenti-tomentosis. Folia 5-15 centim. longa, 2-7 centim. lata, serraturis parvis acutis obtusiusculisve, juxta petiolum plerumque obsolete, acumine acuto integro vel serrulato, nervis basilaribus ad quartam partem superiorem limbi productis supra parum impressis, intermedio reticulato, lamina exsiccata submembranacea, petiolis in eodem jugo ut plurimum valde inaequalibus (longiore limbum dimidium saepe aequante) pubescentibus tomentosive. Stipulae ovato-lanceolatae, 6-8 millimetrales, acuminatae, extus nisi in nervo medio glabrescentes vel mollior pubescentes, subpersistentes. Spicae solitariae, primum erectae, mox folio longiores diffusaeque aut pendulae, saepissime plus quam bidecimetales, masculae et feminae subaequilongae vel masculae paulo longiores, rachis pubescente, glomerulis plerumque discretis subdiscretisve aut femineis nonnunquam submaturitate confluentibus pisi majoris circiter magnitudine. Flores masc. perigonio 4-partito, lobis ovatis acutis magis minusve pubescentibus, pistilli rudimento subclavato basi parce piloso, alabastro brevissime acuminato. Flores fem. subsessiles, perigonio elliptico apice in collum cylindricum breve attenuato ore 2-4-dentato, stylo basi glabrescente superne latiore villosoque quam perigonium longiore. Perigonium fructiferum membranaceum, quam maxime compressum, subdiscoideum, 2-2½ millim. longum, apice pilis brevibus rec-

tis uncinatisve hispidum, achænium parvum pedicellatum arcte includens. (v. s.)

β, *arborescens*, foliis lanceolatis subtus parce pubescentibus haud rugosis. 5 In Brasilia (Gardner). — *B. arborescens* Gardner, in Hook. Journ. of bot. 1. p. 541. (v. s.)

γ, *arguta*, foliis eadem forma ac in typo sed subtus albidotomentosis. 5 In Brasilia (Martius). — *B. arguta* Martius mscr. Miq. l. c. p. 186. — *B. peruviana* Bl. l. c. p. 224 ad hanc varietatem verisimiliter referenda est (v. s.).

13. *B. CYLINDRICA* (Willd. Spec. 4. p. 340), monoica, foliis oppositis alternisque ovato-vel elliptico-lanceolatis lanceolatisve acuminatis basi rotundatis subcordatisve sæpius breviter petiolatis serratis vel crenato-serratis rugulosis supra sparsim piloso-hispidis demumve glabris lævibusve subtus ramulisque villosis vel subtiliter pubescentibus, glomerulisque spicatis, nonnullis axillaribus, spicis simplicibus erectis apice nudis aut foliiferis, perigonio fructifero lenticulari-compresso crasse marginato superne pilis uncinatis sparsim instructo. 2-5 In Americæ borealis sylvis umbriosis saxosis et circa fluvios a sinu mexicano usque in Canadam frequens. — Wedd. Monogr. p. 362. t. 11. f. 18-23; Bl. Mus. bot. lugd. bat. 2. p. 212; Griseb. Fl. Br. W. Ind. p. 159. *Urtica cylindrica* Linn. Spec. 1396 et auct. U. mariana Miller Dict. Duretia cylindrica Gaud. Bot. Voy. Uran. p. 499. — Forma spicis apice subaphyllis: *B. cylindrica* Muehlenb. Cat. pl. Am. sept.; Pursh Fl. Am. sept. 1. p. 112. *B. dasypoda* Miq. in Mart. Fl. bras. 12. p. 187. — Forma spicis apice foliiferis: *B. lateriflora* Muehlenb. l. c.; Pursh l. c. *B. phyllostachya* et *B. florida* Miq. l. c. — Caulis herbaceus vel suffrutescens, erectus, 6-10 decim. altus, subsimplex vel parce ramosus, obtuse quadrangulus aut teretiusculus, glabratus vel apice (ramulisque) villosus, puberulus hispidulusve. Folia 3-15 centim. longa, 1-7 centim. lata, in caule opposita, in ramulis autem fere semper alterna, serraturis nunc triangularibus acutis obtusiusculisve nunc utrinque convexis et interdum subacuminatis parvis grossiusculis, acumine ipsius limbi lanceolato acuto integro, nervis basilaribus circiter ad trientem superiorem limbi productis, supra vix impressis subtus parum prominentibus, lamina exsiccata membranacea vel rigidula juniore ciliolata, petiolo gracili longitudine maxime variabili in typo sæpe vix centimetrali in var. β contra limbo haud raro æquilongopuberulo glabrato. Stipulæ lanceolatæ, 3-6 millim. longæ, in nervo dorsali pilosulæ, subsistentes. Glomeruli in ramulis lateralibus gracilibus inferne aphyllis apiceque foliiferis alterni vel in axillis superioribus ipsius caulis sessiles, parvi, unisexuales aut androgyni. Flores masc. mox decidui, perigonii 4-lobi lobis sub apice gibbosis pilisque quibusdam uncinatis instructis, pistilli rudimento ellipsoideo glaberrimo. Flores fem. fere sessiles, perigonio obovato vel elliptico deorsum cuneato compresso apice pilis uncinatis ciliato et magis minusve attenuato quam stigma longiuscule pilosum superne incurvum fere dimidio longiore, ore 4-dentato, fructifero valde compresso 1-1½ millim. longo latoque in parte superiore solum aut in toto ambitu late crasseque marginato faciebus impressis, achænium ellipticum arcte includente eoque plus minus adhærente. (v. s.)

β, *Drummondiana*, foliis supra asperimis subtus molliter pubescenti-tomentosis, cæteris ut in typo. — In Texas (Drummond, n. 267). *B. Drummondiana* Wedd. in Ann. Sc. nat. 4<sup>e</sup> sér. 1. p. 201; Bl. l. c. p. 214. — Spicæ breves, apice plus minus foliosæ.

γ, *littoralis*, foliis latius ovatis basi cordatis rotundatisve longe petiolatis supra plerumque lævibus, spicis apice foliatis. — In Jamaica, Peruvia (Matthews n. 2039), Brasiliæ prov. Minas Geraes (Pohl), Alagoas (Gardner, n. 1405), et Bahia (Salzmann), in America boreali rarior (Pal. Beauv.). *B. littoralis* Swartz, Prodr. p. 34, Fl. Ind. occ. 1. p. 281. *Procris littoralis* Poir. Encycl. 5. p. 628. *Urtica palustris* Juss.; Pers. Synops. 2. p. 552. *Duretia palustris* Gaudich.

δ, *brachystachys*, glomerulis breviter spicatis sive subcapitatis. — In ins. Antillis Martinica et Portorico (Plée), necnon in Brasiliæ prov. Piahy et Ceara (Gardner, n. 2004) et Rio Grande (St. Hilaire, n. 1727).

14. *B. ELONGATA* (Bl. l. c. p. 218), ramulis petiolisque et nervis subtus pube subtilissima subvelutinis, foliis ovalis v. ovato-ellipticis acuminatis basi rotundatis v. subcordatis dentato-serratis membranaceis supra pilis raris obsitis dein glabriusculis, spicis simplicibus solitariis v. fasciculatis foliis æquilongis v. sæpe longissimis, floribus in glomerulis subconfluentibus androgynis, perigonio fructifero ovoideo lenticulari-compresso hirtulo ore bidentato. § In horto Schœnbrunensi culta et forsitan ex America introducta. (Blume.)
15. *B. BULLATA* (Kunth in H. et B. Nov. gen. et sp. 2. p. 35), monoica, foliis alternis amplius elliptico-ovatis ovatisve, aliis attenuatis obtusiusculisque, aliis hisce alternantibus majoribus acuminatisque, omnibus basi rotundatis aut subcordatis minute serratis supra creberrime bullato-rugosis et asperulato-hispidis subtus lacunosus et pubescenti-hispidis, spicis folio æquilongis longioribusve simplicibus flexuosis aphyllis, perigonio fructifero ovato-lanceolato vix compresso piloso libero. § In temperatis regionis quitensis, prope Hambato et Lactacunga, altit. 2700-3000 metr. (Humb. et Bonpl.). — Wedd. Monogr. p. 379. Ramuli crassiusculi, hirtotomentosi, internodiis modice elongatis. Folia 3-15 centim. longa, 2-6 centim. lata, patula, ante insertionem petioli magis minusve emarginata, nervis basilaribus ad quartam partem superiorem limbi productis, coriacea, pagina superiore eximie verrucosa, verrucis subpyramidalis lobulatis apiceque sæpe piliferis, petioli 1-2 centimetralibus tomentosis. Stipulæ ovatæ, 6-10-millimetrales, acuminatæ, dorso pilosæ, deciduæ. Spicæ solitariæ, patulæ, 10-15 centim. longæ, rhachi crassiuscula tomentosa, glomerulis distantibus pisi magnitudine vel paulo majoribus bracteis ovatis semicentimetrum longis subsistentibusque suffultis. Flores masc. numerosissimi, pedicellati, mox decidui, pedicellis 1-3-millimetralibus inter bracteolas creberrimas persistentibus, perigonii 4-lobi segmentis acutis hispidis. Flores fem. (saltem in specim. suppet.) inter masculos rarissimi, perigonio stylo subæquilongo achænium sessile laxè includente (v. s.).

## § 2. *Stirpes gerontogæ.*

### A. — *Glomeruli axillares* (1). *Folia alterna aut opposita.*

16. *B. MALABARICA* (Wedd. Monogr. p. 355), monoica, foliis alternis subisomorphis in nodis vicinis nunc valde inæqualibus nunc fere eadem magnitudine, majoribus ellipticis oblongisve, minoribus potius ovatis, acuminatis basi rotundatis aut acutatis excepta basi crenulato-serratis planiusculis rugosulisve supra glabriusculis subtus breviter pubescentibus, glomerulis axillaribus, masculis plerumque inferioribus, perigonio fructifero late elliptico vel rotundato obtuse marginato pubescente. § In regionibus tropicis Indiæ orientalis frequens, altit. 0-1000 metr., in Zeylania, in prov. Carnatic, Concan, Silhet et in montibus Khasiæ et Himalayæ orientalis. — Urtica malabarica Wall. Cat. n. 4610. — Frutex 2-3-metralis, ramulis teretiusculis vel apice obtusè quadrangularibus villosulis basi sæpe aphyllis, internodiis longitudine variantibus sæpe vix centimetralibus. Folia 7-20 centim. longa, 3-12 centim. lata, in ipsa petioli insertionem haud raro emarginata, acumine angusto ut limbus reliquus serrulato vel subintegro, dentibus parvis obtusissimis vel acutis nonnunquam fere obsoletis, nervis basilaribus tenuibus trientem superiorem folii attingentibus, lamina exsiccata submembranacea sæpiusque fusco virente, petiolo gracili longitudine admodum variabili in foliis quibusdam dimidium limbum æquante s. fere decimetrati in aliis (minoribus alternisque) vix centimetrali villosulo vel demum glabrato. Stipulæ parvæ, extus puberulæ, mox deciduæ. Glomeruli pisiformes, diametro 3-8-millimetrali, densiflori, floribus plerisque subsessilibus, masculis (inferioribus) et femineis fere semper ut videtur in eodem ramo, florum masc. perigonio lobis acutiusculis prope apicem pubescentibus, perigonio fructifero 1-1½ millim. longo quam stylus multo brevior apice in collum brevissimum acuminato achænium ovatum submillimetrale ferrugineum arcte includente. (v. s.)

[1.] Vid. etiam, inter species glomerulis spicatis dictas, *B. Zollingerianam* et *B. acuminatam*.

- β, *depauperata*, fruticulosa, ramulis petiolis foliisque subtus scabriusculè pubescentibus, foliis longiusculè petiolatis supra basim grossiusculè serrato-dentatis (Miquel). ♂ In sylvulis humidis ins. Javæ, prope Tjikoya (Zollinger. n. 977). — *B. depauperata* Wedd. in Ann. Sc. nat. 4<sup>e</sup> sér. 1. p. 202; Bl. l. c. p. 199. *B. glomerulifera* Miq. in Zoll. System. Verzeichn. p. 104. (v. s.)
17. *B. HETEROPHYLLA* (Wedd. Monogr. p. 351, non Bl.), monoica, foliis alternis dimorphis, aliis majoribus lanceolatis inæquilateralibus longe et angustè acuminatis basi obtusatis subemarginatis petiolatisque, aliis (multo minoribus) oblique ovato-cordatis sessilibus, omnibus acute serratis supra exasperato-hispidis subtus sparsim hispido-pubescentibus, glomerulis axillaribus androgynis, perigonio fructifero compresso marginato apice attenuato achænio non adhærente. ♂ In ins. Manilla Philippinarum (Cuming, n. 1561). — *B. Cumingiana* Mus. bot. lugd. bat. 2. p. 199. — Ramuli graciles, subtiliter pubescentes, apice canescentes, internodiis brevibus. Folia minora bicentimetralia; majora 10-12 centim. longa, 2½-3½ centim. lata, ima basi nonnihil angustata, integraque, trinervia, serraturis reliqui limbi triangularibus margine exteriorè convexis sæpissimeque mucronulatis superioribus sæpe majoribus, nervis basilaribus hinc s. in latere latiore limbi vix usque ad mediætam hujusce productis at illic basim acuminis linearis et bi-tri-centimétralis fere attingentibus pilosulis, lamina exsiccata membranacea plana asperima, petiolo subcentimétrali gracili pubescente. Stipulæ lanceolato-subulatæ, extus pilosæ, deciduæ. Glomeruli magnitudine pisi; floribus masc. quam feminei numerosioribus, perigonii 4-partiti segmentis extus strigosus infra apicem mucronatis, mucronibus majusculis in alabastro conniventibus. Perigonium fem. tubulosum, ore acute bidentato, ad margines superne ciliatum, stigmate quam perigonium longiore persistente, fructiferum elongato-ellipsoideum lenticulari-compressum. — *B. ramifloram* habitu nonnihil refert (v. s.).
18. *B. BLUMEI*, dioica, foliis alternis, dissimilibus inæquilateralibus, aliis oblongo-vel ovato-lanceolatis acuminatis basi rotundatis petiolatis, aliis alternis multo minoribus late ovatis rotundatisve acutis basi oblique cordatis subsessilibus, omnibus dentato-vel crenato-serratis supra rugosis sparsimque pilosis et asperis subtus pubescenti-tomentosis nervis pilosis. ♂ In ins. Philippinis (Cuming, n. 731). — *B. heterophylla* Bl. Mus. bot. lugd. bat. 2. p. 204, non Wedd. *Margarocarpus heterophyllus* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 204 — Ramuli petiolique villosi. Stipulæ persistentes. Folia majora 8-14 centim. longa, 3¼-5½ centim. lata, petiolo 6-12-millimétrali. Flores masc. (solum visi) bracteis majusculis ovatis vel subrotundis involucriati, perigonii segmentis lanceolatis subulato-acuminatis dorso supra basin singulis appendice elongata recta lineari acuminata membranacea pilosa præditis (Blume). — Habitus præcedentis a qua tamen prima facie distinguitur foliis latioribus admodum rugosis necnon stipulis persistentibus et ramis villosis. (v. s. absque fl.)
19. *B. HELFERII* (Bl. Mus. bot. lugd. bat. 2. p. 201), monoica, foliis alternis subisomorphis cordato-ovatis acuminatis argute dentatis, petiolis elongatis in eodem pare inæquilongis, limbo plano supra sparsim piloso nervis subtus hirtulis, glomerulis axillaribus confluentibus unisexualibus, perigonio fem. tubuloso parum ventricoso ore bidentato. ♂ In Indiæ orientalis provincia Tenasserim (Helfer). — Ramuli apice adpresse puberuli et mox glabrati, internodiis elongatis. Folia ejusdem nodi nonnihil inæquimagna, 6-12 centim. longa, 4-6½ centim. lata, basi cordata vel subtruncata integraque, dentibus reliqui limbi triangularibus vix incurvis, acumine angusto l-sesqui-centimétrali basi serrato, nervis basilaribus ultra medium limbum in margine desinentibus, nervisque secundariis et venulis pubescenti-pilosis, lamina exsiccata membranacea læviscula, petiolis 4-8-centimétralibus gracilibus puberulis demumque glaberrimis. Stipulæ lanceolato-subulatæ, subpersistentes. Glomeruli fem. (solum visi) magnitudine pisi, floribus congestis, perigonio circiter millimétrali superne parce hirtulo stigmate recto circiter æquilongo.
20. *B. DIDYMOGYNE*, monoica, foliis alternis isomorphis ovatis acuminatis basi obtusis rotundatis longiusculè petiolatis supra quartam vel dimidiam partem inferiorem crenato-serratis præsertim supra sparse pilosulis, glomerulis axil-

laribus androgynis, perigonis fem. plerisque ovaria duo foventibus usque plus minus adnatis.  $\frac{5}{3}$  In Indiæ orientalis districtu Moulmein (Parish). — *Didymogyne boehmerioides* Wedd. ined. — Caulis herbaceus vel inferne plus minus lignescens, in specimine unico suppetente 5 decim. altus, inferne longe denudatus et crassitudine pennæ anatinæ apice parce ramoso et vix puberulo cæterum glaberrimus. Ramuli graciles, patentes, angulati, puberuli vel glabrati, internodiis plus minus elongatis. Folia 6-10 centim. longa, 3-5 centim. lata, serraturis æqualibus inæqualibusve, numero harumce in duobus lateribus limbi diverso, acumine lineari-lanceolato centimetrali, nervis basilaribus in margine ultra medium limbi desinentibus costaque et venulis subtus pilosulis glabratissive, lamina exsiccata membranacea lævi, petiolis 2-5 centim. longis patentibus glabriusculis. Stipulæ lanceolatæ, acuminatæ, subsistentes. Glomeruli parvi, floribus sessilibus congestis, masculorum perigonio 3-4-partito pubescente, segmentis sub apice mucronatis; perigonio fem. primum lanceolato dein oblongo v. elliptico compresso superne angustato pilisque rectis et hamatis hirtulo ore bidentato, stigmatibus linis parallelis approximatisque perigonio circiter æquilongis colloque hujusce basi constrictis. — Jam prope fui ut genus *Didymogynen* olim a me pro specie *Droguetiae* erratum conditum ad hanc stirpem recipiendam revocarem, sed indoles floris fem. adhuc aliquanto dubia videtur et fructus nullos maturos examinare licuit. Revera perigonium et ovaria speciminis indagati fere omnino accreta occurrunt, at vera natura forsitan propter compressionem mutata est. (v. s. in h. Kew.)

21. *B. MONTICOLA* (Bl. Mus. bot. lugd. bat. 2. p. 227), monoica, foliis alternatim magnitudine et petiolorum longitudine dissimilibus ovatis vel ovato-ellipticis acuminatis basi rotundatis infra medium ad apicem crenato-serratis membranaceis supra glabriusculis subtus petiolisque et ramulis tomentosis, glomerulis axillaribus et lateralibus androgynis, perigonio fructifero ellipsoideo-compresso pubescente superne attenuato quadridentato (Blume).  $\frac{5}{3}$  In sylvis Myrapi insulæ Sumatræ (Korthals). — Miquel, Fl. Ind. bat. 1 (2). p. 251.

22. *B. PARVIFOLIA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. 203; Monogr. p. 359), dioica?, foliis parvis oppositis isomorphis ovatis acuminatis basi acutis rotundatisve acute serratis planis supra hispido-pilosis subtus subtiliter pubescentibus nervis pilosulis, glomerulis axillaribus, perigonio fructifero ovato hirtello haud marginato.  $\frac{5}{3}$  In graminosis humidis ad Waliran, in ins. Java (Zollinger, n. 2185). — Miq. Fl. ind. bat. 1 (2). p. 250. — Caulis debilis, elongatus, prostratus, parce ramosus, ramis filiformibus ascendentibus pubescenti-pilosis. Folia vix sesquicentimetralia, dentibus acutis, acumine brevi, nervis basilaribus medium limbum circiter attingentibus subtus prominentibus, lamina exsiccata membranacea, petiolis in eodem jugo fere æquilongis bimillimetralibus pilosis. Stipulæ pro foliorum magnitudine majusculæ, ovatæ, petiolis circiter æquilongæ, nervo dorsali pilosulæ. Glomeruli ut videtur pauciflori, omnes in speciminibus visis feminei, floribus sessilibus, perigonii ovati quam stylus vix brevioris ore latiusculo 4-dentato, fructifero sesquimillimetrali achænium late ellipticum atrum laxiuscule includente. (v. s.)

23. *B. COMOSA*, monoica, foliis alternis oppositisque fere isomorphis lanceolato-ovatis anguste acuminatis basi acutis obtusatisve etiamque rotundatis æqualiter acuteque serratis planis rugosulisve utrinque ramulisque hirtis, glomerulis axillaribus spicatisque (spicis apice comosis), perigonio fructifero lanceolato v. elliptico v. obovato compresso deorsum cuneato hispido.  $\frac{5-3}{3}$  In Indiæ orientalis regione tropica montium Khasiæ et Himalayæ prov. Sikkim et Nepalia alit. 900-1500 metr. (J. D. Hooker), Mishmi (Griffith), Pegu et Rangoon (Mac Clelland), necnon in ins. Java (de Vriese). *B. diffusa* Wedd. Monogr. p. 356. *Urtica comosa* Hamilt. in Wall. Cat. — Rami graciles, sæpe diffusi, teretiusculi, raro glabrati. Folia ramorum sæpe opposita, ramulorum vero (præsertim floriferorum) fere semper alterna, vulgo lanceolato-ovata sed interdum (etiam in eodem specimine) ovata, alterna vulgo majora, in ramulis floriferis 2-6-centimetralia, in sterilibus vero sæpe duplo majora, nempe 10-12 centim. longa et circiter 4-6 centim. lata, acumine lineari nunc integerrimo nunc ut limbus reliquus serrato, nervis basilaribus usque ad trientem superiorem laminæ excurrentibus, pagina superiore plana vel rugosula asperula

inter pilos molles et patentes (quum adsunt) pilos alios adpressos sæpe exhibente, pagina inferiore in nervis præsertim pilosa, lamina exsiccata membranacea vel rigidula, petiolo brevi vel 1-2-centimetro hirsuto vel strigoso. Stipulæ lineari-lanceolatae, 3-5 millim. longæ, dorso pilosæ. Glomeruli utriusque sexus conformes, multiflori, floribus sessilibus v. pedicellatis, perigonii masc. lobis superne strigoso-hispidis. Glomeruli fructiferi discreti vel (præsertim aphylli) magis minusve confluentes, pisiformes, sæpius valde multiflori, perigonis vix millimetralibus compressis diversiformibus anguste acutiusculeque marginatis achænio arcte adpressis adhærentibusque, ore bi-quadri-dentato. — Hujus forma oppositifolia est *B. sidæfolia* Wedd. in *Ann. sc. nat. l. c.* (v. s.)

β, *strigosa*, foliis rugosis supra vel utrinque (ramulisque) strigosis.

γ, *canescens*, foliis majoribus quam in typo longiusque petiolatis plerisque ovatis (dentibus subacuminatis) planis supra ramulisque strigosis subtus breviter asperato-pubescentibus canescentibusque.

δ, *neglecta*, ramulis petiolis foliisque subtus tomentosis, foliis basi obtusis v. subrotundatis serrulatis rugosis supra glabris nervis subtus incano-hirtulis, glomerulis dioicis (Blume). § In Java occidentali. — *B. neglecta* Bl. Mus. bot. lugd. bat. 2. p. 200; Miq. Fl. Ind. bat. 1 (2), p. 249.

24. *B. AUSTRALIS* (Endl. Prodr. fl. Norfolk, p. 38; Icon. gen. pl. t. 86), monoica, foliis alternis isomorphis ovatis sensim acuminatis basi cordatis argute serratis supra retrorsum scabris subtus subtilissime cinereo-tomentosis, glomerulis axillaribus androgynis, perigonio fructifero elliptico compresso membranaceo-marginato pilis rigidis hispido. § In ins. Norfolk (Ferd. Bauer). — Wedd. Monogr. p. 357; Bl. l. c. p. 205. *Procris splendens* Lindley, Veget. Kingd. ed. 4. f. 175. — Ramuli crassitie pennæ anserinæ, pilis brevissimis rigidis cinerascensibus scabri, internodiis brevibus. Folia patula, 6-15 centum. longa, 4-9 centum. lata, basi late rotundata sinuque angusto cordata, serraturis margine exteriori convexiusculis, nervis basilaribus ad trientem superiorem limbi circiter pertingentibus hispidulis, petiolis foliorum superiorum tricen-timetralibus brevioribus inferiorum multo longioribus crassiusculis incano-tomentosis. Glomeruli nuclei cerasini magnitudine, floribus masc. et fem. intermixtis, perigonii masc. 4-partiti segmentis oblongis acutis pilis erectis rigidiusculisque hispidis, femineorumve brevissime pedicellatorum perigonio inflato-tubuloso ore brevissime 4-dentato quam stylus villosus persistens pilis longiore achæniisque crustaceum læve includente. — Specimen nulum vidi. Affinis videtur *B. excelsæ* insulæ Juan Fernandez.

#### B. — Glomeruli spicati vel cymoso-paniculati.

\* Folia alterna.

25. *B. NIVEA* (Hook. et Arn. Bot. Voy. Beech. p. 214), dioica aut monoica?, foliis amplis alternis isomorphis æquilateralibus late ovatis vel elliptico-rotundatis acuminatis basi cordatis sæpiusve juxta petiolum breviter cuneatis rarius basi attenuatis truncatisve crenato-serratis (dentibus mucronulatis) planis supra scabro-hispidis subtus (nervis exceptis) niveo-tomentosis, stipulis liberis, glomerulis laxè cymoso-paniculatis, perigonio fructifero elliptico aut oblongo-compresso piloso. 2-5 In Asia tropica et præsertim temperata et subtemperata, scilicet regione sinensi, fere tota, indigena et culta in India orientali, in ins. Philippinis, Mariannis et cæt. — Hook. Journ. of Bot. ann. 1851. t. 8; Miq. Pl. Jungh. 1. p. 33; Fl. Ind. bat. 1 (2). p. 253; Wedd. Monogr. p. 380. t. 11. f. 10-17; Bl. Mus. bot. lugd. bat. 2. p. 210. *Urtica nivea* Linn. Spec. 1398; Jacq. Hort. Vindob. t. 166. *Procris nivea* Gaudich. Bot. Voy. Uran. p. 499. — Cæspitosa, caulis metralibus simplicibus v. ramosis crassitudine pennæ ramulisque olivaceis aut gracilioribus obtuse angulatis superne ramulisque molliter hirtis canescentibusque inferne glabris, nodis magis minusve distantibus. Folia patula, 6-30 centum. longa, 4-18 centum. lata, sensim vel subabrupte acuminata (acumine ut plurimum lineari et acuto), dentibus sæpius utroque margine convexis in dimidia parte superiorè

limbi majoribus in ejusdem basi sensim imminutis ac juxta petiolum plerumque nullis, nervis basilaribus ad quartam partem superiorem limbi excurrentibus et in margine desinentibus subtus sicut costa hirsutis, lamina exsiccata membranacea, petiolo longitudine maxime variabili limbo interdum subæquilongo at vulgo fere dimidio brevioris magis minusve hirsuto. Stipulæ liberræ, lanceolatæ, subulato-acuminatæ, 8-12-millimetrales, ciliatæ, nervo dorsali pilosæ. Inflorescentiæ in singulis axillis solitariæ vel geminatæ, amplitudine quam maxime variantes, modo petiolis breviores, modo limbum ipsum superantes, pedunculo brevi vel longiusculo suffultæ, ramosissimæ, pedunculo ramulisque gracillimis hirtis; glomerulis discretis; femineis initio semen sinapinum et mox cannabinum magnitudine æmulantibus aut paulo superantibus, florum sessilium perigonio elliptico (stylo hinc villosillo illic sparsim piloso subæquilongo) pilis rigidis undique hirtis ore parum ampliato 4-dentato; perigonium fructiferum vix semimillimetrale, angustissime marginatum pilis superioribus dentes fere occultantibus, achænio nunc discreto nunc contra cum perigonio arcte cohærente. — Fibræ corticales B. niveæ, robore et pulchritudine nullis secundæ, per Asiam orientalem præsertim, in usu hodierno sunt ad textilia conficienda. Caules stirpis in Europa cultæ frequenter ob hyemem annuæ sunt. Planta vulgo dicitur Chuma s. Tchou-ma apud Sinos; in Java et Sumatra : Rami et Klooi s. Caloi; in Anglia : China Grass. (v. s. et v. cult.)

β, *candicans*, foliis concoloribus vel imprimis junioribus subtus albicantitomentosis. δ Cum præcedente culta, in ins. Java, Sumatra et Borneo frequentior. — B. tenacissima Gaudich. Bot. Voy. Uran. p. 500; Bl. in Ind. By. 1. p. 483; Mus. bot. lugd. bat. 2. p. 211. t. 56. b. B. candicans Hassk. Cat. Bog. p. 79 et Pl. jav. rar. p. 203 in Obs. Rami majus Rumph. Hort. Amb. 5. p. 214. t. 79. *Urtica candicans* Burm. ? Fl. Ind. p. 297. Bl. Bijdr. p. 503. U. tenacissima Roxb. Fl. Ind. 3. p. 590; Wight, Icon. pl. Ind. 2. t. 688. U. utilis hort. — Caules sæpe robustiores quam in typo; folia majora et longius petiolata. (v. s. et v. cult.)

26. B. COMPACTA (Bl. Mus. bot. Lugd. bat. p. 210), dioica?, foliis alternis ovatis acuminatissimis basi rotundatis v. truncatis rarissime acutiusculis grosse dentato-serratis membranaceis supra asperis subtus incano-tomentosis, glomerulis fem. in axillis sessilibus aut in spicis brevissimis conglomeratis, perigonio fructifero turbinato lenticulari-compresso inferne anguste marginato puberulo ore quadridentato. ♀ In Java occidentali. — Ramuli petiolique 3½-8 centim. longi) caescenti-hirtuli. Folia 5-14 centim. longa, 5-10 centim. lata. Flores fem. dense fasciculato-glomerati, brevissime pedicellati, pedicellis bracteolis minutissimis subulatis ferrugineis circumvallatis. Perigonium ventricosum-tubulosum, dentibus oris ovatis sæpe inæqualibus. Ovarium liberum. Stigma apice sæpe incurvum, ad apicem unilateraliter villosulum. Achænium ovoideum, sublenticulari-compressum, perigonio membranaceo obovoideo v. turbinato inclusum (Blume). — Ex hac stirpe, mihi prorsus incognita, et e B. nivea conficitur in opere Blumeano supra laudato sectio propria generis sub nomine *Tilocnide*.

27. B. RUGULOSA (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 200; Monogr. p. 373), dioica?, foliis alternis ovatis vel elliptico-ovatis apice attenuatis subacuminatisve basi rotundatis vel obtusis crenato-serrulatis planis aut rugulosis supra glabris subtus brevissime velutino-pubentibus, stipulis ejusque folii in unam axillarem apice bidentatam coalitis, spicis fem. quam folia multo brevioribus pendulis simplicibus, perigonio fructifero elliptico valde compresso submembranaceo margine longe ciliato. δ In regionibus tropicis Himalayæ centralis, altit. 1200 metr. prov. Bhotan (Griffith), Nepalia (Wallich), Kumaon (Thomson) et prope Nahau (Jacquemont). — Bl. Mus. bot. lugd. bat. 2. p. 212. *Urtica rugulosa* Wall. Cat. n. 4597. U. venosa ejusd. l. c. n. 4602 B. — Arbuscula 3-4-metralis, ramis suberectis teretiusculis siccitate rugosulis ferrugineisque, junioribus quadrangulis et apice præsertim puberulis. Folia 6-12 centim. longa, 2½-4½ centim. lata, obtusiuscula, basi plerumque integra, ad summum apicem usque denticulata, denticulis inæqualibus; nervis basilaribus arcuatis limbo ipsi æquilongis, primariis cæteris brevibus obliquis arcuatimque anastomosantibus, supra impressis subtus pube tenui vestitis;

lamina exsiccata coriacea discolore; petiolo sesqui-tri-centimetro initiali puberulo dein glabrato. Stipulæ ovatæ, 5-6-millimetricæ, dorso puberulæ. Spicæ femineæ (solum visæ) in singulis axillis 1-3, rhachi pubescente, receptaculo s. axi glomerulorum pilis albis dense hirsuto, perigonio longiuscule attenuato quam stylus paulo longiore bifido, fructiferæ 4-6 centim. longæ, glomerulis approximatis pisum magnitudine fere æmulantibus. Achænium ellipticum, utrinque attenuatum, pedicellatum, ventricosum, nitidum, perigonio  $1\frac{1}{2}$ -2-millimetrici arcte inclusum. (v. s.)

\*\* Folia opposita.

28. B. BILOBA (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 199; Monogr. p. 373), monoica, foliis oppositis heteromorphis ellipticis basi acutis plerisque apice magis minusve profunde bifidis (lobis sensim acuminatis) cæterum fere in toto margine crenulatis rugosis supra hispidis subtus pubescenti-tomentosis, stipulis liberis, spicis unisexualibus vel androgynis erectis quam folia brevioribus, perigonio fem. elliptico compresso pubescenti-hirto libero.  $\zeta$  In Japonia sylvestris, nec non in hortis botanicis frequenter culta. — Miq. in Zolling. Syst. Verzeichn. 2, p. 100. B. bifida Bl. Mus. bot. lugd. bat. 2. p. 222. Urtica biloba Siebold, Kruidd. Naamlijst, p. 38 et hort. U. bifida Hassk. Cat. Hort. bogor p. 79. Splitgerbera japonica Miq. Comment. phyt. p. 139. t. 14. — Frutex parce ramosus, ramulis teretiusculis, novellis pubescentibus. — Folia patula, in eodem jugo fere æquimagna, 6-10 centim. longa, 3-6 centim. lata, lobis terminalibus plerumque duobus 5-30-millimetra longis nunc subæqualibus (limbo scilicet costaque simul bifurcis s. diremptis) nunc valde inæquimagnis altero nempe in dentem lateralem reducto vel subnullo, nervis basilaribus costæ vix non parallelis et fere ad basin loborum productis, venis secundariis valde obliquis subparallelisque prope marginem arcuatum anastomosantibus supra impressis subtus autem prominulis; laminæ exsiccata coriaceæ pagina superiore aspera, inferiore lacunoso-reticulata, petioli foliorum oppositorum inæquilongis  $\frac{1}{2}$ -2 centim. longis robustis hispido-pubescentibus. Stipulæ oblongo-lanceolatæ, 10-12-millimetricæ, basi lata ramo insertæ, nervo dorsali pubescente, cæterum glabrata vel puberulæ. Spicæ masc. in axillis inferioribus ramulorum nascentes, in speciminibus observatis quam folia paulo breviores, glomerulis discretis 6-10-floris, floribus sessilibus, perigonii lobis extus pubescentibus summo dorso mucrone brevi obtusiusculoque instructis, pistilli rudimento obovato-clavato basi brevissime lanato. Spicæ fem. ex axillis superioribus ortæ, masculis conformes, sed floribus numerosioribus, perigonio pilis rectis hirto quam stylus paulo brevior, fructifero achænium membranaceo-marginatum arcte obvestiente. — Folia rarissime inæqualiter tri-quinque-fida. (v. s. et v. cult.)

29. B. ZOLLINGERIANA (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 201; Monogr. p. 372), monoica, foliis oppositis alternisve late ovatis vel lanceolato-ovatis acuminatis basi rotundatis rarius (superioribus) acutatis crenato-serratis planis supra sparsim hispidulis demum glabratissimis nervis subtus pilosulis hispidulisque, stipulis liberis, floribus fem. in spicis paniculatas terminales masculisque (5-andris) in glomerulos axillares digestis, perigonio fructifero elliptico parum compresso breviter hispido.  $\zeta$  In ins. Java, ad rupes prov. Bandawosso (Zollinger, n. 2765). — B. diversiflora Miq. in Zolling. System. Verzeichn. p. 101. — Ramuli graciles, striato-sulcati, apice minute hispiduli, mox glaberrimi, epidermide siccitate fusco-rubente. Folia inferiora ramulorum opposita aut fere opposita, superiora alterna, in nodis singulis vel inter se proximis valde inæquimagna, 3-15 centim. longa,  $1\frac{1}{2}$ -8 centim. lata, acumine fere lineari integro aut basi serrato, dentibus reliqui limbi obtusiusculisve, nervis basilaribus dimidium limbum paulo superantibus, lamina exsiccata membranacea, petioli gracillimis valde inæquilongis 1-7-centimetricis glabris lævibusque. Stipulæ lanceolatæ, semicentimetricæ, in nervo dorsali pilosulæ. Glomeruli masc. in axillis inferioribus ramulorum solitarii, oppositi, laxè confluentes, magnitudine pisi majoris, floribus pedicellatis, perigonii 5-partiti lobis ovatis breviter acuminatis extus glabriusculis, pistilli rudimento globoso basi pilis lanatis crebriusculis stipato. Glomeruli fem. in



parte superiore ramulorum orti, parvi, alii axillares, alii et plerique interrupte spicati, spicis simplicibus paniculatis, perigonio elliptico-lanceolato vel obovato apice abrupte attenuato (ore bi-quadri-dentato) achænium arcte includente styloque gracili villosa paulo brevior. (v. s.)

30. *B. ACUMINATA* (non Pers.), monoica v. dioica?, foliis oppositis disparibus subinæquilateralis ovato-lanceolatis longe et anguste acuminatis basi rotundatis v. subcordatis toto margine crenulato-serrulatis rugosis supra sparsim hispidulis demumque glabratibus subtus pubescentibus, altero minore brevius petiolato sæpiusque magis cordato, glomerulis unisexualibus, masculis spicatis, femineis axillaribus pisiformibus, perigonio fructifero (immaturo) elliptico strigoso.  $\S$  In Nova Caledonia, ad ripas, prope Pueblo (Vieillard). Ramuli graciles, superne subangulati et appresse puberuli, mox glabrati, internodiis 1-3-centimetralibus. Folia majora 15-20 centim. longa,  $4\frac{1}{2}$ -6 $\frac{1}{2}$  centim. lata, acuminata falcato haud raro 5-centimetrali, nervis basilaribus usque ad basim huius productis, petiolo 2-3 centim. longo pubescente; minora 6-8-centimetralia, petiolo circiter centimetrali, lamina exsiccata rigidula. Stipulæ lanceolatæ, deciduæ. Spicæ masc. 4-8-centimetrales, graciles, erectæ, in singulis axillis solitariæ, breviter pedunculatæ, glomerulis approximatis parvis, uno majore ad basin pedunculi s. in ipsa axilla sessili, perigonii parce hispidi segmentis sub apice acuminato breviter mucronatis. Glomeruli fem. in ramo distincto vel ad basim spicarum masc. confluentes masculisque subæquimagni, perigonio oblongo hispido superne sensim attenuato, ore suboblique bidentato, stigmatibus initio basi glabra denudato sed mox perigonio accreto usque ad partem superiorem inflexam vaginato. — Species sedis subdubiæ ob defectum fructus maturi. Stigma Cypholophi, sed habitus pro floribus spicatis cæt. omnino Boehmeriæ. (v. s. in hb. Mus. par.)

31. *B. MACROPHYLLA* (Don, Prodr. fl. Nepa! p. 60, non Sieb. et Zucc.), monoica, foliis oppositis lanceolatis longe attenuato-acuminatis basi acutis serrulatis rugosis supra sparsim hispidis demumque glabratibus subtus pubescentibus, stipulis liberis, spicis elongatis pendulis aphyllis unisexualibus simplicibusque aut basi breviter ramulosis (ramulos masc. plerumque gerentibus), perigonio fructifero anguste obovato-cuneato compresso margine ciliato apice in collum brevem 2-4-dentatum abruptissime angustato.  $\S$  In regione tropica montium Khasiæ et Himalayæ orientalis, Assam et Kumaon, altit. 0-1200 metr. — Wedd. Monogr. p. 375. t. 11, f. 1-9; Bl. l. c. p. 217. *B. penduliflora* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 199. *Urtica penduliflora* Wall. Cat. n. 4595, A, B, C. — Ramuli teretiusculi, parce pubescentes, exsiccati obtuse angulati. Folia 1-3 decim. longa,  $1\frac{1}{2}$ -4 centim. lata, patula, in eodem jugo vix inæquimagna, æquilateralis, basi sensim attenuata, ad apicem usque serrulata, nervis basilaribus margine parallelis usque ad trientem superiorem limbi vel ultra productis dein cum nervis lateralibus arcuatim anastomosantibus, lamina exsiccata rigida nervis supra impressis subtus prominulis, petiolis in eodem jugo parum inæquilongis 1-3-centimetralibus sparsim puberulis. Stipulæ anguste lanceolatæ, 6-8 millim. longæ, acutissimæ, in nervo medio puberulæ. Spicæ sæpissime unisexuales; masculæ foliis breviores, graciles, glomerulis subconfluentibus, floribus subsessilibus, perigonii 4-partiti lobis ovatis acutis extus puberulis, pistilli rudimento subgloboso glabro, alabastro brevissime acuminato. Spicæ fem. foliis demum longiores brevioresve, interrupte glomerulatæ, rhachi pubescente, glomerulis floriferis seminis cannabini circiter magnitudine discretis, fructiferis pisiformibus approximatis, perigonio membranaceo demum valde compresso  $1\frac{1}{2}$ -2 $\frac{1}{2}$  millim. longo, achænio lanceolato utrinque attenuato pedicellato compresso crassiuscule subfungosæque marginato perigonio laxo incluso. (v. s.)

32. *B. STIPULARIS* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 200; Monogr. p. 376), monoica vel dioica?, foliis oppositis amplis late ovatis ellipticis rotundatisve acutis vel breviter acuminatis subcordatis serratis utrinque sparsim pilosis hispidulis vel subtus præsertim in nervis pubescentibus interdum rugosis, stipulis (majusculis) cujusque folii in unam axillarem integram binervem coadunatis, spicis unisexualibus androgynisve longe paniculato-ramosis quam folia brevioribus erecto-patulis, perigonio fructifero elliptico compresso mem-

branceo-marginato superne hispidulo.  $\frac{3}{4}$  In insulis Sandwicensibus (Beechey, Remy, etc.), Madagascaria (Petit-Thouars), Borbonia (Gaudichaud, Boivin) et Mauritio. — *B. amplissima* Bl. l. c. p. 219. *Urtica grandis* Hook. et Arn. Bot. Beech. Voy. p. 95. — Ramuli robusti, obtuse quadranguli, adpresse hispidulo vel strigilloso-pubescentes. Folia 1-2  $\frac{1}{2}$  decim. longa, 8-20 centim. lata, basi late rotundata et juxta petiolum emarginata, toto margine acute serrata, serraturis margine exteriori sæpe convexusculis, nervis basilaribus supra costaque sæpe pilosulis subtus prominulis ad partem quartam superiorem limbi circiter productis, venulis ante marginem arcuatim anastomosantibus. Lamina exsiccata rigida vix asperula, petiolis 2-8 centim. longis robustis in eodem jugo persæpe valde inæqualibus magis minusve pubentibus. Stipulæ intraaxillares ovatæ, acuminatæ, 1  $\frac{1}{2}$ -3 centim. longæ, obtuse bicarinatæ, margine glabræ, dorso et præsertim carenis adpresse subincano-pubescentes vel pilosæ, subpersistentes. Inflorescentiæ patulo-ramosæ, late ovatæ, in singulis axillis solitariæ, pedunculo subcentimetro et rhachi strigoso-pubescentibus, glomerulis parvis plerumque discretis bractea late ovata brevi suffultis, floribus masc. brevissime pedicellatis, perigonii lobis 4 ovatis extus breviter strigosos apice subacuminatis, pistilli rudimento clavato basi pilosulo, perigonio fem. elliptico apice hispido (ore 4-dentato) quam stylus superne longiuscule villosus brevior, fructifero sesquimillimetrali faciebus ante marginem membranaceum convexusculis achænio sublenticulari pedicellato arcte adhærente. Stipulis concretis a sequente bene distincta. (v. s.)

33. *B. PLATYPHYLLA* (Don ex Hamilt. Prodr. fl. Nepal. p. 60, non Sieb. et Zucc.), monoca vel dioica? foliis oppositis vel superioribus haud raro alternis amplius fere isomorphis late ovatis in eodem jugo æqualibus vel inæquimagnis acuminatis basi rotundatis cordatisve margine toto (etiam acumine) serratis vel crenato-serratis planis aut magis minusve rugosis exsiccatis rigidis supra sparsim hispidis subtus hispidulis, stipulis liberis, spicis unisexualibus androgynisve simplicibus aut (masculis præsertim) simpliciter ramosis fere semper aphyllis, femineis folia æquantibus superantibusve erecto-patulis glomerulis pilosioribus discretis, perigonio fructifero elliptico obovato vel rotundato plus minus compresso aut inæqualiter angulato sæpius obtuse marginato pulisque rectis vel quibusdam uncinatis hispido.  $\frac{3}{4}$  In Indiæ orientalis regione tropica et subtropica montium Khasiæ et Himalayæ totius, altit. 0-1500 metr., Concan, Zeylantia et Bahar. — Wedd. Monogr. p. 364. *B. macrostachya* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 199; Bl. Mus. bot. lugd. bat. 2. p. 224. *Urtica macrostachya* Wall. Cat. n. 4582. A. B. C. — Caulis 1-3-metralis, magis minus ramosus, ramis junioribus obsolete angulatis nunc pube obductis nunc fere glabris. Folia 1-3-decimetralia, nervis basilaribus parum arcuatis ad trientem superiorem limbi nec ad apicem ejusdem productis, limbo supra plano aut venis et venulis plus minus impressis, petiolis ejusdem paris inæquilongis pube eadem ac ramulorum. Stipulæ triangulari-lanceolatæ, 6-12 millim. longæ, sensim acuminatæ, extus in nervo dorsali præsertim pilosulæ, vix persistentes. Spicæ masc. interdum fasciculatæ, rhachi puberula vel villosa, glomerulis quam feminei vulgo minoribus, illis utriusque sexus sæpius confluentibus rarius distantibus; floribus masc. brevissime pedicellatis, lobis perigonii sub apice acuto sæpissime gibbosulis, pistilli rudimento obovato pilis paucis stipato. Glomeruli fem. plerique pisiformes, centimetrum sæpe diametro metientes, floribus breviter pedicellatis, perigonio primum fere lanceolato et subæqualiter hispido, fructifero heteromorpho plus minus compresso et in dimidio inferiore glabrato apice breviter attenuato seu rostrato ore quadri-dentato faciebus planiusculis achænio arcte adhærente. — Stirps quam maxime polymorpha, ideo, velut *Urtica dioica*, cui optime comparatur, in species permultas ab auctoribus consecta. Hasce in Monographiâ mea pro varietatibus tantum habui, et hodie, re iterato perpensa, in eadem sententiâ hæreo. Omnes etenim formæ infra enumeratæ non sunt meo sensu nisi annuli ejusdem catenæ. Formæ extremæ quidem sat absimiles videntur, sed has inter gradibus fere imperceptis transitur; et nihil dubito quin numerus talium varietatum fere usque ad infinitum augescere possit, quoniam vix ullum sit specimen quod caractere quodam magis minusve notabili enitescat. (v. s.)

$\beta$ , *virgata*, foliis (8-10 centim. longis) sæpius late ovatis acuminatis basi

rotundatis vel subcordatis toto margine crenato-serratis planis aut rugosulis membranaceis supra sparsim puberulis et mox glabris nitidisque subtus in nervis puberulis aut interdum hirtis, spicis sæpius ramosis folia parum superantibus patulis, glomerulis pisiformibus. § In ins. Taiti, Fiji, Nov. Hebrid. etc. — *Urtica virgata* G. Forst. Prodr. n. 345. B. *virgata* Guillem. in Ann. sc. nat. 2<sup>e</sup> sér. 7. p. 182. B. interrupta ejusd. l. c.; Bl. l. c. p. 218. B. *Taitensis* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 200. (v. s.)

- γ, *Sumatrana*, foliis (8-12 centim. longis) nonnunquam ternis rotundato-ovatis vel subdeltoideis grosse dentato-serratis supra hispidulis subtus sparsim strigoso-pilosulis, spicis simplicibus sæpiusve ramosis conferte glomeruliferis folio multo brevioribus. § In ins. Sumatra — B. *Sumatrana* Miq. Pl. Jungh 1. p. 32; Bl. l. c. p. 219. (v. s.)

δ, *macrostachya*, foliis (6-12 centim. longis) ovatis vel interdum fere rotundatis sensim vel subabrupte acuminatis toto margine crenato-dentatis serratisve aut basi acumineque integris planis aut vix rugosulis membranaceis utrinque sparsim et subadpresse hispidis pubescentibusve rarius glabratiss, spicis fem. plerumque simplicibus longissimisve pendulis, glomerulis quam in typo minoribus (nempe 4-5 millim. tantum latis) distantioribusque. § Passim in Himalaya, in montibus Nilghiri, in ins. Java et Zeylantia (Thwaites), Mascarenis, præsertim Mauritio, necnon ad Fernando Po Africæ occidentalis (Mann). — *Urtica caudata* Poir. Encycl. 4. p. 640. B. *caudata* ejusd. in Bonpl. Pl. rar. Malm. t. 15; Dict. sc. nat. t. 283, non Sw. nec Vahl. B. *mauritiana* Wedd. l. c. p. 200; Bl. l. c. p. 216. B. *Wightiana* ejusd. l. c. p. 217. Splitgerbera *macrostachya* Wight, Icon. Pl. Ind. 6. t. 1977. — Occurrit forma (subvar. *elliptica*) hujus varietatis foliis ellipticis abrupte breviterque acuminatis quam olim (l. c. p. 200) sub nomine B. *elliptica* descripti. Altera (subvar. *phylllostachya*) spicis apice comosis gaudet. Duas sequentes refert beatus Blumius: 1<sup>o</sup> (subvar. *rugosa*), foliis limbo rugoso et valde aspero toto margine acute serrato, ex India orientali prov. Garhwal, ad Masuri (Hugel, n. 49); hæc est B. *masuriensis* Bl. l. c. p. 216. 2<sup>o</sup> (subvar. *molliuscula*), foliis subtus molliter subvelutinis, ex ins. Mauritio (Sieber, n. 375) et Borbonia. (v. s.)

- ε, *microcarpa*, foliis quam in typo multo minoribus ovatis vel lanceolato-ovatis lanceolatisve sensim acuminatis basi rotundatis obtusatisve toto margine serratis aut serrulatis sæpe rugosulis magis minusve asperato-hispidis subtus (in nervis præsertim) pubescentibus rigidis, spicis ramosis sæpe androgynis erecto-patulis quam folia brevioribus, glomerulis approximatis. § In insulis moluccanis et oceanicis. — B. *microcarpa* Wedd. l. c. p. 201; Miq. Fl. Ind. bat. 1 (2). p. 254. (v. s.)

ζ, *Zeylantica*, foliis elliptico-ovatis ellipticisve anguste acuminatis basi rotundatis vel subcordatis toto margine crenato-serratis (crenis brevissime mucronato-acuminatis) planiusculis supra sparsim hispidis subtus in nervis pilosulis membranaceis, spicis simplicibus erecto-patulis unisexualibus androgynisve quam folia vulgo brevioribus, glomerulis fructiferis fere confluentibus, perigonis heteromorphis angulatis vel irregulariter alatis. § In Zeylantia (Thwaites, n. 2197). (v. s.)

- η, *Celebica*, foliis (8-20 centim. longis) ovato-vel elliptico-oblongis oblongisve basis obtusis acutiusculisve inæqualiter dentato-serrulatis planis aut vix rugosulis membranaceis supra sparsim strigoso-hispidis subtus pubescenti-hirtulis, spicis simplicibus erecto-patulis quam folia brevioribus aut vix longioribus. § In ins. Celebes et Java. B. *platyphylla* var. *moluccana* Wedd. Monogr. p. 366, pro parte. *Urtica celebica* Bl. Bijdr. p. 492; Mus. bot. lugd. bat. 2. p. 217. (v. s.)

θ, *scabrella*, foliis (quam in typo minoribus) ut plurimum late ovatis rarius ellipticis obovatisve subabrupte acuminatis basi rotundatis vel obtusis acute dentatis sæpe rugosulis supraque asperato-hispidis rigidis, spicis subsimplicibus erectis quam folia brevioribus. § In Indiæ orientalis prov. Assam, Nepalia et Khasia, in montibus Nilghiri et in regione tropica

- Zeylanæ et Javæ. B. scabrella Gaudich. Bot. Voy. Uran. p. 499; Miq. Fl. Ind. bat. 1 (2) p. 252. B. ourantha Miq. Pl. jungh. 1. p. 33; Bl. Mus. bot. lug. bat. 2. p. 221. Urtica caudata Nic. Burm. Fl. Ind. p. 198; Bl. Bijdr. p. 492, non Poir. nec Vahl. U. scabrella Roxb. Fl. Ind. 3. p. 581; Wall. Cat. n. 4581; Wight, Icon. pl. Ind. 2. t. 691. U. uragera Steud. Nomencl. 2. p. 736. (v. s.)
- ι, *rugosissima*, foliis amplis (10-25 centim. longis) late ovatis vel subrotundis acuminatis basi sæpe subcordatis margine toto serratis coriaceis valde rugosis supra hispidis subtus in nervis præsertim pubescenti-hirtis, spicis subfasciculatis quam folia brevioribus patulis. § In Zeylanæ et Javæ occidentalis sylvis montanis et in Sumatra (Korthals). B. rugosissima Miq. Pl. Jungh. 1. p. 32; Fl. Ind. bat. 1 (2). p. 251; Bl. Mus. bot. lugd. bat. 2. p. 222. Urtica rugosissima Reinw. in Bl. Bijdr. p. 490. — Folia intervenio supra verrucoso sive concinne rugoso, rugis numerosissimis conicisque, pagina inferiore limbi lacunosa. — Pro loci natura, ait Blumius, admodum variat foliorum magnitudine, quæ modo textura satis crassa, rugosissima, subtus præsertim in nervis, sicuti in petiolis et ramulis, satis dense pilosa, modo textura membranacea minus rugosa, subtus pube parca et in nervis, sicuti in petiolis et ramulis, pilis appressis substrigosa. (v. s.)
- κ, *clidemioides*, characteribus plerisque præcedentis, sed foliis minoribus (5-15 centim. longis) sæpe ovatis vel oblongis et nonnihil disparibus valdeque inæquimagnis, spicis sæpe solitariis. § B. clidemioides Miq. Pl. Jungh. 1. p. 34; Fl. Ind. bot. 1 (2). p. 252. B. pilosiuscula Hassk. Cat. bogor. p. 79; Bl. Mus. bat. lugd. bot. 2. p. 223. Urtica pilosiuscula Bl. Bijdr. p. 491. — Proxime accedit ad var. *rugosissimam*. — Spicæ interdum apice comosæ. (v. s.)
- λ, *humilis*, omnibus partibus quam in præced. minoribus, foliis fere isomorphis at plerumque inæquimagnis late ovatis ellipticisve basi rotundatis vel subcordatis rugosis supra sparsim hispidis subtus in nervis pilosis aut densiuscule pubescentibus, spicis simplicibus quam folia brevioribus erectis. § In declivibus montium Javæ ad Medim et Ugaram (Junghuhn). B. humilis Miq. Pl. Jungh. 1. p. 33; Fl. Ind. bat. 1 (2). p. 253, Wedd. Monogr. p. 374. Statura multo minore humili decumbente a B. rugosissima longe distare videtur, singularum partium structura autem arctius cognata est (Miq.) (v. s. in hb. h. bot. lugd. bat.).
- μ, *tomentosa*, foliis ovatis acuminatis basi rotundatis vel subcordatis toto margine nisi prope petiolum acute serratis planis subcoriaceis supra molliter hirsuto-pubescentibus. spicis ramosis vel simplicibus, femineis patulis vel pendulis folia æquantibus superantibusve. § In regione subtropica montium Khasiæ Indiæ orient. et Javæ et in Madagascaria (Commerson, etc.) B. tomentosa Wedd. in Ann. sc. nat. 1. c. p. 200. (v. s.)
- ν, *holosericea*, foliis (8-14 centim. longis) subrotundo-ovatis acutis vel breviter cuspidatis basi rotundatis vel subcordatis grosse crenato-serratis subcoriaceis supra dense appresse pubescentibus subtus velutino-tomentosis, cæt. ut in præced. a qua differt præcipue dentibus foliorum obtusis. § In Japonia (Burger). B. holosericea Bl. l. c. p. 221. (v. s.)
- ξ, *rotundifolia*, foliis (6-12 centim. longis) rotundatis vel late ellipticis ovatisve abrupte acuminatis s. cuspidatis (cuspidate 1-2-centimetrali anguste lanceolato integro) basi plerumque late cuneatis integrisque cæterum grosse crenatis vel serratis planis membranaceis utrinque breviter pubescentibus vel supra hispidulis demumque glabratibus, spicis ut plurimum unisexualibus simplicibus vel (masculis præsertim) basi ramosis s. fere fasciculatis patulis folio brevioribus. § In India orientali, regione tropica et subtropica montium Khasiæ et Himalayæ, altit. 0-1500 metr., prov. Concan et Nepalia, necnon in Zeylania. B. rotundifolia Hamilt. et Don Prodr. Fl. Nepal. p. 60. B. caudigera Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 199; Bl. Mus. bot. lugd. bat. 2. p. 219. U. blanda et U. caudigera Wall. cat. Rami novelli petiolique sericeo-velutini. (v. s.)

- o, *canescens*, foliis in toto margine excepto acumine serratis canescentibus, cæt. ut in var.  $\xi$ .  $\S$  In India orientali, cum præcedente. B. ternifolia Don, Prodr. Fl. Nepal. p. 59. B. canescens Wedd. l. c. p. 28; Bl. l. c. p. 224. Urtica macrostachya var. canescens Wall. Cat. n. 4582, D. (v. s.)
- π, *Japonica*, foliis (4-8 centim. longis) rhombeo-ovatis in cuspidem linearem productis basi subcuneatis integrisque cæterum grosse et acute subinciso-serratis planis membranaceis glabriusculis vel subtiliter hispidulo-pubescentibus, spicis fem. simplicibus aut basi ramulosis erectis folium sæpissime superantibus interrupte glomeruliferis.  $\S$  In Japonia. B. spicata Thunb. in Trans. Linn. Soc. 2. p. 330; Bl. l. c. p. 220. Urtica spicata Thunb. Fl. jap. p. 69; Bl. Bijdr. p. 492. U. japonica Linn. Suppl. p. 418. (v. s.)
- p, *macrophylla*, foliis (8-20 cent. longis, 8-25 cent. latis, ex Bl.) orbiculari-ovatis ellipticisve in cuspidem linearem apice productis basi integra rotundatis vel interdum subcordatis inæqualiter grosse et profunde dentato-serratis rugulosis supra pubescenti-scabris subtus holosericeis, spicis masc. ramosis, femineis simplicibus rariusve basi ramulosis elongatis interrupte glomeruliferis.  $\S$  In Japonia et Sina. — B. macrophylla Sieb. et Zucc. in Abhandl. der math. phys. Kl. der baier. Ak. 4(3), p. 215. B. longispica Steud. in Bot. Zeit. 33. p. 260; Bl. Mus. bot. lugd. bat. 2. p. 221. B. grandifolia Wedd. l. c. p. 199. Urtica macrophylla Thunb. Fl. jap. p. 70; Poir. Encycl. 4. p. 642. U. grandifolia Thunb. mscr. in herb. Pourret. — Folia nonnunquam basi cuneato-obtusa, rarissime palmatifissa, serraturis sat variabilibus rectis vel subfalcatis acutis vel acuminatis dorso interdum 1-2-dentatis raro obtusis. Forma altera a Blumio relata (var. appendiculata Bl. l. c.) culturæque tribuenda folia habet apice in cuspidem longam linearem basi subbidentatam terminata superne grosse inciso-dentato-serrata, serraturis acutis vel acuminatis ad marginem deorsum spectantem interdum unidentatis. (v. s.)
- σ, *Sieboldiana*, foliis (6-20 centim. longis) ellipticis vel ovato-lanceolatis cuspidatis basi obtusis vel rotundatis grosse argute serratis membranaceis glabriusculis vel ad marginem et nervos utrinque puberulo-scabridis, spicis masc. (solum visis) simplicibus vel ramosis folio brevioribus interrupte glomerulatis.  $\gamma$ - $\S$  in Japonia. B. Sieboldiana Bl. l. c. p. 220. Ramuli petiolique pube brevissima scabridi. Spicæ masc. graciles, pubescentia densiore quam petioli obsitæ. (v. s. typ. cl. Miq. communic.)
- τ, *Loo-Chooensis*, foliis (8-10 centim. longis) oblongis lanceolatisve acuminatis basi obtusa acutatave excepta serratis submembranaceis supra adpresse et sparsim puberulis nervis et venulis paginæ inferioris adpresse canescenti-pilosis, spicis fructiferis simplicibus patulis pendulisve folium æquantibus, glomerulis approximatis, perigorio fructifero lanceolato vel obovato-lanceolato.  $\S$ ? In ins. Loo-Choo (Wright, n. 303). Ramuli novelli petiolique canescentes, internodiis brevibus. Folia in triente inferiore integra, serraturis reliqui limbi parvis subacuminatis. Rhachis spicæ breviter pedunculatæ puberula. (v. s. in hb. kew.)
- υ, *Nigeriana*, foliis (9-12 centim. longis) oblongo-ovatis cuspidato-acuminatis basi obtusis et ad insertionem petioli emarginatis excepta quinta parte inferiore dentato-serratis (acumine lanceolato integro) supra hispidopilosis nervis venulisque paginæ inferioris hirtis, spicis fem. simplicibus folio longioribus patulis, glomerulis discretis, perigonio fructifero obovato.  $\S$ ? In Africa occidentali, ad Aboh (Barter) — Ramuli novelli et petioli hirsuti, internodiis elongatis. Serraturæ foliorum quam in var. præcedente duplo majores. Spicæ longiuscule pedunculatæ, rhachi villosa. (v. s. in hb. Mus. par. et kew.)
- φ, *Hamiltoniana*, glabriuscula, foliis (8-16 centim. longis) lanceolatis sensim acuminatis basi acutis aut obtusis crenulato-serratis planis nervis subtus adpresse pilosulis glabrativis membranaceis (exsiccatis læte viridibus), spicis simplicibus folio longioribus brevioribusve solitariis aut subfasciculatis fere pendulis.  $\S$  In India orientali, regione tropica Hima-

layæ orientalis a Bhotan ad Sikkim, et montibus Khasia, altit. 600-1500 metr. (Griffith etc.) B. Hamiltoniana Wedd. in ann. sc. nat. 4<sup>e</sup> sér. I. p. 199; Monogr. p. 371. Urtica Hamiltoniana Wall. Cat. n. 4590. Frutex bi-tri-metralis, ramulis elongatis et vulgo glaberrimis. Folia in dimidia parte superiore longe attenuato-acuminata, acumine ut reliquis limbus serrato, serraturis in foliis quibusdam fere obsoletis, petiolis ejusdem jugi sæpius valde inæquilongis. Spicæ fem. (solum visæ) illis var.  $\delta$  maxime similes sed breviores. (v. s.)

Obs. — Stirpes sequentes, pleræque a beato Blumio conditæ atque mihi non aut vix cognitæ, forsitan ad varietates quasdam B. platyphyllæ referendæ sunt.

34. B. HETEROIDEA (Bl. Mus. bot. lugd. bat. 2. p. 216), ramulis petiolisque et nervis subtus parce pilosiusculis, foliis ovatis acuminatissimis basi rotundatis crenato-serratis membranaceis supra glabriusculis, floribus in axillis et spicis paniculatis terminalibus longissimis glomerulis spicarum interruptis, perigonio fructifero bialato ore bidentato.  $\delta$  In India orientali (Helfar). Folia 6-10 centim. longa,  $3\frac{1}{2}$ -5 centim. lata. Caryopsis non ut in B. caudata alata, sed margine incrassato cincta et perigonio membranaceo-alato obsolete pubigero ore constricto vestita (Blume).
35. B. CUSPIDATA (Bl. l. c. p. 216, non Wedd.), ramulis petiolisque et nervis subtus pube parca brevissima strigosulis, foliis ovatis ellipticisve cuspidatis dentato-serratis membranaceis supra pilis brevibus albis hispidulis subtus puberulis, spicis fem. solitariis simplicibus filiformibus interrupte glomeruliferis folia longitudine adæquantibus, perigonio fructifero ovoideo lenticulari-compresso superne parce pubigero, ore quadridentato,  $\delta$  In Nepalia. Folia 4-7 centim. longa, 2-4 $\frac{1}{2}$  centim. lata. Petioli 1-4-centimetreales.
36. B. SPICIFLORA (Bl. l. c. p. 217), ramulis petiolisque et nervis subtus appresse puberulis, foliis magnitudine et petiolorum longitudine sæpe inæqualibus ovatis v. ovato-oblongis acuminatis basi suboblique rotundatis v. subcordatis dentato-serratis chartaceis supra pilis raris conspersis mox glabratiss, spicis fem. solitariis subramosis folia longitudine superantibus interrupte glomeruliferis, perigonio fructifero ellipsoideo compresso anguste marginato puberulo ore subbidentato.  $\delta$  In India orientali (fide specim. Mus. Vindob.) Caturus spiciflorus herb. Jacq. fil. non Linn. Folia 10-14 centim. longa, 6-9 centim. lata, petiolis 2-5-centimetralibus.
37. B. HUGELIANA (Bl. l. c. p. 218), ramulis folisque subtus et petiolis pilis brevibus substrigosis, foliis subrotundo-ovatis v. ovato-ellipticis acuminatis basi rotundatis obtusisve aut subcordatis argute dentato-serrulatis membranaceis supra pilosiusculis, spicis solitariis longitudine foliorum v. longioribus monoicis interrupte glomeruliferis femineis simplicibus v. subramosis, masculis fasciculatis, perigonio fructifero ovoideo lenticulari-compresso hirtulo ore bidentato.  $\delta$  In India orientali (Hügel, n. 1935, in Mus. Vindob.) — Folia 10-20 centim. longa, 5-14 centim. lata, petiolis 1-7-centimetralibus (Blume).
38. B. HISPIDULA (Bl. l. c. p. 223), ramulis petiolisque hirtulis, foliis subrotundo-ovatis acuminatis basi rotundatis grosse dentatis membranaceis supra pilis sparsis hispidulis subtus dense pubescentibus, spicis fem. solitariis pedunculatis simplicibus cylindraceis folio brevioribus, perigonio fructifero elongato-turbinato compresso membranaceo-marginato hispidulo ore bidentato.  $\delta$  In provincia Palembang insulæ Sumatræ. — Affinis B. holosericeæ Bl. et B. pilosiusculæ Hassk. (Blume).
39. B. JACQUINIANA (Bl. l. c. p. 219), ramulis petiolisque puberulis, foliis (magnitudine et petiolorum longitudine inæqualibus) ovatis v. ellipticis cuspidatis basi obtusis v. rotundatis v. retusis grosse dentato-serratis membranaceis supra glabriusculis subtus imprimis in margine nervisque appresse puberulis, spicis fem. solitariis simplicibus v. basi parce ramosis longissimis interrupte glomeruliferis, perigonio fructifero (semiadhærente) ellipsoideo-compresso puberulo, ore bidentato.  $\delta$  Hab. ? — Urtica interrupta hb. Jacq. fil. non Linn. — Folia 5-9 centim. longa, 2-5 centim. lata, petiolis 1-2-centimetralibus. —

*B. spicatæ* Thunb. accedens, foliis autem basi obtusioribus et spicis longissimis distincta. Diagn. fide specim. Mus. Vindob. (Blume).

40. *B. POLYSTACHYA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1, p. 200; Monogr. p. 370), dioica, foliis oppositis amplis ovatis sensim acuminatis basi obtusatis rotundatisve argute serratis planis supra pilosulis glabrativise nervis subtus puberulis, spicis quam folia brevioribus basi ramosis sive subfasciculatis erecto-patulis, perigonio fructifero obovato vel elliptico-rotundato valde compresso latiuscule membranaceo-marginato vel in dimidio superiore hispido. ♂ In Nepalia (Wallich). *Urtica polystachya* Wall. Cat. n. 4584. — Hæc stirps a me olim in herb. Wallichiano visa videtur forma s. subvarietas typi *B. platyphyllæ*.
41. *B. DENSIFLORA* (Hook. et Arn. Bot. Voy. Beech. p. 271), dioica, foliis oppositis lanceolatis superne sensim attenuatis utrinque acutis basi integris cæterum acute serratis in pagina superiore sparsim pilosulis glabrativise, spicis fem. simplicibus elongatis erectis solitariis apice nudis glomerulis confluentibus, perigonio fructifero ventricoso apice abrupte angustato lævi glabriusculo. ♂? In prov. sinensi Tche-Kiang (Staunton) et in insula Loo-Choo (Beechey). — Wedd. Monogr. p. 373. — Rami virgati, puberuli. Folia 12–16 centim. longa, 2½–3½ centim. lata, serraturis interdum majusculis, exsiccata coriacea obscure viridia, petiolo gracili 2–4-centimetrali prope limbum adpresse pilosulo. Stipulæ lineari-lanceolatæ, glabræ, mox deciduæ. Spicæ fem. (solum visæ) folio breviores longioresve, cylindricæ, vix crassæ, perigonio fructifero (1½–1¾ millim. longo) quam stylus longiore, in collum breve angustato, ore denticulato. — Cum var. ♂ *B. platyphyllæ* comparanda. (v. s. in hb. Mus. brit.)
42. *B. SUBPERFORATA* (Wedd. Monogr. p. 383), dioica, foliis alternis late ovatis vel ovato-lanceolatis anguste acuminatis basi rotundatis obtusivise acute serratis supra scabro-hispidis subtus in nervis subtiliter pubescentibus, petiolis in nodis proximis sæpe valde inæquulongis ipso limbo interdum longioribus, stipulis subsistentibus, floribus in glomerulos nudos aut folio suffultis dispositis, alabastris florum marum canescentibus, cat. ignotis. ♂? In India orientali (Wallich). *Urtica subperforata* Wall. — Habitus nonnihil *B. Malabaricæ*.
43. *B. ERYTHROPODA* (Miq. in Zolling. System. Verzeichn. 2. p. 101 et 104), ramulis petiolis nervis et inflorescentia (in sicco rubro-fuscis) tenerrime puberis, foliis oppositis longiuscule petiolatis (altero minore) ovatis vel ovalibus acuminatis supra basin rotundatam vel acutiusculam serratis alte trinerviis et paucivenosis transverse venulosis membranaceis supra glabris lævibus, spicis axillaribus solitariis gracilibus folia subæquantibus interruptis. ♂ In sylvis umbrosis, prope Lamadang Tenga, in Ins. Java (Zollinger, n. 534). — Folia 8–15 centim. longa, 3½–6½ centim. lata, saturate viridia, subtus pallidiora et rubrinervia, demum sublacunosa. Stipulæ subsiccæ fuscæ, lanceolatæ, carinatæ, glabræ. Fasciculi florum remoti, bracteis lanceolatis muniti. Perigonium fem. puberum. Stigma elongatum. — Species valde singularis, prope *B. trinervem* inserenda (Miquel). — Hanc stirpem in monographia mea ad *B. Zollingerianam* errore duxi. Monente cl. Miquelio, affinis est *B. trinervia* Miq. (Pl. Jungh.) quæ est species Pipturi; folia autem plantarum hujus generis semper sunt alterna, quum in *B. erythropoda* opposita dicantur.
44. *B. GREWLEFOLIA* (Presl. Botan. Bemerck. p. 110), ramulis foliisque novellis canescentibus, foliis oppositis ovatis acutis dentatis supra puberulis subtus glabris, spicis filiformibus laxis folium vulgo subæquantibus, glomerulis distantibus. ♂? Hab. ? Folia in eodem jugo inæqualia, altero duplo fere majore, petiolis circiter 3-centimetralibus, limbo 10 centim. longo, 4–5 centim. lato, stipulis fugacibus lanceolatis pubescentibus. Spicæ usque 2 decim. longæ, bracteis glomerulos suffulcentibus lineari-subulatis deciduis.
45. *B. ? RIGIDA* (Benth. in Hook. Fl. nigrit. p. 519), dioica, ramis aculeato-setosis, foliis alternis late ovatis obovatisve basi obtusis serratis glabris subtus albicantibus, cymis masc. axillaribus laxè paniculatis folia æquantibus. ♂ In Sierra Leone (Don). — *Urtica rigida* G. Don, mscr. in hb. propr. — Specimen unicum masc. supp. Ramus crassus, glaber, setis crassis aculeiformibus

echinatus. Folia breviter petiolata, 7 cent. longa, 5-6½ centim. lata, a basi ad apicem grosse serrata. Inflorescentia eadem ac in stirpibus masc. *B. nivæ* et *B. Puyæ*, sed laxior. Flores pentameri; ovarii rudimento clavato (Benth.).

46. *B. NUDFLORA* (Willd. Spec. pl. 4. p. 343), foliis alternis oblongis acuminatis integerrimis utrinque scabriusculis, spica solitaria terminali interrupta. 3 Prope Caracas, in sylvis, ad ripas torrentium (Bredemeyer). — Caulis sesquimetralis, debilis. Folia 4 centim. longa, triplinervia. Spica sesquidecimetralis, filiformis, floribus glomeratis, masculorum perigonio tetraphyllo. Flos fem. in glomerulis unicus, perigonio et squamis 4 adpressis constante, stylo filiformi (Willd.). — An *Boehmeria*? Stirps conferenda cum *Pouzolzia acalyphoidi*.

47. *B. MOLLICOMA* (Miq. in Zolling. System. Verzeichn. p. 100 et 105; Fl. Ind. bat. 1 (2), p. 255), innovationibus aureo-tomentosis, foliis alternis longe petiolatis cordato-ovatis acuminatis tenuiter serratis subcoriaceis, adultis subius glaucescenti-griseo-tomentosis supra pubescenti-scabris, spicis fem. axillaribus dichotome ramosis interrupte glomeruliferis, perigonis ovoideis incanulis. 3 In ins. Java (Zollinger, n. 1454). Statura et habitus *B. tomentosæ* (Miquel). — Forte ad genus *Pipturus* referenda.

### *Species exclusæ.*

- B. ACALYPHOIDES* Bl. = *Pouzolzia acalyphoides*.
- B. ACUMINATA* Hook. et Arn. = *Pipturus albidus*.
- B. ALIENATA* Willd. = *Pouzolzia indica*.
- B. ANGUSTATA* Hassk. = *Debregeasia longifolia*.
- B. ANGUSTIFOLIA* Kunth in H. et B. = *Phenax angustifolius*.
- B. ARACHNOIDEA* Walp. = *Pouzolzia macrura*.
- B. ARBOREA* Desf. = *Gesnouiina arborea*.
- B. ARGENTEA* Guillem. = *Pipturus argenteus*.
- B. BALLOTÆFOLIA* Kunth in H. et B. = *Phenax ballotæfolius*.
- B. BOIVINIANA* Bl. = *Pouzolzia Boiviniana*.
- B. BOLIVIANA* Bl. = *Pouzolzia Boiviniana*.
- B. CÆRULEA* Bl. = *Cypholophus cæruleus*.
- B. CANDIDISSIMA* Hassk. = *Leucosyke candidissima*.
- B. CANDOLLEANA* Gaudich. = *Pipturus argenteus*.
- B. CAPENSIS* Spr. = *Droguetia spec.*
- B. CASTANÆFOLIA* Cunningh. = alius ordinis.
- B. CEPHALIDA* Pers. = *Procris pedunculata*.
- B. CHINENSIS* Pers. syn. 2, p. 557. = alius ordinis.
- B. CINERASCENS* Hassk. = *Pipturus argenteus*.
- B. COCHINCHINENSIS* Pers. syn. 2, p. 557. = alius ordinis.
- B. COCHINCHINENSIS* Spr. = *Pouzolzia indica*.
- B. CONGLOBATA* Bl. = *Pouzolzia conglobata*.
- B. CRENATA* Bl. = *Cypholophus macrocephalus*.
- B. DICHOTOMA* Hassk. = *Debregeasia longifolia*.
- B. DISCOLOR* Poepp. = *Pouzolzia discolor*.
- B. DIVERSIFOLIA* Miq. = *Maoutia rugosa*.
- B. ELONGATA* Fisch. = *Phenax Sonneratii*.
- B. FAGIFOLIA* Pers. = *Elatostema fagifolium*.



- B. FORMICARIA Poepp. = Pouzolzia formicaria.  
 B. FRONDOSA Don = Pouzolzia viminea.  
 B. FRUTESCENS Hamilt. et Don = Maoutia Puya.  
 B. FRUTESCENS Thunb. = Villebrunea frutescens.  
 B. GLABRA Steud. = Neraudia melastomæfolia.  
 B. GLOCHIDIATA Hassk. = Pouzolzia indica.  
 B. GUATEMALANA Bl. = Pouzolzia Guatemalana.  
 B. HARVEYI Seem. = Cypholophus heterophyllus.  
 B. HIRTA Swartz = Phenax urticæfolius.  
 B. HYPOLEUCA Hochst. = Debregeasia bicolor.  
 B. INCANA Hassk. = Pipturus incanus  
 B. INTERRUPTA Willd. = Fleurya interrupta.  
 B. JANATSI-ITSIGO Siebold. = Debregeasia edulis.  
 B. JAVANICA Hassk. = Fleurya interrupta.  
 B. LATIFOLIA Bl. = Cypholophus macrocephalus.  
 B. LUTESCENS Bl. = Cypholophus lutescens.  
 B. MACROTHYRSA Miq. = Laportia decumana.  
 B. MACULATA Pers. = Elatostema maculatum.  
 B. MELANOCARPA Bl. = Cypholophus melanocarpus.  
 B. MELASTOMÆFOLIA Hook. et Arn. = Neraudia melastomæfolia.  
 B. MICROCELTIS Miq. = Pouzolzia viminea.  
 B. MOLLIS Wedd. = Cypholophus macrocephalus.  
 B. MOLUCCANA Wedd. = Cypholophus macrocephalus.  
 B. NANA D. Don = Pouzolzia indica.  
 B. NEPALENSIS Wedd. = Villebrunea frutescens.  
 B. NUDIFLORA Willd. = ?  
 B. ODONTOPHYLLA Miq. = Maoutia aspera.  
 B. OVALIS Miq. = Pouzolzia viminea.  
 B. OVATA Seud. = Neraudia melastomæfolia.  
 B. PALMATA Hassk. Cat. h. Bogor. alter p. 79. = ?  
 B. PAUCIFLORA Bl. = Droguetia pauciflora.  
 B. PROCRIDIOIDES Bl. = Pouzolzia procridioides.  
 B. PROPINQUA Decsne. = Pipturus argenteus.  
 B. PROSTRATA Bl. = Cypholophus prostratus.  
 B. PUTA Hook. = Maoutia Puya.  
 B. RACEMIFLORA Forst. = alius ordinis.  
 B. RADICANS Nees et Mart. = Pilea rhizobola.  
 B. REPANDA Hassk. = Pipturus repandus.  
 B. REPENS Nees et Mart. = Pilea hirtella.  
 B. RUBESCENS Hassk. = Villebrunea rubescens.  
 B. RUBESCENS Jacq. = Gesnouinia arborea.  
 B. RUGOSA Pers. = Phenax ballotæfolius.  
 B. SALICIFOLIA Don = Debregeasia bicolor.  
 B. SANGUINEA Hassk. = Pouzolzia ovalis.

- B. SCABRA Hassk. Cat. h. Bogor. alter p. 79 = alius ordinis.  
 B. SERICEA Bl. = Cypholophus macrocephalus.  
 B. SQUAMIGERA Wedd. = Chamaibainia squamigera.  
 B. STRIGOSULA Bl. = Pouzolzia viminea.  
 B. SYLVATICA Hassk. = Villebrunea sylvatica.  
 B. TRINERVATA Pers. = Gyrotænia cephalantha.  
 B. TRINERVIS Miq. = Pipturus repandus.  
 B. URTICÆFOLIA Pers. = Phenax urticæfolius  
 B. VELUTINA Don = Pipturus incanus.  
 B. VESTITA Bl. = Cypholophus vestitus.  
 B. VIALIS Casaret. = Phenax Sonneratii.  
 B. VIMINEA Bl. = Pouzolzia viminea.  
 B. WAITZIANA Bl. = Pouzolzia viminea.

XVIII. CHAMABAINIA Wight, Icon. pl. Ind. or. 6, t. 1981; Wedd. Monogr. p. 387. — Urticæ spec. Wall. — Boehmeriæ sectionis spec. Bl. Mus. bot. lugd bat. 2. p. 207.

Flores monoici vel dioici?, in glomerulis axillaribus conferti, bracteis brevibus scariosis stipati. *Masc.* Perigonium 4-lobum, segmentis sub apice acutissimo gibbosis mucronatisve, in præfloratione valvatis. Stamina 4. Pistilli rudimentum clavatum glaberrimum vel pilis paucissimis stipatum. *Fem.* Perigonium tubulosum, compressum, erostre, ore valde contracto minute 4-dentato. Ovarium ovulumque ut in Boehmeria. Stigma breve, ovatum s. ellipticum, ovarii apice acuminato continuum. Achaenium ellipticum, læve, perigonio persistente et marcescente (?) inolusum, pericarpio crustaceo. Embryo ut in genere memorato. — Herba indica, caule repente demumque lignescente. Folia opposita, isomorpha, serrata, trinervia, pilosula. Stipulæ insignes, liberæ, axillares, lateralesque, inter petiolos sæpe nonnihil concretæ. — Genus monotypum, a Boehmeria conformatione stigmatidis bene distinctum.

C. SQUAMIGERA, foliis ovatis vel elliptico-ovatis superne attenuatis acuminatisve (acumine obtusiusculo) excepta basi acuta aut obtusa acute serratis, stipulis glomerulos involucrentibus. ♂ ♀ In sylvis humidis, ad radices arborum et locis udis reg. temp. et subtrop. montium Khasiæ, Himalayæ a Sikkim ad Kumaon (Wallich, Strachey et Winterbottom, n. 8; Hook. f. et Thomson) et in montibus Nilghiri peninsulae indicæ (Wight). — C. cuspidata Wight, l. c.; Wedd. Monogr. p. 387, t. 12. B. Urtica squamigera Wall. Cat. n. 4592. Boehmeria squamigera Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 203; Bl. l. c. p. 207. — Diffuse ramosa, caule sæpius longe repente et e nodis radicante, ramis adscendentibus 1-3-decimetralibus gracilibus subsimplicibus quadrangulis magis minusve pilosis, exsiccatis ut plurimum ferrugineis, internodiis (præsertim mediis) elongatis. Folia patula, 1½-5 centim. longa, 2½ centim. lata, basi plerumque late cuneata integraque, serraturis reliqui limbi obtusiusculis acutisve et margine exteriori sæpius convexis, nervis basilaribus ad trientem superiorem limbi excurrentibus subtus pubescentibus aut (sicut in pagina superiore) sparsim pilosis, lamina exsiccata membranacea et haud rara fusciscente, petiolis 5-20 millim. longis atque in eodem jugo nonnihil inæqualibus pubescentibus. Stipulæ 4-5 millim. longæ, ovatæ, acutæ vel rotundatæ et mucronatæ, ejusdem nodi caulem amplectentes sæpeque basi connatæ et perfoliatæ, membranaceæ, erectæ patulæve vel reflexæ, nervo dorsali pilosæ, siccitate ferrugineæ. Glomeruli masc. ex axillis superioribus vel in ramis distinctis orti, floribus laxiuscule congestis, pedicellis 2-3-millimetralibus, perigonii lobis sub apice mucrone brevissimo v. longiusculo

obtusiusculoque instructis extus in dimidio superiore longe pilosis, mucrone interdum glabro, alabastro circiter æsquamillimetrali. Glomeruli fem. inferiores, densiflori, floribus subsessilibus bracteis brevibus lanceolatis stipatis, perigonio obovato compresso (ore valde contracto) in dimidio superiore densiuscule piloso, fructifero fere rotundato millimetrali margine obtuso achæ-nium ellipticum acuminatum laxiuscule includente. (v. s.)

- XIX. POUZOLZIA *Gaudich. Bot. Voy. Uran. p. 503; Wedd. Monogr. p. 389; Benth. Fl. Hongk. p. 331; Miq. Fl. Ind. bat. 1 (2). p. 256. — Pouzolziæ spec. Bennett, Pl. jav. rar. p. 66; Wight, Icon. pl. Ind. 6. p. 36. — Pouzolziæ et Boehmeriæ spec., Leptocnide et Stachyocnide Bl. Mus. lugd. bat. 2. p. 193 et seq. — Lencococcus Liebm. in Kgl. danske Vidensk. Selsk. Skr. 5 Række, naturv. og. math. Afd. 2. p. 311. — Margarocarpus et Pouzolziæ spec. Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 203 et 206.*

Flores monoici, rarius dioici, in glomerulos axillares v. spicatos dispositi, masculi femineique in stirpibus monoicis sæpius commixti, bracteis parvis scariosis suffulti, florum masc. pedicellis (quum adsunt) articulatis. Masc. Perigonium 4-5-lobum v. partitum, raro tripartitum, lobis ovatis acuminatis dorso convexis (nec mucronatis nec transverse infractis), in præfloratione valvatis. Stamina 3-5. Pistilli rudimentum clavatum v. obovatum, glabrum aut plus minus lanatum. Fem. Perigonium tubulosum, sæpius ovatum nervosumque, ore contracto bi-quadridentato. Ovarium inclusum, nunc et frequentissime liberum, nunc (ut in Boehmeria) perigonio plus minus cohærens s. adnatum, sessile, apiculatum. Ovulum erectum aut adscendens. Stigma filiforme, cum apice ovarii articulatum et fere semper ante maturitatem fructus deciduum (1), altero latere villosum. Achæ-nium ovario subconforme, perigonio marcescente subimmutato aut costis alisque aucto vestitum, pericarpio crustaceo nitido modo atro modo fusco aut eburneo. — Suffrutes frutesceve aut herbæ in regionibus tropicis vigentes, alii neogei, alii gerontogei, glabriusculi aut varie pubentes. Folia alterna rariusve opposita plerumque homomorpha rarissime inæquilateralia integerrima aut rarius dentata trinervia, nervis basilaribus ramosis nec unquam usque ad apicem limbi productis, cystolithis punctiformibus. Stipulæ liberæ sæpissime persistentes. — Pouzolziæ nonnullæ integrifoliæ Parietarias habitu mentiuntur; species vero foliis dentatis donatæ Boehmeriis quibusdam adeo similes sunt ut ab illis non nisi stigmate cum apice ovarii articulo (nec continuo) distinguendæ sint. Character desumptus e costis aut alis quibus perigonium fructiferum in Pouzolzia et in Memoriali ornatum est, licet optimus quando sese ostendit, nihilominus tantum abest ut genericus sit ut vix est specificus, quoniam in eadem inflorescentia inveniuntur nonnunquam perigonia alia alata, alia costata aliaque aptera et fere ecostata. Adde quod inter Boehmerias videantur stirpes quorum perigonium fructiferum etiam alatum vel subalatum sit, sed advertatur alam in hisce semper expansionem esse marginis perigonii; alæ autem perigonii Pouzolziarum sunt frequentissime expansiones nervorum lateralium atque ideo sunt vulgo 4 nec binæ.

(1) In speciebus paucis perigonio femineo apice longiuscule attenuato donatis accidit ut stigma intra collum hujusce basi detentum persistens videatur.

I. — Species gerontogææ.

### § 1. — Folia integerrima.

A. — *Glomeruli axillares. Perigonium ab ovario discretum.*

\* Flores masculi tetrandri v. triandri (1).

1. P. INDICA (Gaudich. Bot. Voy. Uran. p. 503), caule sæpius diffuse ramoso ramisque adpresse pubescentibus pilosisve aut fere glabratiss, foliis omnibus aut fere omnibus alternis ovatis lanceolatisve acutis vel acuminatis basi acutis obtusisve aut rotundatis breviter petiolatis integerrimis tri-tripli-nerviis, costa utrinque nervum unicum primarium præter basilares (simpliciter ramosos) emittente, supra parce pubescentibus pilosisve vel glabratiss, nervis subtus adpresse pubescenti-pilosis, margine hispido-spinulosis, glomerulis axillaribus androgynis sæpius paucifloris, floribus masc. tetrandris, perigonio fem. quam stigma brevior, fructifero subæqualiter costato vel costis margini proximis paulo elevatioribus margine ipso nisi in parte superiore exalato. 3-4 In Asia tropica orientali et in insulis maris Indici admodum frequens. — Benn. Pl. jav. rar. p. 67; Wight, Icon. pl. Ind. 6. t. 1980 et 2100. n. 40; Wedd. Monogr. p. 398. t. 13. B. f. 17-16; Bl. Mus. bot. lugd.-bat. 2. p. 233; Miq. Fl. Ind. bat. 1 (2). p. 257. P. glomerata Decsne in Nouv. Ann. Mus. 3. p. 492 in obs.; Bl. l. c. p. 232 (foliis ovatis v. ovato-lanceolatis supra parce subtus den-e adpresse pubescentibus, ramulorum minoribus). P. parietarioides Decsne l. c. Bl. l. c. p. 235 (foliis inferioribus oppositis). P. bornensis Bl. l. c. p. 234 (foliis subtriplinerviis). P. pauciflora Benn. l. c. (caule vix ramoso, foliis longius petiolatis). Parietaria indica Linn. Mantiss. p. 128. P. zeylanica ejusd. l. c. 501. Urtica glomerata Bl. Bijdr. p. 502. U. triplinervis Wall. Cat. n. 4607. U. glomerata et U. repens ejusd. l. c. n. 4619. A et F. Herba Memoriae Rumph. Herb. Amb. 6. t. 12. f. 2. — Caulis ut plurimum basi lignescens, crassitie variabili raro autem ultra eam pennæ corvinæ vel anatinæ abeunte, altitudine s. longitudine modo vix decimetrales, modo plus quam semimetrales, exsiccati sæpe ochracei vel ferruginei, longitudine internodiorum admodum varia. Folia vulgo 1-3 centim. longa, 4-13 millim. lata, sed sæpissime etiam minora et haud raro majora, lamina exsiccata rigidula v. membranacea vix asperula, petiolo sæpius brevi centimetrum raro superante. Stipulæ liberæ, axillari-laterales, a basi dilatata latiuscule ovata, subabrupte et anguste acuminatæ, 1½-3 millim. longæ, scariosæ, ciliatæ, extus vel tantum nervo dorsali et acumine longiuscule pilosæ. Glomeruli pauciflori. Flores sexus utriusque intermixti et sæpe fere æquinumeri, subsessiles, perigonii masc. 4-partiti segmentis breviter acuminatis extus pilis rectis hamatisque strigoso-hispidis. Flores fem. perigonio magis minusve strigoso vel piloso sed demum (nisi ad apicem) sæpius glabrato. Perigonium fructiferum ovatum, ventricosum, superne sensim attenuatum, acipiti-compressum apiceque bidentatum, 1½-2 millim. longum, 1 millim. circiter latum, costis 8-12 ad maturitatem valde prominentibus crassiusculisque, lateralibus raro nonnihil elevatioribus. Achænium ovatum, millimetrale, nitidum, initio nisi in ipso apice eburneum sed demum omnino atrum. (v. s.) — Subvarietates sunt: 1° *Ceramica*, caule erecto a basi subfastigiato-ramoso. foliis (teste Blumio) alternis v. oppositis, floribus masc. triandris. 3 In Ceramia magna Moluccana (Forster). Specimen vidi, mecum ex herbario Lugd. Bat. a cl. Miquelio benigne communicatum, foliis omnibus alternis floribusque masc. omnibus tetrandris. — 2° *suffruticosa*, caule elatiore, ramis virgatis, foliis superioribus lineari lanceolatis, basi subcordatis, perigonio fructifero costis lateralibus nonnihil elevatioribus. 3-5 In India orientali (Roxburgh, Wight). P. suffruticosa Wight, l. c. Urtica suffruticosa Roxb. Fl. Ind. 3. p. 583; Wight, l. c. 2. t. 694. (v. s.) — 3° *procumbens*, caulibus s. ramis procumbentibus v. repentibus, foliis ovatis lanceolatisve obtusiusculis v. acutis. 4-5 In India orien-

(1) Vid. etiam, inter species foliis dentatis, P. heterophyllum.

tali et ins. Java et Hongkong. *P. procumbens* Wight, l. c. 6. t. 2099; Bl. l. c. p. 231. *P. parva* ejusd. l. c. p. 234. Fl. masc. (texte Blumio) tetrandri v. pentandri atque perigonio fructifero passim bi-tri-v. quadri-marginato. (v. s.) — 4° *umbrosa*, foliis amplioribus tenuiter membranaceis, exsiccatis læte viridibus. 3 In India orientalis prov. Silhet, etc. (Wallich n. 4617. c.) (v. s.)

β, *angustifolia*, caule basi præsertim ramoso et vulgo elatiore quam in typo, ramis elongatis, foliis plerisque alternis lanceolatis v. ovato-lanceolatis petiolatis supra glabris nervis subtus pilosulis, perigonio fructifero subcompresso facie utraque costis quinque plus minus prominentibus percursa sæpiusque in toto margine sed præsertim apice membranaceo-alato. 5 In ins. Java et Borneo (Junghuhn, Zolling. n. 52, etc.), Philippinis, et in peninsula Malaccensi (Griffith). — *P. glomerata* Hassk. Cat. hort. Bogor. p. 80, Miq. Pl. Jungh. 1. p. 24; Fl. Ind. bat. 1 (2). p. 238; Wedd. Monogr. p. 401. *P. angustifolia* Wight, l. c. 6. t. 2100. n. 39? — Caulis ramique (4-5-decimetrales) siccitate fusciscentes, internodiis inferioribus 3-10 centim. longis. Folia 3-10 centim. longa, 8 millim.-2½ centim. lata, petiolo 3-15-millimetrali. Perigonium fructiferum oblongum, 2½ millim. circiter longum, 1-1½ millim. latum, ala marginali prope apicem præsertim conspicua, costis sæpe parum elevatis. Achænium quam in typo vix majus. — Figura iconi Wightianæ supra laudatæ ob perigonium fructiferum haud marginatum forsitan ad ipsum typum potius adscribenda. (v. s.)

γ, *alienata*, caule vix ramoso, foliis plerisque oppositis ovatis quam in typo longius petiolatis basi rotundatis v. subcordatis, perigonio fructifero modo subæqualiter costato modo plus minus marginato aut 4-auriculato-alato, costis nempe lateralibus utriusque faciei in auriculas crassas semirotundatas v. triangulares dilatatis. 4-5 In India orientali et Zeylanica. — *P. alienata* Gaudich. l. c. P. Zeylanica Benn. l. c. p. 67; Wight, l. c. t. 2101. n. 45. *Parietaria zeylanica* Linn. Fl. zeyl. n. 371; Spec. pl. 1492, nec Mantiss. *Urtica alienata* Linn. Syst. veg. ed. 13. p. 709; Roxb. Fl. Ind. 3. p. 582. *Boehmeria alienata* Willd. Spec. pl. 4. p. 341. Folia vulgo ampliora quam in typo, pilosula, exsiccata tenuiter membranacea. (v. s.) — Subvarietates sunt: 1° *heterophylla*, procumbens, laxè ramosa, ramis elongatis, foliis plurimis oppositis breviter petiolatis aliis subrotundis v. ovalibus, aliis ovatis v. ovato-oblongis supra sparse pilosis subtus dense appresse pubescentibus, perigonio fructifero costato et interdum 4-marginato. 5? In India orientali. *P. heterophylla* Bl. l. c. p. 237. *Urtica alienata* Roxb. l. c. Wight, l. c. 2. t. 693, ex Bl. P. Forsteriana Bl. l. c. p. 238 (foliis partim magis lanceolatis acuminatis et perigonii fructiferi minus distincte costatis). (v. s.) — 2° *muralis*, sæpius herbacea, opposite ramosa, foliis oppositis ovatis v. ovato-lanceolatis breviter petiolatis supra scabriusculis subtus densiuscule et appresse pubescentibus pilosisve, perigonio fructifero angulato-costato v. sæpius quadrialato, alis rotundatis. 6-5. In ins. Java (Junghuhn). *P. muralis* Miq. Pl. Jungh. 1. p. 24; Bl. l. c. p. 237. *P. glochidiata*? Bl. l. c. p. 245; Miq. Pl. Ind. bat. 1 (2). p. 238. *Boehmeria glochidiata* Hassk. Pl. Jav. rar. p. 206. Folia 1½-3½ cent. longa, siccitate rigidula, inferiora passim ovato-rotundata. (v. s.) — 3° *strigosa*, procumbens, laxè ramosa, ramis elongatis, foliis oppositis ovato-ellipticis breviter petiolatis (infirmis et ramulorum minoribus ovalibus v. obovatis obtusis) supra sparsim pilosis subtus imprimis ad nervos dense pubescentibus, perigonio fructifero obsolete costato v. interdum quadrimarginato. 5? In ins. Java, Borneo, cat. P. strigosa Hassk. Cat. hort. Boq. p. 80; Bl. l. c. p. 237; Miq. l. c. p. 239. (v. s.) — 4° *microphylla*, procumbens, diffusa, ramosissima, foliis ovatis acutis v. obtusis breviter petiolatis supra pilosulis subtus dense appresse pubescentibus, perigonio fructifero costato apterisque v. quadrialato, alis rotundatis. 4-5 In ins. maris Indici, peninsula indica, Sina et Cochinchina. — P. prostrata. Benn. l. c. p. 67? *P. microphylla* (non Wight), P. nana et P. Cochinchinensis Bl. l. c. p. 236 et 245. *Parietaria Cochinchinensis* Luureiro, Fl. cochinch. p. 654. *Urtica nana* Hamilt. mscr. *Boehmeria nana* D. Don, Prodr. Fl. nepal. p. 60. *B. Cochinchinensis*

Spr. 3, p. 844. Folia subconformia v. dissimilia (v. s.) — 5<sup>a</sup> Subvar. *tenuis*, caule erecto v. ascendente simplici v. subramoso, foliis superioribus alternis inferioribus oppositis longius petiolatis ovatis. subrhombico-oblongis obtusis v. obtuse acuminatis utrinque pilosulis, perigonio fructifero costato et inaequali-quadrialato. 4 In India orientali (Hügel). P. *tenuis* Bl. l. c. p. 233. *Urtica tenuis* Steud. in Fl. Regensb. 1850, p. 257.

2, *tetraptera*, caule basi præsertim ramoso, ramis virgatis, foliis plerisque alternis ovatis v. ovato-lanceolatis subæqualibus quam in typo vulgo majoribus breviter petiolatis utrinque pilosis, perigonio fructifero subæqualiter costato vel latiuscule membranaceo-quadrialato, alis membranaceis nucleo ipso æquilatis apicemque perigonii sæpe superantibus. 5 In India orientali (Wight et cat.) — P. *tetraptera* et P. *Johnsoniana* Wight, l. c. t. 2190. n. 42 et t. 2101. n. 47. P. *trichophora* Hobenack. in sched. pl. Ind. or. n. 179. (v. s.) — Subvarietates sunt: 1<sup>a</sup> *pyramidata*, foliis plerisque lanceolatis, superioribus conspicue diminutis. 5 In India orientali (Wight) P. *pyramidata* Wight, l. c. t. 2101. n. 48. (v. s.) — 2<sup>a</sup> *pilosa*, caulibus a. ramis primariis robustis simplicibus elongatis prostratis, foliis inferioribus alternis oppositis ovatis magnitudine superiorum vix diminuta. 5 In India orientali (Wight et cat.) P. *pilosa* Wight, l. c. t. 2100. n. 45; Bl. l. c. p. 234. (v. s.) — 3<sup>a</sup> *minor*, caulibus s. ramis primariis gracilibus diffusis prostratis, foliis parvis omnibus alternis ovatis pilosis. 4 In iisdem locis ac præced. P. *minor* Wight, l. c. t. 2100, n. 43. (v. s.)

2. P. *TUBEROSA* (Wight, l. c. 6, gen. Pouz. n. 44), radice tuberosa fusiformi, caulibus pluribus elongatis prostratis, foliis inferioribus oppositis superioribus alternis ovatis lanceolatisve acutis integerrimis pilosis, floribus masc. tetrandris, perigonio fructifero lævi aut vix profunde costato aptero. 4 In India orientalis locis umbrosis ad sepes et rivulos (Roxburgh). — Wedd. Monogr. p. 402. *Urtica tuberosa* Roxb. Fl. Ind. 3. p. 583; Wight, l. c. 2. t. 697. — Radiciis pars tuberosa plus quam decimetralis, crassitie 4-5 centim. (ex icone Wight.). Caules annui, 3-18 decim. longi, flaccidi, striati, parce pilosi. Folia patula, petiolata; 2-8 centim. longa, inferiora s. caulina majora, ramea lanceolata. Flores congesti, sessiles, masculi et feminei intermixti, perigonio fem. minuto gibboso ore contracto subdentato, stylo elongato piloso (Roxburgh). — *Urticæ alienatæ* (Pouz. indicæ var.) affinis sec. Roxburgh. Radix edulis dicitur. Specimen unicum (floriferum) a me in herb. Bentham. visum P. indicam omnino refert. Addatur radicem ipsius P. indicæ s. P. parietarii dis Decsne nonnihil tuberosam esse (vid. Bl. l. c. p. 235.)

3. P. *VESICARIA* (Wight, Icon. pl. Ind. 6. n. 45), frutescens, erecta, foliis alternis late lanceolatis trinerviis integris pilosis, floribus masc. tetrandris, perigonio fem. basi vesiculis 6-7 obovatis inflatis persistentibusque aucto. 4-5 In montibus peninsulæ Indiæ orientalis Circar dictis (Roxburgh). — Wedd. Monogr. p. 414. *Urtica vesicaria* Roxb. Fl. Ind. 3. p. 587; Wight, l. c. 2. t. 695. — Caules plures, perennes, nonnihil frutescentes, 1-2-metrales, debiles, ramulis pilosis. Folia patula, 6-9 centim. longa, circiter 3 centim. lata. Stipula semilanceolata, caducæ. Flores axillares, sessiles, congesti, masculi et feminei commixti, perigonio fem. ab illo P. tuberosæ non nisi caractere memorato discrepante. — Specimen nullum vidi.

4. P. *ORIENTALIS* (Wedd. Monogr. p. 413), ramis glabrescentibus, foliis alternis ellipticis acuminatis basi cuneatis integerrimis longe petiolatis 5-nerviis (nerviis 2 lateralibus tenuioribus) supra asperis subtus tenuissime et adpressè puberulis, glomerulis axillaribus floribus masc. tetrandris? perigonio florum utriusque sexus incano-villoso. 5 In Indiæ orientalis ditione Tranquebar (Rottler). *Leucococcus orientalis* Liebm. in Kgl. danske Vidensk. Selsk. Skr. Række, naturv. og math. Afd. 2. p. 312. Arborea (teste Liebm.). Rami teretes Folia 8-10½ centim. longa, 4 centim. lata, supra saturate viridia, glaucescentia, opaca, petiolo compressiusculo parce puberulo. Stipula decidua. Glomeruli hemisphærici, stigmatibus florum fem. rufo-villoso. — Numerus staminum non commemoratus in descriptione cl. Liebmannii et nil refertur in illa d. perigonio fructifero. Quantum autem e diagnosi generis *Leucococci* intel

ligi possit flores masc. tetrandri sunt et perigonium fem. post anthesin immutatum permanet.

5. *P. GUINEENSIS* (Benth. in Hook. Niger Flora, p. 518), caule elongato diffuse ramoso pubescente, foliis omnibus alternis ovatis vel ovato-lanceolatis petiolatis integerrimis trinerviis costa utrinque nervum unicum vel duos præter basilares emittente, limbo in utraque facie piloso margineque hispido, glomerulis pauci-multi-floris, floribus masc. tetrandris, perigonio fem. quam stigma longiore, fructifero late ovato acute marginato nec alato in utraque facie nervis filiformibus (nec costis) notata.  $\zeta$  In locis cultis et fruticetis Africæ tropicæ occidentalis, ad Fernando-Po et in insula S. Thomas (Vogel), necnon secus flumen Rio-Nunes dictum in Senegalia (Heudelot). — Wedd. Monogr. p. 403. *Parietaria guineensis* G. Don, mscr. Caulis erectus vel ascendens, 6 dm. et ultra altus, teretiusculus, basi crassitie circiter pennæ anatinæ, superne breviter pubescens aut villosulus et interdum parce pilosus, lævis, inferne glabratus, siccitate pallide ferrugineus. Folia magnitudine admodum variab. 1-5 cm. longa,  $\frac{1}{2}$ -2 $\frac{1}{2}$  cm. lata, acuminata, basi acuta obtusa aut rotundata, nervis subtus prominulis longiuscule pilosis, lamina exsiccata submembranacea in foliis adultioribus asperula, petiolo gracili 2-15 mm. longo, pubescenti-piloso. Stipulæ 1-2 $\frac{1}{2}$  mm. longæ, ovatæ, abrupte subulato-acuminatæ, in nervo dorsali margineque longiuscule pilosæ. Glomeruli densiflori, floribus utriusque sexus sessilibus commixtis, masculorum perigonio 4-partito (rarius 3-partito) extus pilis hamatis hirsuto. Flores fem. perigonio strigoso, stigmatē vix semimillimetrali recurvo longiuscule villoso. Perigonium fructiferum late ovatum aut fere rotundatum, 1 $\frac{1}{2}$  millim. circiter longum, acutum, basi late rotundatum, parce pilosum, nervis 3-4 parum prominentibus percursum. Achænium late ovatum, apiculatum, sesquimillimetrale, acutiuscule marginatum, fusco-albidum, nitidum. — Quasdam formas P. indicæ refert, sed notis designatis et imprimis brevitate stigmati facile distinguitur (v. s. typ. in hb. Benth.)

$\beta$ , *Madagascariensis*, robustior, foliis oblongo-lanceolatis basi rotundatis, perigonio fructifero omnino lævi.  $\zeta$  In summo jugo insulæ Comoræ majoris (Boivin). — Ramuli speciminis suppetentis arcuati, apice subtiliter puberuli, inferne foliis denudati, obscure rubentes. Folia quam in typo nonnihil majora. Perigonium masc. lobis acuminatis pilis rectis parce strigosis. Perigonium fem. lenticulari-compressum, stigmatē haud millimetrali, fructiferum parce strigosum apice in collum breve ore bidentato angustatum, nervis obsoletis. Achænium initio eburneum demumque atrum. — Forsan species distincta (v. s. in hb. Mus. par.)

6. *P. ABYSSINICA* (Bl. Mus. bot. lugd. bat. p. 236), caule herbaceo debili suberecto simpliciter ramoso, foliis anguste acuminatis basi acutis longiuscule petiolatis integerrimis trinerviis (nervis basilariibus ramosis et ante apicem limbi in margine desinentibus, costa utrinque nervum primarium unicum præter basilares emittente) supra hispidis subtus tenuissime arachnoideo-tomentosis, glomerulis androgynis, floribus masc. 3. v. 4-andris, perigonio fem. quam stigma longiore, fructifero nervato.  $\odot$ ? In rupibus vallis abyssinicæ Maimezano (Schimper, n. 1433.). — Wedd. Monogr. p. 404. *Parietaria abyssinica* Hochstett. mscr. in A. Rich. Tent. Flor. Abyss. II. 258. *Margarocarpus Schimperianus* Wedd. in Annal. sc. nat. 4<sup>e</sup> sér. I. p. 205. Caulis gracilis 2-5 decim. longus, inferne denudatus glabratusque, superne diffuse ramosus et pubescenti-pilosus, ramis hirtulis, internodiis 3-5-centimetralibus. Folia 3-8 cm. longa, 2-4 cm. lata, longiuscule acuminata, nervis basilariibus tenuibus simpliciter ramosis usque ad quartam partem superiorem limbi excurrentibus, nervis primariis cæteris infra median costam nascentibus atque ad apicem acuminis arcuatim protensis, lamina exsiccata tenuiter membranacea supra viridi subtus alhicante, petiolo gracili 2-3 cm. longo tenuiter pubescente sparsimque piloso. Stipulæ ovatæ, 3 mm. longæ, subulato-acuminatæ, longiuscule ciliatæ. Glomeruli axillares, pauciflori. Flores masc. femineis intermixti, perigonii 3-4-partiti segmentis breviter acuminatis pilosis. Perigonium fructiferum ovatum, 2-2 $\frac{1}{2}$  mm. longum, superne sensim attenuatum acutum, siccitate interdum irregulariter costatum, superne pilosum, achænium sordide album arcte includens. (v. s.)

7. *P. BOIVINIANA* (Wedd. Monogr. p. 405, t. 13. B. f. 17), monoica, herbacea, caulibus filiformibus repentibus puberulis, foliis lanceolatis vel oblongo-lanceolatis apice attenuatis v. subacuminatis basi obtusiusculis rotundatisve integerrimis concoloribus lævibus ciliolatis et utrinque sparsim pilosis pilis nervorum paginæ inferioris crebrioribus, glomerulis axillaribus paucifloris androgynis unisexualibusve, floribus masc. triandris, perigonio fructifero ecostato piloso. ☉? In convallibus editis umbrosisque insulæ Anjonan, prope Madagascariam (Boivin). *Margarocarpus Boivinianus* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 204. *Bœhmeria Boiviniana* Bl. l. c. p. 227. Caules plures e radice annua? prodeuntes, 1-4 decim. longæ, basi lignescentes demumque denudati, nodis fere omnibus radículas tenues emittentibus, internodiis circiter centimetralibus. Folia 1-4 centim. longa, 4-12 centim. lata, nervis basilaribus tenuibus ad quartam partem superiorem limbi cum nervis primariis sequentibus arcuatim anastomosantibus, lamina exsiccata pallide viridi, petiolo gracili pubescente. Stipulæ ovatæ v. ovato-lanceolatæ, 1-2 millim. longæ, subabrupte acuminatæ, dorso pilosæ, persistentes. Glomeruli in axillis fere omnibus nascentes, e floribus 3-5 (quorum intermedius ut plurimum femineus est) sæpius compositi. Flores masc. breviter pedicellati, perigonii 3-partiti extus densiuscule hirti segmentis valde convexis s. cucullatis apice brevissime acuminatis. Flores fem. sessiles ore inconspicue denticulato, stigmatibus quam ovarium multoties longiore. Perigonium fructiferum ovatum, vix millimetricum, utrinque convexum læveque, inferne glabratum, superne subadpresso pilosum. Achænium eburneum, perigonio conforme. (v. s. in hb. Mus. par.)
8. *P. MIXTA* (v. Solms, in Schweinf. Beitr. z. Fl. Æthiop. p. 889), monoica, ramis lignosis, foliis ovatis acuminatis integerrimis breviter petiolatis supra dense puberulis subtus plus minus albidotomentosis, glomerulis androgynis, floribus masc. 4-v. 5-andris, perigonio fem. ovato superne attenuato piloso quam stigma geniculatum multo brevior. ☿ In districtu Fazogla Abyssiniæ (Cienkow-sky). Fruticulus, ramis nonnihil angulatis reliquiis inflorescentiarum vetustarum conspersis, novellis pube patula albidâ vestitis. Folia 2-3 centim. longa, 15 millim. lata, petiolo 5-millimetrali. Stipulæ e basi ovata lanceolatæ, fuscæ. Glomeruli axillares, floribus pluribus masc. et unico fem. compositi. Florum masc. perigonium subcampanulatum, extus pilosum, segmentis 4 v. 5 apice curvatis barbulatisque, staminibus longe exsertis (filamentis 3-millimetralibus). Fructus ignotus — Affinis, sec. cl. auct., *P. abyssinica* a qua facile tamen distinguitur imprimis caule ramisque lignosis, foliis brevius acuminatis et floribus masc. 4-v. 5-andris.

\*\* Flores masculi pentandri (1).

1<sup>o</sup> Nervi laterales foliorum simpliciter nec dichotome ramosi.

9. *P. LÆVIGATA* (Gaudich. Bot. Voy. Uran. p. 503), ramulis crassis apice petiolisque perigonii et bracteolis subtilissime puberulis cæterum glaberrimis, foliis omnibus alternis elliptico- v. ovato-lanceolatis subacuminatis basi acutis obtusive longiuscule petiolatis integerrimis trinerviis (costa utrinque nervos primarios plures præter basiales emittente) undique etiam margine lævissimis, glomerulis axillaribus, floribus masc. pentandris, perigonio fructifero late dialato utraque facie inter alas majores cristis irregularibus instructa. ♂ In silvis et ad margines torrentium ins. Mauritiæ et Borboniæ (Commerson, etc.) et Timor? — Decsne Herb. Timor. p. 164; Wedd. Monogr. p. 391. t. 13. B: f. 1-6; Bl. Mus. bot. lugd. bat. 2. p. 229; Miq. Fl. Ind. bat. 1 (2). p. 256. *Parietaria lævigata* Poir. Encycl. 5. p. 17. — Arbuscula parce ramosa, ramulis sæpe arcuatis crassitie pennæ cynce v. anserinæ basi denudatis cicatricosisque epidermide obscure rubente vestitis, internodiis superioribus vix semicentimetralibus. Folia ad apices ramulorum vix numerosa, patula, 6-15 centim. longa, 2-5  $\frac{1}{2}$  centim. lata, nervis basilaribus usque ad trientem superiorem limbi productis, primariis cæteris ut plurimum utrinque (sive e

(1) Vid. etiam, inter species floribus masc. tetrandris, *P. mixtam*.



basilaribus nascentibus) numerosis quam ipsi basilaes multoties brevioribus, limina exsiccata membranacea sæpius late viridi, petiolo 1-2 centimetrali. Stipulae liberae, axillari-laterales, lanceolatae, 8-10 millim. longae, subacuminatae, uninerviæ, mox deciduae, siccitate ferrugineo-scariosae. Glomeruli axillares aut in ramuli parte inferiore denudata nascentes, discreti, pauciflori, plerique masculi, paucissimi inter mares florem fem. solitarium exhibentes; floribus masc. bracteolis ovatis ciliolatis stipatis, perigonio 5-loba vel interdum (teste Gaudich.) 6-loba, lobis subacuminatis; floribus femineis sessilibus subsessilibusve, perigonio quam stylus s. stigma multo brevior, fructifero siccitate nigricante transverse oblongo 5-6 millim. lato et dimidium circiter longo s. alto basi subcordato apice bicuspidato, alis majoribus marginalibus subtiliter pubescentibus margine denticulatis et obsolete nervosis. minoribus s. cristis (sæpius utrinque duabus) e costis perigonii ortis quam alae multo minoribus iisque crassioribus forma cæterum irregulari. Achænium ovatum, bimillimetrale, fuscum, nitidum. — Huic stirpi forsitan referenda est sequens. (v. s.)

10. *P. CANESCENS* (Gaudich. l. c.), ramulis crassis, annotinis pubescenti-tomentosis cæteris glabratiss, foliis omnibus alternis ovato-lanceolatis ovatisve attenuato-acuminatis obtusiusculis basi rotundatis petiolatis integerrimis trinerviis utrinque pubescentibus (junioribusque canescentibus) lævibus, nervis simpliciter ramosis cæterumque ut in præcedente, glomerulis axillaribus, floribus masc. (pentandris) femineisque sæpius in axillis distinctis nascentibus, perigonio fructifero ut in specie supra dicta. 5 In montibus altioribus insulae Mauritii, præsertim in illo cui nomen Pouce datur (Commerson, etc.). — Wedd. Monogr. p. 392. *P. mollis* Bl. l. c. p. 230. *Parietaria canescens* Poir. l. c. non Bl. — *Habitus* præcedentis. Rami et folia quod magnitudinem et nervationem ut in eadem, petiolis vero plerumque nonnihil brevioribus crassioribusque pubescenti-tomentosis. Stipulae ovatae, acuminatae, extus excepto margine pubescentes, siccitate ferrugineae. Glomeruli masc. multiflori, brevissime pedicellati, perigonio extus subtiliter pubente 5-loba, lobis subacuminatis; flores feminei pauci, ex axillis superioribus plerumque nascentes, floribus masc. commixti aut sæpissime solitarii; stylo perigonioque fructifero et achænio iis stirpibus præcedentibus simillimis. — Hæc planta et præcedens a sequente facillime distinguendæ sunt nervatione foliorum. (v. s.)

11. *P. AURICULATA* (Wight, Icon. pl. Ind. 6. n. 36. addit. spec.), caule erecto ramoso, ramulis crassis teretibus pubescentibus aut fere tomentosis, foliis omnibus alternis ovatis lanceolatisve aut ellipticis plerumque sensim acuminatis basi acutis obtusisve et interdum rotundatis longiuscule petiolatis integerrimis tri-tripli-subquintuplinerviis costa utrinque nervum primarium unicum præter basilaes (simpliciter ramosos) emittente, limbo discolore supra hispido-puberulo sæpeque asperulo subtus magis minusve pubescenti-tomentoso et in planta juniore canescente margine subtiliter hispido-spinuloso, glomerulis axillaribus androgynis, floribus masc. pentandris, perigonio fructifero sæpius costato apterique aut plus minus alato. 5-5 In montibus australioribus peninsulae Indiae orientalis prov. Carnatic, Mysore et Malabar, necnon in Zeylantia (Wight, etc.). — Wedd. Monogr. p. 393; Wight, l. c. t. 1980 et 2099 (foliis lanceolatis v. ovato-lanceolatis, perig. fruct. auriculato-4-alato); Bl. l. c. p. 232. *P. scabrida* Wight, l. c. t. 2100, n. 41 (foliis lanceolatis, perig. fruct. angulato-4-alato). *P. rotundifolia* Wight, l. c. t. 2098. n. 31 (foliis late ovatis v. ellipticis, perig. fruct. angulato-4-alato). *P. bicuspidata* Wight, l. c. n. 33 (foliis ovato lanceolatis, perig. fruct. marginato). *P. elliptica* Wight, l. c. n. 32 (foliis ellipticis v. oblongo-ellipticis, perig. fruct. aptero); Bl. l. c. p. 230. *P. propinqua* Bl. l. c. p. 230 (foliis ellipticis, perigonio exalato). — Rami elongati, virgati, crassitie pennae anserinae vel cyneae aut etiam crassiores, inferne hispidi asperati v. glabrati epidermideque sæpius castanea vestiti, siccitate plerumque longitudinaliter rugosi, internodiis brevibus. Folia magnitudine admodum variabili, 4-10 centim. longa, 1½-centim. lata, sensim rariusve subabrupte attenuata; nervis basilaribus usque ad basim acuminis vel etiam altius productis, primariis cæteris (e costa scilicet nascentibus) utrinque solitariis limboque dimidio circiter æquilongis, tertiariis (s. e. nervis basilaribus) paucis (3-4) trientem limbi

longitudine fere æquantibus; lamina exsiccata membranacea vel rigida; petiolo 5-15 millim. rariusve 3 aut etiam 4 centim. longo, varie pubescente. Stipulæ liberæ, axillari-laterales, ovatæ vel ovato-lanceolatæ, 4-6 millim. vulgo longæ, anguste acuminatæ, extus majus minusve pubescentes, siccitate ferrugineæ, demum deciduæ. Glomeruli pisiformes, densiflori. Flores masc. quam feminei multo numerosiores, perigonio 5-partito undique pilis brevibus rectiusculis hamatisque strigoso-hispido, lobis breviter acuminatis. Flores fem. sessiles, in axillis superioribus plerumque numerosiores, perigonio stigmati æquilongo vel multo brevior. Perigonium fructiferum 1½-3 millim. latum, superne anceps apiceque bidentatum, nunc sæpissimeque ovatum et subæqualiter costatum, raro in toto ambitu anguste bialato s. tetraptero, frequenter utrinque in triente superiore biauriculatum et cuspidatum, auriculis nempe sive alis (juxta marginem nec ex ipso margine enatis) subtus cuspidem abruptissime orientibus modoque triangularibus modo fere rotundatis aut oblongis. (v. s. typ. Wight.)

β, *Rheedii*, foliis (etiam junioribus) fere concoloribus nempe utraque pagina parce et subæqualiter puberulis, ovato v. ovato-lanceolatis perigonio fructifero auriculato-quadrilato, alis triangularibus s. superne truncatis. ½-¾ In India orientali. P. *Rheedii* Wight, l. c. t. 2099. n. 38. (v. s. typ. Wight.)

γ, *rostrata*, foliis quam in var. α et β amplioribus tenuioribusque ovatis v. rotundatis utrinque hispidulo-pilosis aut fere glabratiss, perigonio fructifero tetraptero alis superne subtruncatis. ¾-¾ In India orientali. P. *rostrata* Wight, l. c. t. 1980 et 2099. n. 34. (v. s. typ. Wight.)

12. P. *DIFFUSA* (Wight, l. c. t. 2099. n. 36); caule gracili prostrato vel repente apice præsertim hirtio, foliis omnibus oppositis ovatis ellipticisve utrinque acutis aut basi obtusatis breviter petiolatis integerrimis pilosis trinerviis, costa in triente inferiore limbi trifurcata s. utrinque nervum unicum primarium præter basiales (simpliciter ramosos) emittente, floribus masc. pentandris, perigonio fructifero costato et anguste subauriculato-4-alato. ¾ In montibus Nilghiri peninsulae Indiæ orientalis (Wight). — Wedd. Monogr. p. 394. — Caulis elongati parce ramosi, crassitie vix pennæ corvinæ, angulati, basi demum glabrati, internodiis 4-5 centim. longis longioribusve. Folia 2-3½ centim. longa, 1-1½ centim. lata, supra sparsim longiusculeque pilosa (pilis paginæ inferioris in nervis præsertim crebris), margine subtiliter spinuloso-hispida, petiolis 3-5 millim. longis hirtio-pilosis. Stipulæ laterales, ovato-lanceolatæ, 3-6 millimetrales, acuminatæ, nervo dorsali pilosæ, siccitate ferrugineæ. Glomeruli vix multiflori, androgyni. Flores masc. pedicellati, perigonii 5-partiti segmentis subacuminatis in dimidio superiore pilis rectis et aliis numerosioribus hamatis instructis. Flores fem. subsessiles, perigonio fructifero ovato bimillimetrali basi obtuso superne ancipiti-compresso apiceque emarginato et hirtio in dimidio inferiore utriusque faciei elevato crasseque 4-5 costato, costis lateralibus plus minus auriculiformibus et in partem superiorem perigonii decurrentibus. — Habitu, ramis angulatis et foliis oppositis a P. auriculata optime distincta. A P. indica (cui affinis) differt imprimis foliis omnibus oppositis et floribus masc. pentandris; videndum si character posterior sufficiens sit ad distinctionem specificam constituendam inter species duas sequentes et eundem typum. (v. s. typ.)

13. P. *ROTHIANA* (Bl. Mus. bot. lugd. bat. 2. p. 231), ramis elongatis angulatis strigosulis, foliis alternis brevissime petiolatis vel subsessilibus integerrimis difformibus, inferioribus subrotundis ovalibusve basi rotundatis, superioribus ovato-lanceolatis acutis v. obtusiusculis basi obtusis supra imprimis ad margines pilosiusculis subtus appresse pubescentibus, glomerulis androgynis paucifloris, perigonio fructifero obsolete costato v. interdum subangulato ore truncato. ¾? In India orientali. — Proxime affinis esse videtur P. minori Wight cujus tamen flores masc. tetrandri necnon perigonia fructifera sæpe alata et in rostrum bidentatum constructa, in hac autem, quos exploravi omnes flores masc. pentandri, perigonia vero fructifera obsolete decem-costata sejunctione duorum dentium orificii in apice brevi plerumque truncata (Blume).

14. *P. CONFINIS* (Bl. l. c. p. 230), laxè ramosa, ramis elongatis angulatis pubescentibus, foliis alternis breviter petiolatis lanceolatis (ramulorum multo minoribus subessilibus ovatis v. ovalibus obtusis) obtuse acuminatis basi acutiusculis integerrimis supra puberulis subtus dense appresse pubescentibus, glomerulis androgynis densifloris, floribus masc. pentandris, perigonio fructifero costato et interdum anguste bi-v. tri-marginato superne subancipiti-attenuato subbidentato. ♂♀ In India orientali. — Perigonia fructifera sublenticulari-compressa, superne in apicem subbidentatum anguste marginatum attenuata, decem-costata, duabus tribusve costis interdum marginum v. cristularum angustarum membranacearum prominentibus. — Habitu et foliorum forma in ramulis tenuioribus *P. glomeratæ* Decsne valde congrua. Etiam *P. scabridæ* Wight affinis, unde ramis angulatis recedit (Blume).
15. *P. HYPOLEUCA*, monoica, caulibus cæspitosis? basi lignosis simplicibus puberulis, foliis alternis amplis oblongis acuminatis basi acutis integerrimis, costa nervos primarios duos præter basilares utrinque emittente, limbo supra glabro et asperulo subtus (nervis primariis costaque exceptis) incano-tomentosis, glomerulis axillaribus androgynis, floribus masc. pentandris, perigonio fructifero ovato superne attenuato ecostato et exalato. 2-5 Ad pedem montis Miramballa, in Africa occid. austr. 14-15 lat. S. (Kirk, exp. Livingston). — Caules semimetrales, teretiusculi, a basi ad apicem foliosi, internodiis 3-4-centimetralibus. Folia 10-12 centim. longa, 4-5 centim. lata, sensim acuminata, nervis basilariibus juxta trientem superiorem limbi in margine desinentibus, venulis parum perspicuis; lamina exsiccata membranacea nervis supra parum impressis, tomento paginæ inferioris tenui et candido; petiolo gracili 2-4-centimetroali puberulo. Stipulæ lanceolatæ, acuminatæ, deciduæ. Glomeruli pisiformes, floribus utriusque sexus intermixtis. Perigonium masc. 5-lobum, segmentis extus puberulis breviter crassiusculeque acuminatis. Perigonium fem. sesquimillimetrale, puberulum, ore bidentato, stigmatibus 5-6-millimetroali; fructiferum late ovatum, bimillimetrale, lævisculum, achæmium conforme eburneumque arcte vestiens. (v. s. in hb. Kew.)
- 2° Nervi laterales foliorum furcati s. subdichotomi.
16. *P. CYMOSA* (Wight, Icon. pl. Ind. t. 1979 bis), caule erecto hirtulo vel glabrato, foliis omnibus oppositis ovato-lanceolatis acuminatis basi rotundatis vel cordatis breviter petiolatis integerrimis trinerviis, costa in triente inferiore trifurcata, nervis lateralibus furcatis s. subdichotomis, floribus laxè cymosis, masculis pentandris, perigonio fructifero aptero. ♂ Ad rivulos, in montibus Nilghiri peninsulae Indiae orientalis (Wight, cæt.). — Wedd. Monogr. p. 396; Bl. l. c. p. 238. *Elkania multinervis* Schlecht. mscr. in sched. Pl. Ind. or. (edit. Hohenack. ann. 1851). Suffrutex simplex aut parce ramosus, 5-8 decim. altus, ramis elongatis gracilibus obtuse quadrangulis pilosis strigosulisve demumque glabratibus et sæpe asperulis, internodiis frequentissime 5-6-centimetralibus. Folia omnia fere æqualia, superiora nempe magnitudine vix diminuta, 3-9 centim. longa, 1½-4 centim. lata, sæpius ovata, sensim acuminata, acutissima, costa nervisque lateralibus (in limbo medio 7-9) fere æqualibus et æquidistantibus, supra (præsertim in intervenio) sparsim et adpresse pilosula, subtus undique pubescenti-pilosula, petiolo 3-5 millim. longo. Stipulæ in nodis singulis quatuor, omnes liberæ, lanceolatæ, subulato-acuminatæ. Cymæ quam petioli ut plurimum plures quam duplo longiores, patule ramosæ, ramulis pilosulis. Flores masc. numerosi, pedicellati, atropurpurei, bracteis ovatis membranaceis suffulti, perigonii 5-partiti segmentis acuminatis extus undique pilosis, pistilli rudimento lanato. Flores feminei pauci interdumque deficientes? ad basim ramorum primariorum cymæ sessiles; perigonio fructifero ovato sesqui-millimetroali subæqualiter altiusculeque 12-costato strigoso. (v. s.)
17. *P. MICROPHYLLA* (Wight, l. c. t. 2098, n. 30), caule debili ramisque pubescentibus, foliis omnibus oppositis ovatis acutis basi rotundatis vel subcordatis breviter petiolatis integerrimis, costa nervisque ut in præcedente furcato-subdichotomis, floribus densiuscule glomerulatis, masculis pentandris, perigonio fructifero costato et anguste tetraptero, ♂♀ In India orientali. —

Wedd. Monogr. p. 397, non Bl. Caulis decumbens, parce pubescens demumque glabratus, ramis omnibus in specimine unico suppetente brevibus et fere aequalibus foliisque canescenti-pubescentibus, internodiis 1-1½-centimetalibus. Folia 10-13 millim. longa, 5-8 millim. lata, subaequalia, non aut vix acuminata, nervis fere ut in specie precedente at subtus magis pilosis, petiolo 1-2-millimetro. Stipulae liberae, ovato-lanceolatae, acuminatae. Glomeruli pisiformes, petiolos superantes. Flores masc. breviter pedicellati, perigonio 5-partito (vel 4-partito, teste cl. Wight), segmentis acuminatis extus aequaliter pilosis, pistilli rudimento lanato. Flores fem. pauci, inter masculos latentes; perigonio fructifero ovato 12-costato, costis quatuor lateralibus quam reliquae majoribus et fere aliformibus (Wight). — A precedente praesertim differt foliis minoribus et glomerulis densis. (v. s. typ. Wight.)

B. *Glomeruli plerique in spicam terminalem elongatam aphyllam dispositi. Perigonium ovario adhærens* (Stachyocnide Bl.).

18. P. ARACHNOIDEA, caule erecto fuscescente ramulis hornotinis pubescentibus, foliis omnibus alternis ellipticis acuminatis basi acutiusculis longiuscule petiolatis trinerviis supra sparsim puberulis asperulisque subtus canescenti-tomentosis, inflorescentia elongata spiciformi terminali aphylla, floribus masc. tetrameris, perigonio fructifero alis duabus marginalibus donato. § In ins. Mindoro Philippinarum (Cuming, n. 1492). P. macrura Wedd. Monogr. p. 396; Miq. Fl. Ind. bat. 1 (2). p. 257. Boehmeria arachnoidea Walp. in Nov. act. Acad. cæs. Leop. Carol. 19. suppl. 1. p. 423. Stachyocnide luzonica Bl. Mus. bot. Lugd. bat. 2. 228. t. 55. — Ramuli crassitie pennae anatinæ vel anserinæ, basi denudati glabratiq, apice contra s. usque ad basim inflorescentiæ crebre foliosi pubescentesque, internodiis 2-3 millim. vix longis. Folia ramea fere æquimagna, 5-8 centim. longa, 2-4 centim. lata, patula; nervis basilariibus ad trientem superiorem limbi productis, cæteris primariis (utrinsecus duobus) e costa supra medium ortis; lamina exsiccata membranacea supra fusco-viridi subtus cinerascens; petiolo gracili 2-5 centim. longo pubescenti tomentoso. Stipulae liberae, axillari-laterales, triangulari-subulatae, persistentes. Spica 2-3 decim. longa, ramulum terminans, demum pendula? rhachi gracili angulata pubescente; glomerulis numerosissimis parvis subdiscretisque plerisque masculis (paucissimis androgynis) infimis ex axillis ipsis foliorum superiorum nascentibus stipulisque involucretis, cæteris bractea parva ovata acuminataque suffultis; floribus masc. sessilibus vel breviter pedicellatis, perigonii 4-lobi hispiduli lobis acuminatis et pilis quibusdam hamatis dorso et margine instructis; floribus femineis in axillis inferioribus praesertim obvis sed in inflorescentiis omnibus quas observare licuit rarissimis atque interdum ut videtur prorsus nullis. Perigonium fructiferum ovatum. 2½-3 millim. longum, basi rotundatum, bi-rariusve quadri-alatum, alis basi angustis superne latioribus orem perigonii superantibus apice sinu parvo discretis, utraque facie inter alas angulato-sulcata et undique hispida. (v. s.)

## § 2. — Folia dentata.

19. P. VIMINEA, monoica, ramis virgatis, foliis ovato-vel oblongo-lanceolatis lanceolatisve acuminatis (acumine integro) basi acutis obtusive dentatis vel serratis supra sparsim hispidis lævibusque aut scabriusculis subtus breviter pubescentibus et asperulis, glomerulis axillaribus multifloris saepius androgynis, floribus masc. tetrandris, perigonio fructifero oblongato-ovato superne angustato angulato et obsolete marginato stigma subaequante. § In India orientali, montibus tropicis et temperatis Khasiæ et Himalayæ, altit. 1300-2200 metr. necnon in sylvis montanis Javæ et Sumatræ (Junghuhn, Zollinger, etc.). — P. ovalis Miq. Pl. Jungh. 1. p. 24; Flor. Ind. bat. p. 2. p. 259. Wedd. Monogr. p. 410. P. borbonica Wight Icon. pl. Ind. 6. t. 2100. n. 44. Urtica viminea Wall. Cat. n. 4616. U. sanguinea Bl. Bijdr. p. 501. Boehmeria viminea Wall. mscr. in sched. pl. exs. hort. bot. calcut.; Bl. Mus. bot. lugd. bat. 2. p. 202. B. sanguinea Hassk. Cat. hort. bogor. p. 367; Bl. l. c. B. ovalis Miq. in Zoll. Syst. Verzeichn. 2. p. 100. Margarocarpus vimineus Wedd. in Ann.

sc. nat. 4<sup>e</sup> sér. 1. p. 203. Leptocnide borbonica Bl. l. c. p. 194. Frutex 1-3-metralis, a basi ramosus, ramis elongatis strictis teretiusculis crassitie pennæ anatinæ vel gracilioribus, merithallis 1-2-centimetalibus, epidermide fusca v. cinerascente demum glabrescente. Folia 3-10 centim. longa, 1½-4 centim. lata, vulgo acute etiamque argute serrata, nervis basilariibus fere usque ad apicem limbi protractis, lamina exsiccata rigida supra sæpius fuscescente, petiolo gracili 6-10-millimetro. Stipulæ ovatæ, vulgo breviter acuminatæ, 3-5 millim. longæ, nervo dorsali pilosulæ. Glomeruli pisiformes, floribus masc. (quam feminei sæpius multo numerosioribus) breviter pedicellatis sessilibus, perigonii 4-lobi segmentis breviter acuminatis extus strigosis. Flores fem. sessiles, stigmatibus filiformi quam perigonium lanceolatum paulo longiore. Perigonium fructiferum 2-2½ millim. longum, subcompressum, ore bi-quadridentato, superne præsertim hispido. Achæmium ovatum vel ellipticum, unispermum, millimetricum, acuminato-apiculatum, nitide eburneum. — Secundum cl. Wight stirps hæc in insula africana Borbonia etiam gignitur, at specimina inde allata e planta culta verisimiliter lecta fuerunt. (v. s.)

β, *glabrata*, foliis supra glabriusculis subtus dense puberulis, cæt. ut in typo. γ In sylvis Javæ et Sumatræ. — *Urtica sanguinea*, var. *glabrata* Bl. Bijdr. p. 501. *Boehmeria sanguinea* β *glabra* Bl. mus. bot. lugd. bat. l. c. (v. s.)

γ, *strigosula*, foliis acute vel obtuse crenato-serratis supra hispidulis, subtus pube subtilissima strigosulis. γ In ins. Sumatra. — *Boehmeria strigosula* Bl. l. c. p. 203. — Folia 5-10 centim. longa, 3-4½ centim. lata, anguste acuminata, laxiuscule serrata, rugosula. (v. s. typ. Blum.)

δ, *Waitziana*, foliis argute serratis supra hispidulis subtus appresse puberulis. γ In ins. Java. B. *Waitziana* Bl. l. c. Ramuli petioli nervique subtus strigosi. Folia 5-15 centim. longa, 2-5 centim. lata. (v. s. typ. Blum.)

ε, *Microceltis*, foliis ovato-lanceolatis argute serratis basi rotundatis supra scaberrimis subtus dense molliterque pubescenti-tomentosis. γ In prov. Bandoeng ins. Javæ. P. *Microceltis* Miq. Fl. Ind. bot. 1 (2). p. 260. *Margarocarpus exasperatus* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 204. *Boehmeria Microceltis* Miq. in Zolling. System. Verzeichn. 1. p. 100. B. *sanguinea* ε, minor Bl. l. c. Ramuli et petioli hirtulo-tomentosi. Folia 3-10 centim. longa, 1½-3½ centim. lata. (v. s.)

ζ, *montana*, foliis quam in var. præcedentibus amplioribus ovatis vel elliptico-ovatis basi subcordatis serratis membranaceis supra asperulis subtus scaberrimis subtus pubescentibus. γ In ins. Java (Junghuhn). P. *montana* Miq. Pl. Jungh. 1. p. 23; Fl. Ind. bat. l. c. B. *sanguinea* γ, *montana* Bl. l. c. Folia 16-18 centim. longa, 9-10 centim. lata, planiuscula, petiolo 3-centimetro. (v. s.)

η, *tomentosa*, foliis cordato-ovatis crenato-serratis supra asperrimis subtus molliter pubescenti-tomentosis. γ In ins. Java. *Urtica rubra* Reinwardt, mscr. in hb. propr. *Boehmeria sanguinea* δ, *rubra* Bl. l. c. — Ramuli quam in typo multo crassiores hirtulo-tomentosi. Folia 3½-10 centim. longa, 2-5 centim. lata, coriacea, rugosa. (v. s.)

θ, *fulgens*, foliis quoad figuram illis typi simillimis sed supra glabris et haud raro sparsim glanduloso-impressis subtusque (nervis exceptis) tomento brevissimo nitido incano vel flavescente vestitis. γ In Indiæ orientalis iisdem fere locis ac typus. P. *fulgens* Wedd. mscr. *Urtica* Chiple Hamilt. in Wall. Cat. n. 4616, D. *Urtica parvifolia* ejusd. l. c. *Boehmeria frondosa* Don Fl. Nepal. p. 59? Ramuli pubescentes. Glomeruli masc. ad apices ramulorum interdum spicati, floribus cano-tomentellis. (v. s.)

20. P. *RUBRICULIS*, monoica, ramulis tenuibus rubentibus pilisque sparsis albidis strigosulis, foliis ovato-lanceolatis cuspidatis præter basin et cuspidem lineari-subulatam grosse acute serratis supra sparsim pilosis subtus pubescentibus, glomerulis axillaribus androgynis stipulis majusculis late ovatis involucreis, floribus masc. tetrandris, perigonio fructifero (achænio adnato) ovoideo compresso apice in collum brevissimum contracto obsolete striato

stigma subæquante.  $\frac{5}{8}$  In Java occidentali, prope Tjanjor. *Leptocnide rubricaulis* Bl. l. c. p. 194. t. 57; Miq. Fl. Ind. bat. 1 (2). p. 256. Herba suffruticosa, ramosa, ramulis obtuse quadrangularibus. Folia 4-6 centim. longa, 2-3 centim. lata, membranacea, supra creberrime punctata, nervis subtus rubentibus, petiolo (etiam rubente) 5-15 millim. longo. Stipulæ latissimæ, membranacæ, rubentes, margine superne nervoque dorsali (in acumen protracto) pilis appressis obsitæ. Flores cymoso-fasciculati, masculis pedicellatis, femineis inter binos masculos sessilibus. Alabastra mascula depresso-subglobosa, in vertice obtuse apiculata, hirtula. Perigonium femineum ovoideum, compressum, ore obsolete dentato, superne pilis appressis conspersum, stigmate crasso curvato mox deciduo. Achænium perigonio conforme apiculatum. — A P. ovali, cui, affinis videtur, stipulis insignibus et collo minimo perigonii fructiferi distinguenda.

21. *P. CINERASCENS*, monoica, fruticosa, ramulis canescenti-strigosulis, foliis (nonnihil dissimilibus) lanceolatis vel ovato-lanceolatis ovatisve acuminatis basi obtusis rotundatisve vel rarius subcordatis acute serratis serrulatisve supra sparsim pubescentibus subtus sericeo-velutinis nervis petiolisque pubescentibus, glomerulis androgynis unisexualibusve, perigonio fructifero ovato subcompresso superne attenuato quam stigma superpersistens multoties brevior.  $\frac{5}{8}$  In Sumatra (Korthals) — *Urtica cinerascens* Korth. nœsc. non Bl. *Boehmeria* (Margarocarpus) *cinerascens* Bl. l. c. p. 203. — Ramuli virgati, internodiis 1-æsequi-centimetalibus. Folia 3-13 centim. longa, 2-4 centim. lata, majora cæteris (sive ovatis) multo longius attenuato-acuminata, trinervia, nervis basilaribus arcuatis usque ad basin acuminis excurrentibus supra costaque impressis, lamina exsiccata rigidiuscula subtus canescente, petiolo quam internodium proximum sæpius brevior. Stipulæ triangulari-lanceolatæ, longe acuminatæ, 4-5-millimetales, persistentes. Glomeruli sat multiflori, floribus sessilibus. Perigonium masc. extus strigosum. Flores fem. stigmate valde elongato (5-6-millim.); perigonio fructifero sesquimillimetro superne præsertim hispido, ore sub-4-dentato. Achænium  $1\frac{1}{4}$  millim. longum, ellipticum, eburneum, apiculatum — Stigmate longissimo a præcedente facile distinguitur. (v. s. in hb. Mus. par. ex hb. Mus. lugd. bat.)
22. *P. CONGLOBATA* (Miq. Fl. Ind. bat. 1 (2). p. 260), monoica, ramulis dense hirsuto-tomentosis, foliis (nonnihil dissimilibus) ovatis aut ovato-ellipticis-oblongisve basi rotundatis v. obtusis apice acuminatis vel acutis supra basin integram dentato-vel crenato-serratis supra rugosulis et hispidulo-scaberrimis subtus molliter canescenti-tomentosis, glomerulis androgynis, perigonio fructifero ovoideo superne vix attenuato stigma deciduum subæquante.  $\frac{5}{8}$  In sylvis montanis Javæ occidentalis (Blume). — *Urtica conglobata* Bl. Bijdr. p. 500. *Boehmeria* (Margarocarpus) *conglobata* ejusd. Mus. bot. lugd. bat. 2. p. 204. — Rami annotini teretes, canali medullari (in specim. viso) vacuo, internodiis ramulorum haud centimetalibus. Folia majora ovato-oblonga v. elliptica,  $4\frac{1}{2}$ -10 centim. longa, 2- $4\frac{1}{2}$  centim. lata; minora ovata, 2-4 centim. longa, 2-3  $\frac{1}{4}$  centim. lata, trinervia, exsiccata subcoriacea, petiolo foliorum majorum sec. cl. Blumium 2  $\frac{1}{4}$ -4 centim. longo, illo minorum 5-20-millimetro. Stipulæ brevès, dense pilosæ, deciduæ? Glomeruli pisiformes, multiflori, floribus masc. et fem. canescenti-tomentosis. Perigonium fructiferum sesquimillimetro, erostratum, ore 4-dentato, dentibus triangulari-lanceolatis acutis, stigmate crassiusculo mox deciduo. Achænium late ovatum, pallide fuscum, apiculatum. — Species optima, imprimis perigonio fructifero ovoideo erostrato et stigmate brevi crasso dignoscenda. (v. s. in hb. Mus. par. ex hb. Mus. lugd. bat.)
23. *P. ELEGANS*, monoica, ramosissima, ramulis patulis puberulis, foliis alternis parvis ovatis vel ellipticis oblongisve non aut vix acuminatis acutis basi rotundatis obtusisve et integris cæterum acute serratis vel crenato-serratis lævibus supra sparsim hispido-puberulis subtus subtilissimè puberulis nervis strigosulis, glomerulis androgynis unisexualibusve, perigonio fructifero ovato subangulato superne attenuato stigma subæquante.  $\frac{5}{8}$  In ins. Formosa, ad ripas rivulorum (Wilford) — Ramuli graciles, teretiusculi, foliosi, internodiis subcentimetalibus. Folia 1-3 centim. longa, 1-1  $\frac{1}{4}$  centim. lata, in eodem ramo sæpe diversiformia, minora interdum fere rotundata, valgo in triente

inferiore integerrima, nervis basilariibus circiter ad trientem superiorem limbi nec ultra protensis, lamina exsiccata rigidula subtus glaucescente; petiolo gracili quam merithallus proximus dimidio brevior. Stipulæ ovatæ, acuminatæ, extus puberulæ. Glomeruli ex axillis omnibus orti, semicentimetrum circiter lati, floribus fem. numerosioribus cum masculis sæpius commixtis. Perigonii masc. 4-lobi segmenta breviter apiculato-acuminata, extus puberula. Perigonium fructiferum sesquimillimetrale, ovatum vel ellipsoideum, sparse hispidum, ore 4-dentato. Achænium millimetrale, elliptico-ovatum, longiuscule apiculatum, eburneum. — Fructus *P. ovalis*, cætera diversa. (v. s. in hb. Kew.)

24. *P. PROCRIDIOIDES* (Wedd. Monogr. p. 412), monoica, caulibus simplicibus apice pubescentibus, foliis amplis ovatis v. ovato-ellipticis acuminatis basi cuneatis aut rotundatis integrisque cæterum grosse crenato-serratis longe petiolatis supra sparsim hispido-pilosis subtus in nervis præsertim parce pubescenti-pilosis, glomerulis axillaribus paucis vel multi-floris androgynis, floribus masc. tetrandris perigonio fructifero ovato lævi.  $\frac{1}{2}$  In Promontorio Bonæ Spei (Drege), Abyssinia (Quartin-Dillon), necnon (fide specim. hb. Bonpland) in America tropica. — *Margarocarpus procridioides* Wedd. in Ann. sc. nat. l. c. p. 204. *Urtica* ? *procridioides* Mey. mscr. in sched. pl. Dreg. n. 4597. *Boehmeria* (*Margarocarpus*) Bl. l. c. p. 204. — Caulis subcæspitosus, circiter metrales, erecti vel adscendentes, basi lignosi teretiusculi crassitie pennæ anserinæ vel cynææ denudati internodiis nonnunquam fere decimetralibus, superne magis minusve angulati et herbacei subtomentosi merithallis centimetralibus. Folia 8-12 centim. longa, 5-7 centim. lata, acumine triangulari-lanceolato uni-sesqui-centimetrali acuto, serraturis margine exteriori convexis acutiusculis, lamina exsiccata membranacea, petiolo gracili 4-6 centim. longo pilosulo vel pubescente. Stipulæ late triangulari-ovatæ, 6-10 millim. longæ, anguste acuminatæ, margine et nervo dorsali ciliato-pilosæ. Glomeruli una cum gemma pisiformes, stipulis involucriati, floribus fem. (stigmatem mox deciduo) inter masculos breviter pedicellatos fere solitariis sessilibusque; perigonii masc. 4-partiti segmentis mucronato-acuminatis extus puberulis. Perigonium fructiferum late ovatum, 2 $\frac{1}{2}$  millim. longum, subcompressum, glabratum membranaceum, achænium conforme et æquilongum arcte includens. (v. s. in hb. Mus. par. cæt.)

$\beta$ , *glabrata*, foliis subtus glabris.  $\frac{1}{2}$  (v. s.)

$\gamma$ , *hirsuta*, ramulis, foliis, floribus et stipulis laxè hirsutis.  $\frac{1}{2}$  In Peruvia (Pavon). (v. s. in hb. Boiss.)

25. *P. HETEROPHYLLA*, foliis ovatis ellipticisve (1 $\frac{1}{2}$ -4 centim. longis, 1-2 $\frac{1}{2}$  centim. latis) acutis v. obtusis integerrimis v. obtuse dentato-serratis membranaceis utrinque pilosiusculis, glomerulis axillaribus, floribus masc. triandris, perigonio fem. ovoideo ovario adnato, ore dilatato bidentato.  $\frac{1}{2}$  In oleraceis Butou. *Leptocnide heterophylla* Bl. l. c. p. 194. — Specimen non vidi.

## II. — Species neogæ (1).

26. *P. OBLIQUA* (Wedd. Monogr. p. 405), dioica, ramis superne hirsuto-pubescentibus, foliis alternis inæquilateralibus oblongis vel oblongo-lanceolatis acuminatis basi rotundatis vel subcordatis, costa utrinque nervos primarios daos præter basilares emittente, limbo utrinque pubescenti-piloso supraque asperulo, glomerulis paucifloris, perigonio fructifero ovato aptero et ecostato apice præsertim hirsuto.  $\frac{1}{2}$  In sylvis prope Panama (Sutton Hayes, n. 350) et Santa Marta Novo Granatensium (Purdie); in Peruvia, prov. Maynas (Poeppig, n. 2032). *Boehmeria obliqua* Poepp. mscr.; Bl. l. c. p. 206, in adnot. — *Margarocarpus obliquus* Wedd. l. c. p. 204. Frutex 3-4-metralis. Rami teretes, basi denudati glabratique et epidermide castanea vestiti, ramulis gracilibus diffusis subdistiche foliosis, internodiis 1-3-centimetralibus, epider-

(1) Vide etiam, inter species sectionis præcedentis foliis dentatis, *P. procridioidem*.

vide subhirsuta etiam castanea. Folia 5-11 centim. longa,  $2\frac{1}{2}$ - $3\frac{1}{2}$  centim. lata, inferiora superioribus multo minora potiusque elliptica quam oblonga aut lanceolata, lamina exsiccata submembranacea fusca hirsutæ paginæ inferioris molliuscula, petiolo 3-6 millim. longo hirt. Stipulæ subulato-triangularis, subcentimetricæ, pilosæ, persistentes. Glomeruli in speciminibus visis omnes feminei, floribus sessilibus, stigmate filiformi 6 millim. longo dense lutescenti-villoso. Perigonium fructiferum  $2\frac{1}{2}$  millim. longum, subcompressum, basi anguste marginatum, achænium ovatum subacuminatum 2 millim. longum sordide eburneum nitidumque arcte includens.

27. *P. OVATA*, monoica, caule s. ramis apice puberulis pilosulisve, foliis alternis ovatis basi obtusis integerrimis nervis primariis præter basilares utriusque pluribus, lamina concolore utrinque glabra lævique margine et nervis paginæ inferioris sparsim et adpresse hispido-pilosulis demumque glabratis, glomerulis axillaribus paucifloris, floribus masc. tetrandris, perigonio fructifero ovato apice attenuato aptero et ecostato parce strigoso.  $\frac{5}{3}$ ? In Nova Granata, prope Cuello (Goudot). *P. glabrata* Wedd. Monogr. p. 406. *Margarocarpus ovatus* Wedd. in Ann. sc. nat. l. c. p. 204. *Boehmeria ovata* Bl. l. c. p. 206, in adnot. Ramus unicus mihi visus bidecimetralis, herbaceus, arcuatus gracilis, in axillis inferioribus ramulis parvis manitus, internodiis 2 centim. longis. Folia 5 centim. circiter longa,  $2\frac{1}{2}$  centim. lata, sensim acuminata, acuta; nervis parum prominentibus, primariis utrinsecus præter basilares 3-5 inæqualibus (nempe 2-majoribus) et sæpe parum conspicuis; lamina exsiccata membranacea læte viridi; petiolo 10-15 millim. longo puberulo. Stipulæ lanceolatæ, 5-8-millimetricæ, subulato-acuminatæ, nervo dorsali et margine parce pilosæ, persistentes. Glomeruli in specimine examinato grani piperini circiter magnitudine, floribus utriusque sexus commixtis. Flores masc. subsessiles, perigonii 4-partiti segmentis breviter mucronato-acuminatis extus sparsim strigosis pilisque quibusdam hamatis hispidis. Flores fem. sessiles, stigmate filiformi bimillimetrici. Perigonium fructiferum 2 millim. longum, acutiusculum, apice penicillato-pilosum, achænium conforme eburneumque arcte vestiens. — Nomen specificum in monographia mea mutaveram propter similitudinem cum nomine *P. ovalis*. (v. s. in hb. Mus. par.)
28. *P. FORMICARIA* (Wedd. Monogr. p. 407), dioica, fruticosa, ramis puberulis, foliis alternis ovatis ellipticisve acuminatis basi obtusis aut subcordatis planis integris vel prope apicem repandis, costa utrinque nervum primarium unicum præter basilares emittente, limbo supra glaberrimo subtus in nervis præsertim pilosulo, glomerulis in spicas axillares dispositis, spicis quam folia brevioribus patulis simplicibus, perigonio fructifero lanceolato carnosulo subangulato apicem versus sensim attenuato pubescenti-hispido.  $\frac{3}{3}$  In prov. Maynas Peruviae (Pöppig). *Boehmeria formicaria* Pöppig, mscr. in sched. pl. exs. n. 2287; Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 201; Bl. mus. bot. lugd. bat. 2. p. 225. t. 56, A. Frutescens, ramulis flexuosis apice puberulis atque hinc et inde pilosulis canali medullari in specimine examinato vacuo. Folia 6-10 centim. longa,  $2\frac{1}{2}$ -5 centim. lata, denticulis parum conspicuis, nervis basilaribus arcuatis usque ad basim acuminis interdum productis subtus prominulis venulisque longiuscule et patentim pilosulis, lamina exsiccata subcoriacea pagina superiore cystolithis punctiformibus conspicuis ornata, petiolo vix centimetrici pilosulo. Stipulæ in specimine viso jam demissæ. Spicæ fem. (solum visæ) 5-10 centim. longæ, rhachi pubescente, glomerulis discretis pisi circiter magnitudine bractea ovato-acuminata puberula suffulta. perigonio quam stylus robustus (in stirpe exsiccata flavescenti-villosus) multo brevior, ore subquadridentato. Ovarium liberum. Fructus desideratur. (v. s.)
29. *P. DISCOLOR* (Wedd. Monogr. p. 408. t. 13. B. f. 18-24, excl. syn. *Margarocarpus asperi* ejusd.), monoica, ramulis apice canescenti-tomentosis, foliis alternis lanceolatis sensim acuminatis basi obtusis v. retusis integerrimis supra adpresse pubescentibus vix asperulis, nervis primariis utrinque 2-3 inclusis basilaribus, glomerulis axillaribus unisexualibus androgynisve, perigonii masc. segmentis lanceolato-triangularibus, perigonio fructifero ovato-lanceolato a medio sensim acuminato s. rostrato, stigmate longissimo.  $\frac{3}{3}$  In Peruvia subandina Pöppig n. 1284. — *Boehmeria discolor* Pöpp. mscr.; Bl.



- l. c. p. 206. *Margarocarpus Poeppigianus* Wedd. in Ann. sc. nat. l. c. p. 204. — Ramuli elongati, herbacei, pubescentes, internodis centimetralibus. Folia majora 12-15 centim. longa, 4 centim. lata, exsiccata chartacea venulis transversalibus distinctis, petiolo centimetr. longo. Stipulae 1 centim. longae, longe acuminatae. Flores utriusque sexus hirtuli, masculi 4-meri. Stigma plus quam centimetrale. Perigonium fructiferum in dimidia parte superiore angustatum, nec angulatum nec marginatum, oris bi-quadridentati laciniis subulatis erectis v. erecto-patentibus. Achænium elongato-ovoideum, nitidum, albidum, perigonio dimidio brevius. (v. s.)
30. *P. ASPERA*, monoica, ramosa, ramulis apice pubescentibus, foliis alternis oblongo-lanceolatis basi acutis obtusiusculisve sensim acuminatis acutisve integerrimis, costa præter nervos basiales utrinque unum vel duos primarios emittente, limbo supra hispido exasperatoque subtus incano-tomentoso, glomerulis axillaribus unisexualibus androgynisve, floribus masc. tetrandris, perigonio fructifero lato ovato apice subtruncato.  $\frac{3}{4}$  in Peruvia (Cl. Gay); in fructicetis provinciae Yungas Boliviae (Wedd.) — *P. discolor* Wedd. Monogr. p. 408, ex parte, t. 13. A. f. 18. *Margarocarpus aspera* ejusd. in Ann. sc. nat. l. c. *Boehmeria aspera* Bl. l. c. p. 206, in adnot. — Frutex metralis, ramis elongatis crassitie calami scriptoris, nodosis, velutinis, cortice fusco siccitate longitudinaliter corrugato, ramulis lateralibus brevibus ligneis adscendentibus. Folia majora, 6 centim. longa, breviter acuminata, nervis basilaribus ad quartam partem superiorem limbi v. altius productis, tertiariis s. venulis transversis tenuissimis, lamina exsiccata rigida nervis supra impressis subtus prominulis pallideque ferrugineis intervenio candido, petiolo 3-6 millim. longo. Stipulae triangulari-ovatae, circiter 3 millimetrales, acuminatae, puberulae, exsiccatae ferrugineae, persistentes. Glomeruli masc. granum piperinum magnitudine æmulantes, floribus subsessilibus bracteis parvis in ramo denudato diu persistentibus suffultis, perigonii 4-lobi segmentis ovatis breviter acuminatis extus strigosus. Glomeruli fem. masculis nonnihil majores, floribus sessilibus, perigonio fructifero fere rotundato 2-2 $\frac{1}{2}$  millim. longo apice subtruncato. Achænium ovatum, acutum, nitide eburneum. (v. v.)
31. *P. LAEVIS*, monoica, caulis caespitosis simplicibus virgatis apice puberulis aut fere omnino glabris, foliis ovato- vel oblongo-lanceolatis ellipticisve basi acutis integerrimis nervis ut in precedente dispositis limbo supra sparsim et breviter pilosulo laevique subtus incano-tomentoso margine vix hispido, glomerulis unisexualibus androgynisve axillaribus aut in apice caulium subspicatis pauci-multi-floris, floribus masc. tetrandris, perigonio fructifero lato ovato erostrato obsolete costato marginato, stigmatè brevi.  $\frac{3}{4}$  in sylvis locisque umbrosis prov. brasiliensis Matto Grosso (Wedd.) — *P. virgata* Wedd. Monogr. p. 409. *Margarocarpus laevis* Wedd. in Ann. sc. nat. l. c. *Boehmeria laevis* Bl. l. c. p. 206. — Caules e rhizomate crasso lignoso radiculis carnosus fusiformibus aucto fasciculatim prodeuntes, circiter metrales, simplicissimi, basi lignescens teretiusculi aphylli crassitieque pennae anserinae vel cycneae, superne herbacei gracilisscentes magisque aut minus angulari, internodiis inferioribus elongatis. Folia forma et magnitudine in speciminibus diversis variabilia, in aliis scilicet elliptica 8-10 centim. longa 3 $\frac{1}{2}$ -4 centim. lata, in aliis autem anguste lanceolata 12-15 centim. longa et vix 2 centim. lata, superiora minima et interdum bracteiformia, omnia sensim acuminata, lamina exsiccata membranacea supra saepe fuscescente nervis subtus nigricantibus, petiolo centimetrali vel breviorè. Stipulae anguste triangulari-lanceolatae, 4-5 millim. longae, sensim subulato-attenuatae, persistentes. Glomeruli inferiores plerumque masculi, superiores androgyni, floribus breviter pedicellatis sessilibusve. Flores masc. perigonio 4-partito, segmentis extus parce strigosulis breviter acuminatis acumine hirtis. Flores fem. pauci, inter masculos fere occulti, perigonio ovato crasse et subadpresse piloso, stigmatè crasso et mox deciduo. Perigonium fructiferum sesquimillimetre, latitudine fere eadem, apice breviter bidentatum subacute marginatum strigosum. Achænium ovatum, millimetre, acutum, eburneum. (v. v.)
32. *P. GUATEMALANA*, monoica, ramosa, ramulis puberulis, foliis lanceolatis acuminatissimis basi obtusis v. acutis integerrimis supra imprimis ad mar-

ginem puberulis v. tandem glabriusculis subtus cano-tomentosis, glomerulis axillaribus et lateralibus androgynis, floribus masc. tetrandris  $\frac{7}{8}$ , perigonio fructifero ovoideo constricto-attenuato obsolete striato strigoso.  $\frac{3}{4}$  In Guatemala (Friedrichstal, n. 1272). — *Boehmeria guatemalana* Bl., l. c. p. 206. — Rami virgati, foliosi s. foliorum cicatricibus tuberculati, inferne glabrati, cortice pubescente vestiti, ad apices subpubescentes. Folia  $6\frac{1}{2}$ –15 centim. longa  $2\frac{1}{4}$  centim. lata, membranacea, nervis subtus petioloque (2-4-centim. metrali) subpubescentibus. Perigonium fructiferum haud marginatum, ore quadridentato, stigmate mox deciduo. — A præcedente imprimis habitu bene distincta.

33. *P. OCCIDENTALIS* (Wedd. Monogr. p. 410), monoica, ramis elongatis dense puberulis vel inferne demum glabratiss, foliis alternis elliptico-v. lanceolato-ovatis acuminatis basi cuneatis aut fere rotundatis integerrimis longiuscule petiolatis supra pilosulis subtus margineque molliter pubescentibus, glomerulis androgynis rariusve unisexualibus, floribus masc. tetrandris segmentis perigonii abrupte aristato-acuminatis, perigonio fructifero lævisculo v. multicostrulato.  $\frac{3}{4}$  In Nicaragua (Orsted), Nova Granata (Orsted, Karsten), et Venezuela, ad Biscaina, altit. 1000 metr. (Fendler, n. 1263). — *Leucococcus occidentalis* Liebm. in Kgl. danske Vidensk. Selsk. Skr. 5 Række, naturv. og math. Afd. 2. p. 311. — Ramuli teretes, inferne sæpius denudati, penna corvina nonnihil crassiores, epidermide in planta exsiccata cinereo-rubente, merithallis superioribus centimetrum circiter longis. Folia 5-12 centim. longa,  $2\frac{1}{4}$ – $6\frac{1}{2}$  centim. lata, anguste acuminata, nervis basilaribus dimidium limbum superantibus, nervis primariis præter basilares utrinque duobus, lamina exsiccata membranacea et fere concolore, petiolo gracili 2-8-centim. metrali longe pubescente. Stipulæ lanceolatae, acuminatae, pilosæ, subcentim. metrales. Glomeruli pauci-multiflori. Flores masc. sessiles v. breviter pedicellati, perigonii 4-lobi segmentis pubescentibus, acumine subulato fere millim. metrali. Perigonium fem. puberulum stigmate mox deciduo paulo brevius; fructiferum late elliptico-ovoideum, sesquimillim. metrale, a basi subtruncata fere usque ad apicem obtusiusculum, striis aut costis 10-15 subæqualibus vel paucis brevioribus interpositis percursum, ore bidentato. Achænium ovatum, perigonio parum brevius, apiculatum, nitide eburneum. — Insignis inter species neogeas perigonio fructifero costulato. (v. s. in hb. DC.)

34. *P. ACALYPHOIDES*, monoica, ramulis pubescentibus hirtulisve, foliis alternis oblongo v. ovato-lanceolatis basi obtusis acutisve acuminatis integerrimis supra parce breviterque hirtulis subtus imprimis in nervis dense pubescentibus hirtulisve, floribus in glomerulos spicatos digestis (spicis apice nudis v. foliatis), masculis tetrandris, perigonio fructifero ovoideo sublenticulari-compresso ecostato.  $\frac{3}{4}$  In Venezuela (Bredemeyer; Fendler n. 1249 et n. 1264). *Boehmeria acalyphoides* Bl. l. c. p. 225. Ramuli teretes, graciles, inferne fere glabrati, internodiis sæpe elongatis. Folia 6-12 centim. longa, 2-5 centim. lata, longiuscule acuminata, nervis basilaribus ad trientem superiorem limbi protractis, lamina exsiccata membranacea concolore nervis nervulisque subtus bene conspicuis, petiolo gracili pubescenti-hirto 1-5-centim. metrali. Stipulæ lanceolatae, subulato-acuminatae, pilosæ. Spicæ terminales, simplices, foliis multo longiores, glomerulis parvis discretis bractea ovato-lanceolata acuminata pubescente suffultis androgynis unisexualibusve, quibusdam ad basin spicæ ex axillis foliorum superiorum etiam enatis. Perigonii masc. 4-lobi lobis acuminatis apice hirtulis. Perigonium fem. strigosum, stigmate paulo brevius; fructiferum bi-millim. metrale, apice constrictum ore bidentato. Achænium perigonio conforme eoque arcte vestitum, nitide eburneum, apiculatum. — Species inflorescentia notabilis in sectione propria (Urocnide) generis *Boehmeria* a cl. Blumio collocata. (v. s. in hb. DC.)

35. *P. LATIFOLIA*, dioica? ramulis elongatis crassis cinerascensibus, foliis amplis late ovatis acuminatis basi truncato-subcordatis integerrimis supra puberulis subtus (nervis exceptis) cano-tomentosis, glomerulis axillaribus, floribus masc. tetrandris.  $\frac{3}{4}$  In imperio mexicano (herb. Pavon). — Ramulus suppetens teres, crassitie calami scriptorii, epidermide cinereo-rubente, dense puberulus et sparsim pilosulus, merithallis bi-tri-centim. metralibus. Folia 6-10 centim.

longa, 4-6½ centim. lata, subabrupte acuminata, nervis basilaribus ad partem sextam superiorem limbi productis, nervis primariis præter has utrinque 3 sicut costa pubescentibus et in sicco rufescentibus, lamina exsiccata membranacea supra viridi, petiolo 2-4½ centim. longo pubescenti-hirtulo. Stipulæ lineari-lanceolatæ, subpersistentes. Glomeruli visi omnes masculi, vix multiflori, floribus sessilibus, perigonii quadrilobi horumce lobis pubescentibus breviter acuminatis. Cætera ignota. (v. s. in hb. Boiss.)

*Species exolusæ.*

- P. ACUTA Wight = Memorialis ambigua.
- P. AMBIGUA Wight = Memorialis ambigua.
- P. ASPERA Wight = Memorialis scabra.
- P. BENNETTIANA Wight = Memorialis hirta.
- P. BRACTEATA Benn. = Memorialis bracteata.
- P. CAUDATA Benn. = Memorialis caudata.
- P. CONCINNA Wight = Memorialis concinna.
- P. COURTALLENsis Wight = Memorialis caudata.
- P. DALZELLII Wight = Memorialis pentandra.
- P. FLACCIDA A. Rich. = Australina Schimperiana.
- P. GARDNERI Wight = Memorialis hirta.
- P. GLABRA Wight = Memorialis aquatica.
- P. HETEROCARPA Wight = Memorialis hirta.
- P. HIRTA Hassk. = Memorialis hirta.
- P. HISPIDA Benn. = Memorialis hirta.
- P. HYPERICIFOLIA Bl. = Memorialis pentandra.
- P. INTEGRIFOLIA Dalz. = Memorialis integrifolia.
- P. LONGIFOLIA Wight = Memorialis ambigua.
- P. LUCIDA Hassk. = Memorialis triandra.
- P. MYSORENSIS Wight = Memorialis hirta.
- P. NEILGHERRENSIS Wight = Memorialis Nilghirensis.
- P. ORLONGIFOLIA Wight = Memorialis Nilghirensis.
- P. OVATIFOLIA Wight = Memorialis hirta.
- P. OVATA Wight = Memorialis Nilghirensis.
- P. QUINQUENRIS Benn. = Memorialis quinquenervis.
- P. PARVIFOLIA Wight = Memorialis parvifolia.
- P. PENTANDRA Benn. = Memorialis pentandra.
- P. QUADRIALATA Wight = Memorialis hirta.
- P. RAMOSISSIMA Wight = Memorialis pentandra.
- P. REPENS Griseb. = Boehmeria repens.
- P. REVOLUTA Bl. = Memorialis hirta.
- P. SCABRA Wight = Memorialis scabra.
- P. SONNERATII Gaudich. = Phenax Sonneratii.
- P. SPONIEFOLIA Schlechtend. = Memorialis hirta.

- P. STOCKSII Wight = *Phenax pentandra*.  
 P. TERNATA Benn. = *Phenax concinna*.  
 P. TOMENTOSA Wight = *Phenax hirta*.  
 P. TRIALATA Wight = *Phenax Nilghirensis*.  
 P. TRIANDRA Bl. = *Phenax triandra*.  
 P. WAITZIANA Bl. = *Phenax hirta*.  
 P. WALKERIANA Wight = *Phenax aquatica*.  
 P. WALLICHIANA Wight = *Phenax Nilghirensis*.  
 P. WIGHTII Benn. = *Phenax Nilghirensis*.

XX. MEMORIALIS Hamilt. mscr. in Wall. Cat. n. 1598 et n. 4601 (anno 1831); Wedd. Monogr. p. 415; Benth. Fl. hongk. p. 332. — Pouzolzia spec. Benn. Pl. jav. rar. p. 66; Wight Icon. pl. Ind. 6. p. 36; Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 205; Bl. Mus. bot. lugd. bat. 2. p. 239 et seq.; non Gaudich. — Gonostegia Turcz. in Bullet. Soc. imp. nat. Mosc. 19. (anno 1846) n. 4. p. 509. — Hyrtanandra Miq. Pl. Junghuhn. 1. p. 25. (anno 1851); Fl. Ind. bat. 1 (2), p. 260.

Flores monoici rariusve dioici, utriusque sexus sæpius commixti, bracteis scariosis parvis stipati glomerulosque axillares aut spicatos constituentes. Masc. Perigonium 4-5-rariusve 3-partitum, segmentis (in præfloratione valvatis) ovatis breviter acuminatis juxta tertiam v. quartam partem superiorem abrupte infractis cristaque s. plica transversa sæpissime ciliata munitis, alabastro ut plurimum campanulato s. turbinato apiceque conjunctione acuminum segmentorum mucronato. Stamina tot quot perigonii segmenta. Pistilli rudimentum lineari-conicum, glabrum. Fem. Perigonium, ovarium, stigma et achænium ut in Pouzolzia, sed pericarpium semper atrum alæque perigonii (quum adsunt) plerumque marginales. — Herbæ perennes v. suffrutescentes, omnes in India orientali v. in insulis maris indici obviæ, glabræ aut varie pubentes, sæpe asperatæ. Folia opposita v. ternata aut sæpissime in eodem caule superiora alterna, homomorpha et æquilatera, integerima, tri-tripli-quintuplinervia, nervis basilaribus a basi ad apicem limbi productis simplicibusque, cystholithis punctiformibus. Stipulæ axillares v. axillari-laterales et liberæ aut utrinque cum illis folii oppositi connatæ s. interpetiolares.

A. — Folia floralia (saltem superiora) cæteris multo minoræ sæpissimeque bracteiformia.

\* Flores masculi pentandri.

1. M. PENTANDRA, caule plerumque erecto glabrato aut in angulis villosulo sæpius valde ramoso, foliis inferioribus semper oppositis lineari-lanceolatis lanceolatisve basi rotundatis vel subcordatis, superioribus ramulorumque s. floriferis ut plurimum alternis et magnitudine valde diminutis, omnibus trinerviis utrinque glabris (rariusve subtus puberis) margine nervisque subtus brevissime et subadpresse spinuloso-setulosis, floribus masc. pentandris, perigonio fructifero late bialato vel subtrialato. 4-5 In locis humidis lapidosisque Indiæ orientalis tropicæ (præsertim borealis et occidentalis) frequens necnon in ins. Java orientali et Philippinis. — M. ciliaris Hamilt. in Wall. Cat. n. 4598; Wedd. Monogr. p. 417. Urtica pentandra Roxb. Fl. Ind. 3. p. 583; Wall. Cat. n. 4598. excl. B; Wight Icon. pl. Ind. 2. p. 696. Pouzolzia pentandra Benn. Pl. jav. rar. p. 66. t. 14; Wight l. c. 6. t. 2096. n. 20; Bl. Mus. bot. lugd. bat. 2. p. 242. Gonostegia oppositifolia Turcz. l. c. p. 510. Hyrtanandra pentandra et H. javanica Miq. Pl. Jungh. 1. p. 25; Fl. Ind. bat. 1 (2). p. 26.

— Caulis erectus v. adscendens, altitudine et crassitie admodum variabilibus, magis minusve angulatus, exsiccatus sæpissime ferrugineus, ramis virgatis aut diffusis angulatis ut caulis ipse hirtellis hispidulisve (in angulis præsertim) vel glabris, merithallis floriferorum brevissimis. Folia caulina magnitudine admodum varia, modo decimetralia et centimetrum vix lata, modo quintuplo minora, acuta vel acuminata rariusve obtusa, sessilia vel breviter petiolata (petiolo 1-3 millim. longo), margine nervisque subtus minute spinulosi asperatisque, lamina exsiccata rigida pallideque viridi. Stipulæ breves, ovatae, liberae vel inter petiolos connatae. Glomeruli pauci-multi-flori, bractea multo breviores, floribus utriusque sexus intermixtis. Flores masc. pedicellati, perigonii 5-partiti segmentis breviter acuminatis (acumine pilosulo), margine acuto plicæ transversalis s. infractionis pilis hamatis magis minusve ciliato, alabastro campanulato. Flores fem. sessiles aut pedicellati. Perigonium fructiferum ut plurimum bi-tri-alatum (raro apterum) apiceque et basi emarginatum, alis lateralibus semiorbicularibus 2-3 millim. longis membranaceo-subcoriaceis, intermedia (quum adest) minore. (v. s.)

β, *Stocksii*, foliis inferioribus oblongo-lanceolatis aut ellipticis sæpe obtusis, floralibus lanceolatis, perigonio fructifero bi-tri-alato. 4-5 In India orientali. Pouzolzia Stocksii Wight, l. c. t. 2097, n. 28, sub P. ramosissima. Caulis debilis, prostratus aut inter fruticeta scandens, sæpius valde ramosus. (v. s.)

γ, *hypericifolia*, foliis inferioribus lineari-lanceolatis acuminatis basi subcordatis, floralibus ovalibus oblongisve obtusis, perigonio fructifero aptero nervoso-striato apice attenuato edentulo. 4-5 In Sina australiori. Pouzolzia hypericifolia Bl. l. c. p. 242. — Folia inferiora 2½-4 centim. longa, 12 millim. lata, superiora sensim minora sed cœnformia, petiolo 1-2-millimetrali.

δ, *ramosissima*, foliis inferioribus s. caulinis ovato-lanceolatis, floralibus cordatis, perigonio fructifero sæpe trialato, ala intermedia incrassata. 4-5 In India orientali. Pouzolzia ramosissima Wight, l. c. t. 2095, n. 17. Caulis erectus v. decumbens aut subscandens, ramosissimus. Folia quam in typo pro longitudinis ratione latiora. (v. s.)

ε, *Dalzellii*, caule procumbente parce ramoso, cæt. ut in var. præced. 4-5 In India orientali. Pouzolzia integrifolia β Dalz. mscr. in hb. Kew. P. Dalzellii Wight, l. c. t. 2096, n. 21. Perigonium fructiferum bi-tri-alatum aut (sicut in præcedente) fortuito apterum et nervoso-costulatum. (v. s.)

2. M. AQUATICA (Wedd. Monogr. p. 418), caule erecto teretiusculo apice pubescente parce ramoso, foliis inferioribus oppositis lanceolatis vel ovato-lanceolatis acuminatis basi rotundatis breviterque petiolatis trinerviis glabris (margine tantum hispido-pilosulis) vel utrinque sparsim pilosis lævibus aut asperulis, floralibus superioribus bracteiformibus lineari-lanceolatis, floribus masc. pentandris, perigonio fructifero bialato aut aptero costatoque. 4 In Zeylaniam, ad ripas et in locis demissis (Moon, Walker, Thwaites, Macrae, Gardner). Urtica pentandra Wall. Cat. n. 4598. B. U. aquatica Moon, Cat. pl. Ceyl. p. 62. Pouzolzia Walkeriana Wight, l. c. t. 2095, n. 16. — Caulis erectus aut adscendens, elongatus, parce ramosus, superne gracilescens, inferne teretiusculus glaberrimusque, crassitie pennæ anserinæ v. anatinæ, nonnunquam glaucescens, angulatus hispidulusque aut pubescens, internodiis inferioribus frequentissime semidecimetrum longis superioribus brevissimis. Folia sterilia (s. inferiora) 5-18 centim. longa, 1½-5 centim. lata, sæpius sensim longiusculeque acuminata, inflorescentiam versus gradatim vel subabrupte diminuta; floralia a basi ad apicem spicæ (sæpe longissimæ) plerumque sensim imminuta, majora 5-8 centim. longa centimetrumque fere lata, minora vix centimetralia, exsiccata læte viridia; petiolis foliorum inferiorum 5-8 millim. longis, illis bractearum circiter bimillimetralibus. Stipulæ ovatae, liberae vel connatae, amplexicaules, caducæ. Glomeruli pauci-multi-flori, petiolos paulo superantes, floribus utriusque sexus in glomerulis distinctis congregatis (staminiferis inferius sitis) vel commixtis. Flores masc.

lævisque subtus in nervis appresse pilosulis intervenio glabro, glomerulis axillaribus pisiformibus.  $\frac{3}{4}$  In silvis montanis ins. Javæ, alt. 1000–1600 metr. (Zollinger, n. 3190 et 3480, cæt.) *C. rufescens* Wedd. Monogr. p. 436; Miq. l. c. p. 263. *Urtica lutescens* Bl. Bijdr. p. 494. *Dendrocnide* ? *cærulea* Miq. Pl. Jungh. 1. p. 31. *Boehmeria lutescens* Bl. Mus. bot. lugd. bat. 2. p. 209. — Ramuli obtuse tetragoni, superne strigosuli, mox glaberrimi, exsiccati fusco-rubrescentes. Folia 4–10 centim. longa,  $1\frac{1}{2}$ –3 centim. lata, breviter acuminata, acumine totoque limbo (ima basi excepta) serraturis triangularibus æqualibus dentata, nervis basilaribus usque ad trientem superiorem limbi solum productis cum sequentibus aruatim anastomosantibus, lamina exsiccata rigide coriacea valde rugosa, nervis primariis supra impressis, intervenio subtus læu-noso-foveolata, petiolis longitudine subæqualibus 5–20-millimetralibus ad-pressis pilosulis. Stipulæ lanceolatæ, 10–15 millim. longæ, vix dorso pilosule, deciduæ. Glomeruli masc. alabastris apice pilosulis (teste Miquel). Glomeruli fem. pisiformes, ramulis quam in *C. macrophylo* brevioribus floribusque igitur paucioribus, perigonio elliptico superne angustato haud gibboso stylo fere a basi longiuscule piloso. Perigonium fructiferum rotundatum s. lenticulare, margine obtuso, ore 4-dentato. Achænium millimetrale, perigonio conforme eodemque arcu inclusum, pericarpio nucamentaceo superne præsertim incrassato. (v. s.)

6. *C. CÆRULEUS*, monoicus, foliis subæqualibus aut disparibus oblongo-lanceolatis ellipticisve acutis v. acuminatis excepta basi (acuta v. obtusa) toto margine (acumine incluso) crenulato-serrulatis planis supra glaberrimis et lævis subtus in nervis pubescentibus intervenio glaberrimula, glomerulis utriusque sexus axillaribus unisexualibus pisiformibus.  $\frac{3}{4}$  In Moluccis (Reinwardt, Blume). — *C. Blumeanus* Wedd. Monogr. p. 437; Miq. l. c. *Urtica cærulea* Bl. Bijdr. p. 495. *Boehmeria cærulea* Bl. Mus. bot. lugd. bat. 2. p. 209. — Ramuli graciles, obsolete angulati v. teretiusculi, superne pubescentes, exsiccati epidermide fusca obtecti, in vivo autem (teste cl. Blumeo) cærulei, internodiis 1–2-centimetralibus. Folia in eodem ramo (saltem in specimenibus visis) asymmetrica, subheteromorpha et inæquimagna, 3–8 centim. longa, 15–30 millim. lata, acumine sæpe obliquo, dentibus obtusiusculis et sæpe inæqualibus, nervis basilaribus ut in præcedente, lamina exsiccata membranacea fusco-virente nervis supra vix impressis, petiolis plerumque inæquilonis gracilibus pubescentibus. Stipulæ lanceolatæ, 5-millimetrales, nervo dorsali et margine ciliolatæ. Glomeruli fem. (solum visi) fere ut in specie præcedente sed perigonio achænioque ut videtur sæpius obovatis nec lenticuliformibus, pericarpii pariete superiore crassissimo et dimidia scilicet parte achæni totius constante. (v. s.)

7. *C. MELANOCARPUS* (Miq. l. c. p. 263), dioicus, foliis æqualibus vel paullo disparibus lanceolatis v. oblongo-lanceolatis acuminatis supra basim acutam integramque obtuse serratis planis utrinque glabris v. in nervis subtus parce appresse puberulis, glomerulis utriusque sexus axillaribus pisiformibus.  $\frac{3}{4}$  In Nova Guinea. — *Boehmeria melanocarpa* Bl. l. c. p. 210. — Aspectus præcedentis. Ramuli glabrati aut apice sparsim appresse puberuli. Folia 4–13 centim. longa,  $1\frac{1}{2}$ –3 $\frac{1}{2}$  centim. lata, acumine angusto sicut reliquis limbus (basi excepta) serrato, lamina exsiccata membranacea, petiolis gracilibus inæqualibus  $\frac{3}{4}$ –2 $\frac{1}{2}$  centim. longis glabris v. sparsim appresse puberulis. Stipulæ lanceolatæ, deciduæ. Glomeruli subconfluentes, illis *C. Blumei* aspectu similes; masculi ignoti. Perigonium fructiferum obovoideo-cuneatum, subangulatum aut varie compressum. (v. s.)

8. *C. ELLIPTICUS*, monoicus, foliis vix disparibus ellipticis v. obovato-ellipticis obtusis acutiusculisve basi subcuneatis integrisque reliquo margine inæqualiter et obtusiuscule serrulato rugosulis supra glabris lævisque subtus in nervis appresse pilosulis intervenio puberulo v. glabrato, glomerulis unisexualibus axillaribus pisiformibus.  $\frac{3}{4}$  In insula Java (de Vries). Ramuli graciles, angulati, superne appresse puberuli, internodiis sesqui-bi-centimetralibus. Folia 2–8 centim. longa, 13–35 millim. lata, nervis basilaribus ad partem quintam superiorem limbi productis margini parallelis, lamina exsiccata rigida, petiolis gracilibus inæqualibus 1–2 centim. longis appresse pilosulis

puberulisve. Stipulæ lanceolatæ deciduæ. Glomeruli illis specierum duarum præcedentium aspectu simillimi, florum masculorum perigonio parce hirtulo, segmentis sub apice acuminato haud mucronatis. Perigonium fructiferum obovatum compressum ore bidentato achænium fere lenticulare millimetrale arcte vestiens, stigmatem quam perigonium circiter dimidio brevius. (v. s.)

9. *C. PROSTRATUS*, ramulis dense hirsutis, foliis valde disparibus argute dentato-serratis rugosis supra hirtis-asperis subtus hirsuto-pubescentibus, majoribus ellipticis v. oblongo-ellipticis acutis v. acuminatis basi obtusis breviter petiolatis, minoribus ovatis ellipticisve obtusis v. acutis basi rotundatis sessilibus, stipuliæ profunde bipartitis longissima subulato-acuminatis.  $\delta$  Ad rupes in Nova Guinea (Zippel). — *Boehmeria prostrata* Bl. l. c. p. 209. — Caulis prostratus, ramis subflexuosis sæpe radicanibus. Folia majora 7-1½ centim. longa, 4-8 centim. lata, exsiccata submembranacea, dentibus acuminatis, petiolo dense hirsuto semicentimetrali; minora 1½-3 centim. longa. Stipulæ dorso v. persistentes. Flores ignoti, (v. s.)

**XXII. TOUCHARDIA** Gaudich. Bot. Voy. Benite, t. 94; Wedd. Monogr. p. 441.

Flores dioici, densissime cymoso-capitati, capitulis pedunculis axillares subsimplices furcatosve v. dichotome ramosos terminantibus. *Masc.* Perigonium 5-partitum, segmentis fere æqualibus ellipticis apice membranaceis puberulis obtusiusculis in præfloratione imbricatis, alabastro umbilicato. Stamina 5. Pistilli rudimentum obovato-clayatum, glabrum. *Fem.* Perigonium subcampanulatum, 4-lobum, lobis sæpe inæqualibus magis minusve angulatis interdum cucullatis carnosulis glaberrimis. Ovarium ovoideum, rectum, perigonio fere æquilongum. Ovulum plerumque obliquum, in funiculum basi loculi affixum adscendentemque plus minus inflexum. Stigma spathulatum, facie altera marginibusque longiuscule papilloso-ciliatis, stylo brevi suffultum. Achænium ovatum, ventricosum, læve, perigonio carnosum et accreto diu vestitum. Embryo cotyledonibus latiuscule ovatis subcordatis, radícula cylindrica longioribus, albumine mediocri. — Frutex sandwicensis, glaber. Folia majuscula, alterna, crenata, trinervia, cystolithis punctiformibus. Stipulæ magnæ, axillares, integerrimæ, subpersistentes. Flores pedicellati, bracteis linearibus intermixti. — Inflorescentiæ capituliformes illas Cypholophorum quorundam referunt, sed structura perigonii fem. junioris sat similis est illi perigonii fem. *Sarcochlamydis*.

- T. *LATIFOLIA* (Gaudich. l. c.), foliis magnis ovatis crenulatis supra glaberrimis subtus in nervis præsertim pubescentibus.  $\delta$ ? In insulis Sandwich (Gaudichaud). — Wedd. Monogr. p. 442. t. 13. *C.* — Ramuli crassi, cicatricibus rugosis (in sicco pallidis) foliorum delapsorum late signatis, cæterum epidermide lævi sparsimque hispida glabratave vestiti, internodiis modo brevissimis modo elongatis. Folia 1-3 decim. longa vel majora, 6-20 centim. lata, circiter a tridente inferiore attenuata, acutiuscula, basi rotundata aut attenuata acutaque et plerumque subintegra, cæterum minute et subinequaliter dentato-crenata, nervis primariis utrinsecus 15-18 ante marginem reticulatim anastomosantibus subtus prominulis jugo inferiore medium limbum attingente, nervis secundariis valde conspicuis inter primarios transverse confluentibus, lamina exsiccata rigide coriacea cystolithis punctiformibus creberrimis obducta; petiolo robusto dimidium limbum æquante vel paulo breviori, juniore pilis quibusdam rigidis acutissimisque hinc et inde insperso. Stipulæ 4-6 centim. longæ, acutæ. Inflorescentiæ masc. pedunculatæ, e capitulis 2-5 globulosis didymisve sessilibus vel longiuscule pedunculatis et centimetrum fere crassis constantes, perigonio puberulo, alabastris 1½ millim. circiter latis. Inflorescentiæ fem. breviter pedunculatæ, capitulis quam in inflorescentiis masc. numerosioribus et paulo minoribus omnino globosis oblongisve aut didymis, ramis pubescentibus hispidisque. (v. s. in hb. Mus. par.)

XXIII. LAUREA Gaudich. Bot. Voy. Bonite, t. 88; Wedd. Monogr. p. 443.

Flores dioici, masculi (solum cogniti) densissime spicati, spicis simplicibus subsimplicibusve in singulis axillis geminis. Perigonium profunde 4-partitum, segmentis obtusiusculis puberulis in præfloratione imbricatis? Stamina 4. Pistilli rudimentum lanceolatum, superne longiuscule acuminato-attenuatum hispidumque, basi pilis rectis stipatum. — Frutex guyanensis, glabriusculus. Folia opposita, in eodem jugo nonnihil inæquimagna et subinæquilatera, integerrima, trinervia, plana, petiolata, cystolithis ut videtur destituta. Stipulæ axillares, integræ, deciduæ. — Genus nimis imperfecte cognitum, inter Sarcocochlamydeas collocatum a similitudine quadam Sarcocochlamydis.

L. TILIÆFOLIA (Gaudich l. c.), foliis amplis rotundato- v. elliptico-ovatis breviter acuminatis basi cordatis supra glabris et lucidis subtus in nervis petioloque subtilissime puberulis, spicis erectis breviter pedunculatis teretiusculis. ♂ In Guyana gallica. — Wedd. l. c. t. 12. D. Piper tiliæfolium Desv. in Ham. Prodr. fl. Ind. occ. p. 4. Folia exsiccata subcoriacea. (v. s. in hb. Mus. par.)

XXIV. SARCOCHLAMYS Gaudich. Bot. Voy. Bonite, t. 89; Wedd. Monogr. p. 439. — Urticæ spec. Roxb. et auct.

Flores dioici, masculi laxiuscule feminei densissime glomerato-spicati, bracteolis brevibus denticulatis stipati, spicis simpliciter ramosis in singulis axillis sæpius geminis. Masc. Perigonium 5-partitum, segmentis ovatis obtusis in præfloratione imbricatis. Stamina 5. Pistilli rudimentum parvum, conoideum, lana densa obtectum. Fem. Perigonium breviter et inæqualiter 4-lobum, sub anthesi campanulatum, tubo denique (latere altero incrementum majus capiente) elevate gibboso, limbo simul omnino laterali lobisque subconniventibus. Ovarium late ellipticum, obliquum, pedicellatum, perigonio inclusum. Ovulum suberectum, funiculo brevi adscendente juxta basim loculi affixum. Stigma sessile, penicillato-capitatum, persistens. Staminum rudimenta nulla. Achænium ovario conforme, læve, perigonio carnoso et accreto laxè inclusum. Embryo... — Frutex indicus, foliis alternis trinerviis serrulatis subtus incanis, cystolithis punctiformibus vix conspicuis, stipulis in unam axillarem v. axillari-lateralem bifidam connatis. — Genus distinctissimum characteribus floris fem. et imprimis perigonii quod, initio rectum et fere symmetricum, ad maturitatem fructus quodam modo (more ovulorum) semianatropum fit.

S. PULCHERRIMA (Gaudich. l. c.), foliis anguste lanceolatis acute serratis serrulatisve supra glaberrimis rugosulisque subtus (nervis nervulisque exceptis) tenuiter niveo-tomentosis. ♂ In rupestribus et ad oras rivulorum regionis tropicæ Indiæ orientalis prov. Silhet, Assam, Chittagong et montium Khasiæ, altit. 0-1000 metr. necnon in ins. Sumatra. — Wedd. Monogr. p. 440. t. 16. C. Miq. Fl. Ind. bat. 1 (2). p. 264. — Frutex bi-tri-metralis. Ramuli elongati, inferne teretes, glabrati corticeque fusco v. rubente vestiti, superne angulato-sulcati puberulique. Folia 8-20 centim. longa vel etiam longiora, 1-3 ½ centim. lata, sensim acuminata, basi acuta, toto margine æqualiter serrata, nervis basilaribus margini parallelis eique proximis sæpe usque ad apicem limbi excurrentibus costaque et venulis (majoribus transverse parallelis) subtus in intervênio toto æqualiter elegantissimeque reticulatis puberulisque supra magis minusve impressis, lamina exsiccata rigide membranacea supra fuscescente subtus fusco-reticulata, petiolo gracili 1-5 centim. longo puberulo. Stipulæ latiuscule amplexicaules, usque ad medium circiter bifidæ (laciniis linearibus ciliolatis), mox deciduæ. Spicæ floriferæ utriusque sexus fere ejus-



dem magnitudinis, petiolo demum circiter dimidio longiores, initio erectæ, mox patulæ vel recurvæ, subdistiche ramosæ, ramis alternis brevibus æqualibusque aut inæqualibus, glomerulis subunilateralibus e ramulis atque ipso axi inflorescentiæ simul nascentibus approximatis, femineis demum (in spica fructifera) omnino confluentibus atque 3-4 millim. crassis. Flores sessiles, bracteis parvis denticulatis stipati. Perigonium utriusque sexus glaberrimum, femineum sub anthesi minimum, fructiferum autem  $\frac{1}{2}$  millim. circiter latum, subglobulosum, basi in pedicellum brevem attenuatum, ore laterali, lobis denticulatis. Ovarium oblique sphæroideum, pedicellatum, eleganter marmorato-reticulatum, perigonio carnosio lutescenteque inclusum vel subexsertum. Achænium... (v. s.)

**XXV. POIKILOSPERMUM** Zippel mscr.; Miquel Ann. Mus. bot. Lugd. bat. 1. p. 203.

Flores dioici, dichotome cymosi. *Masc.* Perigonium campanulatum, 4-fidum, lobis ovatis æstivatione valvatis. Stamina 4. Pistilli rudimentum obconicum, glabrum. *Fem.* Perigonium obconico-subcampanulatum, ore minutissime 4-denticulato, ovarium totum arcte includens, in fructu carnosum achænium basi cupulæ specie amplectens. Stigma sessile, capitellato-peltatum, minute papillosum. Achænium baccatum. Semen conforme, albumine parco. — Frutex, insulas Amboina et Ceram incolens. Folia alterna trinervia. Stipulæ axillares. Flores numerosi. — Monente cl. Miquelio, differt hoc genus a *Sarcochlamyde* perigonio fem. regulari, a *Leucosyke*, præter alia, inflorescentia atque stigmatem non penicillato. Perigonio pro cupula sumpto, Villebrunæ accederet, sed sub accuratiori examine ab hoc genere longe recedere constat. Flore femineo ad *Urereas* etiam haud parum spectat.

**P. AMBOINENSE** (Zipp.; Miq. l. c.), foliis magnis e basi rotundata vel obtusa oblongo- v. obovato-ellipticis breviter obtuse v. acute apiculatis (perraro acute acuminate) integerrimis subcoriaceis, junioribus supra in costa et subtus sparse piliferis. ♂? In ins. Amboina (Zippel, Teysmann) et Ceram (de Vriese). — Rami lignosi cicatricibus petiolorum nodosi, novelli apice sparse et appresse puberi. Folia  $1\frac{1}{2}$  - 3 $\frac{1}{2}$  decim. longa, passim obtusa aut apice rotundata, adulta glabra, lævia, exsiccata supra ad lentem puncticulata subtusque cystolithis striolata, costulis præter basilares utrinque 6-11; petiolo 3-10-centimetrali antice sulcato, juniore parce pilifero. Stipulæ lanceolatæ, 3-5 centim. longæ, dorso bicarinatæ, diu persistentes. Inflorescentiæ breviter pedunculatæ, quam maxime ramosæ, foliis ac sæpius petiolis breviores, densifloræ, floribus cymose sessilibus et subpedicellatis, haud vere capitulatis, axibus tenuibus hirtellis, bracteis ovalibus coloratis sub axibus primariis diutius persistentibus majusculis, masculæ ut videtur densiores et breviores quam femineæ, ramulis ultimis in cymulas capituliformes excurrentibus, perigonii basi parce pilosuli lobis patentibus. Inflorescentia fem. usque 3-decimetralis, ramis primariis longis subæquilongis, floribus in ultimis ramulis per cymulas contractas dispositis exilibus, bracteolis parvis linearibus, perigonio colorato roseo obpyramidato-campaniformi sub-4-angulato margine subtruncato, ovario sub anthesi ovoideo sursum leviter attenuato, fecundo magis lenticulari stigmatem diu coronato carnosulo verruculoso-punctato e perigonio persistente et arcte amplectente  $\frac{1}{2}$  demum exserto. (Descript. ex op. cit. cl. Miq.)

**XXVI. NERAUDIA** Gaudich. Bot. Voy. Uran. p. 500; Wedd. Monogr. p. 437.

Flores dioici, in axillis foliorum glomerati, glomerulis vulgo paucifloris. *Masc.* Perigonium 4-partitum, segmentis ovatis acutis carnosulis in præfloratione valvatis. Stamina 4. Pistilli rudimentum anguste conicum, longe et dense lanatum. *Fem.* Perigonium tubuloso-ventricosum, apice attenuatum, ore contracto sub-4-dentato. Ovarium ovoideum, in-

clusum, sessile, a perigonio discretum. Ovulum imo loculo funiculo brevi affixum, erectum. Stigma elongato-filiforme, cum ovarii apice articulatam, deciduum, basi (colli perigonii inclusa) glabrum, cæterum villosum. Achænium depresso-conicum, basi sæpe lobulatum, perigonio evoluto globuliformi carnosoque laxe inclusum, pericarpio crasso nucamentaceo. Embryo... — Frutices oceanici, succo lacteo (teste Gaudichaud), glabri aut pubentes. Folia alterna, homomorpha et æquilatera, integerrima, trinervia, petiolata, cystolithis punctiformibus. Stipulæ parvæ axillares. — Genus ab omnibus Sarcoclamydeis notis permultis distinctissimum.

1. *N. MELASTOMÆFOLIA* (Gaudich. l. c. p. 500. t. 117), ramulis glaberrimis, foliis oblongo- v. lanceolato-ellipticis ovatisve basi acutis s. cuneatis aut obtusatis sensim acuminatis utrinque glaberrimis v. nervis subtus adpresse pilosis.  $\frac{5}{8}$  In ins. Sandwich (Gaudichaud, Beechey, Remy, etc.). — Wedd. *Monogr.* p. 438. t. 12, A. N. glabra Meyen, Reise, 2. p. 124. *Boehmeria melastomæfolia* Hook. et Arn. Bot. Voy. Beech. p. 96. *Boehm. glabra* Steud. nom. 2. p. 192. — Ramuli nonnihil arcuati, crassitie circiter calami scriptorii, floriferi sijnati et ad insertionem foliorum inflorescentiarumque subnodosi, exsiccati pallide fusco-rubentes glabri aut apice puberuli, internodiis brevibus. Folia 5-15 centim. longa, 2-6 centim. lata, juniora supra sparsim pilosula, nervis primariis ad quartam partem superiorem limbi excurrentibus cumque sequentibus arcuatim anastomosantibus, petiolo gracili (saltem in sicco) 1-5 centim. longo sæpius glaberrimo. Stipulæ inconspicuæ, glomeruli utriusque sexus conformes, 3-10-flori. Flores masc. subsessiles, perigonii ovati 3 millim. fere longi segmentis crassis extus puberulis, pistilli rudimento lanatissimo. Flores fem. etiam fere sessiles, perigonio ovato circiter bimillimetrali superne longiuscule angustato s. rostrato extus puberulo ore acute 4-dentato, ovario millimetrali quam stylus quadruplo brevior; perigonio fructifero sphaeroides aut subdepresso globuloso 3-4 millim. lato in collum breve abrupte acuminato achæniumque basi dilatatum et magis minusve lobulatum laxissime includente. (v. s.)

$\beta$ , *ovata*, ramulis pubescentibus, foliis ovatis basi late rotundatis supra puberulis subtus subtiliter pubescentibus.  $\frac{5}{8}$  Cum typo in ins. Sandwich. *N. ovata* Gaudich. l. c.

2. *N. SERICEA* (Gaudich. l. c.; Bot. Voy. Bon. t. 133), foliis late ellipticis v. elliptico-ovatis sæpius abrupte acuminatis supra sparsim puberulis subtus undique tenuiter molliterque pubescentibus.  $\frac{5}{8}$  In fisdem locis quam spec. præced. (Gaudichaud). Ramuli et flores ut in *N. melastomæfolia* cujus forte varietas est. Folia quam in illa ut plurimum pro longitudine latiora et abruptius acuminata magisque pubentia, cæterum similia. (v. s.)

Subtrib. III. *VILLEBRUNÆ* Wedd. *Monogr.* p. 342. Perigonium femineum ovario semper adnatum, fructiferum siccum v. subhæcatum, limbo minimo dentato v. subintegro, Stigma varium, nunc cum apice ovarii continuum et diu persistens, nunc articulatam et caducissimum.

XXVII. *PIPTURUS* Wedd. in *Ann. sc. nat.* 4<sup>e</sup> sér. 1. p. 196; *Monogr.* p. 444; *Miq. Fl. Ind. bat.* 1 (2). p. 267; *Seem. Fl. Vit.* p. 243. — Nothocnide *Bl. Mus. bot. Ind.* bat. 2. f. 14. — Urticæ et *Boehmeriæ* spec. auct.

Flores dioici, arcte glomerulati s. capitati, bracteis parvis hirtis stipati, glomerulis s. capitulis axillaribus v. interrupte spicatis, spicis interdum distiche ramosis, receptaculo capitulorum fem. demum carnosum. *Masc.* Perigonium 4-5-lobum, lobis ovatis acutis. Stamina tot quot perigonii segmenta. Pistilli rudimentum clavatum, lanatum. *Fem.*

*Perigonium* ovatum, ventricosum, sensim attenuatum, ore contracto minime 4-5-denticulato. Ovarium perigonio conforme eique adnato. Ovulum e basi loculi erectum, funiculo brevi sustentatum. Stigma elongato-filiforme, cum apice ovarii articulatam et caducissimum, altero latere villosum, basi glabrum oreque perigonii brevissime insertum. Fructus ventricosus, e pericarpio nucamentaceo perigonio haccante (?) vestito constitutus. Embryo cotyledonibus ellipticis ovatisve radícula paulo longioribus. Albumen parvum. — Arbusculæ v. frutices (raro scandentes) malaici, oceanici et mauritiani, pubescentes v. tomentosi rariusve glabriusculi. Folia alterna, homomorpha et æquilatera, integra aut dentata, subtus sæpe canescentia, trinervia, petiolata, cystolithis punctiformibus. Stipulæ axillares, cujusque folii in unam profunde bifidam connatæ. — Genus a cæteris omnibus distinctissimum notis floris fem. Fructus s. capitulum fructiferum a peregrinatoribus moriformis dicitur, at specimina exsiccata hunc characterem non retinent.

#### A. — *Glomeruli axillares.*

1. *P. ALBIDUS*, ramulis superne pubescenti-tomentosis, foliis ovatis acuminatis basi acutis v. rotundatis v. subcordatis crenatis serratisve supra puberulis v. glabris lævibusque aut asperulis subtus subtiliter incano-tomentosis, glomerulis omnibus axillaribus, perigonio fem. quam stigma vix dimidio brevior. ♂ In ins. Sandwich. — *P. taitensis* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 197; Monogr. p. 449. *Boehmeria albida* Hook. et Arn. Bot. Voy. Beech. p. 96. — Ramuli obtuse angulati, modice crassi, internodiis brevibus (5-10-millim.). Folia magnitudine admodum variabili, vulgo 4-15 centim. longa, 3-10 centim. lata et haud raro ampliora, acumine triangulari sæpius ut limbus reliquus serrato, serraturis acutis obtusiusculisve, nervis basilaribus usque ad quartam partem superiorem limbi protensis cum primariis sequentibus (plerumque paucis) ante vel juxta marginem anastomosantibus, lamina exsiccata rigide membranacea supra sæpissime fusciscente, petiolis longitudine variabilibus (etiam in eodem ramo) dimidium limbum autem raro excedentibus. Stipulæ fere liberæ, triangulari-lanceolatæ, 3-4-millimetrales, subulato-acuminatæ, extus adpresse pilosulæ. Glomeruli masc. et fem. conformes, pisiformes, ramulum amplectentes, receptaculo femineorum bracteolisque dense hirtis, floribus utriusque sexus sessilibus densiuscule albidopubescentibus. Perigonium fructiferum ovatum, millimetrale. (v. s.)

β, *Gaudichaudianus*, foliis amplioribus quam in typo elliptico- v. rotundato-ovatis subtus vix incanis serraturis s. crenis plerumque majoribus. ♂ In iisdem insulis. — *P. Gaudichaudianus* Wedd. in Ann. sc. nat. 1. c.

γ, *Meyenianus*, ramulis subtiliter puberulis, foliis subacuminatis fere ab insertione petioli usque ad apicem crenatis (crenis semirotondis obtusissimis) supra subtilissime adpresseque hispidis et minute rugosulis subtus in intervenio subtiliter canescenti-tomentosis nervis puberulis. ♂ In ins. sandwicensi Ovahu (Meyen). — Ramulus visus crassitie calami scriptorii, obscure rubens. Folia 5-10 centim. longa, exsiccata subcoriacea. (v. s. in hb. Mart.)

2. *P. ASPER* (Wedd. in Ann. sc. nat. 1. c.; Monogr. p. 450), ramulis inferne puberulis glabrativè superne breviter tomentosis, foliis ovatis sensim longiuscule acuminatis basi obtusatis rotundatisve crenulato-serratis aut subintegris supra hispидulis demumque asperimis subtus velutinis, glomerulis omnibus axillaribus, perigonio fem. quam stigma triplo brevior. ♂ ? In ins. Philippinis (Cuming, n. 724; Lobb, n. 463). — Ramuli ut in *P. albido*, sed internodiis plerumque paululum longioribus. Folia sæpius longius et minus abrupte acuminata quam in eodem, æqualis autem fere longitudinis, nervis quoque basilaribus et primariis ut in specie illa, reticulatione intervenii minus conspicua interdumque ob tomentum vix manifesta, lamina exsiccata in speciminibus omnibus visis rigide membranacea supra olivacea subtus pallide

viridi, petiolis gracilibus breviter denseque pubescenti-tomentosis. Stipulæ liberæ, ovatæ, 3-4 millim. longæ, subabrupte et anguste acuminatæ, extus dense pubescentes. Glomeruli pisiformes, feminei præsertim densiflori. Perigonium masc. pubescens, femineum cano-tomentosum, fructiferum quam in specie antecedente angustius. (v. s.)

B. — *Glomeruli plerique spicati.*

3. P. ROTUNDIFOLIUS (Wedd. in Ann. sc. nat. l. c.; Monogr. p. 445), ramulis apice subtilissime puberulis, foliis elliptico- v. rotundato-ovatis subacuminatis basi rotundatis subcordatisve integerrimis supra glaberrimis nervis subtus subtilissime adpresseque pilosulis, glomerulis in spicas elongatas fasciculatas s. ima basi ramosas superne foliiferas dispositis. ☉ In insula Mauritio (Commerçon, cæt.). — *Urtica rotundifolia* Poir. Encycl. 4. p. 644. *U. monilifera* Smith herb. — Frutex scandens. Ramuli elongati, teretiusculi s. obtuse angulati, crassitie pennæ anserinæ v. cynæ, siccitate fistulosi epidermideque fusco-rubente vestiti, internodiis superioribus 3-5 centim. longis. Folia ramorum sterilibus in speciminibus examinatis 8-10 centim. longa, 5-8 centim. lata, sensim acutata vel acuminata, nervis basilariibus circiter ad sextam partem superiorem s. fere ad apicem limbi productis et ante marginem cum primariis sequentibus (supra medium folium ortis) arcuatim anastomosantibus, intervenio subtus tenuiter reticulato, petiolis 2-4-centimetralibus foliorum juniorum sparsim puberulis sed mox glabratiss. Stipulæ axillares, ovato-lanceolatæ, 3-5 millim. longæ, bifidæ, puberulæ. Spicæ fem. (solum visæ) erecto-patulæ, in singulis axillis plerumque ternæ (duabus lateralibus e basi ipsa spicæ mediæ ortis illaque sæpius brevioribus), 5-15 centim. longæ, e glomerulis 4-16 sessilibus alternis discretis inferioribus nudis superioribusque ex axilla folii parvi (1-2 centim. longi) elliptici et petiolati ortis compositæ, floribus coarctatis. Perigonium puberulum, quam stigma multoties brevius, fructiferum ovoideum et fere millimetrale. (v. s.)
4. P. INCANUS, ramulis breviter incano-tomentosis, foliis amplis late ovatis subabrupte acuminatis basi cordatis v. rotundatis crenulatis serrulatisve discoloribus supra hispidis aut glabratiss. subtus incanis canescentibusve, glomerulis in spicas laxè paniculatas dispositis. ♂ In ins. Malaicis, Societatis, cæt. — P. velutinus Wedd. in Ann. sc. nat. l. c.; Monogr. p. 446. t. 15. D. f. 4-8; Miq. Fl. Ind. bat. 1 (2). p. 268. *Urtica incana* Bl. Bijdr. p. 497. *Boehmeria velutina* Decne Herb. Timor. p. 163. *B. incanana* Hassk. Cat. hort. bogor. p. 79; Pl. jav. rar. p. 207; Miq. Pl. Jungh. 1. p. 36. — Ramuli apice obtuse angulati, crassitie calami scriptorii aut tenuiores, internodiis nunc 1-1½ centimetralibus nunc (præsertim in var. β) etiam brevioribus. Folia patula, magnitudine admodum variabili, alterna nempe valde inæquimagna, majora in speciminibus examinatis 2 decim. circiter longa, 15 centim. lata, minora quadruplo v. quintuplo breviora et angustiora, acumine modo triangulari modo fere lineari acutissimo rarius nullo (limbo scilicet gradatim acutato), sinu basilari profundo vel interdum (ac non raro in folio nodi proximi) prorsus nullo, dentibus semper parvis obtusissimis aut acutis interdumque fere omnino obsoletis, nervis basilariibus circiter ad trientem superiorem limbi excurrentibus et ante marginem cum nervis primariis sequentibus (utrinsecus pluribus) arcuatim anastomosantibus, lamina exsiccata membranacea vel rigidula supra sæpius fusca et magis minusve hispidula etiamque aspera subtus glauco-canescente eleganterque fusco-reticulata, petiolis foliorum majorum sæpissime dimidium limbum longitudine superantibus, foliorum minorum autem (s. cum præcedentibus alternantium) canescenti-tomentosis aut pubescentibus. Stipulæ ovatæ, 3-5-millimetrales, acutæ, extus incanulæ v. glabræ, cito deciduæ. Inflorescentiæ masc. et fem. conformes, quam petioli vulgo breviores, simpliciter pauci-ramosæ, ramis distichis patentibusque, glomerulis in planta exsiccata granum piperinum magnitudine fere æmulantibus discretis, floribus sessilibus, perigonio utriusque sexus ramulisque et pedunculo densiuscule breviterque tomentoso. Fructus ovatus, obtusiusculus, vix millimetralis. (v. s.)

β, *glabrior*, ramulis glaberrimis, cæt. ut in typo.

γ, *pellucidus*, ramulis hirtis-tomentosis, foliis deltoideo-ovatis lineari-acuminatis repando-dentatis fere concoloribus nervis subtus petiolisque hirtis-tomentosis. ♂ In ins. Boutou (de Labillardière). — *Urtica pellucida* Billard. Sert. austr. caledon. p. 79. t. 80. (v. s.)

δ, *asper*, ramulis breviter tomentosis, foliis supra asperis subtus velutinis.

5. *P. ARGENTUS*, ramulis ut in præcedente, foliis minoribus breviusque petiolatis rarius cordatis acuminatis vel sensim attenuatis acutisque discoloribus, adultis supra glabris lævibusque, glomerulis axillaribus aut plerisque in spicas simplices interruptas et nonnunquam apice foliosas dispositis. ♂ In ins. Java (Blume, cæt.), Timor, Nova Hibernia, Taiti (Riedle), Mariannis (Gaudichaud, cæt.), Nova Hollandia, ad Moreton Bay (Leichard) et Queen's Land (F. Mueller). — *P. propinquus* Wedd. Monogr. p. 447. t. 15. D. f. 1-3; Miq. l. c. *P. propinquus* et *P. Candolleanus* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 196. *Urtica argentea* Forst. Prodr. n. 343. *U. cinerascens* Bl. Bijdr. p. 497. *U. albidopunctata* Steud. Repert. 2. p. 734. *Boehmeria propinqua* Decne l. c. B. *Candolleana* Gaudich. Bot. Voy. Uran. t. 148. — Ramuli sæpius nonnihil flexuosi, crassitie calami scriptorii, obtuse angulati, internodiis 5-10-millimetralibus. Folia quam in præcedente sæpius dimidio v. triente minora, interdum elliptico-lanceolata, basi ut plurimum cordata, lamina exsiccata rigida et nonnunquam subcoriacea, petiolis longitudine admodum variabili, altero longiore cum altero breviori alternante. Stipulæ 2-5 millim. longæ. Spicæ (raro nonnullæ ramosæ) petiolos superantes, e glomerulis 5-10 discretis constantes. Cæt. ut in *P. velutino* cui valde affinis est. (v. s.)
6. *P. PLATYPHYLLUS*, ramulis hirtis demumque glabris, foliis amplis ovatis acuminatis basi rotundatis crenato-serratis concoloribus supra asperato-hispidis subtus molliter pubescentibus, glomerulis in spicas simplices interruptas digestis. ♂ In ins. Ovaloho archip. Fidji (Vieillard). — Ramuli teretiusculi, crassitie calami scriptorii, epidermide exsiccata sub hirsutie brunnea. Folia eadem circiter magnitudine ut in *P. velutino*, pleraque sensim acuminata, basi interdum subcordata, serraturis paulo majoribus quam in specie memorata obtusis v. brevissime apiculatis, petiolis (dimidii limbi circiter longitudine) costaque et nervis basilaribus in planta exsiccata subcinereo-pubescentibus. Spicæ in specimine examinato paucæ, graciles, petiolis breviores, glomerulis (fem.) parvis; floribus ut in speciebus præcedentibus. (v. s. in hb. Lenormand.)
7. *P. REPANDUS* (Wedd. Monogr. p. 448), ramulis apice puberulis pubescentibusve, foliis ellipticis v. oblongo-ellipticis subabrupte acuminatis basi obtusis rotundatisve integerrimis v. repando-denticulatis utrinque glabris et lævibus aut nervis pubescentibus, glomerulis fem. in spicas simplices digestis. ♂ ? In ins. Java (de la Billardière, Blume, Zolling. n. 1574 et 3540), in Manilla (Cuming, n. 759). — Miq. Fl. Ind. bat. 1 (2). p. 268. *P. repandus* et *P. ellipticus* Wedd. in Ann. sc. nat. l. c. *Urtica repanda* Bl. Bijdr. p. 501. *Boehmeria repanda* Hassk. Cat. hort. bogor. p. 79; Miq. in Zolling. Syst. Verzeichn. 1. p. 101. *Nothocnide repanda* Bl. Mus. bot. lugd. bat. f. 14. — Ramuli eadem crassitie ac in præcedentibus, superne obtuse angulati, breviter hirsuto-pubescentes puberulive aut omnino glabri, internodiis circiter centimetralibus. Folia in eodem ramulo nonnihil inæquimagna, 5-13 centim. longa, 2½-6 centim. lata, acumine brevi angusto et acuto, nervis basilaribus usque ad quintam v. sextam partem superiorem limbi elongatis, primariis cæteris paucissimis nempe utrinque tantum uno vel rarius duobus (superiore minimo) sæpe (costaque) in pagina utraque pubescentibus, lamina exsiccata rigide membranacea aut coriacea fusca, petiolis 1-2-centimetralibus glabris pubescentibusve. Stipulæ 3-5 millim. circiter longæ, bifidæ, nervo dorsali adpressæ pilosæ. Inflorescentiæ, floresque et fructus ut in præced. sed spicis masc. sæpe ramosis, perigonio masc. glabriusculo, femineo puberulo ore tomentoso. (v. s.)

β, *oblongifolia*, foliis oblongis v. oblongo-lanceolatis, nervis primariis (s. adscendentibus) præter basilares nullis. ♂ ? In ins. Java (de Vriese).

8. *P. MOLLISSIMUS* (Wedd. Monogr. p. 449), ramulis puberulis, foliis obovato-

ellipticis obtusissimis apiculatisve basi obtusis vel rotundatis v. subcordatis integerrimis utrinque pubescentibus, glomerulis spicatis. ♂ ? In ins. Java (blume, cæt.) et Pulo-Pinang (Wall. n. 4609). — *Urtica mollissima* Bl. Bijdr. p. 501. — Ramuli teretes. Folia 10-14 centim. longa, 5-6 centim. lata, nervis basilaribus arcuatis fere usque ad apicem limbi productis, primariis cæteris utrinque 1-2, pube in pagina inferiore limbi brevior et densior, lamina exsiccata rigide membranacea olivacea, petiolis 2-3-centimetralibus pubescentibus. Spicæ masc. (solum visæ) in singulis axillis sæpius 3, petioli circiter longitudine vel eo breviores, glomerulis discretis. (v. s. in hb. DC. Desesert, cæt.)

*Species mihi nomine tantum cognita.*

*P. GRACILIPES* Asa Gray, in litt. — In ins. Fidji.

**XXVIII. VILLEBRUNEA** Gaudich. *Bot. Voy. Bôn. t. 91 et 92; Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 195; Monogr. p. 451; Bl. Mus. bot. lugd. bat. 2. p. 166; Benth. Fl. hongk. 332. — Oreocnide* Miq. *Pl. Jungh. 1. p. 39; Fl. Ind. bat. 1 (2). p. 269. — Morocarpi* spec. *Sieb. et Zuccar. — Urticæ et Boehmerie* spec. *auct.*

Flores dioici, arcte glomerati s. capitati, glomerulis s. capitulis pedunculato modo simplices fasciculatosque modo furcatis dichotomosve (et tunc sæpius geminos) terminantibus. *Masc.* Perigonium 4-partitum, segmentis ovatis acutis extus hispidulis in præfloratione valvatis, alabastro acutiusculo v. obtuso. Stamina 4. Pistilli rudimentum obovato-clavatum, basi pilis lanatis stipatum. *Fem.* Perigonium tubulosum, ventricosum, ovario adnatum, limbo minimo 4-5-dentato. Ovulum e fundo loculi ovarii erectum, funiculo brevi. Stigma fere discoideum, subpeltatum, margine longe ciliato-papillosum. Fructus ventricosus obsolete angulato-sulcatus, strato extimo (e perigonio adnato constante) aliquantum carnosus, intimo s. pericarpio subnucamentaceo. Semen pericarpio fere conforme. Albumen satis copiosum. Embryo cotyledonibus ellipticis radiculæ tereti subæquilongis. — Frutices v. arbusculæ indici, malaici et oceanici. Folia integra v. crenulata, penninervia aut trinervia, glabra v. pubescentia. Stipulæ in unam axillarem bifidam concretæ. Flores fem. cupula carnosae e bracteis pluribus connatis constante suffulti, hacce demum valde accrescente fructumque maturum more perigonii *Ureræ* vestiente. — Genus admodum insignis floris fem. et fructus indole.

A. — *Folia penninervia.*

1. *V. APPENDICULATA*, foliis oblongis v. oblongo-lanceolatis cuspidato-acuminatis basi obtusis v. acutis rariusve rotundatis integris penninerviis planis supra glabris lævibusque nervis et præsertim costa subius hirsuto-pubescentibus. pedunculis bis terve dichotomis. ♂ In regionibus tropicis montosisque Indiæ orientalis, prov. Concan et Chittagong, necnon in montibus Khasiæ et Himalayæ orientalis, altit. 1520 metr. — *V. integrifolia* Gaudich. l. c. t. 91; Bl. l. c. p. 166; Wedd. Monogr. p. 452. *Urtica urophylla* et *U. appendiculata* Wall. Cat. n. 437 et n. 4604. Ramuli obtuse angulati, crassitie variabiles, graciles aut robusti, superne hirsuto-pubescentes, epidermide rubente vel fusco-cinerascente vestiti, internodiis 1-3-centimetralibus. Folia patula, 8-25 centim. longa, 3-12-centim. lata, nunc a medio utrinque attenuata, nunc deorsum fere cuneata, acumine lineari 1-3-centimetrali acutissimo, nervis primariis tenuibus utrinsecus 8-12 (inferioribus quam cæteri brevioribus) fere in ipso margine arcuatis anastomosantibus, lamina exsiccata rigida vel submembranacea fusciscente, petiolo 1-8-centimetrali ut costa pubescenti-hirto. Stipulæ lanceolatae, 5-15 millim. longæ vel etiam longiores, subulato-acuminatæ, extus densiuscule pubescenti-pilosæ. Inflorescentiæ petiolo breviores, sessiles.

utriusque sexus conformes, ramulis breviter hirtulis diametrum glomerulorum longitudine circiter æquantibus. Flores masc. perigonio 4-partito extus hispido, pistilli rudimento lineari-conico lanato. Capitula fem. 10-15-flora, initio 5 millim. circiter crassa demumque (cupula basilari evoluta) centimetralia, cupulis inæqualibus 5-6-angulis  $\frac{1}{2}$ -3 millim. latis totidemque fere altis basi subcuneatis margine obtusis in planta viva albedo-subhyalinis fructus ovatos sesquimillimetrales subacuminatos obtusiusculosque (semina vitis referentes) laxè continentibus. (v. s.)

$\beta$ , *pubescens*, ramulis superne velutinis, foliis subtus undique pube densa molliuscula que vestitis. (v. s.)

2. *V. SEMICRENATA* (Bl. l. c. p. 167), ramulis petiolis folisque subtus ad nervum medium sericeo-puberulis, foliis ellipticis acuminatis basi rotundatis inferne integerrimis superne crenulatis, pedunculis furcatis v. dichotomis brevissimis.  $\frac{3}{4}$  In umbrosis Amboinæ. — Wedd. Monogr. p. 453. Oreocnide semicrenata Miq. Fl. Ind. bat. 1 (2). p. 270.

3. *V. RUBESCENS* (Bl. Mus. bot. lugd. bat. 2. p. 166), foliis oblongis ellipticisve subabrupte acuminatis basi rotundatis sæpiusve subcordatis circiter in dimidio superiore crenulatis v. crenulato-serratis cæterum subintegræ penninerviis planis supra glabris lævibusque subtus præcipue in nervis pubescentibus, pedunculis subsimplicibus (s. ima basi divis) v. furcatis.  $\frac{3}{4}$ - $\delta$  In sylvis montosis ins. Javæ. — Wedd. Monogr. p. 454. t. 15. C. f. 12 et 13. *V. scabra* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 195. *Urtica rubescens* Bl. Bijdr. p. 506. *U. scabra* Herb. Zolling. n. 577 et 581, non Bl. Oreocnide major Miq. Pl. Jungh. 1. p. 41; Fl. Ind. bat. 1. c. Boehmeria rubescens Hassk. Cat. hort. bogor. — Ramuli exsiccati crassitie pennæ anatinæ v. anserinæ, obtuse angulati v. teretiusculi, epidermide fusco-cinerea vestiti, superne pubescentes, internodiis 1-2-centimetalibus. Folia 6-20 centim. longa, 3-8 centim. lata, fere in toto margine crenulata, acumine 1-2-centimetro acutiusculo, nervis primariis in utroque latere costæ 5-10 cæterum ut in *V. integrifolia*, intervenio subtiliter puberulo, lamina exsiccata rigide membranacea fusciscente, petiolo 1-4 centim. longo densiuscule pubescente. Stipulæ lanceolatæ, 6-8 millim. longæ, excepto nervo dorsali glabræ, mox deciduæ. Inflorescentiæ ut plurimum parvæ, pedunculis patulis fasciculatis quam petioli circiter dimidio brevioribus, glomerulis floriferis ut in specie supra memorata sed paululum minoribus, fructiferis non visis. (v. s.)

4. *V. SYLVATICA* (Bl. Mus. bot. lugd. bat. 2. p. 167. f. 15), foliis ellipticis v. elliptico-oblongis lanceolatisve anguste acuminatis basi acutis v. obtusis penninerviis superne obsolete crenulatis supra glabris nervis subtus pilosis, pedunculis simplicibus v. furcatis fasciculatis.  $\delta$  In sylvis montosis ins. Javæ et Zeylanæ. — Wedd. Monogr. p. 453. t. 15. C. f. 1. *Urtica sylvatica* Bl. Bijdr. p. 506. Boehmeria sylvatica Hassk. Hort. bogor. p. 79. Oreocnide sylvatica Miq. Pl. Jungh. 1. p. 40; Fl. Ind. bat. 1. c. — Arbor 6-9-metralis, ramulis glabris aut apice puberulis gracilibus, epidermide exsiccata pallide fusca et corrugata. Folia 6-15 centim. longa, 2-7 centim. lata, acumine lineari centimetrum v. sesquicentimetrum longo, nervis primariis utrinque 5-8 cæterum ut in speciebus præced. saltem jugo inferiore nonnunquam (sed raro) sequentibus longiore (folio tunc vere trinervio), lamina exsiccata membranacea et obscure viridi v. rigida et fusciscente, petiolo gracili  $\frac{1}{2}$ -2 centim. longo et interdum fere triplo longiore. Stipulæ ut in præcedente. Inflorescentiæ et flores ut in eadem, pedunculis autem fortasse longioribus. (v. s.)

$\beta$ , *integrifolia*, foliis integerrimis. (v. s.)

$\gamma$ , *Sumatrana*, foliis repandis v. repando-crenato-serratis, costæ subtus ramulisque et petiolis pubigeris. (Blume.)

$\delta$ , *serrulata*, foliis superne acute serrulatis, subtus ad nervos ut ramuli et petioli pubescentibus, pedunculis plerisque furcatis. — *V. glomerata* Bl. l. c. p. 167. Oreocnide pilosula Miq. l. c. p. 41. (v. s.)

#### B. — Folia trinervia.

5. *V. FRUTESCENS* (Bl. Mus. bot. lugd. bat. 2. p. 168), foliis late ellipticis ova-

tisve v. oblongis v. oblongo-lanceolatis cuspidato-acuminatis basi rotundatis obtusatisve acute serratis rarius crenato-serratis trinerviis, junioribus rugosis concoloribusque v. subtus in intervenio cinereo-tomentellis s. arachnoidis nervis pilosis pagina superiore sparsim hispida, pedunculis brevissimis simplicibus v. furcatis. 5 In Japonia (Thunberg; Oldham, n. 771, etc.) et in Himalayæ regione tropica, ad Sikkim et Kumaon, altit. 600-1500 metr. (J. D. Hooker et Thomson), Assam (Griffith), Nepalia (Wallich, n. 9091) et Concan (Ralph, n. 97). — Rami ramulique graciles, obsolete angulati, puberuli, internodiis florum interdu vix centimetralibus. Folia in eodem ramo sæpe inæquimagna, 2-12 centim. longa, 1-5 centim. lata, basi integra, acumine lineari 1-2-centimetrali acuto integro aut laxe serrato, nervis basilariibus sæpissime supra medium limbum sive usque ad tertiam v. quartam partem superiorem illius elongatis, lamina exsiccata folii adulti membranacea et obscure viridi, ea folii junioris nonnunquam subtus niveo et nigro reticulata. Stipulæ lineari-lanceolatæ, apice subulatæ, 6-8 millim. longæ, in nervo dorsali pilosulæ. Inflorescentiæ primo aspectu sessiles, pedunculis nempe inter glomerulos occultis; masculæ illas Boehmeriæ referentes, glomerulis confluentibus glomerulumque unicum 5-8 millim. latum efformantibus, floribus sessilibus, perigonii tripartiti segmentis ovatis extus superne hirtulis. Capitula fem. longitudine ramulorum inflorescentiæ latiora, 8-12-flora; fructifera fere rotundata et 5-6 millim. lata, solitaria aut sæpissime didyma, ea nodorum proximorum haud raro confluentia et subsessilia, cupula fructum suffulciente iisdem forma et colore ut in V. integrifolia sed minore scilicet 2-2½ millim. lata fructuque glabriusculo sesqui-bi-millimetrali nonnihil superata. — Magnitudine foliorum, ait Blumius, quæ modo subtus niveo-tomentosa, modo in eodem subinde ramulo utrinque concolora et nullo subtus tomento et ut serraturæ plerumque satis magnæ modo acuta modo obtusa, adeo variat ut facile ea non minus esse speciei existimes quamquam nihil certius esse plurimis comparatis speciminibus constat. (v. s.)

β, *pauciserrata*, foliis ovato-lanceolatis, serraturis utrinque grossis acutisque.

6. V. TRINERVIS (Wedd. in Ann. sc. nat. l. c.; Monogr. p. 456), foliis ellipticis v. ovato-ellipticis abrupte acuminatis basi obtusis v. late cuneatis acutatisve integerrimis trinerviis supra glabris lævibusque nervis subtus puberulis, pedunculis bis terve dichotomis. 5? In insula Manilla Philippinarum (Cuming, n. 777). — Oreocnide trinervis Miq. Fl. Ind. bat. l. c. — Ramuli obtuse angulati, graciles, superne puberuli, exsiccati fuscii. Folia in speciminibus visis 7-12 centim. longa, 4½-6 centim. lata, acumine centimetrali vel breviori acuto vel obtusiusculo, nervis basilariibus usque ad quartam partem superiorem limbi excurrentibus juxtaque marginem cum primariis sequentibus (utrinsecus 2-3) arcuatim anastomosantibus, lamina exsiccata membranacea fusca, petiolo gracili 1½-5 centim. longo puberulo glabratoque. Stipulæ ovatæ, acutæ, glabriusculæ, cito deciduæ. Inflorescentiæ fem. (masculæ mihi ignotæ) quam petioli fere dimidio breviores, laxe ramosæ, ramulis glabriusculis, glomerulis floriferis 3 millim. circiter crassis 12-14-floris cæterum ut in præcedentibus, fructiferis non visis. (v. s.)

### *Species dubiæ.*

7. V. ? *RUFESCENS* (Bl. l. c. p. 166), foliis ellipticis subabrupte acuminatis basi obtusatis integris aut superne obsolete crenulato-dentatis penninerviis rugosis supra glabris lævibusque nervis subtus pubescentibus pedunculis furcatis v. dichotomis brevissimis, capitulis fem. 10-15-floris, floribus bracteolis scariosis suffultis. 5 In Nova Guinea (Hind). — Wedd. Monogr. p. 157. Oreocnide? rufescens Miq. Fl. Ind. bat. l. (1). p. 271. Arbor 6-10-metralis, ramis ramulis-

(1) Vid. quod de hoc nomine necnon de nomine Morocarpi dicitur in Monographia mea, p. 460. Tabulæ botanicæ operis dicti *Voyage de la Bonite* inter annos 1839 et 1846 editæ sunt; inscriptio autem hujus iconographiæ post mortem Gaudichaudii s. anno 1851 data est.



que exsiccat is epidermide pallide ferruginea vestitis, novellis villosis. Folia (alternatim minora ac brevius petiolata) in speciminibus examinatis 12-18 centim. longa,  $4\frac{1}{2}$ -8 centim. lata, a medio deorsum sensim attenuata, acumine apicali sesquicentimetrali lineari acutoque, nervis primariis utrinsecus 8-10 prope marginem inter se arcuatim conjunctis (jugo inferiore brevissimo et fere basilari), secundariis et tertiariis in intervênio rete subtile formantibus, laminæ exsiccatæ rigidæ pagina superiore bullulata (venulis impressis) cystolithisque crebre albo-punctulata, petiolo (in vivo rubente)  $1\frac{1}{2}$ -3 $\frac{1}{2}$  centim. longo pubescente. Stipulæ lanceolatæ, 6-8-millimetrales, acutæ, dorso puberulæ. Inflorescentiæ valde contractæ, e capitulis 2-4 ut plurimum constantes, pedunculis villosulis diametri capitulorum circiter longitudine. (v. s. in hb. Benth.)

β, *pubescens* (Blume), ramulis petiolisque et foliis subtus in nervo medio vel infra medium ad apicem obsolete denticulatis.

In floribus fem. speciminis unici suppetentis desiderari videbatur cupula characteristica fructus plantarum hujus generis. Species duæ sequentes mihi omnino incognitæ sunt.

8. V. ? *MURINA* (Bl. l. c. p. 160), foliis ovatis v. oblongo-ovatis acutis v. acuminatis ad basim integerrimis a triente inferiore ad apicem obtuse serrulatis supra scabridis subtus murinis et molliter tomentosis, capitulis in pedunculo brevi paucis subramosis. ♂ ? In Nova Guinea (Zippel). — Wedd. l. c. p. 158. — Folia 4-7 centim. longa, 2-4 centim. lata. (Blume.)
9. V. ? *RHODOPLEURA* (Bl. l. c.), foliis lanceolatis acuminatissimis basi acutis denticulato-serratis subtus niveo-tomentosis, capitulis breviter pedunculatis solitariis v. paucis racemosis. ♂ ? In Nova Guinea (Zippel). — Wedd. l. c. Oreocnide rhodopleura Miq. l. c. Folia 3-10 centim. longa, petiolis 5-15-millimetralibus (Blume). — Conferenda cum *Missiessya corymbulosa*.

### *E genere exclusa.*

V. *CRENULATA* Gaudich. = *Urera sandwicensis*.

XXIX. *DEBREGEASIA* (1) *Gaudich. Bot. Voy. Bon. t. 90; Wedd. Monogr. p. 459.* — *Morocarpus* Sieb. et Zucc. in *Munch. Abhandl. d. math.-phys. Kl. Akad. Wissensch. 4. 3. Abh. p. 218. n. 765; Bl. Mus. bot. lugd. bat. 2. p. 155; Miq. Fl. Ind. bat. 1 (2). p. 272.* — *Leucocnides* spec. *Miq. Pl. Jungh. 1. p. 36.* — *Missiessya* spec. *Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 195.* — *Urticæ* et *Boehmeriæ* spec. *auct.*

Flores monoici v. dioici, masculi glomerulati, feminei capitellati, bracteis scariosis stipati, glomerulis capitulisque ut plurimum in apicibus ramulorum pedunculi furcati v. dichotomi solitariis didymisve. *Masc.* Perigonium 4- rariusve 3-partitum, segmentis ovatis breviter acuminatis dorso convexis in præfloratione valvatis. Stamina 4 v. 3. Pistilli rudimentum ellipsoideum, apiculatum, basi densiuscule lanatum. *Fem.* Perigonium ventricosum-tubulosum, ovario obovato-oblongo subadnatum, ore contracto 4-dentato. Ovulum e fundo loculi suberectum, funiculo brevi. Stigma sessile, penicillato-capitatum. Fructus bacciformis obovatus, strato extimo (s. carnosum) et perigonio et pericarpium parte exteriori constante, intimo s. endocarpio nucamentaceo. Semen endocarpio conforme. Albumen copiosum. Embryo cotyledonibus parvis et fere rotundatis, radiculæ conoidæ circiter æquilongis. — Frutices, regiones tropicas subtropicasve Indiæ orientalis et insularum malaicarum necnon Abyssiniæ incolentes. Folia alterna, serrata, sæpe rugosa, subtus cinereo- v. niveo-tomentosa. Stipulæ axillares, bifidæ. Pedunculi sparsim bracteolati, in quaque axilla gemini. Receptaculum fructiferum vix carnosum.

1. *D. LONGIFOLIA*, ramulis cinereo- vel fusco-tomentosis hirtisve, foliis lanceolatis v. oblongo-lanceolatis sensim acuminatis basi ut plurimum obtusis rotundatisque rarius subcordatis argute serrulatis supra rugosis et molliter pubescentibus aut demum asperulis subtus breviter cinereo-tomentosis nervis hirtis, capitulis s. glomerulis unisexualibus androgynisve in apice ramulorum pedunculi furcati v. dichotomi solitariis aut didymis, perigonio masc. 4-partito bracteolas receptaculi superante. ♀ Copiosa in reg. trop. et subtrop. Indiae orientalis, in montibus Himalaya et Khasia, ad altit. 2000 metr. adscendens, in prov. Carnatic, Canara, Concan, cæt., et in ins. Zeylantia et Java. — *D. velutina* Gaudich.; Wedd. Monogr. p. 460. t. 15. A. f. 19. *Urtica longifolia* Nic. Burm. Fl. Ind. p. 297. *U. angustata* Bl. Bijdr. p. 499. *U. muricata* Heyne, in Wall. Cat. n. 4612. A. C. D. *Boehmeria angustata* Hassk. Hort. bogor. p. 79. *B. dichotoma* ejusd. l. c. *Conocephalus nireus* Wight, Icon. pl. Ind. 6. p. 7. t. 1959. *Mississya velutina* Wedd. in Ann. sc. nat. l. c. *Morocarpus velutinus*, M. longifolius et M. angustatus Bl. Mus. bot. lugd. bat. 2. p. 156. M. longifolius Miq. Fl. Ind. bat. l. c. — Frutex 2-3-metralis, ramulis gracilibus magis minusve elongatis dense pubescenti- vel tomentoso-hirtis demumque glabrescentibus, epidermide sicca sub hirsutie obscure fusco-rubente. Folia 4-15 centim. longa vel multo longiora, vulgo 12-30 millim. lata, ima basi integra et in insertione petioli plus minus emarginata, cæterum æqualiter serrata vel serrulata, nervis basilaribus ultra medium limbum productum cum sequentibus arcuatim aut reticulatim anastomosantibus in sicco costaque fuscescentibus, lamina exsiccata rigide membranacea, petiolo 5-20-millimetrali aut in var. δ multo longiore indumento ut in costa. Stipulæ oblongo-lanceolatae, 6-10 millim. longæ, apice bifidæ, extus puberulæ, nervis dorsalibus pilosis, siccitate pallide ferrugineæ, mox deciduæ. Glomeruli masc. aut fem. vel androgyni, pedunculo longitudine variabili furcato aut bis v. ter dichotomo pubescenti-tomentoso, ramulis cymarum fem. quam in inflorescentia masc. plerumque longioribus parce bracteolatis. Flores masc. sessiles aut breviter pedicellati, alabastro circiter sesquimillimetrali extus puberulo siccitate pallide ferrugineo, bracteis suffulcrantibus concoloribus ovatis ciliolatis. Capitula fem. floribunda, floribus bracteolis parvis ciliolatisque stipatis, fructifera pisiformia 6 millim. circiter lata, bacculis in vivo rubentibus v. anreis, siccitate fuscis. (v. s.)

β, *virens*, foliis obtuse crenato-serrulatis subtus cinereo-virentibus. — *Morocarpus virens* Bl. l. c. (v. s.)

γ, foliis elliptico-lanceolatis basi obtusis rotundatisve acute serratis, (v. s.)

δ, foliis late ovato-ellipticis basi subcuneatis vix rugosis acute serratis. (v. s.)

2. *D. EDULIS* (Wedd. Monogr. p. 462), ramulis pube brevi adpressa vestitis demumque glabratiss, foliis oblongo-lanceolatis ellipticisve sensim breviterque acuminatis basi sæpius obtusis rotundatisve in dimidia parte superiore acute et inæqualiter serrulatis supra rugosis lævibusque et glabratiss subtus tenuiter subcanescenti-tomentosis s. puberulis nervis pubescentibus, capitulis unisexualibus (masculis et femineis in distinctis ramulis), aut androgynis in apicibus ramulorum pedunculi furcati v. subdichotomi solitariis didymisve. ♀? In Japonia ubi Janatsi-itsigo s. Toon-itsigo dicitur. — *Boehmeria Janatsi*-(s. Janagi-) itsigo Siebold, in Act. Batav. 12. p. 70. *Morocarpus edulis* Sieb. et Zucc. l. c. excl. pl. masc.; Bl. Mus. bot. lugd. bat. 2. p. 156. t. 16. *Mississya parvifolia* Wedd. in Ann. sc. nat. l. c. — Ramuli graciles, breves, laterales, pennæ corvinæ circiter crassitie, apice foliosi, lateraliter florigeri, e ramo annotino glabrato fusco prodeuntes. Folia 2-4 centim. longa v. longiora, interdum cuspidata, supra sparsim pilosa denique glabrata, tomento paginæ inferioris tenuissimo virente. nervis ut in præcedente, petiolo 2-3-millimetrali. Stipulæ ovato-lanceolatae petiolis longiores in apice ramulorum imbricatæ, breviter bifidæ, nervis et margine pilosulæ. Capitula globulosa, 2-4-millim. lata. pedunculis plerisque furcatis puberulis bracteolas ovatas quam in spec. cæt. paulo majoribus hinc et inde gerentibus. (v. s. in hb. Monac.)
3. *D. DICHOTOMA* (Wedd. l. c.), ramulis breviter fusco-pubescentibus puberulisve, foliis ovato- v. elliptico-oblongis lanceolatisve anguste acuminatis basi

obtusis rotundatisve v. cordatis serratis v. crenato-serratis supra rugosis et asperulis (junioribus pilosulis) subtus in intermedio cinereo-tomentosis nervis pubescentibus, capitulis unisexualibus androgynisve in apicibus ramulorum pedunculi dichotomi solitarii didymisve, perigonio masc. tripartito. 3 In sylvis montosis ins. Javæ. — *Urtica dichotoma* Bl. Bijdr. p. 499. *Leucocnide dichotoma* Viq. pl. Jungh. 1. p. 38. L. sororia et L. affinis ejusd. 1. c. p. 38 et 39. *Morocarpus dichotomus* et M. affinis Bl. Mus. bot. lugd. bat. 2. p. 157; et M. sororius Miq. Fl. Ind. bat. 1. c. — Ramuli juniores pube brevi subadpressa et siccitate fusciscenti vestiti, demum glabrati, epidermide sicca fusca aut nigricante. Folia 6-14 centim. longa, 3-7 centim. lata, lamina exsiccata subcoriacea supra fusca subtus fusco-reticulata intermedio nonnunquam albidovirente, petiolo 1-6 centim. longo pubescenti-tomentoso. Cætera ut in D. velutina, a qua differt præcipue floribus masc. trimeris. (v. s.)

4. D. BICOLOR, ramulis superne breviter canescenti-tomentosis, foliis anguste lanceolatis sensim acuminatis basi acutis obtusatisve acute serrulatis v. crenulato-serratis supra planis aut demum rugosis et magis minusve asperatis (necnon junioribus parce pilosulis) subtus niveo-tomentosis, capitulis s. glomerulis unisexualibus androgynisve in quavis axilla solitariis geminis aut quaternis sessilibus vel breviter pedunculatis, perigonio masc. 4-partito quam bracteolæ receptaculi brevior. 3 In reg. trop. Himalayæ boreali-occidentalis, altit. 1500 metr. a Kumaon ad Marri; in Afghanistan (Griffith) et in montibus Abyssiniæ, altit. 1700-2600 metr. (Schimper, n. 862, cæc.). D. hypoleuca Wedd. Monogr. p. 463. t. 15. A. f. 10 et 11. *Urtica bicolor* Roxb. Fl. ind. 3. p. 589. *Urtica salicifolia* Don Prodr. fl. nepal. p. 60. *Boehmeria* (Procris) hypoleuca Hochst. in A. Rich. Tent. fl. Abyss. 2. p. 264. *Missiessya hypoleuca* Wedd. in Ann. sc. nat. 1. c. *Morocarpus salicifolius* Bl. Mus. bot. lugd. bat. p. 157. — Frutex, ramulis gracilibus vimineis superne tomentosis canescentibusque inferne puberulis vel omnino glabris et epidermide siccitate fuscorubente vestitis. Folia parum patentia, 3½-15 centim. longa, 1-3 centim. lata, toto margine dentata, nervis basilaribus tenuibus vix ultra medium limbum productis cum nervis primariis sequentibus eleganter arcuatum anastomosantibus costaque incano-tomentellis, intermedio transversim paralleleque reticulato, indumento paginæ inferioris undique (costa interdum excepta) adpresso et nervis exceptis candidissimo, lamina exsiccata rigide membranacea v. subcoriacea, petiolo 4-20 millim. longo ut costa pubescente v. tomentoso. Stipulæ lanceolatae, 5-8-millimetrales, apice bifidæ, in nervis et lacinis adpresse pilosæ, siccitate ferrugineæ, cito deciduæ. Glomeruli (s. capitula) pisiformes, in eodem ramo nunc omnes unisexuales, nunc superiores masculi, inferiores feminei et intermedii androgyni: masculi feminei sæpius fere duplo majores, pauciflori, floribus majusculis (alabastro nempe 2 millim. circiter lato), singulis bractea ampla siccitate ferruginea puberula involutis, segmentis late ovatis extus niveo-tomentosis. Capitula fem. ut plurimum subsessilia, valde multiflora, floribus bracteolis parvis ciliatisque suffultis; fructifera globulosa, interdum confluentia s. didyma, 5-6 millim. lata, bacculis circiter millimetrum longis stigmatis vestigio apiculatis siccitate lutescentibus. (v. s.)

5. D. LEUCOPHYLLA, ramulis glabris, foliis amplis ovato-rotundatis breviter acuminatis basi sæpius subcordatis inæqualiter sinuato-serratis rarius integris planis supra hispidulis sed mox glabris subtus niveo-tomentosis nervis pubescentibus, capitulis in apicibus ramulorum pedunculi ter quaterve dichotomi solitarii, perigonio masc. 4-partito bracteolas receptaculi superante. 3 In reg. trop. Himalayæ orientalis prov. Sikkim (Wallich) et in montibus Khasia usque ad altit. 1200 metr. (J. D. Hooker et Thomson), necnon in Zeylan (Walker). — D. Wallichiana Monogr. p. 464. t. 14. *Urtica leucophylla* Wall Cat. n. 4067. *Missiessya Wallichiana* Wedd. in Ann. sc. nat. 1. c. Ramuli præsi, siccitate longitudinaliter rugosi, epidermide cinereo-v. fuscorubente oblecti, apice tantum foliati, internodiis (saltem in ramulo florifero) brevissimis. Folia vulgo 10-20 centim. longa, 8-17 centim. lata, vel majora, juniora basi fere acuta, adulta rotundata v. subcordata, acumine triangulari centimetrali, nervis basilaribus ad trientem v. quartam partem superiore limbi excurrentibus cæterisque primariis in apicibus acuminatis serraturarum limbi desinentibus, secundariis rectiuscule transversis et mire invicem paral-

lelis in planta exsiccata fuscescentibus, lamina ipsa membranacea, petiolis 3-15 centim. longis puberulis. Stipulæ late ovatæ, 10-18 millim. longæ, bifidæ, extus puberulæ vel glabræ, subpersistentes, lacinii pubescentibus. Cymæ quam petioli breviores v. paulo longiores, geminæ, longiuscule pedunculatæ, pedunculo ramisque divaricatis demum subcentimetralibus villosis. Capitula florifera utriusque sexus conformia, globulosa, multiflora, floribus masc. sessilibus glabris in vivo rubentibus, femineis illis D. velutinæ fere conformibus. Capitula fructifera 8-10 millim. crassa, bacculis rubentibus. (v. s.)

Subtrib. IV. *MAOUTIÆ* Wedd. *Monogr.* p. 342. Perigonium fem. nullum. v. brevisimum liberumque (calyculus) et in fructu siccum. Stigma varium cum apice ovarii continuum.

XXX. LEUCOSYKE Zoll. et Moritz. in *Moritzii Verzeichn. d. Pflanz. Moritz.* p. 76; *Miq. in Zoll. System. Verzeichn.* p. 100; *Fl. Ind. bat.* 1 (2). p. 264. — *Mississya Gaudich. Bot. Voy. Bonite, t.* 93; *Bl. Mus. bot. lugd. bat.* 2. p. 162; *Wedd. Monogr.* p. 465; *Seem. Fl. Vit.* p. 244.

Flores sæpius dioici, masculi glomerulati, feminei receptaculo globuloso sessiles v. breviter pedicellati denseque aggregati, capitulis in axillis foliorum plerumque geminis raro corymbulosis aut irregulariter cymosis. *Masc.* Perigonium 4- v. 5-partitum, segmentis ovatis acutis in præfloratione valvatis, alabastro acutiusculo. Stamina 4 v. 5. Pistilli rudimentum conicum, lanatum v. glabrum. *Fem.* Perigonium breve, cupuliforme, ab ovario discretum, breviter et vulgo obtuse 4-5-dentatum. Ovarium oblique elliptico-ovatum, subcompressum, glabrum vel superne et margine strigillosum. Ovulum imo loculo funiculo brevi affixum, micropyle sæpius infundibuliformi-ampliata fimbriatæque et summo loculo cohærente. Stigma penicillato-capitulum v. subpestatum longe papillosum. Fructus subexsuccus, pericarpium nempe strato extimo vix carnosulo, intimo chartaceo. Semen pericarpium cavitati conforme, testa tenuissima. Albumen parcum. Embryo cotyledonibus ellipticis, radícula cylindrica crassiusculæque vix longioribus. — Arbusculæ v. frutices malaici et oceanici. Folia distiche alterna, subæquilateralia, conspicue v. subobsolete serrata, trinervia, subtus canescenti-tomentosa, cystolithis punctiformibus. Stipulæ in unam axillarem bifidam connatæ. Glomeruli masc. bracteati. Pedicelli florum fem. teretes v. superne incrassati, bracteolati. Receptaculum fructiferum carnosum. — Plantæ hujus generis ab omnibus aliis Boehmeriis distinguendæ sunt foliis plane distichis.

A. — *Capitula in singulis axillis solitaria v. gemina.*

1. *L. CANDIDISSIMA*, foliis late elliptico-ovatis acuminatis basi rotundatis aut obtusis ultra basim integram acute serratis supra minutissime strigosis asperulisque subtus niveis, stipulis magnis oblongis subpersistentibus, capitulis solitariis v. geminis majusculis pedunculatis. *♂* Ad ripas et in sylvis montanis ins. Javæ, altit. 1000-1300 metr. (Zollinger, n. 692, etc.). — *L. javanica* Zoll. et Moritz l. c.; *Miq. Fl. Ind. bat.* 1 (2). p. 265. *L. javensis* *Miq. in Zoll. Syst. Verzeichn.* p. 100. *Urtica candidissima* Bl. *Bijdr.* p. 498. *Boehmeria candidissima* Hassk. *Cat. hort. bogor.* p. 79. *Leucocnide candidissima* *Miq. Pl. Jungh.* 1. p. 37. *Mississya javanica* Wedd. in *Ann. sc. nat.* 4<sup>e</sup> sér. 1. p. 195. *M. candidissima* Bl. *Mus. bot. lugd. bat.* 2. p. 165; *Wedd. Monogr.* p. 466. *Touchardia candidissima* Bl. l. c. t. 24. — Arbuscula, ramulis pennæ cyneæ circiter crassitie vel etiam crassioribus teretiusculis adpresse subcinereo-pubescentibus prætereaque interdum patentim pilosis ad petiolorum insertiones cicatricibus stipularum annulatis notatis, internodiis superioribus 1-3-centi-

metralibus. Folia 1-2 decim. longa, 7-12 centim. lata, sæpius elliptica, ultra medium sensim attenuata, apice autem tantum acuminata, acumine subcentimetrali ut reliquis limbus serrato acutoque, serraturis in parte inferiore limbi plerumque obsoletis in parte superiore contra sæpius acutis aut etiam acuminatis, nervis basilaribus subtus prominentibus ad quartam partem apicalem limbi productis et in margine desinentibus, primariis cæteris numerosissimis parallele patentibus s. transversis necnon a triente superiore ad basim costæ gradatim imminutis et in medio intervenii cum venis (fere duplo numerosioribus) nervorum basilarium arcuatim connexis, lamina exsiccata subcoriacea pagina superiore nervisque inferioribus fuscescentibus, petiolo valido 2-6 centim. longo ut ramuli ipsi et costa adpresse pubescente et interdum magis minusve hirta. Stipulæ quam in cæteris speciebus majores, basi late amplexicaules, 2-4 centimetra longitudine metientes, more solito bifidæ, dorso molliuscule hirta-pilosæ, magis minusve persistentes. Capitula masc. magnitudine nucis avellanæ minoris, pedunculo sesqui-bi-centimetrali suffulta, floribus numerosissimis pedicellatis in vivo pallide rubellis bracteolisque ovatis et apice pilosis stipatis, perigonii 5-partiti segmentis extus et superne strigosius, pistilli rudimento brevissimo vix lanato. Capitula fem. masculis paulo minora, nempe 8-10 millim. diametro vix excedentia breviusque pedunculata, floribus creberrimis confertissimisque breviter pedicellatis (pedicellis teretibus), perigonio inæqualiter 4-5-dentato, achænio oblique ovato sub stigmate altero latere gibboso. (v. s.)

2. *L. HISPIDISSIMA* (Miq. Fl. Ind. bat. l. c.), foliis late ellipticis v. elliptico-ovatis breviter acuminatis excepta basi rotundata acute serratis v. crenato-serratis supra setoso-hispidis asperimisque subtus cinerascensibus nervis hispidis, stipulis oblongo-lanceolatis deciduis, capitulis masc. geminis majusculis pedunculatis. ♂ In ins. Manillæ montibus Igorrotes (Callery). — Missiessya *Hispidissima* Wedd. in Ann. sc. nat. l. c.; Monogr. p. 468. — Ramulus speciminis unici examinati crassitie circiter calami scriptorii, teres, puberulus, lævis, cinerascens, ad insertionem petiolorum annulatus, internodiis elongatis. Folia 10-14 centim. longa, 6½-8 centim. lata, serraturis parvis interdum apiculatis, nervis basilaribus ut costa tenuibus subtus prominulis juxta partem quartam superiorem limbi in margine evanescentibus, venis cæteris fere ut in specie præcedente dispositis, lamina exsiccata rigida supra pallide fusca et nonnihil glaucescente nervis impressis subtus cinerascens nervis secundariis tertiariisque fuscescentibus reteque prominulum fingentibus, petiolo 3-6-centimetrali puberulo sparsimque et adpresse pilosulo albedo. Stipulæ sesquicentimetrales, extus adpresse pubescentes. Capitula masc. pisi majoris circiter magnitudine, pedunculo centimetrali adpresse pubescente, floribus breviter pedicellatis, perigonii 5-partiti segmentis ovato-lanceolatis acutis extus superne strigosius, pistilli rudimento dense lanato. Cætera ignota. (v. s. in hb. Mus. par.)
3. *L. CAPITELLATA*, foliis oblongo- v. elliptico-lanceolatis lanceolatisve acuminatis basi obtusis aut acutis in toto fere margine v. in parte tantum ejusdem acute serratis subchartaceis supra subtiliter hispidulis subtus cinereo- v. albi-to-tomentosis et sæpius utrinque asperulis nervis basilaribus vulgo ad partem quintam superiorem limbi extensis venulis e costa ortis plerisque rectiusculis parallele transversis et plus minus reticulatis, stipulis lanceolatis cito deciduis, capitulis sæpius geminis breviter pedunculatis. ♂ Ad ripas et in sylvis montanis ins. Javæ (Junghuhn; Zollinger n. 3540), ins. Sumbawa (Zollinger, n. 3423), Manilla (Cuming n. 1766 et 1842; Callery, n. 7), necnon in Moluccis (Gaudich.). — *L. alba* et *L. bimensis* Zoll. et Moriti l. c.; Miq. in Zoll. System. Verzeichn. p. 100; Fl. Ind. bat. l. c. *Urtica capitellata* Poir. Encycl. suppl. 4. p. 244. *U. sphærocephala* Roxb. Fl. ind. 3. p. 589. *Missiessya fagifolia* Gaudich. l. c.; Bl. l. c. p. 161; Wedd. Monogr. p. 468. t. 15. B. f. 2. *M. fagifolia*, *M. alba* et *M. bimensis* Wedd. in Ann. sc. nat. l. c. *Leucocne alba* Miq. Pl. Jungh. 1. p. 37. — Ramuli strigillosi sæpeque superne hirtelli vel pubescentes albidique, crassitie variabili, demum glabrati, epidermide cinerascens aut castanea, ad insertionem petiolorum linea s. cicatrice annulari notati, merithallis 1-3 centim. longis. Folia magnitudine admodum variabili, 6-16 centim. longa, 2-6 centim. lata, sæpius sensim et longiuscule acuminata,

serraturis haud raro acuminatis, nervis basilaribus eleganter arcuatis margini subparallelis ad partem quartam vel quintam superiorem limbi productis et in margine desinentibus, nervis cæteris e costa ortis plerisque parallele transversis æqualibusque, petiolo circiter centimetrali ut costa ipsa et nervi pube fuscula brevissima obito pilisque longioribus albis plus minusve consperso, lamina exsiccata rigida supra fuscescente nervis impressis subtus concinne fusco-reticulata. Stipulæ sesqui-bi-centimetricæ petiolisque ut plurimum longiores, extus puberulæ et in nervis adpresse pilosæ. Capitula (interdum teste Miquelio androgyna) pedunculis centimetralibus aut paulo brevioribus suffulta, utriusque sexus valde congestiflora. Flores masc. ut in specie precedente. Flores fem. pedicellati, pedicellis sub perigonio obtusæ dentato magis minusve incrassatis. Achenium oblongo-ovatum, sub stigmate vix gibbosum, in altero margine strigillosum. (v. s.)

β, *ochracea* (Blume), foliis elliptico-oblongis v. oblongo-lanceolatis basi obtusis v. acutiusculis adpresse serrulatis subtus pallide ochraceis. ♂ ? In ins. Borneo.

γ, *angustata* (Blume), foliis oblongo-lanceolatis basi suboblique attenuatis serrulatis uno latere versus basim vulgo altius integerrimis. ♂ ? In ins. Banda et Celebes.

δ, *tenera* (Blume), foliis tenere membranaceis oblongo-lanceolatis basi obtusis v. subattenuatis venulis transversalibus obsolete reticulatis. ♂ ? In ins. Banda.

ε, *retusa* (Blume), foliis oblongo-lanceolatis basi obtusis v. retusis venulis transversalibus reticulatis. ♂ ? In ins. Borneo.

ζ, *Blumei*, foliis ellipticis utrinque subacutis v. sæpius oblongis v. oblongo-lanceolatis (7-20 centim. longis) basi obtusis v. acutiusculis medio v. paulo infra medium ad apicem acute et sæpe obsolete serrulatis subcoriaceis venulis transversalibus rectiusculis quam in typo distinctius reticulatis. ♂ ? In littore australi ins. Javæ, necnon in ins. Moluccis, cæt. — *Missiessya mutabilis* Bl. l. c. p. 163; Wedd. l. c. p. 471. *Leucosyke mutabilis* Miq. l. c. p. 265. (v. s. typ. Bl. in hb. Mus. par.)

η, *villosa*, foliis oblongo-lanceolatis acuminatissimis basi obtusis v. acutiusculis supra basim ad apicem argute serrulatis subcoriaceis supra asperis subtus albidis nervis petiolisque sicut ramuli villosis venulis transversalibus rectiusculis approximatis subreticulatis. ♂ ? In ins. Borneo, Celebes, cæt. — *Missiessya villosa* Bl. l. c.; Wedd. l. c. *Leucosyke villosa* Miq. l. c. p. 265. — Folia 7-20 centim. longa, 3-5 centim. lata, petiolo 4-6-millimetrali. (v. s. typ. Bl. in hb. Mus. par.)

θ, *elongata*, foliis lanceolatis rectiusculis acuminatissimis basi obtusis v. acutiusculis inferne integerrimis medio v. supra medium ad apicem obsolete serrulatis membranaceis venulis transversalibus rectiusculis obsolete reticulatis, pedunculis fem. petiolo longioribus (Blume). ♂ ? In ins. Celebes, Banda, cæt. — *Missiessya elongata* Bl. l. c. p. 164; Wedd. l. c. p. 472. *Leucosyke elongata* Miq. l. c. — Folia 9-17 centim. longa, 3-5 centim. lata, petiolis (4-6-millimetralibus) ramulisque et nervis subtus pilis appressis puberulis. — Foliis magis elongatis tenuioribus supra crebre punctatis et capitulis ♀ longius pedunculatis a var. præced. diversa.

ι, *falcata*, foliis lanceolatis subfalcatis superioribus acuminatissimis (infimis minoribus acutis v. obtusis) basi obtusis v. acutiusculis inferne integerrimis a medio ad apicem obsolete repando-serrulatis chartaceis supra asperulis subtus cæcio-albis venulis transversalibus rectiusculis obsolete reticulatis, pedunculis fem. petiolum æquantibus (Blume). ♂ ? In provincia celebica Menado. — *Missiessya falcata* Bl. l. c.; Wedd. l. c. *Leucosyke falcata* Miq. l. c.

κ, *celtidifolia*, foliis parvis oblongo- v. lanceolato-ovatis sensim acuminatis basi rotundatis obtusis acutisve in toto fere margine v. a tertia aut etiam a dimidia parte inferiore ad apicem argute rariusve subobsolete serrulatis

supra asperimis. ♂? In montibus ins. Philippinarum (Cuming, n. 1360 et 1842). — *Missiessya celtidifolia* Gaudich. Bot. Voy. Bonite, t. 93; Bl. l. c. p. 163; Wedd. l. c. p. 470. t. 15, B. f. 1. *M. celtidifolia* et *M. aspera* Wedd. in Ann. sc. nat. l. c. *Leucosyke celtidifolia* Miq. l. c. — A typo differt omnibus partibus minoribus foliis multo asperioribus et capitulis (interdum omnino sessilibus), e floribus multo paucioribus compositis. (v. s.)

4. *L. SUMATRANA* (Miq. l. c.), foliis amplis ovato-v. elliptico-oblongis acuminatis basi obtusis v. acutiusculis integrisque cæterum acute dentato-serratis subchartaceis supra hispidis asperulisque subtus tenuiter canescenti-tomentosis, nervis basilaribus ad sextam partem superiorem limbi extensis, venulis inter hosce et costam angulato-curvatis et distincte reticulatis, stipulis deciduis, capitulis majusculis sessilibus. ♂? In ins. Sumatra (Korthals). — *Missiessya sumatrana* Bl. l. c. p. 165; Wedd. l. c. — Ramuli crassi, rufescenti-tomentosi et pilis albidis villosi. Folia 15-28 centim. longa, 7-12 centim. lata, nervis basilaribus præter basiales pluribus costaque et petiolo (3-8-centimetrali) indumento eodem ut in ramuli vestitis. Stipulae non visæ. Capitula fructifera (solum suppetentia) globulosa, plus quam centimetrum lata, acheniis confertissimis, calyculo minimo disciformi omnino sessili margine subintegro. — A *L. alba* imprimis distinguitur foliorum venis transversalibus sursum in angulum connexis necnon rectiuscule anastomosatis. (v. s. in hb. Mus. par. communicat. ex hb. lugd. bat.)

β, *sericea* (Blume), pubescentia ramulorum magis sericea et appressa. ♂? In ins. Sumatra.

5. *L. OCHRONEURA*, foliis elliptico-oblongis v. oblongo-lanceolatis utrinque acutis v. apice breviter acuminatis basi excepta laxe et minute serrulatis chartaceis supra scabridis subtus canescenti-tomentosis venulis transversalibus inter nervos basiales et costam curvatis et distincte reticulatis, stipulis deciduis, capitulis geminis pedunculatis. ♂? In ins. Moluccis. — *Missiessya ochroneura* Bl. l. c.; Wedd. l. c. p. 474. — Ramuli petioli nervique subtus ochraceo-v. fuscascenti-tomentelli et pube sericea canescenti vestiti. Folia 12-20 centim. longa, 5-8 centim. lata, nervis basilaribus ut in præcedente. Stipulae majusculæ (petiolos excedentes), lanceolatae, mox deciduæ. Capitula fem. magnitudine pisi, pedunculis 5-6 centimetralibus canescenti-puberulis. — *L. sumatranae* affinis, sed foliis brevius petiolatis serraturis ad margines multo minoribus et interdum fere inconspicuis, venulis transversalibus quoque inter nervos minus curvatis. (v. s. in hb. Mus. par. communicat. ex hb. lugd. bat.)
6. *L. CELEBICA* (Miq. l. c.), foliis oblongo-lanceolatis acuminatissimis basi inæquali obtusiusculis coriaceis paulo supra basim ad apicem acute serrulatis supra scabriusculis subtus incano-tomentosis triplinerviis venulis inter nervos basiales et costam transversalibus rectiusculis v. leviter curvatis et distincte reticulatis. ♂? In ins. Celebes. — *Missiessya celebica* Bl. l. c.; Wedd. l. c. — A præcedente distinguitur foliis triplinerviis longius acuminatis basi obtusis et distinctius serrulatis. (Blume.)
7. *L. BORNEENSIS* (Miq. Fl. Ind. bat. l. c. p. 267), foliis ovatis v. oblongo-ovatis acuminatis basi rotundatis v. obtusis ima basi excepta acute serratis membranaceis v. chartaceis supra asperato-hispidis subtus canescenti-tomentosis, nervis basilaribus ad trientem superiorem limbi productis, venulis inter hosce et costam distincte ramosis reticulatisque, stipulis lanceolatis persistentibus, capitulis pedunculatis. ♂? In ins. Borneo (Korthals). — *Missiessya borneensis* Blume Mus. bot. lugd. bat. 2. p. 164; Wedd. Monogr. p. 473. — Ramuli (in specim. unico viso), graciles, superne subadpresse pubescenti-pilosi. Folia 7-11 centim. longa, 3-5 centim. lata, nervis primariis præter basiales utrinque 2-3 venulisque pubescentibus et in sicco fuscis, petiolo 4-6-millimetrali. (v. s. in hb. Mus. par. communicat. ex hb. lugd. bat.)
8. *L. RUBIGINOSA* (Miq. l. c.), foliis oblongo-ellipticis breviter acuminatis basi acutis integrisque cæterum laxe serrulatis coriaceis supra hispidis scabriusculisque et demum glabris subtus canescentibus et rubiginoso-venosis (in sicco), nervis basilaribus fere ad apicem limbi protensis venulis inter hosce

et costam transverse arcuatis simplicibusque nec reticulatis, stipulis deciduis, capitulis subsessilibus. ♂? In ins. Borneo (Korthals, Müller). — *Missiessya rubiginosa* Bl. l. c.; Wedd. l. c. — Ramuli tomento rubiginoso denso pilis albidis appressis immixto vestiti. Folia 4-8 centim. longa, 2-3 centim. lata, nervis primariis præter basiales ut plurimum utrinque solitariis, petiolo 4-12-millimetrali sicut nervi tomento rubiginoso et pilis albidis vestito. Stipulæ lanceolatæ, extus adpresse pilosæ. Capitula fructifera (solum visa) parva, solitaria, brevissime pedunculata v. sessilia, achæniis valde compressis, calyculo minimo. — Species distinctissima nervatione foliorum. (v. s. in hb. Mus. par. ex hb. lugd. bat. a cl. Miquelio benigne communic.)

B. — *Capitula in singulis axillis plura.*

9. *L. CORYMBULOSA*, foliis ovato-lanceolatis ovatisve acuminatis acutissimis supra basim obtusam rotundatamve argute serrulatis supra adpresse puberulis lævisculisque aut asperulis subtus cinereo- vel albedo-tomentosis, stipulis mox deciduis, capitulis sæpius pluribus in singulis axillis pedunculatis corymbulosis. ♂? In ins. Viti s. Fidji (Cunningham, Milne, Wilkes, Seemann). — *Missiessya corymbulosa* Wedd. l. c. p. 475. t. 15. f. 8; Seem. Fl. Vit. p. 244. Ramuli graciles, flexuosi, adpresse cinereo-pubescentes demumque glabrati, ad insertionem foliorum cicatrice annulari notati. Folia 4-12 centim. longa, 1½-5 centim. lata, a medio sensim attenuata, ima basi aut tertia vel quarta parte inferiore integra, nervis basilaribus leviter arcuatis ad trientem superiorem limbi vel etiam ultra extensis subtus prominulis, secundariis tertiariisque eximie reticulatim connexis, pagina superiore vix asperula adpresse puberulo-hispida, inferiore brevissime cano-tomentosa nervis petiolisque (semicentimetralibus) molliter pubescentibus. Stipulæ lanceolatæ, 8-12 millim. longæ, involutæ, pubescentes rariusve glabrata, mox deciduæ. Capitula petiolos plerumque superantia, plasi circiter magnitudine, pedunculis filiformibus 5-10-millimetralibus nunc geminis axillaribusque nunc pluribus atque in ramulo abbreviato corymbulosis adpresse puberulis, floribus masc. perigonii laciniis extus breviter hispidis, femineis quam in cæteris speciebus longius pedicellatis, perigonii fructiferi lobis acutiusculis cillolatis interdumque hispidulis, achænio sesquimillimetrali. (v. s.)

XXXI. *MAOUTIA* Wedd. in *Ann. sc. nat.* 4<sup>e</sup> sér. 1. p. 193; *Monogr.* p. 476; *Miq. Fl. Ind. bat.* 1 (2). p. 272; *Seem. Fl. Vit.* p. 244. — *Boehmeria* sect. *Miq. Pl. Jungh.* 1. p. 34. — *Lecanocnide* Bl. *Mus. bot. lugd. bat.* 2. t. 12. — *Urticæ* spec. auct.

Flores monoici v. dioici glomerulati s. capitellati, glomerulis laxè irregulariterque cymosis, cymis in singulis axillis sæpius geminis. *Masc.* Perigonium 5-partitum, segmentis ovatis subacuminatis extus hispidulis in præfloratione valvatis, alabastro brevissime acuminato. Stamina 5. Pistilli rudimentum obovoideum, in lana densa nidulans. *Fem.* Perigonium deficiens (nisi in unica specie). Ovarium ovoideum, rectum, setosum aut subadpresse hispidum. Ovulum imo loculo affixum, suberectum. Stigma stylo brevi suffultum, sublaterale, lanceolatum v. subcapitatum, persistens, papillis brevibus elongatisve instructum. Achænium ovatum, subcompressum v. interdum obtuse trigonum, hispidum v. longiuscule setosum, pericarpium strato extimo carnosulo, intimo osseo s. nucamentaceo magis minusve compresso et nonnunquam crasse marginato. Semen ellipticum, compressum, pericarpium cavitatem explens. Albumen tenue. Embryo cotyledonibus ellipticis oblongisve radícula tereti et gracili paulo longioribus. — Frutices in insulis malais et oceanicis necnon in India orientali obvii. Folia alterna, crenulata v. serrata, trinervia, subtus cano-tomentosa, cystolithis punctiformibus. Stipulæ axillares, sæpius profunde bifidæ, glomerulis basi



bracteatis. Flores fem. omnino nudi v. bracteolis paucis parvisque stipati, in *M. ambigua* autem perigonio brevi calyculiformi illo Mississyæ simillimo muniti. — Genus distinctissimum et habitu et characteribus florum.

1. *M. PUYA* (Wedd. in Ann. sc. nat. l. c.; Monogr. p. 477. t. 16, B. f. 1-8), monoica, foliis lanceolato- v. ovato-ellipticis anguste acuminatis basi acutis s. cuneatis siccitate membranaceis supra hispido-pilosis asperisque subtus nervis exsertis niveo-tomentosis, achænio erostri subadpresse hispido. ♂ In Indiæ orientalis regionibus tropicis montium Himalaya, a Sikkim ad Garwhal, altit. 0-1200 metr., et in montibus Khasia, ad oras sylvarum præsertim frequens. — *Urtica Puya* Wall. Cat. n. 46C5. *Boehmeria frutescens* Hamilt. et Don Prodr. Fl. nepal. p. 59. Bæhm. *Puya Hassk.* in Journ. of bot., ann. 1851, t. 7. Frutex s. arbuscula 2-3-metralis, ramulis superne magis minusve hirtis et sub hirsutie breviter strigosis. Folia 5-20 centim. longa, 2-6 centim. lata, acumine lineari 10-15-centimetrali, inferne integra, serraturis reliqui limbi triangularibus et in margine exteriore sæpe nonnihil convexis, nervis basilaribus ad trientem vel quartam partem superiorem limbi productis ut et ipsa costa tenuibus pubescentibus, intervenio vix reticulato, lamina exsiccata supra subtusque in nervis fusciscente, petiolo gracili 1-5-centimetrali canescenti-pubescente. Stipulæ profunde bifidæ, segmentis lanceolatis apice fere subulatis margine et nervo dorsali pilosis. Inflorescentiæ unisexuales aut e glomerulis aliis androgynis aliis masculis constantes, masculæ et androgynæ majores, raro autem petiolos superantes, ramulis strigosos pilosulisque. Glomeruli masc. multiflori, sæpe 4-5 millim. lati s. pisiformes, floribus subæsilibus v. breviter pedicellatis, perigonii lobis subadpresse hispidis. Glomeruli fem. masculis plus quam dimidio minores, achæniis pyramidato-ovatis 1-1½ millim. longis obtuse 3-angulatis stigmate sessili ovato v. rotundato coronatis. (v. s.)
2. *M. ODONTOPHYLLA* (Miq. Fl. Ind. bat. 1 (2). p. 272), monoica, foliis late ellipticis v. elliptico-ovatis breviter acuminatis basi obtusis rotundatisve grosse dentato-serratis subcoriaceis supra rugosis hispido-pilosis asperatisque subtus niveo-tomentosis nervis pubescentibus, achæniis ut in præcedente. ♂ In locis graminosis dumosisque, ad pedes collium ins. Javæ (Junghuhn; Zollinger, n. 861). — *M. aspera* Wedd. in Ann. sc. nat. l. c.; Monogr. p. 478. *Boehmeria odontophylla* Miq. Pl. Jungh. 1. p. 35; in Zoll Syst. Verzeichn. p. 100. — Frutex circiter bimetralis, ramulis elongatis crassitie pennæ anserinæ v. cynææ superne dense tomentoso-hirtis inferne glabratissimis epidermide fusca lævique vestitis. Folia 7-11 centim. longa, 3½-7 centim. lata, subabrupte acuminata, toto margine (ima basi excepta) acute serrata, serraturis quam in *M. Puya* paulo minoribus, nervis basilaribus usque ad sextam partem superiorem limbi excurrentibus et cum sequentibus arcuatim anastomosantibus, lamina exsiccata supra fusca nervis impressis subtus fusco-cinereis, petiolis 2-3-centimetralibus cinereo-tomentosis v. hirtis. Stipulæ profunde bifidæ, laciniis 6-10-millimetralibus apice subulatis extus pubescentibus margine et nervo pilosis. Inflorescentiæ in speciminibus quæ videre licuit omnes androgynæ, ramulis inferioribus hirtulis, floribus achæniisque cæterum ut in specie supra memorata. (v. s.)
3. *M. RETICULATA*, dioica, foliis ovatis acuminatis basi cuneatis integrisque cæterum serratis (serraturis subacuminatis) membranaceis supra planiusculis hispidulisque subtus cinereo-tenuiter canescenti- s. cinereo-tomentosis intervenio distincte reticulato. ♂ ? In insulis Mariannis (Gaudichaud) et Philippinis (Née). — Ramuli superne hirtotomentosi canescentesque, epidermide partis glabræ obscure brunnea. Folia 10-15 centim. longa, 5-8 centim. lata, acumine angusto sicut reliquis limbus serrato, parte cuneata integraque limbi sextantem hujusce circiter æquante, nervis basilaribus ut in *M. odontophylla*, venulis supra impressis subtus in lamina exsiccata rete nigricans quam in specie citata magis distinctum fingentibus, petiolis 2-8 centim. longis indumento fere ut in ramulis. Stipulæ laciniis subulato-lanceolatis pilosis. Inflorescentiæ masc. (solum visæ) petiolis majoribus breviores, divaricato-ramosæ, ramulis pubescenti-hirtis, glomerulis parvis (nempe plerisque 3 millim. vix

latis), floribus breviter pedicellatis v. subsessilibus, perigonii segmentis puberulis. — Habitus speciei præcedentis, sed affinitates veræ ob defectum florum fem. adhuc dubiæ. (v. s.)

4. *M. DIVERSIFOLIA*, monoica v. dioica ?, foliis ovatis v. ovato-lanceolatis ~~menais~~ longiusculeque acuminatis basi oblique obtusatis rotundatisve minute crenulato-serratis coriaceis supra rugosis asperulisque subtus nervis exceptis niveo-tomentosis, achæniis trigonis breviter rostratis undique sed præsertim in angulis subadpresse setulosi, stigmatibus obliquo recurvo. ♂ In ins. Java (Jung-huhn; Zollinger, n. 2532). — *M. rugosa* Wedd. Monogr. p. 479; Miq. Fl. Ind. bat. 1 (2). p. 274. *Urtica diversifolia* Bl. Bijdr. p. 498. *Boehmeria diversifolia* Miq. Pl. Jungh. 1. p. 34; et in Zolling. System. Verzeichn. p. 100. *Lecanocnide diversifolia* Bl. Mus. bot. lugd. bat. 2. t. XII. — *Arbuscula* 3-6-metralis, ramulis dense breviterque lucano-tomentosis crassitie calami scriptorii v. crassioribus obtuse angulatis. Folia 8-16 centim. longa, 3-5 centim. lata et ut in hoc genere solet nonnihil invicem dissimilia, altero nempe paulo minore basi que magis rotundata et brevius petiolato, toto margine serrulata aut juxta petioli insertionem integra, nervis basilaribus fere usque ad basim acuminis excurrentibus subtus costa que pubescenti-tomentosis et in lamina exsiccata fusciscentibus intervenio eleganter reticulato foveolatoque, petioli 1-4 centim. longis breviter canescenti-tomentosis. Stipulæ subcentimetricales atque ut in congeneribus plerisque profunde bifidis, lacinii a basi sensim attenuatis nervo dorsali et marginibus pilosis. Inflorescentiæ unisexuales aut superiores masculæ v. androgynæ et inferiores femineæ, ramulosissimæ, petiolos superantes, ramis tenuibus vix divaricatis puberulis, glomerulis parvis laxis, floribus breviter pedicellatis. Flores masc. pentandri, perigonii segmentis adpresse hispidis. Achænia ovata, vix millimetralia, acutiuscule trigona, stigmatibus majusculo unilaterali s. in rostrum achæniis decurrente et illo longiore. (v. s.)

β, foliis ovatis quam in typo magis serratis submembranaceis, achæniis longius rostratis. ♂ In ins. Java (Zollinger, n. 1252). — *Maoutia Zollingeriana* Wedd. in Ann. sc. nat. l. c.; Monogr. p. 480. — Folia subtus magis minusve foveolata, nervis albidis. (v. s.)

5. *M. AUSTRALIS* (Wedd. Monogr. p. 480), sæpius dioica, foliis ovatis vel elliptico-ovatis lanceolatisve anguste acuminatis basi acutis v. obtusatis rotundatisve acuminis (interdum) basi que exceptis grossiuscule crenato-serratis serratisve siccitate plerumque subcoriaceis planis rugosulisve supra sparsim pilosulis et mox fere omnino glabratis lævibusque subtus niveo-tomentosis nervis adpresse pubescentibus, achæniis breviter rostratis inferne et in angulis præsertim patule v. subadpresse setosis, stigmatibus obliquo subrecurvo. ♂? In ins. Fidji s. Viti (Milne, Jacquinot, Leguillou, Seemann, etc.), Societatis (Bidwill) et Taiti (hb. Jaubert). — Ramuli elongati, teretiusculi v. obtuse angulati, crassitie circiter pennæ anserinæ, superne breviter canescenti-pubescentes, inferne glabrati, siccitate longitudinaliter rugosi fusciscentisque s. fusco-cinerei. Folia 8-20 centim. longa, 3½-7 centim. lata, acumine sub-integro v. serrato, serraturis reliqui limbi obtusiusculis acutis apiculatisve, nervis (basilaribus ut in præcedente) supra parum impressis, intervenio subtus vage reticulato, petioli breviter incanulo-pubescentibus 1-4 centim. longis. Stipulæ 18-20-millimetrales, usque ad medium circiter bifidis, extus pubescentes, nervo dorsali et margine pilosæ. Inflorescentiæ petiolos æquantibus v. parum superantes, divaricatis ramosæ; masculæ quam femineæ longius pedunculatæ, ramis puberulis, glomerulis seminibus cannabini circiter magnitudine, floribus subsessilibus, perigonio adpresse hispido; femineæ glomerulis (raro androgynis) quam masculi paulo minoribus, floribus sessilibus, achæniis millimetralibus, stigmatibus subunilateralibus rostrum excedente. — Præcedenti affinis. (v. s.)

6. *M. PLATTYSTIGMA* (Wedd. in Ann. sc. nat. l. c.; Monogr. p. 481. t. 16. B. f. 10), dioica, foliis oblongo-vel ovato-lanceolatis acuminatis basi rotundatis crenulato-dentatis siccitate membranaceis planis supra glabris lævibusque subtus nisi in nervis niveo-tomentosis, achæniis breviter rostratis inæqualiter patentem setosis, stigmatibus majusculo subcapitato quam rostrum achæniis longiore. ♂? In ins. Manilla Philippinarum (Cuming, n. 1441). — Miq. Fl. Ind.

bat. l. c. — Ramuli graciles, obtuse angulati, breviter canescenti-pubentes, laxe foliosi. Folia 8-15 centim. longa, 3-5 centim. lata, toto fere margine (etiā acumine) obtusiuscule dentata, nervis basilaribus ut in præcedente, lamina exsiccata supra fusca nervis subtus breviter pubescentibus cinereis, petiolis gracilibus patentibus 2-5 centim. longis. Stipulæ 10-12-millimetrales, ad medium circiter bifidæ, laciniis superne subulatis nervo medio cano-pubescente. Inflorescentiæ fem. (solum visæ) petiolos superantes, divaricato-ramosissimæ, ramulis adpresse incanulo-pubescentibus, achæniis subsessilibus plerumque quam in cæteris speciebus minoribus (id est stigmatē exempto semimillimetrum vix metientibus) fuscis inferne et in angulis præsertim setosis. (v. s.)

7. *M. SETOSA* (Wedd. in Ann. sc. nat. l. c.; Monogr. p. 482. t. 16. B. f. 9), monoica, foliis elliptico-ovatis breviter acuminatis basi rotundatis obtusiuscule crenulato-dentatis coriaceis supra in nervis sparsim pilosis subtus niveo-tomentosis, achæniis patule longiusculeque setosi rostro quam stigma longiore. ♂? In Manillæ montibus Igorrotes (Callery, n. 30). — Miq. Fl. Ind. bat. l. c. — Rami ramulique molliter canescenti-pubescentes, apice incani. Folia 3½-10 centim. longa, 2-5 centim. lata, sensim acuminata (acumine acuto), fere ab ipsa basi crenulata, nervis basilaribus usque ad partem quartam quintamve superiorem limbi protractis supra parum impressis subtus prominulis interveniisque fere concoloribus, lamina exsiccata rigida supra læviscula, petiolo 1-3-centimetrali tomentoso. Stipulæ profunde bifidæ, 5-10 millim. longæ, basi exteriore nervoque dorsali canescenti-pilose. Inflorescentiæ petiolis subæquilongæ, divaricato-ramosæ, pedunculis brevibus ramulisque densiuscule canescenti-pubescentibus, glomerulis parvis discretis aut fere confluentibus androgynis unisexualibusve. Flores masc. sessiles perigonii lobis late ovatis adpresse hispidis, pistilli rudimento obovato et ut in stirpibus præcedentibus densissime lanato. Achænium late ovatum, pedicellatum vel sessile, apice in rostrum longiusculum subabrupte acuminatum setisque longis rigidis patentibus obsitum abiens, stigmatē brevi. (v. s.)

8. *M. AMBIGUA* (Wedd. Monogr. p. 483), dioica, foliis ovatis acuminatis basi rotundatis v. subcordatis toto margine serratis coriaceis rugosis supra hispidis asperisque subtus incanis nervis hirtis intervenio distincte reticulato, achæniis compressis erostratis calyculatis fere glabris. ♂? In ins. Amboina. — Miq. Fl. Ind. bat. l. (2). p. 274. — Ramuli breviter cinereo-hirti demumque omnino (?) glabrati. Folia in specimine unico viso decimetralia, 5-5½ centim. lata, anguste acuminata, æqualiter serrata, serraturis parvis acutis aut in sicco obtusiusculis (apice inflexo), nervis subtus pubescenti-hirtis, basilaribus arcuatis ad partem quintam superiorem limbi productis, secundariis tenuibus et fere transversis, tertiariis reticulatis, petiolis valde inæqualibus, 1½-5 centim. longis, indumento ut in nervis. Stipulæ profunde bifidæ, laciniis 6-8-millimetralibus lineari-subulatis hirtisque. Inflorescentiæ fem. (masculis non visis) petiolis majoribus breviores, pedunculis circiter centimetralibus parce ramosis pubescentibusque, glomerulis (seminis cannabini magnitudine) sessilibus vel breviter pedunculatis, achæniis crebris oblique ovatis ¼ millim. circiter longis compressis superne in angulis parce et subadpresse setulosis, calyculo ut in genere *Mississya* brevissimo sub-5-denticulato et ad apices dentium setuloso suffultis, stigmatē parvo subcapitato crasse papilloso. — Stirps admodum insignis, *Boehmeriam* niveam habitu referens sed ad genus *Mississyam* spectans floribus femineis calyculatis. (v. s. in hb. Mus. par.)

XXXII. *MYRIOCARPA* Benth. Bot. Voy. Sulph. p. 168; Miq. in Mart. Fl. bras. fasc. 12. p. 197; Wedd. Monogr. p. 484.

Flores dioici aut rarius monoici, masculi (glomerulati) et feminei (calyculo 2-4-phyllo suffulti) in spicas racemosve gracillimos subunilaterales pendulos dense aggregati. Masc. Perigonium 4- aut rarius 3-5-partitum, segmentis ovatis obtusis ciliolatis. Stamina 3-5. Pistilli rudimentum minutum conicum glabrum. Fem. Perigonium nullum. Ovarium ellipticum, compressum, basi attenuatum, apice in stylum

longiusculum subabrupte acuminatum productum, margine styloque et pedicello setoso-ciliatis. Stigma laterale, semilunare, villosopapillosum, persistens. Ovulum funiculo gracili imo loculo affixum, erectum. Achænium ovario conforme. Embryo cotyledonibus rotundatis radícula conica dimidio longioribus. — Arbusculæ v. frutices in regionibus calidioribus Mexici, Novæ Granatæ et Peruvix præsertim obvii, facie propria, omnes magis minusve pubescentes. Folia alterna, sæpe majuscula, dentata, petiolata, trinervia, limbo exsiccato cystolithis linearibus a basi pilorum paginæ superioris radiantibus sub lente eximie ornato. Stipulæ axillares, integræ. Flores masc. et fem. breviter pedicellati sessilesve, feminei bractea minima infra calyculum sæpe muniti, calyculi folioli longiuscule ciliatis aut glabriusculis. — Genus admodum naturale, speciebus (genuinis forte paucissimis) pro speciminibus mancis non omnibus satis cognitis et pluribus inter se difficile dignoscendis.

1. *M. STIPITATA* (Benth. l. c. t. 55), dioica, ramulis pubescentibus et mox glabratiss, foliis ovato-vel obovato-ellipticis rariusve elliptico-lanceolatis breviter acuminatis basi rotundatis v. obtusis rarissime subcordatis inæqualiter serrato-denticulatis crenulatisve planis aut parum rugosulis siccitate membranaceis supra sparsim hispido-pilosis demumque glaberrimis nervis subtus adpresse pilosulo-pubescentibus, inflorescentiis fructiferis longitudine circiter foliorum, calyculi sæpius bifoliolati foliolis ovatis et fere glabris, achænio stipitato longiuscule setoso-ciliato. ♂ In imperio mexicano (Botteri, Bourgeau), Nova Granata (Goudot), Venezuela (Fendler, n. 1213), Bolivia (Wedd.) et Brasilia (Widgren). — Miq. l. c.; Wedd. Monogr. p. 485. t. 16, D. f. 7. *M. humilis* ejusd. in Ann. sc. nat. 3<sup>e</sup> sér. 18. p. 231. — Arbuscula 2-3-metralis v. frutex. Ramuli novelli subincano-pubescentes, sed demum glabrati, epidermide tunc (saltem in planta exsiccata) ferruginea. Folia magnitudine et forma variabilia, vulgo 10-18 centim. longa, 4-10 centim. lata, acumine circiter centimetrali et haud raro (ut limbus ipse) subobliquo, margine toto serrata, dentibus plerumque acutis rarius obtusiusculis, nervis basilaribus non infrequenter paulo supra insertionem petioli ortis ultra medium limbum vix productis ante marginem cum primariis sequentibus (utrinsecus 6-8) anastomosantibus subtus parum prominentibus, lamina exsiccata membranacea sæpius læte viridi, cystolithis paginæ superioris tenuibus creberrimisque, petiolo 1-3-centimetrali indumento ut in costa, illo foliorum juniorum sæpe incanulo. Stipulæ ovatæ, subcentimetales, acuminatæ, extus adpresse puberulæ, mox deciduæ. Inflorescentiæ ut plurimum ter dichotomæ, ramis primariis brevibus, ultimis quam folia nunc brevioribus s. 10-18 centim. longis nunc iis multo longioribus nempe fere 3-decimetralibus. Inflorescentiæ fem. (plerumque masculis majores) pedunculo circiter centimetrali primum puberulo at demum ut ramuli glabrato. Flores masc. glomerulati, subsessiles, perigonii segmentis extus hispidulis; feminei plus minus conferti, foliolis calyculi basi subconcretis siccitate recurvis. Achænium ellipticum, margine patule setosum, faciebus breviter hispidis, cum stipite et rostro æquilongis pariterque ciliatis 1½ millim. longum. (v. v. et s.)

β, *polystachya*, calyculo vulgo 4-phylo, foliolis glabriusculis. ♂ In provincia Maracaybo Novo-Granatensium (Plée). — *M. polystachya* Wedd. in Ann. sc. nat. l. c. (v. s.)

γ, *ambigua*, calyculi foliolis lineari-spathulatis magis minusve ciliatis. ♂? In sylvis et pratis provinciæ mexicanæ Vera Cruz, altit. 1300 metr. (Galettii, n. 317). (v. s.)

2. *M. DENSIFLORA* (Benth. l. c. p. 169), dioica, ramulis novellis tomentosiss, foliis ovatis sæpius breviter acuminatis basi rotundatis vel subcordatis acule serrulatis denticulatisve, junioribus supra planiusculis et hispido-pubentibus, adultis rugosulis et glabratiss subtus in venis præsertim pubescenti-tomentosis siccitate rigidis, inflorescentiis (saltem femineis) quam folia longioribus dichotomis, calyculo florum fem. sæpius bifoliolato, foliis spathulatis achæniisque crebre setoso-ciliolatis. ♂ In Peruvia (Mathews, n. 2040). — Wedd. Mo-

nogr. p. 487. t. 16. D. f. 1-5. — Ramuli crassi, epidermide exsiccata sub hirsutie ferruginea. Folia 10-15 centim. longa, 5-9 centim. lata, superne nonnunquam attenuata, nervis basilaribus fere ut in stirpe præcedente dispositis sed subtus paulo magis prominentibus supraque sæpe (costaque et primariis cæteris) plus minus pubescentibus, lamina exsiccata obscure viridi, cystolithis paginæ superioris creberrimis et in limbo adulto e vertice rugarum (tum pilis destitutarum) radiantibus. petiolo  $1\frac{1}{2}$ -3 centim. longo tomentoso. Stipulæ 10-15-millimetrales, ovatæ, acuminatæ, extus dense pubescentes, cito deciduæ. Inflorescentiæ fructiferæ (solum visæ) bi-tri-decimetales, pedunculo circiter centimetrali pubescente, floribus confertis, folioli calyculi dorso et margine setosis. Achæniun ad maturitatem ellipticum aut fere rotundatum, margine quam in stirpe præcedente crebrius setoso, faciebus lævibus, cum stipite brevi et rostro  $1-2\frac{1}{2}$  millim. metiens. (v. s.)

β, *Dombeyana*, foliis oblongo-ovatis valde rugosis subtus petiolisque et ramulis dense tomentosis, inflorescentiis bifurcis. δ In Peruvia (Dombey). — M. Dombeyana Wedd. in Ann. sc. nat. l. c.; Miq. l. c. *Urtica spiralis* Domb. mscr. (v. s.)

3. M. CORDIFOLIA (Liebm. in Vidensk. Selsk. Skr. 5 Række, naturv. og math. Afd. 2 Bind, p. 306), dioica, ramis crassis puberulis, foliis suborbiculatis cordatis acutis crassis dentatis bullato-rugosis supra sparsim setosis subtus imprimis ad nervos venasque pubescentibus demum glabrescentibus, inflorescentiis fem. axi dichotoma longissima, floribus pedicellatis, perigonii diphylli foliolis patulis acutis dorso achænioque ciliatis. δ In sylvis primævis prov. mexicanæ Vera Cruz (Liebmann). — Wedd. Monogr. p. 488. — Arbor 4-5-metralis. Folia 20-25 centim. longa, 18-20 centim. lata, trinervia (venis utrinsecus 5-6 subtus prominulis); exsiccata supra obscure viridia, bullato-pustulata, sparsim setosa pilisque minimis adnatis (cystolithis) eleganter stellato-lacunosa, fusco-viridia; petiolis 3-5 centim. longis teretibus striatis incano-pubescentibus demum glabris. Stipulæ  $2\frac{1}{2}$  centim. circiter longæ, convolutæ, incano-puberulæ, deciduæ. Inflorescentiæ fem. circiter 6-decimetales, bractea bifida pubescente suffultæ, floribus minutissimis pedicellatis bracteis minimis rufis suffultis, achænio ovato acuto marginato fusco. (Descript. ex Liebmann.)
4. M. LONGIPES (Liebmann, l. c.), dioica, ramis crassis teretibus puberulis mox glabrescentibus, foliis ellipticis v. late ellipticis acutis basi obtusatis margine denticulatis supra lacunoso-rugosis et immerso-glandulosis subtus ad costam nervosque incano-pubescentibus ad venulas tenuissime puberulis mox glabrescentibus, inflorescentiis fem. dichotomis longissimis, floribus sessilibus, perigonii diphylli foliolis breviter patentibus achænio margine breviter ciliato duplo brevior. δ In ditone Costa Rica prope Aguacate (Orsted). — Wedd. Monogr. p. 489. — Folia 1-3 decim. longa, 8-18 centim. lata, trinervia, venis utrinsecus 6-7 citra marginem arcuato-anastomosantibus subtus prominulis, pagina superiore pilis minutissimis adpressis (cystolithis) fusiformibus circa glandulas immersas stellato-radiatis obsita, petiolis 5-centimetralibus parum compressis supra canaliculatis puberulis. Inflorescentiæ masc. 3-10 centim. longæ, e spicis pluribus 5-7 centim. longis compositæ, floribus sessilibus densissime confertis, perigonio tenui membranaceo glabro. Inflorescentiæ fem. 6 decim. circiter longæ, axi filiformi puberulo, pedunculo  $1\frac{1}{2}$ -3 centim. longo, bractea bifida amplectente basi circumvallato. (Descript. ex Liebmann.)

β, *Colipensis*, petiolis quam in typo longioribus nonnihil compressis et in facie superiore sulcatis, floribus fem. quam in eodem majoribus pedicellatisque. δ Prope Colipa, in prov. mexicana Vera Cruz (Liebmann) — M. colipensis Liebmann l. c. M. cordifolia β colipensis Wedd. Monogr. p. 488.

5. M. INÆQUILATERA (Liebmann, l. c. p. 307), dioica?, ramis verrucosis cinereo-pubescentibus mox glabris, foliis ellipticis acutis parum obliquis basi obtusatis vel obtuse cuneatis late serratis (serraturis obtusiusculis) supra demum glabris subtus ad costam nervos venulasque subtilissime puberulis. δ In eisdem locis ac præcedens. — Wedd. Monogr. p. 489. — Folia 18-22 centim. longa,

ciliatis. Flores masc. breviter pedicellati, perigonii campanulati vel subinfundibuliformis 4-lobi lobis extus in nervo medio acuminque fere millimetrali pilosis, filamentis staminum (interdum pilosis) inferne cum basi pistilli rudimentarii (fere linearis lanatique) in columnam teretem circiter semimillimetralem coactis. Achænia oblique ovato-elliptica s. dimidiato-rotundata,  $\frac{1}{2}$ - $\frac{3}{4}$  millim. longa, margine minutissime verruculosa, glaberrima, stylo circiter quadruplo breviora. — Insignis longitudine acuminis lorum perigonii masculi, cæt. (v. s.)

β, *velutinus*, foliis ovatis acuminatis basi rotundatis v. obtusis rugosulis subtus velutinis. ♂ Cum typo (Fendler, n. 1268).

γ, *rotundatus*, foliis rotundato-ovatis non aut vix acuminatis crenato-serratis supra vix hispidis subtus breviter pubescenti-tomentosis. ♂ In provincia Bogota Novo-Granatensium, altit. 1900 metr. (Linden, n. 810). Phenax tomentosa Wedd. in Ann. sc. nat. l. c., Bl. l. c. p. 170. Filamenta staminum pilosa. (v. s.)

3. *P. rugosa*, monoicus v. dioicus, foliis ovatis acuminatis basi rotundatis cordatisve acute et frequenter grossiuscule crenato-serratis supra rugosulis et magis minus pilosulis glabratise lævibusque floribus masc. tetrandris androceo sæpius sessili. ♂ In Nova Granata, prope urbem Bogota locoque dicto Peña Vieja (Goudot, etc.), necnon in temperatis Andium quitenium prope Hambato et Llacungua, alt. 2700-3000 metr. (Humb. et Bonpl., Jameson, etc.). *P. ballotæfolius* Wedd. in Ann. sc. nat. l. c.; Monogr. p. 494. Procris rugosa Poir. Encycl. 5. p. 628. Boehmeria rugosa Pers. Synops. 2. p. 556. *B. ballotæfolia* Kunth in H. B. Nov. gen. et sp. 2. p. 35. Ramuli fere ut præcedente. Folia ut in eadem specie magnitudine admodum variabilia, cæterum basi sæpe cordata, serraturis juniorum ut plurimum majusculis obtusisque, acumine s. dente apicali haud raro lanceolato v. lineari, petiolo 1-4-centimetrali. Stipulæ ovatae, 3-10-millimetrales, acuminatae, magis minusve pubescentes. Glomeruli sphaerici, diametro 6-10-millimetrali, bracteis rotundatis obtusis parce ciliatis. Flores masc. breviter pedicellati, pedicello piloso, perigonii campanulati 4-lobi lobis convexis in acumen  $\frac{1}{2}$  millim. longum pilosulumque intus nonnunquam lanatulum productis, filamentis glabris in fundo perigonii e basi pistilli rudimentarii linearis lanatique nascentibus. Achæniæ ut in *P. uliginoso* sed margine dorsali superne pilis rectis aut hamatis numerosioribus paucioribusve ornatum fere glaberrimum. (v. s.)

β, *Mandoni*, foliis late ovatis acuminatis basi truncato-cordatis integrisque cæterum crenulato-serratis supra rugoso-hispidis subtus velutino-tomentosis. ♂ In Bolivia (Mandon). (v. s.)

γ, *serratus*, foliis ovatis acuminatis basi cordatis ab ipsa basi acute multi-serratis supra rugosulis et pubescenti-hispidis subtus dense hispidopubescentibus. ♂ In Bolivia (Mandon, n. 110-4). (v. s.)

δ, *minor*, foliis multo minoribus quam in typo potiusque ellipticis quam ovatis. ♂ ? Prope Tarma, in Peruvia. (v. s.)

4. *P. hirtus*, monoicus, foliis subcordato-ovatis acuminatis acute serratis planiusculis aut rugosis lævibus supra sparsim hispidis nervis subtus hirtis, nervis primariis inæqualibus, floribus masc. tetrandris, androceo sæpius breviter stipitato. ♂ In locis humidis umbrosisque vel subapertis Mexici (Schiede; Galeotti, n. 310; Botteri, n. 274, 275, 276 et 277), Antillarum (Swartz, Purdie, etc.), Colombiae et Peruviae. — Bl. l. c. p. 169. t. 54.; Grisebach, l. c. *P. urticæfolius* Wedd. in Ann. sc. nat. l. c.; Monogr. p. 495. *Urtica hirta* Swartz, Fl. Ind. occ. 1. p. 285. Procris urticæfolia Poir. Encycl. 5. p. 628. Boehmeria urticæfolia et *B. hirta* Pers. Synops. 2. p. 556. Ramuli graciles, obtuse angulati, sparsim pilosuli v. puberuli, epidermide ramorum veterum castanea. Folia 3-7 centim. longa,  $1\frac{1}{2}$ -4 centim. lata, acumine s. dente apicali sæpius lineari acutissimoque, serraturis grossiusculis, nervis ut in *P. uliginoso*, lamina exsiccata rigida, petiolo gracili raro plusquam bicornu- metrali sæpissimeque brevior hirta v. puberulo. Stipulæ ovatae, anguste acuminatae, nervo dorsali apiceque pilosae. Glomeruli androgyni (floribus

utriusque sexus vulgo numero fere æquali), sphaerici, majores centimetrum diametro metientes, bracteis rotundatis ovatisve ciliatis. Flores masc. breviter pedicellati, perigonii campanulati segmentis convexis acuminatis, acumine pilosulo semimillimetræli, filamentis glabris inferne cum basi pistilli rudimentarii (lineari et lanati) in stipitem brevem connatis. Achænia magis minusve verruculosa, glabra v. superne pilosula, stylo circiter quadruplo breviora. — Hæcce stirps et præcedentes duæ affines sunt; specimina vidi, nec rara, quæ ad typum proprium ægre referre potui. (v. s.)

β, *Pentlandianus*, foliis crenato-serratis nervis subtus pubescentibus siccitate membranaceis. ♀ In Bolivia (Pentland). — *P. Pentlandianus* Wedd. in Ann. sc. nat. l. c. (v. s.)

γ, *petiolaris*, foliis ovatis v. rotundato-ovatis sensim acuminatis inferne cuneatis acute serratis petiolo brevissimo aut limbum excedente. ♀? In Peruvia (Pavon). — *P. petiolaris* Wedd. l. c.; Monogr. p. 497. — Folia ima basi obtusata v. rotundata integraque, siccitate membranacea. Segmenta perigonii masc. sparsim uncinato-hispida. Achænia læviuscula, glabra. (v. s.)

δ, *minor*, foliis quam in typo minoribus crenato-serratis planis nervis subtus puberulis. ♀ In Peruvia. (v. s.)

5. *P. ASPER*, monoicus, foliis ellipticis acutis basi rotundatis acute angulato-dentatis utrinque sed præsertim supra asperato-hispidis demumque rugosis, nervis primariis paucissimis, floribus masc. tetrandris, androceo sessili. ♀ In ins. Cuba orientali, prope villam Monte Verde (Wright, n. 532). — *P. urticifolius* Griseb. in Mem. Acad. Am. sc. et art. n. ser. 8 (ann. 1860), Cat. pl. Cub. p. 60; non Wedd. Fruticulus 2-3-decimetrælis, caulibus s. ramis primariis gracillimis subsimplicibus basi procumbentibus radicantibusque puberulis hispidulisve, epidermide exsiccata obscure ferruginea. Folia  $1\frac{1}{2}$ -3 centim. longa, 1-2 centim. lata, dentibus utrinque 4-7 mucronato-acuminatis, dente apicali brevi triangulari, nervis primariis s. adscendentibus præter basiliares tantummodo 2 in utroque latere costæ, intervenio paginæ inferioris in folio adulto conspicue reticulato, lamina exsiccata rigida, petiolo quam dimidius limbus brevioris hispido. Stipulæ lanceolatæ, acuminatæ, superne parce hispidopilosæ. Glomeruli androgyni, vix multiflori, petiolo breviores, bracteis glabris. Flores masc. breviter pedicellati, perigonii lobis convexis acumine lineari semimillimetræli pilosulo subabrupte terminatis, filamentis glabris. Achænia (imatura solum visa) elliptica, puberula, stylo multoties breviora. — Habitu a congeneribus omnibus distinctissimus. (v. s.)

6. *P. LEVIGATUS* (Wedd. in Ann. sc. nat. l. c.; Monogr. p. 497. t. 16. A. f. 1-7), monoicus, foliis oblongo-ellipticis ovatisve apice attenuatis vel subacuminatis acutis basi rotundatis obtusisve et integris cæterum laxè grossiusculeque crenato-serratis (crenis submucronatis) junioribus supra etiam in costa nervisque puberulis (intervenio mox glabrato) venis paginæ inferioris inæqualiter reticulatæ sparsim et adpresse pilosis, glomerulis androgynis, floribus masc. tetrandris. ♀ In Peruvia (Dombey; Mathews, n. 1556). — *P. urticifolius* Griseb. Fl. Br. W. Ind. l. c. — Suffrutex basi præsertim ramosus, ramis simplicibus aut parce ramulosis 2-4 decim. longis flexuosis diffusis striatis puberulis v. apice pubescentibus foliosis (internodiis sesqui-bi-centimetralibus) adultis epidermide fusca v. castanea vestitis raro basi omnino glabris. Folia 3-7 cm. longa,  $1\frac{1}{2}$ -3  $\frac{1}{2}$  cm. lata, trinervia, superiora plerumque sensim acuminata, inferiora sæpe late elliptico-ovata vel etiam rotundata obtusaque, dentibus in diversis foliis nunc acutis nunc obtusis et abrupte mucronato-acuminatis, nervis primariis pluribus (8-10) subæqualibus supra parum impressis densiusculeque puberulis subtus vix prominulis sed in lamina exsiccata submembranacea conspicuis, venulis lateralibus oblique transversis, petiolo gracili 1-3-centimetræli puberulo aut (in foliis junioribus) villosulo. Stipulæ lineari-lanceolatæ, semicentimetrales, extus pubentes. Glomeruli circiter crassitudine pisi majoris vel etiam crassiores, densiflori. Flores masc. breviter pedicellati, perigonii lobis extus et superne hispidulis mucrone lineari pubescente instructis, androceo breviter stipitato, filamentis glabris, pistilli rudimento millimetræli acuto densiuscule lanato. Achænium late ovatum  $\frac{1}{4}$  mm. circiter longum, mi-

nute granulosum superne præsertim in altero margine pilosulum, quam stylus multoties brevis. (v. s. in hb. Mus. par. Kew. etc.)

β, *Haitiensis*, foliis late ovatis acuminatis basi subcordatis breviter petiolatis supra basim crenatis v. crenato-serratis (crenis non mucronatis) planis, supra sparsim breviterque pilosulis nervis subtus adpresse pilosulis intervenio venulis secundariis reticulato. 5 In insula S. Domingo (Nectoux). — P. haitiensis Wedd. in Ann. sc. nat. l. c.; Monogr. p. 496. (v. s. in hb. Mus. par.)

γ, *serratus*, fruticosus, foliis utrinque acutis v. basi obtusis dentibus acutis numerosioribus quam in typo limbo sæpius plus minus rugosulo. 3 In Peruvia et Bolivia (Mandon, n. 1110).

7. P. *LAXIFLORUS* (Wedd. Monogr. p. 499), monoicus, foliis ovatis ellipticisve obtusis mucronulatis basi late cuneatis acutisque vel obtusis grosse et inæqualiter crenatis supra pilosulis subtus subtiliter pubescentibus, glomerulis unisexualibus (femineis superioribus), floribus masc. longiuscule pedicellatis, tetrandris. 5 In Peruvia (Pavon). Suffrutescens, ramulis gracilibus puberulis internodiis 1-2-centimetralibus. Folia late patentia, 1½-4 cm. longa, 1-2 cm. lata, dentibus valde inæqualibus obtusis in dimidio inferiore limbi sæpiissime nullis, limbo toto nonnunquam integro, nervis basilaribus tenuibus ad 1 superiorem folii productis, petiolo 5-10 mm. longo subtilissime pubescente. Stipulæ lanceolatæ, 2-3-millimetrales, extus pubescentes. Glomeruli masc. in axillis inferioribus ramulorum laxiflori, femineis multo majores, perigonii lobis dorso pilis paucis hamatis instructis, pistilli rudimento conico lanato. Glomeruli fem. pauciflori, ex axillis superioribus orti, bracteis ovatis ciliolatis pilis longis hamatis puberulis. Achænia margine densiuscule pilosula, pilis albis rectis adpressisque albis hamatis, stylo quam achænium triplo circiter longiore hinc villosillo illic adpresse piloso. (v. s. in hb. Kew. et Boiss.)

8. P. *INTEGRIFOLIUS* (Wedd. in Ann. sc. nat. l. c.; Monogr. p. 499), monoicus, foliis parvis lanceolatis utrinque acutis integerrimis quintuplinerviis supra in intervenio pilis longis adpressis lineatim ornatis nervis subtus pilosis, glomerulis densifloris plerisque androgynis 3 In Peruvia (Pavon), in Nova Granata, prope Pasto (Friana). Suffrutex 3-6-decimetralis, caule basi radicante dein procumbente vel ascendente diffuse breviterque ramoso villosillo internodiis brevibus. Folia conferta, 1-2 cm. longa, 3-8 mm. lata, nervis primariis subparallelis cæteris inconspicuis, pagina superiore pilorum lineis 5-6 ornata, intervenio paginae inferioris glabro, lamina exsiccata rigida, petiolo 1-2 mm. vix longo. Stipulæ subsistentes, latiuscule ovatae, 3-4-millimetrales, acutæ, nervo dorsali piloso. Glomeruli conferti demumque confluentes, circiter magnitudine pisi, androgyni vel interdum in axillis superioribus feminei. Flores masc. brevissime pedicellati, lobis acuminatis superne longiuscule pilosis, alabastro sesqui-bi-millimetrali. Achænia elliptico-ovata, 3 mmp. longa, glaberrima, minute granulosa. (v. s. in hb. Webb. et Boiss.)

9. P. *MEXICANUS* (Wedd. Monogr. p. 500), monoicus, foliis elliptico-vel ovato-lanceolatis lanceolatisve sensim acuminatis acutiusculis excepta basi acuta integra crenulato-serratis tri-triplinerviis supra glabris et lævibus nervis subtus puberulis s. breviter pilosulis, glomerulis unisexualibus aut androgynis, floribus masc. triandris. 5 In imperii mexicani provinciis Oaxaca, alut. [200 metr. (Galeotti, n. 325 aut 345?), et Vera-Cruz, monte Orizaba (Botteri, n. 278 et 279; Fr. Müller, cæt.). — P. Mexicanus et P. Gaudichaudianus Wedd. in Ann. sc. nat. l. c. p. 193. P. Galeottianus Bl. l. c. p. 227. — Ramuli plus minus elongati, erecto-patuli, epidermide pallide castanea v. interdum cinerascens vestiti. Folia etiam in uno et eodem ramo sæpe valde inæquimagna, 2½-12 cm. longa, 1-3½ cm. lata, serraturis obtusiusculis in acumine (acuto aut obtusato) sæpius obsoletis, nervis basilaribus circiter usque ad trientem superiorem limbi excurrentibus et ante marginem cum primariis sequentibus arcuatum connexis subtus parum prominentibus, venulis reticulatis, lamina exsiccata membranacea supra obscure subtus pallide viridi et nonnunquam subcyanescente, petiolo 3-30 mm. longo gracili puberulo. Stipulæ lanceolatæ, hispidae. Glomeruli magnitudine variabiles, scilicet pisum interdum vix æmulantes aut interdum illo duplo majores, siccitate obscure ferruginei, aut raro unisexuales s. e floribus fem. tantum compositi, floribus masc.



in quovis glomerulo androgyno ut plurimum paucis, bracteis ovatis acutis obtusisve glabris vel superne breviter et inæqualiter fimbriato-ciliatis. Flores masc. breviter pedicellati, perigonii lobis triangularibus acutiusculis fere glabris. Achænia elliptica, læviuscula, superne in margine parce hispida, stylo villosa sextuplo breviora. (v. s.)

10. P. ANGUSTIFOLIUS (Wedd. in Ann. sc. nat. l. c.; Monogr. p. 501, t. 16, A. f. 8), dioicus, foliis anguste lanceolatis sensim acuminatis acutissimisque basi rotundatis v. obtusis v. acutis apice et basi exceptis minute laxeq. serrulatis trinerviis supra glabris et lævibus subtus subtiliter strigosis, floribus masc. tetrandris. 3 Prope Aguacate, in diti ne Costa-Rica (Ersted); in temperatis Novæ Granatæ, prope Mariquita, Ibagué, etc. (Humb. et Bonpl. Goudot), in Peruvia subandina (Poeppig, n. 1315) et in Brasiliæ prov. Minas Geraes (sep. specim. hb. imp. bras.). *Boehmeria angustifolia* Kunth in H. et B. Nov. gen. et sp. 2. p. 34. Ramuli graciles, teretiusculi, epidermide obscure castanea vestitis pubesque brevi adpressa alba magis minusve conspersis, internodiis ut plurimum uni-sesqui-centimetralibus. Folia 5-15 cm. longa, 6-25 mm. lata, erecto-patentia, longe angusteq. acuminato-attenuata, nervis basilariibus usque ad summum limbum productis margini subparallelis eique quam costæ multo propioribus, lamina exsiccata rigida fusciscente interdum rugosula nervis supra impressis subtus prominulis, petiolo gracili 5-15 millimetrali strigosulo. Stipulæ lanceolatæ, 2-4 mm. longæ, in nervo dorsali adpresse pilosulæ. Glomeruli fem. (solum visi) in axillis omnibus ramuli nascentes, grani Zeæ circiter magnitudine, siccitate obscure ferruginei, e spiculis innumera (spiculas graminis quodammodo referentibus) densissime aggregatis compositi, bracteis ovato-lanceolatis acutis apice ciliolatis arcte involucriatis. Achænia elliptica, vix semimillimetralia, margine oculo armato sparsim grappulo-papillosa summoque dorso pilis paucis minutis rectis hamatisque munita, stylo longissimo sub lente subtiliter villosulo atque hic illic adpresse pilifero. (v. s.)

Tribus IV. *PARIETARIÆ* Wedd monogr. p. 502. — *Parietariae*-  
rum gen. *Gaudich.*

Herbæ annuæ aut perennes, rarius suffrutices v. frutices, rarissime arbusculæ, inermes. Folia alterna, homomorpha, æquilateralia aut rarissime (nempe solum in *Helxine*) inæqualiteralia, integerrima (specie unica gen. *Hemistylis* excepta), cystolithis punctiformibus v. rarissime lineari-oblongis inspersa. Stipulæ omnino deficientes aut axillares aut petiolares. Inflorescentiæ (saltem femineæ) bracteis herbaceis duabus v. tribus aut pluribus s. involucri munitæ. Flores declini aut polygami. *Masc.* : perigonio 4-partito, segmentis acutis muticis in præfloratione valvatis, pistilli rudimentò vario basi sæpe lanato. *Fem.* : perigonio libero tubuloso-ventricosò, ore magis minusve contracto 4-fido aut 2-v. 4-dentato, stigmate spatulato v. subcapitato longeq. penicillato aut filiformi, staminum rudimentis nullis. *Hermaphr.* : perigonio et staminibus ut in flor. masc., ovario et stigmate ut in flor. fem. Fructus siccus. Embryo cotyledonibus elliptico-v. quadrato rotundatis radiculæ æquilongis v. illa longioribus, albumine vario sed nunquam deficiente.

XXXIV. *PARIETARIA* Tournef. Instit. p. 289; Linn. Gen. pl. n. 1152 (excl. sp.), Bt. Mus. bot. lugd. bat. 2 p. 245; Wedd. Monogr. p. 503. — *Parietaria*, Thaumuria et Freirea Gaudich. Bot. Voy. Uran. p. 501 et 502; Lindl. Veget. Kingd. ed. 2. p. 262. — *Parietariæ* sect. a, b, et c, Endlich. Gen. pl. p. 284.

- Flores polygami (hermaphroditi et feminei), dense v. laxiuscule cymosi, cymis s. glomerulis axillaribus sæpius geminis 3-multifloris bi-

paris v. ultra basim scalari-uniparis, bracteis herbaceis 1-3 singulos flores fulcientibus liberisque aut inter se aut cum illis florum vicinorum basi plus minus coalitis. *Hermaphr.* : perigonium 4-partitum, intus glabrum, tubo breviter campanulato post anthesin sæpe accreto; laciniis ovatis breviter acuminatis uncinato-hispidis pubescentibusve. Stamina 4. Ovarium et stigma ut in floribus femineis. Stylus brevis aut nullus. *Fem.* : perigonium tubuloso-ventricosum, intus glabrum lanatumve, limbo 4-fido, laciniis conniventibus extus hamato- v. rarius glanduloso-pilosis. Ovarium rectum, ovoideum v. oblongum. Ovulum vulgo obliquum, funiculo brevi insidens. Stigma spatulatum, valde recurvum et propter papillas elongatas latus alterum vestientes aspergilliforme, stylo elongato-filiformi vel brevissimo suffultum, mox deciduum. Achænium rectum, ovoideum, nitidum, perigonio evoluta (in floribus hermaphroditis sæpe elongato-cylindraceo) marcescente nervoso v. striato rufo v. fuscescente inclusum. Embryo cotyledonibus oblongo-ovatis radicula subæquilongis. — Plantæ herbaceæ, annuæ v. perennes, rarius suffrutescentes, utriusque orbis incolæ, plerumque pubescentes pilisque hamatis frequentes obsitæ. Folia alterna, integerrima, trinervia aut triplinervia, cystolithis punctiformibus inspersa. Stipulæ prorsus nullæ. Cymæ nunc symmetrice dichotomæ, nunc abortu asymmetricæ s. scalares; bracteis inflorescentiæ cum ramulis suis (s. ex axillis suis natis) a basi ad apicem usque coadunatis ideoque in ipsis angulis s. sinubus dichotomiæ manifestis. — Genus admodum naturale e typis paucis genuinis sed pro habitatione et cœlo valde mutabilibus constans.

*A. — Stirpes pleræque perennes. Folia triplinervia.*

1. *P. OFFICINALIS* (Linn. Spec. pl. p. 1492), perennis, caespitosa, caulibus herbaceis aut lignescensibus sæpius pubescentibus, foliis polymorphis acutis acuminatisve basi nunc longe angustatis acutisque nunc rotundatis obtusive triplinerviis, inflorescentia glomerata dichotoma pauci-multi-flora, floribus aliis femineis aliis (numerosioribus) hermaphroditis, bracteis ovatis oblongisve, perigonio fl. hermaphr. post anthesin sæpissime elongato.

*α, diffusa*, caulibus prostratis v. diffusis rarius erectis mox ramosis, foliis lanceolatis v. ovatis ellipticisve, glomerulis densiusculis mox ramosis, foliis basi plus minus coalitis.  $\frac{1}{2}$  Ad muros et vias et in ruderalis, in Europa tota, frigidioribus excepta, in insulis Canariis, Africa boreali, Syria, Anatolia, Mesopotamia et Persia. — *P. officinalis* Linn. herb. 1 Smith Engl. bot. t. 879, Fl. brit. 1. p. 189; Curtis Fl. Lond. 3. t. 521; Bertol., Fl. ital. 2. p. 213; Coss. et G. de St.-P. Fl. env. par. ed. 1. p. 475, ed 2. p. 582; Wedd. Monogr. p. 506. t. 17. f. 1-14. *P. judaica* Vill. Fl. Dauph. 2. p. 346; G. F. Hoffm. Deutschl. Fl. 2. p. 276; Lam. Illustr. t. 853. f. 2; DC. Fl. fr. 3. p. 824; Schkuhr Handb. t. 346. f. inf.; Webb Phytogr. canar. 3. p. 502; non Linn. *P. diffusa* M. et K. Deutschl. Fl. 1. p. 827; Koch Synops. ed. 1. p. 636; Guss. Fl. sic. synops. p. 640; Reichb. Ic. fl. germ. 12. t. 651, f. 1318; Gren. et Godr. Fl. fr. 3. p. 109; Bl. Mus. bot. lugd. bat. 2. p. 247. *P. assurgens* Poir. Encycl. 5. p. 15; Pers. Synops. 2. p. 555. *P. maderensis* Reichb. in Bot. Zeit. ann. 1830. p. 131? — Caules magis minusve numerosi, erizomate lignoso ramoso nodosoque (siccitate fusco) nascentes, sæpissime adscendentes basique radicantes, crassitie varia, subsimplices vel simpliciter rariusve composite ramosi, superne patule pubescentes vel puberuli, inferne glabrati, exsiccati olivacei v. fusci v. fuscorubentes. Folia quoad formam et magnitudinem quam maxime variabilia, majora 8-10 cm. longa, minora vix centimetralia, acumine nullo vel elongato et angusto nunc acuto nunc obtuso, nervis basilaribus in limbo deorsum longe attenuato procul ab insertione petioli (nempe 5-10 mm.), in

limbo autem basi rotundato proxime (id est vix 1-2 mm. supra eandem insertionem) orientibus, limbo pubescenti-piloso margineque ciliato, pilis paucis ut plurimum rectis hamatisque intermixtis, lamina exsiccata membranacea læte viridi vel fuscescente subtus ut in vivo pallidiore, petiolo (foliorum caulinorum præsertim) limbum dimidium sæpe æquante et haud raro excedente ramealium contra plerumque brevi pubescente. Cymæ in axillis foliorum caulis ramorumque simul nascentes, pisiformes aut interdum nucem avellanam minorem magnitudine æmulantes, ramulis abbreviatis, bracteis ut plurimum ovatis obtusisque pilos uncinatos crebros (paucis rectis intermixtos) gerentibus. Flores hermaphr. in apicibus ramorum cymæ obvii, singuli bracteis tribus basi connatis involucriati, perigonii lobis acuminatis extusque pilis uncinatis rectis crebre ornatis demum inflexis, tubo brevi campanulato et fere glabro vel sparsim pilosulo post anthesin plus minus elongato cylindrico pallideque rubente intus lanulato. Flores fem. vulgo in angulis dichotomiæ sessiles, bractea unica stipati, perigonio ovato demumque pyramidato s. superne 4-angulo, lobis anguste triangularibus acutissimis arcte conniventibus, stylo filiformi exserto stigma aspergilliforme læteque purpureum fulciente, stigmatum florum hermaphroditorum plerumque minore pallidiore styloque brevi suffulto. Achenium ovatum, compressum, obtusiusculum, non aut vix apiculatum, in floribus fem. fere semper atrum et fertile pericarpio crustaceo, in floribus hermaphr. contra haud raro olivaceum et effictum pericarpio submembranaceo. — Ad hanc formam (cum *P. judaica* L. sæpe confusam) referenda est etiam *P. punctata* Willd. (Sp. pl. 4. p. 953) quæ ab illa perigonis fl. hermaphr. post anthesin minus elongatis vix differt. Magis distinctæ, præcipue quoad habitum, sunt varietates sequentes (v. v. et s.)

β, *microphylla*, caule prostrato v. adscendente a basi diffuse ramoso sæpe canescenti-hirtulo, foliis multo minoribus quam in typo ovatis lanceolatisve, inflorescentiis parvis, perigonis fl. hermaphr. post anthesin subcampanulatis. ¼ Passim, in locis aridioribus quam typus. *P. canescens* Bl. l. c. p. 249. *P. judaica* hb. Willkomm lt. hisp. secund. n. 39. (v. s.)

γ, *populifolia*, caule sat elato erecto v. suberecto parce pubente, foliis late subtriangulari-ovatis basi late subtruncatis v. leviter cordatis. ¼ In ins. Melita (Nyman) et verisimiliter alibi. *P. populifolia* Nym. in Linnaea, 18. p. 661. Folia (teste Nyman) illis *Populi balsamiferæ* v. *nigræ* majora.

δ, *erecta*, caulibus simplicibus v. subsimplicibus erectis, foliis ovato- v. elliptico- v. oblongo-lanceolatis, inflorescentiis plerumque majoribus laxioribusque quam in typo bracteisque basi minus coalitis, floribus hermaphr. post anthesin vulgo minus elongatis s. campanulatis. ¼ In iisdem locis ac typus sed rarior, præsertim in Europa australi. *P. erecta* Mart. et K. l. c. p. 827; Koch l. c. p. 636; Schkuhr l. c. t. 346; Gren. et Godr. l. c. p. officinalis DC. l. c.; Reichb. l. c. f. 1317; Bl. l. c. p. 246. *P. officinalis* β *longifolia* Coss. et Germ. l. c. ed. 1. p. 475. Cymæ quam in var. magis ramosæ laxioresque. (v. v. cult. et s.)

2. *P. JUDAICA* (Linn. Sp. pl. p. 1492), perennis, cæspitosa, caulibus suffruticosis perennantibusque puberulis rigidis siccitate pallide ferrugineis cinereisve, foliis ovatis rariusve lanceolatis acutis v. breviter lanceolatis basi rotundatis v. subcuneatis triplinerviis, inflorescentia dichotoma pauciflora, flore inferiore cymulæ femineo cæteris hermaphroditis, bracteis late ovatis basi plus minus connatis, perigonio fl. hermaphr. post anthesin elongato oblongo. ¼-½ In Anatolia (Boissier et Heldreich, Botta, Tchihatcheff), Syria monte Libano (Labillardière), Kurdistania (Kotschy, pl. Mesop. Kurd. n. 329), Armenia (Hohenacker), Georgia caucasica (Un. it. ann. 1834), Persia (Olivier et Bruguère; Kotschy, pl. Pers. n. 257 et 510), Belutschistania, alt. 1520 metr. (Stocks) et regione temperata Thibetiae occid. alt. 2430-3040 metr. (J. D. Hooker). — Willd. Spec. pl. 9. p. 954; Wedd. in Bullet. Soc. bot. fr. 2. p. 663; Monogr. p. 508. t. 17. f. 15; Bl. l. c. p. 250, non auct. *P. multicaulis* Boiss.

et Heldr. in sched. pl. Anatol. ann. 1846; Boiss. Diagn. pl. or. nov. fasc. 12. p. 106: P. nitens et P. cryptorum C. Koch in Linnæa, 22. p. 606 et 607; Bl. l. c.: P. tibethana Bl. l. c. — Caules numerosi, e rhizomate lignoso (siccitate fusco v. ferrugineo) fasciculatim prodeuntes, 15-25 centim. alti. crassitie circiter pennæ corvinæ vel paulo crassiores, erecti vel subdiffusi, basi non radican- tes, novelli simplicissimi, annotini simpliciter breviterque ramosi, inferne sæpe nudi, siccitate plerumque fusco- v. cinereo-ferruginei. Folia majora 10-20 millim. longa, 8-15 millim. lata, nervis basilaribus ut plurimum paulo supra insertionem petioli nascentibus, limbo supra sparsim hispido-puberulo demumque glabrato in margine nervisque subtus pilis brevibus hispido, ex- siccatu ut plurimum rigidulo viridi aut sæpissime ferrugineo-fuscescente pa- gina superiore cystolithis crebris verruculosa, petiolo brevi sive partem quartam v. tertiam limbi longitudine raro excedente puberulo. Cymæ axillares, in caulibus hornotinis aut ramis caulium annotinorum diversis temporibus (nec unquam simul hic et illic) nascentes, quam petioli breviores, ramulis brevissimis, bracteis ovatis ellipticisve sæpius obtusis pilis brevibus arcuatis vel paucis hamatis hispidis. Flores hermaphr. quam feminei multo numero- siores, perigonio post anthesin elongato-cylindrico 2 millim. circiter longo rubente tubo sparsim pilosulo lobis abrupte inflexis conniventibusque et pilis aliis lanatis aliisque hamatis munitis. Flores fem. ut in præcedente, sed con- stanter paucissimi. (v. s.)

β, *velutina*, caulibus dense pubescentibus, foliis subtus velutino-pubescentibus, bracteis basi discretis. ♀ In monte altissimo Kuh-Delu Persiæ australis. — P. velutina Bl. l. c. p. 251.

3. P. CRETICA (Linn. Spec. pl. p. 1492), perennis, caulibus diffusis pubescentibus, foliis parvis ovatis v. rhomboideo- aut rotundato-ellipticis triplinerviis, cymulis trifloris, bracteis in involucrium quinquepartitum demumque induratum coalitis, exterioribus allformibus, floribus lateralibus hermaphroditis intermedio femineo, perigonio hermaphroditorum post anthesin elongato. ♂-♀ In Peloponneso, insulis Archipelagi et Creta; necnon in ins. siculis Favignana, Pantellaria, Lampedusa, cæt. — Sibth. Fl. græc. 2. p. 44. t. 154; Guss. l. c. p. 642; Bl. l. c. p. 250; Wedd. Monogr. p. 511. t. 17. f. 16 et 17. P. bracteata Moench, sec. Steud. P. affinis Hornem. Hort. Hafn. (1850), ex Bl. P. cretica minor capsulis seminum alatis Tournef. Instijt. p. 38. Thau- maria cretica Gaudich. Bot. Voy. Uran. p. 502; Bonite. t. 98, B. — Caules numerosi, 1-3-decimetrales, simplices vel parce simpliciterque ramosi, gra- ciles, undique diffusi, inferne demum glabrati et lignescens. Folia 5-10 mil- lim. longa, 3-8 millim. lata, obtusa, basi sæpius cuneata, nervis basilaribus e costa paulo supra petioli insertionem nascentibus, limbo utrinque sparsim pilosulo ciliato, exsiccatu membranaceo viridi vel fuscescente cystolithis parce punctulato, petiolo gracili quam limbus brevior. Cymæ in singulis axillis geminæ, petiolis breviores aut paulo longiores, demum patulæ. Involucrium post anthesin valde auctum rubens aut violaceum, tubo campanulato florem femineum amplectente, bractearum valde inæqualium intermedia lineari, cu- jusque paris lateralium (flores hermaphr. involucriantium) altera exteriori oblonga v. spatulata 2-4 millim. longa, altera triente vel dimidio brevior et angustior, omnibus rigidis pilosque breves arcuatos et hamatos gerentibus. Flores hermaphr. pallide rubentes, pilis ut in involucrio. Flores fem. illos P. officinalis quoad formam referentes, sed perigonio intus longe et densissime piloso extusque pilis clavatis obsito, lobis apice breviter penicillatis. Achenium elliptico-ovatum, sesquimillimetrale, acutiusculum, fuscum, perigonio tubo- que involucri indurato circumvallatum. (v. v. cult. et s.)

β. — *Stirpes annuæ. Folia trinervia s. nervis basilaribus ex ipsa insertionem petioli nascentibus, rarissime triplinervia.*

4. P. MAURITANICA (Durieu in Duchartre Rev. bot. 2. p. 427), annua, caule erecto v. diffuso glabriusculo, foliis late ovatis v. triangulari-ovatis trinerviis, cymis laxis pauci-multi-floris, bracteis ovatis v. oblongo-ovatis liberis post anthesin subimmutatis, floribus infimis (paucissimis) femineis, cæteris herma- phroditis perigonio post anthesin elongato;

α, *erecta* (Wedd. mon. p. 513. t. 17. f. 18). caule elato ramisque erectis, foliis amplis basi obtusis rotundatis v. subcordatis. ☉ In fissuris rupium, locis humidis umbrosisque convalliis et ad ora speluncarum, in Lusitania prope Cintram et Olyssiponem (Ant. de Jussieu; Welwitsch, it. lusit. n. 158); in Hispania prov. Andalusia, Murcia et Valencia (Bourgeau, n. 449. a. 1485 et 1669); ins. Majorca (Cambessedes); Mauritania, prope Algeriam, Oran, Tlemcen, cæt. (Durieu, Munby, Cosson; Balansa, n. 174 et 348, cæt.); Tunisia, prope Gabès (Kralik) et in monte Clarence ins. Fernando Po, alt. 2430 metr. (Gustav Mann). P. Ocymifolia Bocc. Ic. et descr. pl. rar. Sic. p. 47. f. a (?). P. ocymifolia Poir. Encycl. 5. p. 15 (?). P. circeifolia Pourret mscr. in hb. Mus. par. P. Welwitschii Bl. l. c. p. 251. P. latifolia trimestris Grisley Virid. lusit. n. 67. Freirea ocymifolia Gaudich. Bot. Voy. Uran. p. 502 (?). — Caulis simpliciter ramosus (ramis gracilibus), superne sæpe puberulus, cæterum glaberrimus, striatus, lævis, succulentus, raro inferne lignescens. Folia majora 5-8 centim. longa, 3-6 centim. lata, sæpius subabrupte acuminata (acumine triangulari obtuso), rarius sensim attenuata, basi ut plurimum late cuneata, utrinque sparsim pilosula, margine ciliata, raro omnino glabrata, lamina exsiccata tenuiter membranacea late viridi cystolithis minutissimis punctulata, petiolo gracili quam limbus longiore aut breviori pubescente vel glabrato. Cymæ laxiusculæ, basi dichotomæ s. biparæ dein scalari-uniparæ, ramis divaricatis pubescentibus, bracteis primum flores vix superantibus sed quam illi demum brevioribus ciliatis. Flores hermaphr. albi aut rubentes post anthesin 3-millimetrales, perigonio lobis angustis erectis superne ciliatis, in nervo dorsali pilis paucis hamatis munitis, tubo extus hispidulo intus glabrato. Flores fem. perigonio ovato post anthesin parum aucto, segmentis lanceolatis scariosis nervo dorsali et margine fuscis, extus pilis clavatis dense obsitis, stylo elongato. Achænium ovatum, bimillimetrale, rectiuscule et subconspicue apiculatum, fuscum, nitidum. — Stylus floris fem. elongatus est ut in plantis divisionis præcedentis; e contrario in speciebus sequentibus e quibus conficitur genus Freirea Gaudich. stylus floris ♀ brevis est. (v. v. et s.)

β, *diffusa* (Wedd. mon. l. c.), caule quam in typo breviori diffuse ramoso, foliis parvis basi obtusis acutisve. ☉ In iisdem locis ac forma typica. — P. mauritanica γ minor Wilk. et Lange, Prodr. fl. hisp. p. 252. — Folia 1-2-centimetralia. (v. v. et s.)

5. P. DEBILIS (G. Forst. Fl. ins. austr. prodr. n. 387), annua, caule erecto v. diffuso, foliis ovatis v. rhomboideo-ovatis subrotundatisve trinerviis, cymis paucifloris bracteis vulgo linearibus lanceolatisve post anthesin vix acutis, floribus paucis hermaphroditis perigonio post anthesin sæpe immutatis, cæteris femineis perigonio post anthesin plus minus evoluto lobis acuminatis.

α, *micrantha*, caule erecto v. adscendente rarius prostrato parce ramoso, foliis sæpius ampliusculis late ovatis cordatisve obtusis v. obtuse acuminatis tenuiter membranaceis sparsim hispidulis late virentibus petiolo elongato, bracteis linearibus flores vix aut non superantibus, perigonio fl. fem. post anthesin ovato lobis obtusiuscule acuminatis. ☉ Per totum fere orbem extra-europæum (partibus frigidioribus exceptis) diffusa: in Sibiria et Dahuria, Mongolia, Himalaya a Sikkim ad Kistwar, altit. 1800-3900 metr., Belutschistan, Arabia, Abyssinia (Schimper, n. 753), ins. Gorgoniceis, Canariis, Madeira, Fernando Po, Cotonifloris, Oceani australis, in Nova Hollandia, Tasmania, Nova Zelandia, Buëbs-Ayrès, Monte Video, Brasilia australi, Chilè, Juan Fernandez, Peruvia, Nova Granata, ins. Gallapagos, Mexico, cæt. P. debilis Hook. fil. Fl. Nov. Zel. p. 226; Fl. Tasm. l. p. 344; Wedd. Monogr. p. 517. t. 17. f. 21-25. P. micrantha Ledeb. Fl. alt. 4. p. 303; Icon. t. 22. P. appendiculata Webb Phyt. canar. 3. p. 265. P. debilis, P. micrantha, P. carnosula et P. australis Bl. l. c. p. 254 et seq. Urtica debilis Endlich. Prodr. fl. Norf. p. 37. Freirea australis N. ab E. Pl. Preiss. l. p. 638. F. erecta Philippi Florul. atacam. p. 49. — Caulis simplex aut simpliciter ramosus, 1-5 decim. altus, ramis sæpius diffusis gracilibus puberulis pilosiusculisve. Folia 2-6 centim. longa, 1½-5 centim.

lata, obtusa v. obtuse acuminata, raro omnino glabrata, lamina exsiccata membranacea cystolithisque crebriusculè punctulata, petiolo longitudine quam maxime variabili. Cymulæ 3-7-floræ, bracteis liberis longiuscule ciliatis. Flores hermaphr. in angulo dichotomiæ præsertim obviæ, perigonio (deflorato campanulato) pilis parvis uncinatis rectisque hispido. Flores fem. perigonio extus pilis brevibus uncinatis rectisque (paucis numerosisve) atque nonnullis glanduloso-capitatis munito intus glabro, lobis obtuse v. acutiuscule acuminatis et apice ciliatis; perigonio fructifero 1½-2 millim. longo submembranaceo fusco-rubente lobis nonnunquam apice obscure virentibus vulgoque plus minus patentibus nec (ut in *P. lusitanica*) exacte conniventibus nervo dorsali nonnihil prominente lateralibus plerumque obsoletis. Achænium ¼-1¼ millim. longum, oblique breviterque aut rarius rectè apiculatum, fuscum aut saturate olivaceum. — Stirps quoad staturam et amplitudinem foliorum necnon perigonii florum fem. et hermaphr. incrementum admodum variabilis. Formæ sequentes præsertim distinguendæ sunt. — Subvar. *micrantha*, perigonii florum fem. et hermaphr. post anthesin immutatis. ☉ In regionibus frigidioribus præcipue obvia. *P. micrantha* Ledeb. ad hanc formam forte referenda potius quam ad typum. (v. v. cult. et s.) — Subvar. *macrantha*, perigonii omnibus florum femineorum hermaphroditorumque post anthesin ampliatis. ☉ In regionibus calidioribus Novæ Granatæ (Goudot) et verisimiliter in aliis locis. Caules in specimine viso late procumbentes. Folia late ovata, subcordata, late breviterque acuminata, petiolo centimetro. Flores hermaphr. post anthesin late campanulati circiter bimillimetrales. (v. s.)

β, *gracilis*, caulibus s. ramis primariis vulgo numerosis procumbentibusque, foliis parvis ovatis v. rhombeo-ovatis subrotundatisve sparsim pilosis pubescentibusve obscure aut pallide viridibus ciliatis, cæt. ut in typo. ☉ Passim, cum typo. — *P. gracilis* Lowe, Primit. fl. mader. in Trans. Camb. Phil. Soc. p. 16. *P. maderensis* Rchb. in Bot. Zeit. ann. 1830. p. 131. ? quæ forsân *P. officinalis*. *P. appendiculata* var. *Vulcani* Webb l. c. *P. humifusa* Bl. l. c. (forma tenuior). *Urtica parietariæfolia* hb. Berter. U. freireæformis Steud. in Bot. Zeit. 33. p. 257. *Freirea fernandeziana* Steud. Nomencl. bot. p. 734. *F. humifusa* Gay Fl. chil. 5. p. 366. — Caules plus minus elongati, prostrati, ramis ascendentibus. Folia vulgo vix centimetralia, sed interdum majora, nervis rarissime omnino inconspicuis, limbo nonnunquam (sicut caules filiformes) subhyalino, petiolo brevi aut elongato. Cymæ axillarum inferiorum haud raro ad florem unicum redactæ. — Occurrunt specimina pusillima foliis vix semicentimetralibus cymisque fere omnibus depauperatis; specimina autem staturæ mediæ *P. lusitanicam* habitu omnino referunt.

γ, *lanuginosa*, ramis foliis bracteisque lanatulo-hirtis, cæt. ut in præced. ☉ In ins. Canariis. *P. appendiculata* var. *lanuginosa* Webb l. c.

δ, *ceratosantha*, perigonio fem. in dorso loborum verrucis cylindricis inæqualibus onusta, achænio quam in var. diffusa longius apiculato, charact. cæt. similibus. ☉ In Peruvia.

ε, *squalida* (Hook. f. Fl. tasm. 1. p. 344), caule ramisque erectis aut adscendentibus ramosis, foliis ovatis acuminatis subacuminatisve, cymis triplici-floris, bracteis oblongis v. ellipticis flores non aut vix superantibus, perigonii florum fem. et hermaphr. post anthesin auctis oblongis lobis acutiusculis. ☉ In ins. Tasmania et Australia (Hook. fil., cæt.). *P. squalida* Hook. fil. in Lond. Journ. of bot. 6. p. 285; Bl. l. c. p. 256.

6. *P. LUSITANICA*, annua, caule ramisque procumbentibus puberulis glabratissive, foliis parvis ovatis v. rhombeo- aut rotundato-ovatis trinerviis, cymis paucimultifloris, bracteis oblongis lanceolatisve liberis æqualibus post anthesin subimmutatis, floribus paucis hermaphroditis sterilibus perigonio post anthesin immutato, cæteris femineis perigonio fructifero vix aucto sed valde indurato lobis ovatis acutis exacte conniventibus. ☉ In Gallia meridionali prope Telo-piam, Perpiniacum, etc., Lusitania, Hispania (Bourgeau), Corsica, Sicilia,

Peninsula italica, Peloponneso, Rossia australi ?, ins. Archipelagi, Creta, prov. Constantina Mauritaniz, Syria et Asia minore (Balansa, pl. or. n. 344). — DC. Fl. fr. 5. p. 356; Ledeb. Fl. ross. 3. p. 639; Gren. et Godr. Fl. fr. 3. p. 110; Bl. l. c. p. 252; Wedd. Monogr. p. 517. t. 17. f. 19 et 20. — Caulis s. rami primarii pauci v. numerosi, plerumque filiformes, 1-2-decimetales fere simplices, siccitate flavescenti-virides vel fuscæscentes. Folia 5-10 millim. longa, 4-8 millim. lata, superne sensim attenuata rariusve subacuminata, acutiuscula aut obtusa, basi sæpius obtusa vel rotundata, rarius acuta, utrinque sparsim pilosula demumque ut caulis ipse glabrata, limbo exsiccato membranaceo læte vel obscure viridi cystolithis crebris minutissimis punctulato, petiolo dimidii limbi circiter longitudinis vel paulo longiore pubescente aut glabriusculo. Cymulæ 3-7-floræ, bracteis pubescentibus ciliatisque, pilis rectis aut quibusdam hamatis intermixtis. Flores hermaphr. dilute virentes, perigonio pilis paucis uncinatis brevibus hispido. Flores fem. perigonio ovato extus æqualiter hispido intus glabro, fructifero  $1\frac{1}{2}$ - $1\frac{1}{4}$  millim. longo incrassato saturate castaneo capsulamque quadrialvalem quodammodo referente, valvis s. lobis sensim attenuatis apice conniventibus glabriusculis nervo triplici prominente percursis. Achænium elliptico-ovatum, vix millimetrale, recte apiculatum, dilute fuscum. — Affinis præcedenti, sed facile distinguenda characteribus perigonii fructiferi. Bractææ semper (teste cl. Gussone) perigonio adpressæ. (v. s.)

β, *Chersonensis*, foliis amplis late ovatis subacuminatis obtusis longe petiolatis læte viridibus exsiccatis hyalinis, cymis glomeruliformibus, bracteis linearibus perigonio superantibus. ☉ In locis umbrosis saxosis, prope Odessa. P. lusitanica β chersonensis Lange et Szovics. Caulis suberectus. Folia 3-4-centimetralia basi obtusa v. acuta. (v. s. in hb. Monac.)

γ, *filiformis* (Wedd. mon. p. 518), caule ramisque quam in typo elongatioribus, foliis paulo angustioribus, bracteis lineari-lanceolatis subæqualibus demum squaroso-patulis perigoniumque fructiferum superantibus. ☉ In saxosis aridis vulcanicis ins. Ustica (Gussone), Capri (Tenore) et in Græcia (Heldreich). P. filiformis Ten. Syll. p. 77; Guss. l. c. p. 642.

δ, *decipiens* (Wedd. mon. p. 518), caule ramisque diffusis v. erectiusculis elongatis, foliis majusculis lanceolatis longe petiolatis, cymis paucifloris, bracteis lineari-lanceolatis flores longe superantibus. ☉ In Georgia caucasica (Hohenacker, Un. it. ann. 1838). — Specimina visa aspectum P. pensylvanicæ omnino habent; in hac autem folia triplinervia sunt nec trinervia et perigonium fructiferum valde diversum.

7. P. PENSYLVANICA (Mühlenb. Cat. pl. Am.), annua, caule erecto elato subsimplici v. ramoso basi interdum lignescente, foliis lanceolatis v. rarius ovato-lanceolatis majusculis triplinerviis sparsim puberulis diluteque viridibus, cymis multifloris, bracteis linearibus flores superantibus post anthesin subimmutatis, floribus paucis hermaphroditis perigonio post anthesin haud mutato, cæteris femineis perigonio fructifero elongato cylindraceo lobis breviter acutecque acuminatis. ☉ In America septentrionali temperata frequens, Texas, Mexico, cæt. — Willd. Spec. pl. 4. p. 955; Asa Gray, Man. bot. U. St. p. 400; Bl. l. c. p. 254. P. debilis e pensylvanica Wedd. Monogr. p. 516. t. 17. f. 28-31. — Sicut species præcedentes magnitudine omnium partium sat variabilis. Caulis 2-5 decim. altus, subpubescens, fere semper erectus et basi interdum sat robustus. Folia  $2\frac{1}{2}$ -7 centim. longa, 1-2 centim. lata, vulgo sensim acuminata, acumine acuto v. obtuso, basi attenuata v. plus minus abrupte cuneata, triplinervia v. subpenninervia, nervis basilaribus supra petioli insertionem petioli plus minus distanter ortis; limbo imprimis in margine pubescente exsiccato tenuiter membranaceo, petiolo gracili  $\frac{3}{4}$ -2 $\frac{1}{2}$  centim. longo ut limbus magis minusve pubescenti-piloso. Cymæ quam in speciebus duabus præcedentibus sæpius majores, basi dichotomæ dein scalari-uniparæ, subaxillares, bracteis 3-4 millim. longis obtusis patulis puberulis ciliatisque. Flores hermaphr. vix nisi in apice ramulorum cymæ nascentes parvi, perigonii segmentis pilis brevibus rectis uncinatisque hispulis. Flores fem. numerosi, singuli bracteis duabus elongatis stipati, perigonio ellipsoideo lobis hispidis post anthesin oblongo 2 millim. circiter longo fusco et plus minus indurato,

lobis crasse breviterque mucronato-acuminatis et medio dorso nervo sat prominente percursis. Achænium ellipticum 1-1½ millim. longum fusco-olivaceum oblique breviterque apiculatum. — Stirps a me olim cum *P. debilis* conjuncta, sed, quantum hodie opinor, foliis triplinerviis necnon cymis multifloris et perigonis fem. post anthesin magis elongatis satis distincta. (v. v. cult. et z.)

β, *Floridana*, caulibus s. ramis primariis diffusis v. suberectis sæpe pubescentibus, foliis minoribus quam in typo elliptico-lanceolatis ovatisve subabrupte lateque acuminatis (acumine obtuso) tri-triplinerviis magis minusve pubescentibus ciliatisque sæpius obscure viridibus, petiolo limbo breviori, cymis pauci-multifloris bracteis flores parum superantibus. ☉ In imperio mexicano et Florida, Ludoviciana, Texas, etc. — *P. floridana* Nutt. gen. 2 p. 208. *P. Orizabæ* et *P. linguæfolia* Liebm. in Vidensk. Selsk. 5 R. naturv. og math. Afd. 2. p. 313. *P. debilis* & *floridana* Wedd. l. c. — Forma inter *P. pennsylvanicam* prototypam et *P. debilem* fere intermedia. (v. s.)

γ, *microphylla*, caule humili filiformi ramoso erectiusculo, foliis parvis ovatis ellipticisve non aut vix acuminatis obtusissimis opacis vage triplinerviis aut sæpius nervis basilariibus omnino inconspicuis pubescentibus obscure viridibus, petiolo quam limbus breviori, cymis paucifloris congestis bracteis flores parum superantibus. ☉ Inter Laredo et Bejar in imperio mexicano (Berlandier, n. 1536). — Caules in specimenibus visis 5 centim. alti, ramique pubescentes. Folia superne vix aut non producta, majora haud centimetralia. — Forma pro foliis etiam sub lente sæpe subnerviis typo bene distincta. (v. s. in hb. Mus. par.)

8. *P. ALSINIFOLIA* (Delile, Bot. Egypte, p. 137. t. 50. f. 2), annua, caulibus ramisque diffusis pubescentibus, foliis parvis late ovatis trinerviis, cymis trifloris, bractea inferiore lanceolata florem hermaphroditum sterilem post anthesin nonnihil elongatum suffulciente, cæteris (utrinque ternatis) floribus femineis involucrentibus et denique late cordato-ovatis membranaceisque, perigonio fructifero parum aucto segmentis acutis arcte conniventibus. ☉ In Ægypto (Delile) et in umbrosis rupium Arabiæ et Syriæ (Aucher-Eloi, Schimper, Unio it. ann. 1835. n. 121; Botta, Boissier, etc.). — Bl. l. c. p. 256; Wedd. Monogr. p. 519. t. 17. f. 26 et 27. *Freirea alsinifolia* Gaudich. Bot. Voy. Uran. p. 502. — Caules s. rami primarii graciles, 1-3-decimetrales, nonnunquam fasciculati, simplices vel simpliciter ramosi, debiles, patule pubescentes, exsiccati albido-fuscescentes, ramulis sæpius elongatis filiformibusque. Folia illa *Alsines* mediæ quoad formam bene referentia, sed minora, nempe plerumque vix centimetralia interdumque multo minora, basi rotundata vel obtusa, rarius acuta, superne attenuata vel subacuminata sæpiusque obtusa, in utraque pagina margineque subtiliter pubescentia, limbo exsiccato tenuiter membranaceo saturate viridi cystolithis creberrimis sed subinconspicuis obdito, petiolo tenui limbo ipso æquilongo aut paulo breviori puberulo. Cymulæ petiolis breviores, illæ nodorum superiorum interdum confluentes, bracteis inferioribus illas *P. debilis* figura referentibus persistentibus siccitate ut et folia ipsa obscure viridibus, superioribus forma diversa insignibus accretis ad maturitatem fructus 3 vel etiam 4 millim. latis et longis acutis aut obtusis tenuiter membranaceis venosis siccitate dilute viridibus vel subscariosis subtiliter puberulis una cum flore involucreto demum cadentibus. Flores hermaphr. perigonio intus parce lanulato, segmentis post anthesin elongatis lineari-spathulatis ciliolatisque castaneis apice virentibus. Perigonium floris fem. ovatum extus pilis clavatis majusculis obditum, segmentis obtusiusculis, fructiferum sesquimillimetrale, dilute castaneum vel subscariosum, achænium elliptico-oblongum 1½ millim. longum fusco-rubens suboblique apiculatum arcte amplectens. (v. s.)

β, *elongata* (Blume Mus. L. B. 2. p. 256), involucri foliolis obtuse acuminatis. ☉ In monte Sinai.

### *Species non satis notæ.*

9. *P. ELLIPTICA* (C. Koch, in Linnæa, 22. p. 606), erecta ramosa, caulibus et ramis glabris, foliis tenuibus ellipticis glabris margine puberulis supra tu-



herculis (sub lente) albidis obsitis obscure viridibus subtus subtiliter bullulatis pallidis nervis minime eminentibus petiolo triplo brevior instructis, floribus glomerulatis, involucri 1-2-floro 4-6-fido puberulo, perigonio fructifero elongato vix puberulo quadrilobo staminibus pæne triplo longiore lobis ciliatis faucem includentibus, achæniis nigrescentibus. ☉? Crescit in Græcia.

10. *P. PARVIFLORA* (Friedrichsth. mscr.; Bl. Mus. bot. lugd. bat. 2. p. 249), caule prostrato filiformi ramoso pubescente, foliis 3-nerviis rhombeo-ovatis v. subrotundis obtusis utrinque pubescentibus, involucri 3-flori foliolis inæqualibus obtusis ima basi cohærentibus, flore medio femineo. ☉? In rupibus Acropolis Athenarum (Friedrichsth., n. 179, in Mus. vindob.); etiam in rupetribus montosis Smyrnæ. — *P. cretica* (haud Linn.) Hb. Un. itin. Fleisch. — Folia 4-8-millim. longa, 4-5 millim. lata. — Transitum facit ad *P. creticam* L., sed multo magis nana ut *Helxine Soleirolii* Req., foliolis involucri triflori (quorum unum duo v. tria majora) fere ad basin usque distinctis post anthesin non induratis. Flos medius femineus, lateraliibus masc. v. hermaphr. (Blume.)

*Species exclusæ.*

- P. ARBOREA* L'Hérit. = *Gesnoulia arborea*.  
*P. ASPERA* Leschen. = alius ordinis.  
*P. CANESCENS* Poir. = *Pouzolzia canescens*.  
*P. CAPENSIS* Thunb. = ign.  
*P. COCHINCHINENSIS* Lour. = *Pouzolzia indica*.  
*P. CUNEATA* E. Mey. = *Didymodoxa cuneata*.  
*P. CUNEATA* Eckl. et Zeyh. = *Droguetia urticifolia*.  
*P. DECAISNEI* Steud. = *Pouzolzia indica*.  
*P. FILAMENTOSA* Webb = *Gesnoulia filamentosa*.  
*P. GUINEENSIS* G. Don = *Pouzolzia guineensis*.  
*P. LÆVIGATA* Poir. = *Pouzolzia lævigata*.  
*P. LANCEOLATA* Thunb. = ? *Didymodoxa debilis*.  
*P. LUSITANICA* Viv. = *Helxine Soleirolii*.  
*P. MICROPHYLLA* Linn. = *Pilea microphylla*.  
*P. ORIENTALIS* Tournef. = alius ordinis.  
*P. PROSTRATA* Spr. = *Pilea serpyllifolia*.  
*P. RAMIFLORA* Moench = ign.  
*P. RECLINATA* Moon = ign.  
*P. SERPYLLIFOLIA* Poir. = ? *Pilea serpyllifolia*.  
*P. SOLEIROLII* Spr. = *Helxine Soleirolii*.  
*P. SONNERATII* Poir. = *Phenax Sonneratii*.  
*P. URTICÆFOLIA* Linn. fil. = *Pilea rupipendia*?  
*P. VERBASCIFOLIA* Poir. = *Pilea verbascifolia*.  
*P. ZETLANICA* Linn. = *Pouzolzia indica*.

XXXV. *GESNOULIA* Gaudich. Bot. Voy. Uran. p. 502, Wedd. Monogr. p. 521, — *Parietariæ* sect. *Gesnoulia* Endlich. Gen. pl. p. 285, Webb Phytogr. Carnar. 3. p. 264. — *Urticæ*, *Boehmeriæ* et *Parietariæ* sp. auct.

Flores monoici, in cymulas 3-floras involucrias axillares aut interrupte spicatas paniculatasve digesti, floribus lateraliibus cymulæ masculis, intermedio femineo, involucri gamophylli 7-fidi laciniis linearibus

sæpius inæqualibus. *Masc.* : perigonium 4-partitum, segmentis ovatis longiuscule acuminatis ciliatis. Stamina 4. Pistilli rudimentum ovato-oblongum, subsessile vel stipitatum, acuminatum, basi lanatum. *Fem.* : perigonium ovoideo-tubulosum, limbo acute 4-dentato, tubo cum basi involucri arcte cohærente. Ovarium rectum, ovoideum. Ovulum erectum, imo loculo funiculi gracilis subobliqui ope affixum. Stylus elongatus, filiformis, apicem versus sensim incrassatus, inferne glaber, cæterum altero latere piloso-stigmatosus, deciduus. Achænium ovoideum, rectum, nitidum, perigonio vix accreto involucrique marcescentibus vestitum. Embryo cotyledonibus rotundatis radícula crassa longioribus. Albumen tenue. — Frutices suffruticesve canarienses. Folia alterna, integerrima, tri-tripli-nervia, cystolithis punctiformibus inspersa. Stipulæ omnino deficientes. — Genus a *Parietaria* distinguendum imprimis floribus haud polygamis et conformatione stigmati.

1. *G. ARBOREA* (Gaudich. Bot. Voy. Uran. p. 502), arborea rariusve fruticosa, ramulis novellis canescentibus, foliis lanceolatis v. oblongo-lanceolatis triplinerviis, glomerulis in paniculam amplam terminalem dispositis. 5-3 In sylvis umbrosis ins. Canariæ, Teneriffæ (Bourgeau, pl. can. n. 1506) et Palmæ. — Wedd. Monogr. p. 521. t. 18. B. f. 17. *Urtica* arborea Linn. Suppl. p. 417; L'Hérit. Stirp. nov. p. 39. t. 20. *Parietaria* arborea L'Hérit. in Journ. de Roz. 33. p. 55; Hort. Kew. ed. 1. 3. p. 429; Poir. Encycl. 5. p. 16; Webb Phytogr. Canar. 3. p. 264. t. 213. *Boehmeria* rubescens Jacq. Fragm. t. 5. f. 1. *B. arborea* Desf. Cat. pl. hort. reg. par. ed. 3. p. 348. — *Arbuscula* interdum 6-metralis, patule ramosa; ramulis teretibus rubentibus cano-pilosis, exsiccatis fusco-cinereis. Folia patentia, 4-12 cm. longa, 1-5 cm. lata (floralia cæteris multo minora), a medio vel supra medium ad apicem sensim attenuata, sæpius acuminata acutissimaque, inferne subcuneata imaque basi acutata obtusa rotundata vel subcordata, triplinervia (nervis basilariibus ad trientem superiorem limbi excurrentibus), juniora supra in nervis præsertim puberula subtusque molliter pubescentia, nervis hirtulis sed demum fere omnino glabris, lamina exsiccata membranacea olivaceo-v. fusco-viridi, petiolo gracili limbum dimidium longitudine circiter æquante vel paulo breviori interdumque longiore pubescente vel puberulo. Cymulæ (singulæ involucri proprio munitæ) numerosissimæ, glomerulatæ, glomerulis grani piperini circiter magnitudine in apicibus aphyllis ramorum ramulorumque superiorum (pariter aphyllorum) interrupte spicatis, spicis singulis simplicibus ramosisve una paniculam amplam elegantissimamque magis minusve foliosam et subnutantem efformantibus. Involucrum in vivo late coccineum sed siccitate sæpius obscurare ferrugineum, tubo campanulato millimetrali glabro segmentis linearibus ciliatis et obtusiusculis sæpius paulo breviori. Flores masc. subsessiles, perigonio subcampanulato albicante v. rubello, lobis abrupte et anguste acuminatis margine acumineque ciliatis, pistilli rudimento sessili ovato pariter acuminato in lana nidulante. Flores fem. perigonio tenui inferne glabro et cum tubo involucri cohærente apice exserto breviterque hirtulo, stylo subclavato hirtulo. Achænium vix millimetrale, fuscum, pericarpio tenui. (v. v. cult. et s.)
2. *G. FILAMENTOSA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 207; Monogr. p. 523. t. 18. B. f. 8), suffruticosa, caulibus diffusis, foliis ovato-lanceolatis v. subrhomboideis tri-tripli-nerviis, cymulis axillaribus. 3 In rupestribus convallium calidiorum Teneriffæ (Webb; Bourgeau, pl. canar. n. 520 et 1507; etc.). — *Parietaria* filamentosa Webb. l. c. p. 263. t. 212. — *Habitus* *Parietariae*. Caulis lignosus, tortuosus, rufescens, diffuse ramosus, ramis tenuibus elongatis foliosis pube patula cinerea vestitis, epidermide exsiccata fuscescente. Folia 8-20 mm. longa, 3-15 mm. lata, superne attenuata vel subacuminata, basi acuta aut obtusata, utrinque pubescentia supraque demum glabrata, lamina exsiccata membranacea ut plurimum nonnihil cinerascens, petiolo gracili pubescente longitudine variabili. Cymulæ breviter pedunculatæ, in axillis singulis geminæ. Involucrum in vivo dilute viride, tubo late subinfundibuliformi glabro, lacinjiis elongatis filiformibus acutissimis pilosis initio patulis

sed excusso florum masc. ornatu appressis. Flores masc. breviter pedicellati, perigonii campanulati segmentis subacuminatis acutissimis pilosis, pistilli rudimento elliptico apiculato longe stipitato apice hirtulo basi pilis copiosis stipato. Flores fem. sessiles, perigonio ut in specie præcedente (in involucri tubo autem minus immerso) fere glabro, dentibus ciliolatis. Achænium ovatum, submillimetrale, acutum, fuscum. (v. s.)

XXXVI. *HELKINE* Req. in *Ann. sc. nat.* 1<sup>re</sup> sér. 5. p. 384; *Duby, Bot. gall.* 1. p. 418; *Wedd. Monogr.* p. 529 — *Soleirolia Gaudich. Bot. Voy. Uran.* p. 504; *Bot. Voy. Bonite.* t. 114. B; *Endlich Gen. pl.* p. 285. — *Parietaræ spec. Viv. Append. ad Fl. Cors. prodr.* p. 7; *Spreng. Syst. pl.* 4. pars 2. p. 218, *aliorumque*. — *Parietaræ sect. Mutel, Fl. fr.* 3. p. 172; *Gren. et Godr. Fl. fr.* 3. p. 110.

Flores monoici, in axillis foliorum solitarii, masculi involucrio triphylo, feminei involucrio gamophyllo tubuloso alato apiceque trilobo præditi. *Masc.* : perigonium 4-partitum, segmentis ovatis subacuminatis, in præfloratione valvatis. Stamina 4. Pistilli rudimentum obovatum, basi pilis fasciculato-concretis petala abortiva quodammodo mentientibus stipatum. *Fem.* : perigonium ventricosum-tubulosum, ore contracto 4-laciniato. Ovarium oblongum. Ovulum e fundo loculi erectiusculum. Stigma penicillato-capitatum, stylo brevi suffultum, vix persistens. Achænium ovoideum, nitidum, pericarpio subnucumentaceo, perigonio marcescente et involucrio evoluto crasseque 3-alato inclusum. — *Herbula corsica et sardoa*. Caules filiformes, repentes. Folia alterna et subdisticha, inæquilateralia, integerrima, trinervia, cystolithis oblongis inspersa. Stipulæ nullæ. Flores in quaque axilla singuli una cum gemma foliifera (verisimiliter ejusdem dignitatis) nascentes. — Inflorescentia in hoc genere ab illa *Parietaræ* primo intuitu admodum discrepare videtur, at revera ad illam sat proxime accedit, interposito discrimine quod *Helxines* cymulæ flore unico constent.

H. *SOLEIROLII* (Req. l. c.), foliis oblique subrotundis breviter petiolatis sessilibusve sparsim pilosis. 2. Ad rupes et in fissuris scopulorum in humidis umbrosisque montium et secus torrentes Corsicæ et Sardiniae. — *Duby, l. c.*; *Wedd. Monogr.* p. 530. t. 18. A. *Parietaria lusitanica* Viv. l. c. t. 1. f. 1. P. repens *Soleirol exsicc.*; *Mutel Fl. fr.* 3. p. 172. P. *Soleirolii* *Spreng. l. c.*; *Gren. et Godr. l. c.*; *Mabille, Herb. cors. n.* 387. P. *cretica* *Moris Stirp. sard. elench. fasc.* 1. p. 41. *Soleirolia corsica* *Gaudich. l. c.* *Habitus Menthæ Requienii*. Caules numerosissimi, gracillimi, 3-15 cm. longi longioresve, e nodis inferioribus præsertim radicanter sæpissimeque intertexti, striati, puberuli, internodiis 1-10-millimetralibus. Folia 2-5 mm. lata, longitudine paulo minore, apice rotundata vel obtusa, sæpius in petiolum 1-2-millimetralem attenuata, lamina exsiccata tenuiter membranacea læte viridi supraque (in folio adulto) cystolithis oblongis s. fursiformibus sub lente distinctissimis ornata. Florum masc. involucrium breviter pedicellatum, foliolis linearibus apice parce pilosis, perigonii lobis acutis superne pilosis cæterum glabris, alabastro late turbinato breviterque acuminato. Flores fem. millimetrum circiter mentientes, involucrio infra limbum ob incrementum nervi medii foliolorum anguste et acute tri-alato, alis (pilis rigidis patulis et sursum uncinatis ornatis) ad maturitatem fructus valde incrassatis et semiorundatis, perigonii lacinulis ciliatis, tubo lanceolato superne parce piloso cæterum glabro. Achænium ovatum, millimetrale, acutum, fuscescens. (v. s.)

XXXVII. *HEMISTYLIS* Benth. *Pl. Hartweg.* p. 123; *Miq. in Mart. Fl. bras. fasc.* 12. p. 193; *Wedd. Monogr.* p. 524.

Flores monoici, masculis in glomerulos spicatos, femineis in cymulas bifloras involucratas digestis, involucris fem. ad basim spicæ macs. vel

in axillis foliorum superiorum rami sessilibus rariusve inflorescentiam distinctam efformantibus, e bracteis duabus late ovatis ima basi angustatis in tubum brevem coalitis superiusque lineari-auriculatis constantibus. *Masc.* Perigonium 4-partitum, segmentis acutis in præfloratione valvatis. Stamina 4. Pistilli rudimentum obovatum, basi lanatum. *Fem.* Perigonium ventricosum-tubulosum, ore contracto 4-dentato, fructiferum valde compressum et membranaceo-marginatum, facie altera cum tubo involucri; altera cum perigonio floris contigui connatum s. ut minime dicam admodum adhærens. Ovarium lanceolato-ovatum, liberum. Stylus filiformis, rectiusculus v. incurvus, cum summo ovario articulatus et ideo mox deciduus, latere altero a basi ad apicem stigmatosus. Achænium compressum, perigonio et involucro evoluto marcescenteque diu vestitum. Semen pericarpio conforme. Albumen tenue. Embryo cotyledonibus quadrato-orbicularibus utrinque subemarginatis, radícula ovidea multo longioribus. — Arbusculæ v. frutices colombiani. Folia alterna, ampla, sæpius integerrima, tri-tripli-nervia, cystolithis punctiformibus. Stipulæ axillares, liberæ, caducæ. Glomeruli masc. (interdum valde pauciflori) in axillis bractearum brevium subscariosarum nascentes. Receptaculum involucri fem. nonnunquam inter flores magis minusve productum (ut in genere sequente). Perigonium fructiferum utroque margine a basi ad apicem frequentissime scissum. — Indole perigonii fem. genus *Hemistylis* ad *Rousselliam* nonnihil accedit, cæterum vero admodum discrepat.

1. *H. BOEHMERIOIDES* (Benth. l. c.), foliis late ovatis subabrupte acuminatis basi rotundatis v. fere truncatis integerrimis supra subtiliter hispida asperulisque subtus (nervis venulisque exceptis) albo-tomentosis, spicis androgynis solitariis fasciculatisve erectiusculis quam folia brevioribus. § In sylvis ditionis Equatoris, prope Guayaquil (Hartweg n. 696). — Miq. l. c.; Wedd. Monogr. p. 525. — Ramuli crassi, herbacei, breviter pubescentes, siccitate fuscescentes et fistulosi, internodiis 1-3-centimetalibus. Folia 8-12 cm. longa, 6-8 cm. lata, acumine 1-1½-centimetro, nervis basilariibus ex ima basi laminæ ortis ad partem quartam superiorem ejusdem productis subtus nervisque secundariis et tertiariis (parallele transversis) in limbo exsiccato fuscescentibus valdeque evidentibus, petiolo 2-4 cm. longo puberulo. Stipulæ parvæ, ovato-lanceolatæ, acuminatæ. Spicæ in axillis superioribus ramorum solitariæ sæpiusve geminæ aut ternæ, 3-5 cm. longæ, graciles, in specimenibus examinatis erectiusculæ, breviter pedunculatæ, pedunculo puberulo, glomerulis approximatis. Flores masculi sessiles, perigonio puberulo; feminei in glomerulis inferioribus cum masculis intermixtis aut involucratiss, involucris ad basim spicæ sessilibus solitariis. (v. s.)
2. *H. MACROSTACHYA* (Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 208; Monogr. p. 526. t. 18. C), foliis ovatis sensim acuminatis basi rotundatis integerrimis supra subtiliter pubescentibus demumque glabratis asperulisque subtus molliter pubescentibus junioribusque canescentibus, spicis androgynis pendulis quam folia multo longioribus. § In Venezuela, inter La Guayra et Caracas (Plée). — Ramuli floriferi crassitie pennæ anserinæ vel cycneæ, superne pubescentes demumque glabrati, siccitate longitudinaliter rugosi et fusi, internodiis vix semipentimetalibus. Folia adulta 6-10 cm. longa, 3-5 cm. lata, nervis basilariibus ex ipsa insertione petioli vel paulo supra illam nascentibus usque ad partem quartam v. quintam apicalem excurrentibus, illis limbi adulti subtus cum intervenio concoloribus, petiolo gracili quam limbus dimidiis brevior puberulo. Stipulæ lanceolato-subulatæ, 4-5-millimetrales. Spicæ 1-2-decimetrales, graciles, longiuscule pedunculatæ; glomerulis masc. confluentibus, floribus sessilibus, perigonio pubescente. Involucra fem. ad basim spicæ masc. solitaria vel gemina, rarius plura, foliolis cordatis 4-5 mm. longis latisque hirtulis, lobulis basilariibus circiter sesquimillimetalibus. (v. s. in hb. Mus. par.)
3. *H. VELUTINA* (Wedd. Monogr. p. 526), foliis ellipticis acuminatis basi acutis

- integerrimis supra velutinis subtus (costa et nervis primariis exceptis) incanotomentosis, floribus masc. spicatis (spicis pendulis quam folia brevioribus) involucris fem. axillaribus.  $\S$ ? In Nova Granata (Triana). — Ramulus in specimine unico examinato modice crassus, adpresse cinereo-pubescentis, internodiis centimetralibus. Folia a basi ad apicem ramuli valde imminuta, summa nempe fere bracteiformia sessilia lanceolata et vix centimetralia, inferiora s. majora 10-15 cm. longa, 5-8 cm. lata, anguste acuminata, infra medium sensim attenuata longeque petiolata, nervis basilaribus usque ad quintam vel sextam partem superiorem limbi productis subtus in sicco fusciscentibus, secundariis tertiariisque parum conspicuis, petiolis foliorum inferiorum 3-8 cm. adpresse canescenti-puberulis. Stipulæ ovatæ, 3 mm. circiter longæ, subabrupte attenuatæ. Spicæ masc. in specimine viso 6-10 cm. longæ, pedunculo 1-2-centimetrali suffultæ; graciles, teretes, floribus sessilibus confertis, perigonio tomentosus. Involucra fem. in axillis foliorum superiorum rami vel in ramulis abbreviatis ex iisdem axillis ortis sessilia, foliolis ut in specie præcedente sed lobulis basilaribus longioribus. (v. s. in hb. Mus. par.)
4. *H. ODONTOPHYLLA*, foliis late ovatis acuminatis basi rotundatis v. subcordatis inæqualiter dentatis supra pilosulis subtus pubescentibus concoloribus, floribus masc. spicatis, spicis in parte inferiore rami axillaribus folio brevioribus longioribusve, involucris fem. in paniculam amplam terminalem dispositis.  $\S$  In Venezuela prope coloniam Tovar (Fendler, n. 1827). — Rami ramulique virgati, villosi, in sicco pallide fusco-rubentes, internodiis partis foliiferæ sat elongatis, floriferæ contra brevibus aut brevissimis. Folia inferiora 8-15 cm. longa, 5-9 cm. lata, sensim et anguste acuminata (acumine integro), basi integra, dentibus reliqui limbi acutiusculis obtusisve, nervis basilaribus ex ipsa insertione petioli ortis usque ad trientem superiorem limbi productis, venulis parallelæ inter nervos primarios anastomosatis, petiolo 2-5-centimetrali pubescente. Stipulæ parvæ deciduæ. Spicæ masc. graciles 10-15 cm. longæ, pedunculo bidentimetro tomentoso, floribus in axi canescenti-tomentoso sessilibus glomerulatis congestis, perigonio subincano-pubescente. Panicula florum fem. 2-4 decim. longa, pyramidalis, ramulis gracillimis simplicibus ramosisve, bracteis plerisque parvis lanceolatis cæteris foliaceis. Involucra admodum numerosa sessilia, foliolis cordatis 4-5-millimetralibus pilosulis ciliatisque, lobulis s. laciniis basilaribus 2 mm. longis densius hirtulis. — Stirps inter Parietarleas valde insignis, propter folia dentata. (v. s. in h. Kew.)

XXXVIII. *ROUSSELIA* Gaudich. Bot. Voy. Uran. p. 503; Bot. Voy. Bonite, t. 98. A; Wedd. Monogr. p. 527. — Parietariæ sect. Rousselia Endlich. gen. pl. p. 285. — Urticæ spec. auct.

Flores monoici, in axillis foliorum superiorum cymulosi, cymulis masc. laxiuscule paucifloris exinvolucratis, femineis bifloris involucro sessili e bracteis duabus late ovatis integrisque constante munitis. Masc. Perigonium 4-partitum, segmentis acutis in præfloratione valvatis. Stamina 4. Pistilli rudimentum anguste conicum s. teretiusculum, basi lanatum. Fem. Perigonium ventricosum-tubulosum, ore contracto bi-quadri-dentato, fructiferum compressum angustiusque membranaceo-marginatum, facie altera cum involucri pagina superiore excavata confærens. Ovarium ovatum, liberum. Stylus filiformis, nonnihil incurvus, in latere uno plumoso-papillosus, cum apice ovarii articulatus. Achænium compressum, perigonio involucrique evolutis et marcescentibus diu vestitum. Semen pericarpio conforme. Albumen tenue. Embryo cotyledonibus quadrato-orbiculatis, basi et apice subemarginatis, radícula ovoidea iisdem dimidio brevior. — Herba perennis, in insulis Antillanis hucusque solum lecta. Caules graciles repentes. Folia alterna, integerrima, trinervia. Stipulæ liberæ, petiolares, persistentes. Flores fem. in utraque facie receptaculi communis admodum compressi et disciformis inserti horizontaliterque patentes s. divergentes. — Genus monotypum, præcedenti solo pro involucri fem. conformatione nonnihil affine.

R. LAPPULACEA (Gaudich. l. c.), foliis ovatis v. rhomboideis sparsim pilosis.  $\gamma$  In ins. S.-Domingo, Jamaica, Cuba (Wight, n. 1681), S.-Thomas, Porto-Rico, etc. — Wedd. Monogr. p. 528. t. 18. D; Griseb. Fl. Brit. W. Ind. p. 159. Urtica humilis et U. lappulacea Swartz. — Caules numerosi, filiformes, longe repentes, parce ramosi, glabrati, ramis adscendentibus 1-2-decimetralibus, puberulis, parce foliosis, internodiis scilicet plerumque 2-3 cm. longis vel longioribus. Folia vulgo 5-20 mm. longa, 4-15 mm. lata, interdum fere rotundata, acuta v. obtusata, inferne sæpius cuneata, lamina exsiccata membranacea supra obscure subtus dilute viridi, petiolo gracili quam limbus sæpe longiore ut costa nervique paginæ inferioris pubescente. Stipulæ triangulari-lanceolatae, siccitate ferrugineæ. Cymulæ parum conspicuæ, in axillis superioribus præsertim obviæ; masculæ pedunculatæ, pedunculo filiformi 2-4-millimetrali cum petioli basi connato puberulo, floribus minimis breviter pedicellatis singulis bractea oblonga ciliata suffultis, perigonio extus hirtio. Involucra fœm. sessilia, diametro majore 3-4-millimetrali, foliolis late ovatis acutiusculis obtusisve subinæquimagnis venosis ciliatis supraque pilis hamatis majusculis conspersis. Perigonium fructiferum  $1\frac{1}{2}$ -1 $\frac{1}{4}$  mm. circiter latum, ciliolatum, achænium ovatum compressum fuscum nitidulum arcte includens. (v. s.)

$\beta$ , fruticulosa, caule suffruticuloso ramosissimo. — In iisdem locis ac typus. (v. s.)

### Tribus V. FORSKOHLÆÆ.

Herbæ annuæ aut sæpius perennes vel suffrutices, stimulis veris destituti, sed interdum pilis induratis aculeolati. Folia alterna aut opposita, homomorpha, æquilateralia, varie dentata, rarissime integra, cystolithis punctiformibus inspersa. Stipulæ petiolares et liberæ vel petiolari-laterales et tunc (saltem in speciebus oppositifoliis) connatæ. Inflorescentiæ involucreatæ (involucro gamo-polyphyllo), aut exinvolucreatæ, floribus involucreis in lana copiosa sæpius nidulantibus. Flores unisexuales. *Masc.* : perigonio sæpius monophyllo, basi tubuloso, superne dilatato bracteiformi v. subbilabiato, rarissime diphylo; stamine unico, rarissime stam. duobus; pistilli rudimento nullo, aut (in flore diphylo) lineari et densiuscule lanato. *Fem.* : perigonio tubuloso-ventricosso (ore 4-dentato), aut deficiente; stigmate filiformi aut rarissime subcapitato; staminum rudimentis nullis. Fructus siccus. Embryo cotyledonibus ellipticis, oblongis rotundatisve radícula longioribus; albumine vario, nunquam deficiente.

Subtrib. 1. *EUFORSKOHLÆÆ*. Flores in involucro gamo-polyphyllo aggregati et plerumque densissime lanati.

XXXIX. FORSKOHLÆA Linn. Mant. p. 11. n. 1262; Juss. Gen. pl. p. 403 (Forska-lea); Endlich. Gen. pl. p. 285 (Forskolea); Wedd. Monogr. p. 532. — Caid-beja Forsk. Ægypt. p. 82.

Flores monoici, ut plurimum in involucreis turbinatis 3-6-phyllis intus densissime lanatis arcte congregati, masculi numerosiores in circuito, feminei 1-5 centrales, femineis rarius solitariis in involucro diphylo, involucreis in axillis foliorum bractearumve geminis vel glomeratis. *Masc.* Perigonium monophyllum, basi anguste tubulosum, superne dilatatum s. bracteiforme obtuseque 3-dentatum s. lobatum, florem unilabiato referens, lobo intermedio in alabastro compressiusculo inflexo. Stamen unicum. Pistilli rudimentum nullum. *Fem.* Perigonium nullum.

Ovarium rectum, ovato-vel elliptico-lanceolatum, lanatum, superne in stigma filiforme villosa-hispidum abiens. Ovulum nonnihil obliquum, basi funiculo brevi continuum. Achænium rectum, compressum, ovatum, punctulatum atque (ut ipsius involucri facies interna tubusque perigonii masc.) lana gossypina quam maxime intricata densissimaque obductum. Embryo cotyledonibus rotundatis basi emarginatis radícula cylindrica longioribus. Albumen modicum. — Herbæ suffruticesve africanæ, in Europa mediterranea vero necnon in Arabia et India orientali etiam obvii, novo orbe exsules, tenaces, pilis rigidis basi incrassatis subpungentibusque asperato-hispidi. Folia alterna, crenata aut varie dentata, triplinervia, pilis hamatis adscendentibusve obsita. Stipulæ laterales, liberæ. Involucra extus longe sericeo-pilosa. — Genus admodum naturale. Discrimen specierum non semper facile.

1. *F. TENACISSIMA* (Linn. Mantiss. p. 72) Syst. pl. ed. 13. p. 363, foliis orbicularibus v. obovatis v. rhomboideis apice obtusis rotundatisve, supra basim cuneatam integramque grosse crenato-serratis, pagina superiore viridi pilisque hamatis densiuscule hispida, inferiore (nervis exceptis) bombycino-tomentosa, capitulis geminis v. quaternis, involucri vulgo 4-5-foliolati foliolis lanceolatis acutis prope apicem rigide ciliolatis cæterum præsertim in dimidio inferiore longe denseque sericeo-pilosis. 4-5 In Numidia, prope Biskra, etc.) (Jamin, Balansa, n. 751, etc.), Ægypto, Arabia (Schimp. pl. arab. n. 230, etc.), Aden (Hook. fil.), Beluchistan (Stocks), Afghanistan (Griffith), et Indiæ orientalis provinciis occidentilibus, ad Simla (Stocks) et Punjab (Jacquemont, Fleming). — Jacq. Vindob. p. 48; Lam. Encycl. 2. p. 522, illustr. t. 388; Desfont. Fl. atlant. 2. p. 329. *F. latifolia* Retz. Obs. 3. n. 51. *Caidbeja* adhærens Forsk. Ægypt. p. 82. Herba basi lignescens vel suffrutex 1-5-decimetralis, a basi diffuse ramosus, caule ramisque asperato-s. aculeolato-hispidis et inter spinulas adpresse tomentosus, epidermide ipsa sæpissime rubente, internodiis brevibus. Folia vulgo 1-2 cm. longa, 7-18 mm. lata, dentibus acutiusculis aut obtusis, supra magis minus tomentosa et valde asperata, in vivo viridia, siccitate sæpius fusco-cinerea, nervis paginæ inferioris laxè patuleque albo-pilosis, tomento intervenii molli, petiolo dimidium limbum circiter æquante vel etiam paulo longiore ut limbi margines rigide ciliato parceque tomentoso. Stipulæ ovatæ, acutæ, scariosæ, pilis rigidis longiuscule ciliatæ. Capitula numerosa, immo in ramulis quibusdam ob breviter internodiorum imbricata. 6-10 mm. longa, involucri fere usque ad apicem pilis sericeis vestito, pilis siccitate haud raro virescentibus, foliolis lanceolatis nec apice dilatatis. Flores in singulis involucri numero quam maxime vario, masculi scilicet interdum nulli (in involucri bifoliolato), aut (in involucri plurifoliolato) 8-12; feminei 1-6, omnes lana bombycina densissima albida, aut (siccitate?) ferruginea intertexta obvoluti. Perigonium masc. apice inæqualiter 3-dentato-lobatum, lobo s. dente intermedio majore obtuso et hispido-ciliato. (v. s.)

β, *Cossoniana*, capitulis plerumque geminis, involucri sæpius trifoliolato. 4-5 In arenosis aridis Hispaniæ prov. Almeria (Bourgeau, pl. hisp. n. 1486). — *F. Cossoniana* Webb Otia hisp. p. 49, t. 44; Wilk. et Lange Prodr. fl. hisp. p. 253. (v. s.)

2. *F. PROCRIDIFOLIA* (Webb Spicil. gorgon. in Hook. Fl. Nigrit. p. 179), foliis elliptico-lanceolatis ovatisve superne attenuatis et acutissimis deorsumque in petiolum longiusculum sæpe decurrentibus et supra basim v. partem quartam tertiamve inferiorem serratis v. dentato-serratis, dentibus spinosis, pagina superiore viridi pilisque rectis aut (in folio juniore) hamatis hispido, inferiore inter nervos primarios incano-tomentosa, capitulis geminis, involucri vulgo 3-10-foliolati foliolis lanceolatis acutis membranaceis subcoriaceisve inferne pilosis superne rigide ciliatis. 3 In insulis gorgonicis, ad rupes et in locis montosis. — Schmidt, Beitrag. zur Fl. C. Verd. Ins. p. 169; Wedd. Monogr. p. 535. t. 19. B. f. 1-5. — Suffrutex magis minusve ramosus, ramis nunc elongatis erectisque nunc diffusis pilis rigidis s. aculeolis numerosissimis adscendentibusque hirtis aut etiam fere inermibus, epidermide sæpius fusco-rubente et inter pilos pubescente, internodiis 1-4-centimetralibus. Folia quam

in *F. tenacissima* vulgo majora, 3-6 cm. circiter longa, supra plus minus asperata, pagina superiore folii exsiccati obscure viridi, inferiore cinerea v. nivea, nervis laxè et patentim albo-pilosis, petiolo longitudine varia patule hirsuto v. pubescente. Stipulæ ovato-lanceolatæ, subacuminatæ, margine et basi præsertim scariosæ margineque longiuscule pilosæ. Capitula distincta nec imbricata, 6-10-millimetralia, involucri raro bifoliolato sæpius turbinate et 6-8-foliolato basi tantum et angulis piloso, foliolis membranaceis aut rigidis extus parce hispidis. Flores numero admodum variabiles, perigonio masc. fere ut in *F. tenacissima*, stylo (teste Webb) quam in illa pilosiore, lana albidâ. (v. s.)

β, *umbrosa*, foliis amplioribus grosse obtuseque crenatis membranaceis. γ Cum typo, sed in locis minus apertis. — Folia interdum 10-centimetralia, 4½ centim. lata, capitula ampliora. (v. s.)

γ, *microphylla*, foliis multo minoribus quam in typo argute serratis dentibus spinosis. δ In ins. gorgonicis. — *F. proceridifolia* var. *microphylla* Schmidt l. c.? *F. candida* Ch. Smith, in Tuck. Voy. p. 251; Webb l. c. non Linn. fil. (v. s.)

3. *F. CANDIDA* (Linn. fil. Suppl. p. 245), foliis elliptico-ovatis ovatisve acutis inferne in petiolum attenuatis integrisque cæterum sinuato-dentatis v. grosse serratis supra asperato-hispidis subtus inter nervos incano-tomentosis, capitulis geminis, involucri 2-5-foliolati foliolis obovato- v. lanceolato-spathulatis acutiusculis aut obtusis membranaceis inferne densiuscule pilosis in dimidia parte superiore ampliata nitidulis ciliatisque. δ In arenosis Africae australis (Zeyher, n. 1544; Drege, etc.) — Willd. Spec. pl. 2. p. 475; Thunb. Prodr. p. 77; Wedd. Monogr. p. 536. *F. scabra* Retz. Obs. 3. n. 49. — Caulis magis minusve asperato-hispidus, 2-3 decim. altus. Rami erecti, graciles, parce et subtiliter aculeolato-hispidi, cæterum glabri v. puberuli, siccitate fusci, internodiis 5-10-millimetrales. Folia 1-2-centimetralia, 5-10 mm. lata, dentibus seu serraturis magnitudine variabili utrinque 2-4 acutis obtusisve, tomento paginæ inferioris adpresso niveoque, nervis parce hispidis, pagina superiore siccitate ut plurimum nigricante, petiolo 3-10 mm. longo puberulo asperatoque aut glabrato. Stipulæ parvæ, ovatæ vel lanceolatæ, pilosulæ. Capitula discreta, 5-8 mm. longa, involucri in dimidio inferiore tantum piloso, foliolis inferne linearibus at in dimidio superiore subrhombicis, in sicco scariosis aut rarius subcoriaceis dorso vix hispidulis margineque breviter ciliatis. Flores in singulis involucri ut plurimum pauci, lana gossypina albidâ aut pallide ferruginea intertexta uniti cæterumque ut in *F. tenacissima*. (v. s.)

β, capitulis majoribus s. fere centimetralibus, foliolis lanceolato-spathulatis superne granulosis. (v. s.)

γ, *virescens*, foliis oblongo-spathulatis (dentibus minoribus quam in typo) subtus virentibus. — In Africa australi. (v. s. in hb. Lenormand.)

4. *F. VIRIDIS* (Ehrenb. in hort. bot. Berol., ex Desfont. Cat. hort. bot. par. ed. 3. p. 347), foliis ovato-lanceolatis lanceolatisve utrinque attenuatis acuminatisve longiuscule petiolatis inferne integris cæterum grosse et irregulariter crenato-serratis rariusve subintegris concoloribus aut in pagina inferiore nonnihil pallidioribus s. subcinereo-tomentellis pilis hamatis ut plurimum omnino destitutis supraque parce hispidulis, capitulis aliis bi aliis 4-foliolatis, hisce geminis quaternisve, foliolis obovatis obtusis aut (in involucri bifoliolato) ovatis acutis acuminatisve membranaceis inferne in marginibus præsertim sericeo-pilosis supra medium parce ciliatis hispidisque. γ - δ In vallibus altis, circa portum Praya ins. gorgonæ S. Jacobi (J. D. Hooker); in Abyssinia prope Gurrsofa, altit. 1200 metr. (Schimper). — Webb, Spicil. gorgon. l. c.; Wedd. Monogr. p. 537. t. 19. B. f. 5 et seq. — Caulis et rami basi vulgè lignescentes, graciles, pilis tenuibus hispidi aut glabriusculi, siccitate fuscescentes, internodiis 1-2 centimetralibus longioribusve. Folia 1-5 cm. longa, 6-25 mm. lata, dente terminali s. acumine sæpius obtusiusculo, serraturis lateralibus (numero variabili) nunc obtusis nunc acutiusculis muticis ciliatis, lamina exsiccata membranacea pallide viridi, nervis subtus pilosis, intervenio (in folio juniore) nonnunquam parce tomentello, petiolo longitudine dimidii limbi vel brevioris puberulo vel glabrato. Stipulæ ovatæ, acuminatæ,



subcartilaginea, margine pilosæ. Capitula discreta nec imbricata, 5-8 mm. longa, turbinata; limbo valde ampliato s. patente. tubo in angulis præsertim piloso, foliolis involucri turbinati fere semper quatuor foliaceis, floribus lana albidâ immersis, cæterum ut in præcedente. (v. v. cult. et s.)

5. *F. ANGUSTIFOLIA* (Retz. Obs. bot. fasc. 3. p. 31), foliis lanceolatis v. oblongo-lanceolatis utrinque attenuatis inferne integris cæterum sinuato-dentatis supra viridibus pilisque hamatis parce conspersis subtus nisi in nervis incano-tomentosis, capitulis parvis sed sæpius valde numerosis in singulis axillis geminis, involucri 3-6-foliolati foliolis anguste lanceolatis acutis fere usque ad apicem dense sericeo-pilosis. 3 in rupestribus siccis et in ruderatis regionis inferioris insularum canariensium omnium ubi copiosissima (Bourgeau, n. 479, ann. 1845, et n. 1504, ann. 1855, etc.). — Lam. Illustr. t. 388; Webb Phytogr. canar. 3. p. 266; Wedd. Monogr. p. 538. *F. fruticosa* Willd. Enum. hort. Berol. 2. p. 950; Spreng. Syst. veget. 3. p. 777. — Caulis basi lignosus, diffuse ramosus, ramis pilis rigidis adscendentibus asperato-hispidis, epidermide exsiccata fusca, internodiis sæpe brevissimis. Folia 8-30 mm. longa, 1½-10 mm. lata, acuta, dentibus sæpe spinuloso-mucronatis, marginibus (sæpe maxis minus revolutis) spinuloso-ciliolatis, lamina exsiccata rigide membranacea supra obscure viridi, nervis subtus sparsissime albo-pilosis, petiolo gracili 2-8 mm. longo pubescenti-hirto. Stipulæ parvæ, lanceolatæ, margine pilosæ. Capitula ut plurimum valde numerosa et in ramulis lateralibus dense imbricata s. glomerulata, glomerulis confluentibus. Involucria quam in cæteris speciebus ut plurimum multo minora, scilicet didymum centimetrum longitudine raro excedentia, anguste turbinata et undique longe pilosa, pilis haud raro rufescentibus. Flores quam in speciebus præcedentibus minores, ætæroquin quoad formam similes, in lana albidâ aut pallide ferruginea nidulantes. — Variat caule ramisque plus minus abbreviatis elongatis, nec notâ magnitudine foliorum quæ interdum sed raro utrinque viridia sunt. (v. s.)

XL. *DROGUETIA* Gaudich. Bot. Voy. Uran. p. 505; Wedd. Monogr. p. 538. — Forskohlæ spec. Wight, Icon. pl. Ind. VI. t. 1982. — Didymogyne Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 35, ex parte. — Urticæ spec. auct.

Flores monoici, in involucriis gamophyllis campanulatis aut tubulosis numero admodum variabili (1-20), sessiles, involucriis axillaribus aut racemoso-spicatis ut plurimum androgynis (floribus fem. in involucrio multifloro paucioribus centralibusque), rarius (in involucriis paucifloris) omnino femineis. *Masc.* Perigonium ut in Forskøhlea monophyllum basi que anguste tubulosum, extus lanatum, superne dilatatum, stamen unicum involvens, marginibus in alaba-tro sæpius omnino concretis demumque floris evolutione dilaceratis. Pistilli rudimentum nullum. *Fem.* Perigonium nullum. Ovarium quoad formam et lanæ indumentum illo Forskøhleæ simillimum aut glaberrimum ovariumque Boehmeriæ potius referens, ovariis duobus interdum in unum concretis. Stigma filiforme cum apice ovarii continuum, villosulum. Ovulum erectum aut subadscendens, funiculo brevi suffultum. Achænium ut in Forskøhlea aut glaberrimum nitidissimumque. Embryo cotyledonibus ellipticis radícula crassa paulo longioribus. Albumen magis minusve copiosum. — Herbæ perennes v. suffrutescentes, omnes africanæ (specie unica in India orientali etiam lecta), caulibus lævibus. Folia alterna aut opposita, serrata, pilis hamatis destituta, cystolithis punctiformibus. Stipulæ laterales, liberæ. Involucria dentata, pubescentia v. tomentosa. — Planta feminea cujusdam speciei hujus generis pro pistillis glaberrimis in involucrio tubuloso-ventricosum subsolitariis cum Boehmeria facillime confundenda.

1. *D. LEPTOSTACHYS*, herbacea v. suffruticulosa, rhizomatibus longe repentibus, foliis oppositis ovatis aut elliptico- v. ovato-lanceolatis sensim acuminatis acutisve excepta basi acute serratis asperis, involucriis bi-pluri-floris omnibus in

spicas s. racemos angustos axillares et terminales dispositis aut aliis spicatis alisque (majoribus) axillaribus, achæniis lanatis. 2-3 In sylvis excelsioribus montium insulæ Bourbon (Commerson, cæt.). — *D. ovata* et *D. elliptica* Gaudich. Bot. Voy. Uran. t. 86 l. c. *D. ovata* Wedd. Monogr. p. 540. t. 19. f. 12 et seq. *Urtica leptostachys* Juss. mscr. in Pers. Synops. 2. p. 554. *U. leptostachya* Smith in Rees' Cycloped. 37, *Urtica*, n. 65. — Rhizomata metralia vel etiam longiora. Caules graciles, 2-4 decim. alti, diffuse ramosi, adscendentes, radicales, pubescentes. Folia vulgo 2-5 centim. longa, 1-2½ centim. lata, supra parce pilosa asperataque, nervis subtus pubescentibus et pilosis, lamina exsiccata membranacea sæpiusque viridi, petiolo longitudine varia pubescente. Stipulæ subulatæ, 1-2-millimetrales, parum conspicuæ. Spicæ s. racemi simplices vel parce ramosi paniculamque terminalem efformantes, sæpius aphylli et longitudine admodum variabiles (haud raro decimetrales). Involucra approximata, plerumque anguste campanulata, obtuse dentata, pubescentia. Flores masc. nunc in involucrio tubuloso cum flore fem. unico solitarii, nunc numerosi, involucrio tunc latius campanulato s. fere patelliformi, perigonii limbo obtuse tridentato dentibus lateralibus interdum obsoletis intermedio multico, extus ut flores fem. et involucri facies interna lana densissima obvestiti. Achænia ovata, circiter millimetralia, ventricosa aut plus minus compressa, fusca, pericarpio sub lana punctulato. (v. s.)

2. *D. THOUARSIANA* (Gaudich. Bot. Voy. Bon. t. 87), fruticulosa, subcæspitosa, foliis alternis parvis ovatis ellipticisve acutis v. obtusis acute vel obtusiuscule serratis supra asperulis nervisque subtus sparsim hispidulo-pilosis, involucri bi-pauci-floris axillaribus, achæniis lanatis. 3 In locis humidis montium insularum Mauritiæ et Bourbon (Commerson, cæt.). — Wedd. Monogr. t. 541. — Caules fruticulosi, numerosi, graciles, 1-4 decim. alti, subsimplices aut diffuse ramosi, adscendentes aut fere prostrati, basi sæpe denudati glabratique, superne pubescentes, internodiis brevibus. Folia 3-15 millim. longa, 2½-8 millim. lata, interdum (minora præsertim) fere rotundata, basi plerumque integra obtusaque aut subcuneata, lamina exsiccata rigide membranacea obscure viridi, petiolo 1-5-millimetrali pubescente. Stipulæ parvæ, subulatæ, puberulæ. Involucra campanulata, uni-sesqui-millimetralia, pubescentia, in axillis superioribus ramorum solitaria vel pauca glomerata, sessilia, in speciminibus omnibus visis admodum pauciflora. Cætera ut in specie præcedente. (v. s.)
3. *D. URTICÆFOLIA*, caulibus herbaceis diffusis v. erectiusculis, foliis oppositis rariusve paucis alternis late ovatis supra basim cuneatam sæpius grosse crenatis v. crenato-serratis utrinque sparsim pilosis, involucri axillaribus sæpe glomerulatis bi-pluri-floris, achæniis lanatis. 4 In Africa australi (Drege, Zeyher n. 3866, Gueinzus, n. 174. — *D. ambigua* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 39; Monogr. p. 543. *P. urticæfolia* E. Mey. pl. exs. Dreg., non Linn. fil. *P. cuneata* Eckl. et Zeyh. Bot. Zeit. 28. p. 88. — Caules adscendentes, non aut vix radicales, ramique parce rubentes. Folia 1-5 centim. longa, 1-3 centim. lata, in triente inferiore cuneata integraque, serraturis s. crenis reliqui limbi obtusis, dente apicali ovato, laminæ exsiccatae membranaceæ saturate viridis pagina inferiore sæpe glaucescente, petiolo 3-15 millim. longo gracili pubescente. Stipulæ ovatæ, 2-4-millimetrales, subulato-acuminatæ, ciliatæ, siccitate albo-scariosæ, nervo dorsali viridi. Involucra gemina aut glomerulata, campanulata v. tubuloso-campanulata, 1½-2 millim. alta, 1-3 millim. lata, margine acute dentata v. denticulata, extus pubescentia, intus lanata. Flores masc. in involucrio cum flore unico fem. solitarii, nunquam numerosi, perigonio anguste acuminato inferne præsertim lanato. Flores fem. interdum marginibus geminatim concreti, achæniis cæterum ut in *D. ovata*. (v. s.)
4. *D. PAUCIFLORA*, caulibus herbaceis lignescentibusve diffusis, foliis oppositis ovatis acuminatis crenato-dentatis serratisve supra sparsim hispidulo-pilosis nervis subtus pilosulis, floribus interdum dioicis, involucri axillaribus uni-multi-floris, achæniis glaberrimis nitidisque. 2-3 In Abyssiniæ locis umbrosis regionis mediæ montis Selki, etc., alt. 2600 metr. (Schimper), necnon in peninsula Indiarum orientalis montibus Nilghiri (Wight, etc.) et ins. Java (Zollinger, n. 2185). — *D. diffusa* Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 211.

Monogr. p. 541. t. 19, A. f. 1-11. *Urtica pauciflora* Hochstet. in pl. Schimp. exs. n. 682. *U. rivularis* Schlechtend. in pl. ind. Hohenack. n. 1326. *Pouzolzia pauciflora* A. Rich. Tent. fl. Abyss. 2. p. 259; Walp. Ann. bot. 3. p. 416. *Forskohlea urticoides* Wight, Icon. pl. Ind. t. 1982. *Didymogyne abyssinica* Wedd. in Ann. sc. nat. 1. c. *Boehmeria pauciflora* Bl. Mus. bot. lugd. bat. 2. p. 201. *B. parvifolia* Wedd. Monogr. p. 259. — Caulis longitudine seu altitudine et crassitie admodum variabiles, sæpissime basi prostrati radicantesque et demum adscendentes, obtuse angulati, pubescentes aut demum (præsertim inferne) glabrati, internodiis sæpe valde elongatis. Folia 1-6 centim. longa, 3½-6 centim. lata, basi obtusa v. late cuneata, in triente aut quarta parte inferiore integra, dentibus s. serraturis subæqualibus nonnunquam grossiusculis, acumine integro oblongo-lanceolato acuto v. obtusiusculo, lamina exsiccata membranacea supra obscure subtus pallide viridi, petiolo gracili 5-30 millim. longo pubescente aut patentim piloso. Stipulæ quam in cæteris speciebus evidētiore, ovatæ vel ovato-lanceolatæ, 2-5-millimetrales, sæpe subulato-acuminatæ, siccitate ferrugineæ, nervo dorsali pilosæ, cæterum glabræ v. ciliolatæ. Involucra in singulis axillis sæpius gemina, ventricosotubulosa aut late campanulata s. crateriformia, pubescentia, margine dentata, facie interna lanulata aut glabra, modo in eodem specimine aut ramo flores fem. (in singulis involucris 1-2) solum gerentia et tunc tubulosa perigoniumque *Boehmeriæ* referentia, modo majora floresque masc. et fem. (ut in genere solet) simul continentia. Flores masc. in involucri multifloro haud raro biserialiter interdumque cum flore fem. unico solitarii, perigonio apice anguste acuminato (acumine longiuscule ciliato) intus et inferne copiose lanato. Flores fem. ovario glabro, stylo filiformi villosulo. Achænia ovata ventricosa vel angulata, 1-1½ millim. longa, atra, nitidissima. — Planta fem. involucra 1-2-flora solum exhibens a me atque aliis ad *Boehmeriæ* olim relata fuit. (v. s.)

Suppetunt specimina alterius ut videtur speciei in Africa australi lectæ. Folia et habitus plantam abyssinicam referunt, sed involucrum unicum visum (caule divulsam) achænia lanata continebat.

### Subtrib. II. AUSTRALINEÆ. Flores exinvolucrati lanaque destituti.

XLI. AUSTRALINA Gaudich. Bot. Voy. Uran. p. 505; Hook. fl. Fl. Nov. Zel. p. 225; Wedd. Monogr. p. 543. *Urticæ* spec. auct:

Flores monoici in cymulas s. glomerulos paucifloros axillares exinvolucratos dispositi, cymulis fem. sessilibus, masculis (in distinctis axillis) pedunculatis bifloris. *Masc.* Perigonium monophyllum, infundibuliforme v. subcampanulatum, subhyalinum, limbo irregulariter bilabiato ciliatoque, labio exteriori s. longiore in alabastro inflexo. Stamen unicum. Pistilli rudimentum nullum. *Fem.* Perigonium ventricosotubulosum, liberum, limbo subquinqudentato. Ovarium rectum, ovato-lanceolatum. Ovulum e fundo loculi erectum, funiculo nullo v. brevissimo. Stylus linearis, cum apice ovarii continuus, altero latere præsertim villosulus. Achæniū ovatum, læviusculum, perigonio persistente inclusum. Embryo cotyledonibus late ellipticis radícula conica longioribus, albumine tenui. — Herbæ perennes in Australia, Tasmania, Nova Zelandia et Abyssinia vigentes, caulibus repentibus. Folia alterna aut rarius opposita, crenata v. crenato serrata, trinervia. Stipulæ laterales simulque petiolares, in specie oppositifolia inter petiolos connatæ. Cymulæ ad florem unicum haud raro reductæ. — Genus inter *Forskohleas* facile dignoscendum inflorescentiis unisexualibus et cymulis masc. constanter bifloris.

1. A. PUSILLA (Gaudich. Bot. Voy. Uran. p. 505), caulibus ramisque filiformibus repentibus, foliis alternis parvis rotundatis grosse crenatis longiuscule petio-

latis utrinque sparsim hispido-pilosis. ¼ In sylvis humidis Australiæ fœdici haud raro (F. Mueller); in sylvis obscuris Tasmaniæ et ins. sept. Novæ Zelandiæ (Gunn, Hook. fil., etc.). — Hook. fil. Fl. Tasman. 1. p. 345; Wedd. Monogr. p. 545. A. Novæ Zelandiæ Hook. fil. Fl. Nov. Zel. p. 225. *Urtica pusilla* Poir. Encycl. suppl. 4. p. 224. — Habitus fere Sibthorpiæ. Caulis numerosi, subintricati, radicales, herbacei aut demum nonnihil lignescentes, magis minusve pubescentes, 10-20 centim. longi. Folia 3-8 millim. longa et lata, obtusissima, basi cuneata aut truncata, in utroque margine bi-crenata, lamina exsiccata tenuiter membranacea sæpiusque saturate viridi supra cystolithis oblongis linearibusve conspicue ornatis, petiolo tenui limbi circiter longitudinis vel nonnihil brevioris sicut caulis pubescente. Stipulæ lineares v. subulatae, puberulae, parum conspicuae. Cymulae masc. in axillis superioribus obviae, ut in genere solet bifloræ, pedunculo (petiolum æquante) puberulo, floribus subsessilibus, perigonii fere campanulati hirtuli breviter et inæqualiter bilabiati labio exteriori s. majore late triangulari in alabastro pyriformi inflexo. Flores fem. solitarii aut sæpius in glomerulos 2-4-flosos digesti, sessiles subsessilesve, perigonio tubuloso demum valde compresso et circiter sesquimillimetrali elliptico aut rotundato apice in collum breve constricto marginibus acutis ciliato. Achæmium ellipticum, millimetrale, compressum, utrinque apiculatum, nitidulum, saturate fuscum. — Secundum cl. F. Mueller, hæc species statu majore sequentem includit. (v. s.)

2. A. MUELLERI (Wedd. Monogr. p. 545), caulibus gracilibus longe repentibus, ramis demum adscendentibus, foliis alternis rhombico-ovatis obtusis basi fere cuneatis integrisque cæterum grosse crenato-serratis utrinque sparsim pilosis. ¼ In ditone Victoria Australiæ, ad flumen Delatite (F. Mueller). — *Urtica tasmanica* Mueller, Syst. ind. pl. Vict. p. 18. — Rami subsimplices, 1-5 decim. longi, striati, villosi vel puberuli. Folia 2-4 centim. longa, 12-25 millim. lata sensim attenuata, in dimidio vel sæpius in triente inferiore cuneata aut rotundata, crenis utrinque 4-6 (inferioribus plerumque minoribus), lamina exsiccata membranacea sæpius obscure viridi pagina superiore cystolithis brevibus linearibus secus costam rarissimis ad marginem contra crebrisculis inspersa, petiolo gracili longitudine una quam limbus vulgo breviori pubescente. Stipulæ lanceolatae, inconspicuæ. Flores masc. cymulam bifloram efformantes, pedunculo filiformi petiolum æquante, perigonio sub anthesi bilabiato pilosulo, alabastro ovato obtusissimo. Flores fem. glomerulati, rarius solitarii, perigonio ovato ventricosulo villosulo breviter rostrato ore acute bidentato, stylo s. stigmatem ovario fere æquilongum. Achæmium ovatum, millimetrale, minute granulosum, obscure ferrugineum, pedicellatum, stylo persistente coronatum perigonioque marcescente arcte inclusum. (v. s. in lib. Kew.)
3. A. FLACCIDA (Wedd. Monogr. p. 546. t. 20, C), caulibus gracilibus elongatis repentibus, ramis adscendentibus, foliis oppositis ovato- vel elliptico-lanceolatis acuminatis grosse crenato-serratis brevissime petiolatis in apice ramorum subverticillatis utrinque sparsim pilosis. ¼ In locis umbrosis et in declivibus montium Abyssiniæ, altit. 2000 metr. (Schimper). — A. Schimperiana Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 212. *Pouzolzia flaccida* A. Rich. Tent. fl. Abyss. 2. p. 259. — Caulis longe lateque repentes, haud raro metrales, ut rami ipsi (1-2 decim. alti) hirtuli, internodiis inferioribus 2-4-centimetralibus, superioribus contra brevissimis. Folia magnitudine admodum variabilia (in apice ramorum sæpius majora), 1-3 centim. longa, 5-15 millim. lata, in triente v. quarta parte inferiore integra, superius utrinque crenis s. serraturis 1-5 obtusis munita, dente apicali s. acumine ovato obtuso, lamina exsiccata membranacea sæpiusque pallide viridi cystolithis punctiformibus inspersa, petiolo nullo aut 1-2-millimetrali. Stipulæ conspicuæ, ovatae, liberae vel sæpius inter petiolos connatae, scariosæ, longe ciliatae. Cymulae masc. (in axillis superioribus solum visæ) pedunculo quam petiolus duplo v. triplo longiore suffultæ, floribus sessilibus, perigonii circiter millimetralis tubo infundibuliformi, limbo inæqualiter bilabiato, labio majore fere triangulari-lanceolato acuminato minoreque late triangulari, utroque longiuscule ciliato. Glomeruli fem. subbiflori, floribus sessilibus, perigonio tubuloso demum ovato 2 millim. longo puberulo ore obtuse 4-5-dentato. Achæmium ovatum, fuscum, pericarpio tenui. (v. s.)

XLII. DIDYMODOXA E. Mey. in sched. pl. Dreg.; Wedd. Monogr. p. 547. — Australinæ spec. Wedd. in Ann. sc. nat. 4<sup>e</sup> sér. 1. p. 212.

Flores monoici, in glomerulos axillares pauci-multi-floros exinvolucratos dispositi, masculi et feminei in eodem glomerulo intermixti sessilesque. *Masc.* Perigonium monophyllum, fere bracteiforme, inferne breviter tubulosum, limbi cucullati acuminati marginibus ciliatis in alabastro lanceolato arcte conniventibus. Stamen unicum. Pistilli rudimentum nullum. *Fem.* Perigonium deficiens. Ovarium rectum, ovatum, apice in stigma subcapitatum aut breviter filiforme villosulum sæpeque incurvum desinens. Ovulum e fundo loculi erectum, funiculum destitutum. Achænium suboblique ovatum, compressum, margine altero crassiuscule carinatum. Embryo cotyledonibus rotundatis radicula tereti fere æquilongis. Albumen parvum. — Herbæ annuæ, capenses, diffuse ramosæ. Folia alterna, crenata aut integra, trinervia. Stipulæ petiolares, scariosæ, ciliatæ. Glomeruli sæpius 5-20-flori. Flores fem. interdum geminatim concreti. — Genus a præcedente distinctum glomerulis androgynis et floribus fem. nudis.

1. D. DEBILIS (E. Mey. in sched. pl. Dreg.), foliis late ovatis rotundatisve obtusissimis ima basi s. juxta petiolum sæpius breviter cuneatis grosse crenatis v. subintegris utrinque hispido-pilosis glabrativse longiuscule petiolatis, stipulis ovatis v. lanceolatis plerumque anguste acuminatis, floribus fem. haud raro concretis, stigmate brevissimo incurvo. ☉ In Africa australi (Drege). — Wedd. Monogr. p. 548. t. 20, B. Australina capensis Wedd. in Ann. sc. nat. 1. c. — Caulis gracilis, flaccidus, longe et diffuse ramosus, ramis superne haud infrequenter hirtulis. Folia magnitudine admodum variabili, majora 3 centim. circiter longa, 2 centim. lata, minora vix semicentimetralia, crenis s. serraturis plerumque obtusissimis nec raro omnino obsoletis, lamina exsiccata tenuiter membranacea pallide vel intense viridi cystolithis ut plurimum bene conspicuis, petiolo quam limbus longiore vel brevior glabrato vel hirtulo. Stipulæ nonnunquam sublobatæ, muticæ aut mucronato-acuminatæ. Glomeruli quinque-multi-flori, florum masc. perigonio (expanso) obovato elliptico 2 millim. circiter longo mucronato ciliato virente aut colerato. Achænia compressa, bimillimetralia, margine altero carinata (carina cum mucrone apicali arcuato seu stigmate persistente continua), alteroque cum margine achæni proximi a basi fere ad apicem sæpe connato. (v. s.)

β, lanceolata, foliis ovato-lanceolatis acutis crenato-serratis. ☉ In Africa australi (Drege). Parietaria lanceolata Thunb. Prodr. fl. cap. p. 31. ? Urtica ? radula E. Mey. l. c. — Folia quam in typo nonnihil majora, crenis nonnunquam subacutis. (v. s.)

2. D. INTEGRIFOLIA (Wedd. Monogr. p. 549), foliis late ovatis obtusis acutiusculisve basi subabrupte breviterque cuneatis integerrimis supra nervisque subtus sparsim hispidis, stipulis ovatis, floribus fem. discretis, stigmate filiformi dimidii ovarii circiter longitudinis, alabastris masc. breviter acuminatis. ☉ In Africa australi (Drege). — Parietaria lanceolata Thunb. ? E. Mey. in sched. pl. Dreg. Australina integrifolia Wedd. in Ann. sc. nat. 1. c. — Caulis elongatus, gracilis, flaccidus, adscendens v. subprostratus basi que sæpe nudatus, superne breviter ramosus subtilissimeque puberulus vel glabratus, internodiis 2-5-centimetralibus. Folia in speciminibus visis 5-15 millim. longa, 5-12 millim. lata, haud acuminata, obtusa vel acutiuscula, lamina exsiccata membranacea obscure viridi cystolithis vix conspicuis, petiolo dimidii limbi longitudinis vel brevior puberulo. Stipulæ interdum angulatæ v. sublobatæ. Glomeruli 5-10-flori, alabastris masc. cuneato-lanceolatis subæqualiter ciliatis. Achænia ovata, sesquimillimetralia, compressa, hispida. (v. s.)

3. D. CUNEATA, foliis rhombo-ovatis obtusisque vel sæpius (supremis præser-

tim) ovato-lanceolatis longiusculeque acuminatis (acumine obtusiusculo) basi cuneatis integrisque cæterum grosse crenatis vel crenato-serratis utrinque sparsim pilosulis, stipulis ovatis acuminatis floribus æquilongis, stigmatibus filiformi quam dimidium ovarium brevioribus, alabastris masc. breviter acuminatis. ○ In Africa australi (Drege). — *D. acuminata* Wedd. Monogr. p. 549. *Parietaria cuneata* E. Mey. in sched. pl. Dreg., non Eckl. et Zeyh. *Australina acuminata* Wedd. in Ann. sc. nat. l. c. — Caulis simplex vel a basi diffuse ramosus, 2-4 decim. altus, gracilis, striatus, puberulus. Folia 2½-6 centim. longa, 1-3 centim. lata, superne attenuata vel sensim acuminata, acumine ipso interdum centimetrali lineari integroque, crenis obtusissimis raro subnullis limbo tunc fere integro, lamina exsiccata tenuiter membranacea læte viridi supra cystolithis punctiformibus sparsis ornata, petiolo quam limbus brevioribus gracilibus magis minusve pilosulis. Flores fem. adpresse hispidi, stigmatibus vix semimillimetrali. Achænia 2 millim. circiter longa. (v. s.)

XLIII. DISTEMON *Wedd. Monogr. p. 550. — Urticæ spec. Wall.*

Flores monoici, in glomerulos paucifloros exinvolucratos laxè spicatos dispositi, floribus utriusque sexus sessilibus, femineis sæpe geminatis concretis. *Masc.* Perigonium subinfundibuliforme, bipartitum, segmentis æqualibus extus convexis breviter acuminatis integris. Stamina 2. Pistilli rudimentum lineare, longe lanatum. *Fem.* Perigonium tubuloso-ventricosum ovario adnatum, limbo minimo denticulato, ad maturitatem fructus carnosulum. Ovarium rectum, oblongo-lanceolatum. Ovulum funiculo brevissimo adfixum. Stylus linearis cum apice ovarii articulatus et ideo mox deciduus, altero latere a basi ad apicem stigmatoso-papillosus. Fructus ovatus, simplex aut sæpissime didymus, scilicet e floribus duobus concretis compactus et tunc figuram magis minusve irregularem exhibens, strato extimo (s. e perigonio confecto) carnosulo. Semen pericarpium nucamentacei cavitati conforme, testa tenuissima. Embryo cotyledonibus quadrato-rotundatis, radícula conica dimidio longioribus. Albumen satis copiosum. — Herba perennis, Indiam orientalem et Javam incolens. Caules erecti adscendentesque. Folia alterna, grosse serrata, trinervia. Stipulæ laterales, liberæ. Glomeruli subtriflori, in axilla bractearum ovatarum. — Genus a præcedentibus omnibus ejusdem tribus floribus masc. diandris et stigmatibus articulato distinctissimum.

*D. grossum*, foliis ovatis acuminatis grosse crenato-serratis utrinque sparsim pilosis subtusque glaucescentibus, spicis gracilibus terminalibus axillaribusque. ✕ In regionibus tropicis Indiæ orientalis (Wallich) et ins. Javæ (Zollinger, n. 2785). — *D. indicum* Wedd. Monogr. p. 551. t. 20, A; Miq. Fl. Ind. bat. 1 (2). p. 275. *Urtica grossa* Wall. Cat. n. 4615. — Caulis gracilis, 3-5 decim. altus, simplex v. parce ramosus, obtuse quadrangulus, striatus, inferne sæpius denudatus, haud raro lignescens glabratusque, superne interdum puberulus, siccitate lutescens, internodiis elongatis. Folia pauca, in medio caule majuscula s. 5-7 centim. longa et 3-4 centim. lata, ad basim spicæ autem multo minor, utrinque serraturis obtusis 3-7 ornata, ipsa basi sæpius integra et fere cuneata, dente apicali s. acumine cæteris multo majore triangulari, supra sparsim pilosa, nervis subtus etiam parce pilosis, intervenio subtiliter arachnoideo-tomentoso, lamina exsiccata membranacea fusciscentia subtus pallidior cinerascensque pagina superiore sub lente cystolithis punctiformibus creberrimis ornata, petiolo gracili 1-5 centim. longo adpresse puberulo aut glabrato. Stipulæ 3-5-millimetrales, e basi latiuscula subulatae, pilosulae. Spicæ simplices, solitariae, terminales aut axillares, priores cæteris majores, 2-10 centim. longi, rhachi filiformi angulata glabruscula. Glomeruli parvi, pauciflori discreti, bracteis late ovatis submillimetralibus acutis mucrona-

tisve integris vel basi auriculatis. Flores masc. minimi, breviter pedicellati. perigonii vix semimillimetratis lobis cucullatis summo dorso pilos paucos hamatos majusculos gerentibus cæterum fere glabris, pistilli rudimento (staminibus interposito) perigonii dimidiam longitudinem circiter æquante, lana quam hocce etiam longiore. Flores fem. primum bractea suffulciente multo breviores, sed demum illam superantes, perigonio strigosulo parallele longitudinaliter nervato, nervis demum obsoletis. Fructus regulariter vel irregulariter ovatus, apicem versus sensim attenuatus, unilocularis vel sæpe (præsertim in speciminibus javanicis) ob florum conjunctionem bilocularis, 2-3 millim. longus. (v. s.)

### URTICACEIS ADDENDA.

Pag. 51 δ. Adde : U. Kioviensis Schur, Enum. Transylv. p. 601, nec alior. ! nil est, teste cl. Kanitz, nisi subvarietas typi α.

α. Adde : Kanitz, Bot. Zeit. 1863 p. 54. — Dioica, caule inflorescentia petiolis et foliis parce setosis vel foliorum pagina superiore setis densis intertextis plus minus tomentosis imo cano-tomentosis, stimulis nullis vel paucissimis. (Kanitz.)

— 52. Post var. ξ Sicula, adde : In Styria inf. lecta est, ut videtur, forma hybrida U. dioicam inter et U. urentem, a cl. Koch (in litt. et in *Maly Fl. N. Steiermark*, p. 63) sub *U. oblongata* divulgata. Planta dicitur annua, et, monente cl. Kanitz (in litt.), qui typum in herb. Kochianum frustra quæsit, nil est nisi forma juvenilis U. dioicæ. (Confer ad *Flora*, ann. 1868, p. 183.)

— 57 δ. Secundum cl. Kanitz, synonym. U. atlantica Bl. ad U. urentem absque dubio pertinet.

— 59. U. fastigiata. Species hæc Blumeana, auctore cl. Kanitz, ad U. chamaedryoidem potius quam ad U. magellanicam referenda erit.

— — Inter species exclusas adde : URTICA NAUCLEIFLORA Griff. Not. 4. p. 385, quæ non est Urticacea, et U. RUBRICAULIS Span. quæ verisimiliter est species quædam Boehmeriæ.

— 140, adde : 91 bis. PILEA ULIGINOSA (Philippi, in Linn. 30. p. 199), caule humifuso radicante ramosissimo, foliis exacte ellipticis breviter petiolatis crenato-serratis ad summum 7 lin. longis, glomerulis pedunculatis flores fem. una cum masculis gerentibus, pedunculo folium subæquante. — Frequens in prov. Valdivia circa Puerto Monti in locis sylvaticis uliginosis. — Cum P. elliptica confusa. Folia minora breviter petiolata, petiolo ad summum tertiam folii partem æquante. Caulis longe magis ramosus. Flores fem. non ad basim pedunculi sessiles. Pedunculus denique brevior. (Phil.)

— 235<sup>1</sup>. Linea ante ultimam : Phenax.... lege Memorialis.

— 235<sup>2</sup>. Linea 1-9 : Phenax.... lege : Memorialis.

— 235<sup>3</sup>. M. QUINQUENRERVIS. ....foliis omnibus similibus, adde : oppositis. Adde postea : Ad hanc speciem forsitan adscribenda est Boehmeria melast-

moides Griff. Notul. 4. t. 63. f. 2. — Caulis basi decumbens et suffrutescens, pilis scabris lineatus. Folia (floribus exceptis) opposita, linearia vel lanceolato-linearia. Flores masc. 4-5-andri. Perigonium fructiferum 3-rarius 4-alatum, corollam papilionaceam nonnihil referens.

Pag. 235<sup>19</sup>. Linea 23 et 29, P. velutino.... lege P. incano.



## Ordo CLXXXVI. PIPERACEÆ (1).

*Urticis affinia* Juss. gen. p. 405. — *Urticeæ* § 2 *Piperiteæ* DC. *théor. élém.* (1813) p. 218. — *Piperaceæ* Kunth nov. gen. v. 1. p. 46 (1815); *Rich. obs. fr. transl. by Lindl.* p. 50 (1819); *E. Meyer de Houttuynia et Saurureis* p. 9; *Endl. gen.* p. 205; *Meissn. gen.* p. 335; *Kunth in Linn.* v. 13. p. 561; *Bl. monogr. der Oost. Ind. pep. in Verh. der Bat. Gen.* v. 11; *Miq. syst. Rip. in-8°* 2 fasc. (1843 et 1844), *Ill. in Nov. act.* v. 19 suppl., *Ann. bot. Mus. Lugd. Bat.* v. 1. fasc. 5; *Lindl. veg. Kingd.* p. 514; *Schnützl. iconogr. fam.* 81. t. 81; *Cas. DC. Mém. sur la fam. des Pip.* in *Mém. soc. phys. et hist. nat. Genev.* 18. pars 2; *Decaisne et Lemaout, traité de bot.* p. 502. — *Saurureæ* L. C. *Rich. anal. fr.* p. 41; *Meyer de Houtt.*; *Rich. dict. class.*; *Endlich. gen.*; *Meissn. gen.*; *Lindl. veg. Kingd.*; *Schnützl. icon.*; *Lemaout et Decaisne, traité de bot.* p. 500.

Flos hermaphroditus vel unisexualis, a bractea liber vel cum ea coalitus, sessilis vel pedicellatus vel rhachi impressus vel in ea profunde immersus. Bractea forma variabili, squamosa, sessilis vel pedicellata vel cum rhachi adnata et tum margine libera. Perigonium nullum. Stamina 1-10, rhachi vel basi imbricati vel et raro ovario inserta. Anthera vel cum filamentis articulata et caduca, vel non articulata et persistens, maturitate quadrivalvata (junior quadricellata) vel bivalvata (junior bilocellata), semper longitudinaliter dehiscentes. Carpella 1-4, monosperma vel polysperma, connata vel libera, sessilia vel stipitata vel rhachi impressa vel in ea profunde immersa et tum sæpe cum ea concreta, apice stylifera vel stylo destituta. Stigmata 1-4-plura. Ovula orthotropa. Bacca ovario incrassato formata, sessilis vel stipitata vel rhachi impressa vel in ea profunde immersa et tum sæpe cum ea concreta. Epicarpium membranaceum vel pulposum. Endocarpium durum. Albumen farinosum, aromaticum. Embryo minutus, intra sacculum persistentem vertice albuminis situs. — Herbæ vel frutices vel et arbores humiles terrarum omnium calidiorum. Systema fibro-vasculare modo duplex, id est fasciculis periphericis tubum periphericum lignosum formantibus et fasciculis in medulla sparsis sæpe tubum lignosum internum formantibus semper inter se et cum periphericis quoque nodo anastomosantibus; modo simplex, id est nunc tubo peripherico nunc fasciculis sparsis sistens. Vegetatio continua vel in singulo nodo interrupta. Folia monomera (sensu in opusc. meo *Théorie de la feuille*), simplicia, integerrima. Nervi primarii fasciculos

(1) Auctore Casimir de Candolle.

rami periphericos sine nodo continuantes (hic *nervi* dicti), secundarii, tertiarii, etc. inter se reticulatim anastomosantes (hic *nervuli* dicti), sæpe rari vel et deficientes; secundarii fere semper ante marginem inter se bifurcatim anastomosantes. Stipulæ petiolo sæpe ad limbum usque adnatæ, caducæ vel persistentes, sæpe rudimentariæ (Stipula oppositifolia auct. proprie rami sequentis prophyllus lateralis est). Petiolus stipulis ope vaginam formantibus gemmulam terminalem includens (*vaginans* hic dictus). Limbus, quando vegetatio interrupta, fere semper basi inæquilateralis, consistentia et crassitudine variabili, sæpissime succulentus. Inflorescentia amentacea. Amenta sparsa vel paniculatim spicativè apice ramorum conferta, aut (vegetatione quoque nodo interrupta), oppositifolia.

### CONSPECTUS TRIBUUM ET GENERUM.

Trib. I. *SAURURÆ*. — Systema fibro-vasculare simplex periphericum. Ovarium unicum, uniloculare, e pluribus foliis constans, multiovulatum, vel ovaria 3-4, bi-uni-ovulata. Anthera non articulata.

I. *Gymnotheca* Decaisne. Flos hermaphroditus. Stamina 6 (interdum 4-7), summo ovario inserta. Styli 4. Ovarium unicum. Placentæ parietales, multi-ovulatæ. Ovula cujusve placentæ biserialia.

II. *Anemiopsis* Hook. et Arn. Bractea libera. Flos hermaphroditus. Stamina 5-6. Ovarium unicum, rhachi profunde immersum et cum ea coalitum. Stigmata 3. Placentæ 3, parietales, cum stigmatibus alternæ. Ovula cuique placentæ 6-8, horizontalia, biserialia.

III. *Houttuynia* Thunb. Bractea libera. Flos hermaphroditus. Stamina 3, libera. Ovarium unicum, emersum, uniloculare, e 3-4 carpellis constans. Stigmata 3-4. Placentæ 3, raro 4, parietales, cum stigmatibus alternæ. Ovula cuique placentæ circ. 6, horizontalia, biserialia.

IV. *Saururus* L. Bractea cum flore concreta (raro libera). Flos hermaphroditus. Stamina 6, libera. Ovaria 3-4, libera, basi biovulata. Ovula sessilia, adscendentia.

V. *Saururopsis* Turcz. Bractea cum flore coalita. Flos hermaphroditus. Stamina 6. Ovaria 4.

Trib. II. *PIPERÆ*. — Systema fibro-vasculare duplex. Ovarium unicum, uniovulatum.

VI. *Piper* L. Anthera matura 4-valvata, utroque latere dehiscens.

VII. *Chavica* Miq. Anthera matura bivalvata, extrorsum dehiscens.

Trib. III. *PEPEROMIÆ* — Systema fibro-vasculare simplex sparsum. Ovarium unicum, uniloculare.

VIII. *Verhuellia* Miq. Bractea libera. Stamina 2. Anthera matura quadrivalvata, utroque latere dehiscens. Stigmata 4.

IX. *Peperomia* Ruiz et Pav. Bractea libera. Flos hermaphroditus. Stamina 2. Anthera matura bivalvata, extrorsum dehiscens. Stigma unicum.

Genera dubia : X. *DUGAGELIA* et XI. *Symbryon*.

Trib. I. *SAURUREÆ*. — Naiadearum et Typharum gen. *Juss. gen.* — Ordo Saurureæ *L. Cl. Rich.* (1808) *anal. fr.* p. 41.; *Endl. gen. p.* 266. — Ordo Saururaceæ *Lindl. nat. syst. ed. 2.* p. 184.

Systema fibro-vasculare simplex periphericum. Ovarium unicum, uniloculare, multicarpellare, multiovulatum, vel ovaria 3-4, biuni-ovulata. Anthera non articulata, matura 4-valvata.

I. *GYMNOTHECA* Decaisne in *Ann. sc. nat. ser. 3. v. 3. p.* 100. t. 5; *Decaisne et Le Maout Traité de bot. p.* 500; *Lindl. veg. Kingd. p.* 521 et 747.

Flores hermaphroditi. Stamina 6 (interd. 4-7) summo ovario inserta. Styli 4, recurvati, longitrorsum stigmatosi. Placentæ 4, parietales, multi-ovulatæ. Ovula orthotropa, horizontalia, cuique placentæ biseriata. — Herba perennis, erecta. (Ex Decaisne, l. c.)

G. *CHINENSIS* (Decaisne, l. c.), foliis petiolatis auriculato-cordatis apice obtusis vel breviter acuminatis utrinque glabris rigidulo-membranaceis subtilissime pellucido-punctulatis supra lævibus viridibus subtus pallidioribus reticulato-venosis 5-7-nerviis nervis centralibus 3 subtus prominentibus, petiolo glaberrimo basi vaginante, amentis oppositifoliis pedunculatis, pedunculo petiolum paullo superante amento limbum æquante, bractea ovato-rotunda carnosula, staminibus sæpius 6 summo ovario insertis, anthera oblonga filamentum superante. 4 In China (h. Mus. Par.). Herba glabra, erecta, rhizomatica. Limbi circ. 0,05 longi, 0,40 lati. Petioli 0,04 lati. (Ex Decaisne, l. c.)

II. *ANEMIOPSIS* Hook. et Arn. in *Beech. voy. p.* 390. t. 92. — *Anemia* Nutt. in *Ann. of Nat. Hist. v. 1. p.* 136. — *Anemiopsis* *Endl. gen. suppl. n.* 1825; *Meissn. gen.* 248; *Lindl. veg. King. p.* 521.

Bractea libera. Flos hermaphroditus. Stamina 5-6, libera. Ovarium unicum, rhachi profunde immersum, cum ea coalitum. Stigmata 3. Placentæ 3, parietales, cum stigmatibus alternæ. Ovula cuique placentæ 6-8, horizontalia, biserialia. — Herba americana.

A. *CALIFORNICA* (Hook et Arn. in *Beech. voy. p.* 390. t. 92), foliis alternis radicalibus longe petiolatis obovato-oblongis utrinque glabris et stomatiferris junioribus basi ciliolatis subcoriaceis opacis pellucido-punctulatis laxè venu-

lois penninerviis, centrali nervo validiore subtus prominente nervoq. subtiliores tota fere longitudine utrinque 4-5 subalternos subadscendentes ante marginem anastomosantes mittente, petiolo parce pilosulo basi vaginante angustè alato, amentis ramulos radicales diphyllis terminantibus quam folia multoties brevioribus, involucri 6-phylli phyllis obovato-oblongis glabris subpellucidis subtiliter pellucido-punctulatis, firmo-membranaceis, bractea spatulata apice obtusa glabra, staminibus 5-6, filamento antheram æquante, ovulis cuicque placentæ 6-8 horizontaliter insertis. ♀ In Nova California (Douglas!, in h. DC.), Zacatecas (Hartweg, n. 4, fide Hook. l. c.). — Nutt. in Ann. of nat. hist. v. 1. p. 136, Hook. in Bot. mag. t. 5292. Herba. Ramuli floriferi circ. 0,2 alti, parce pilosi. Limbi 0,065 longi, 0,025 lati; petioli 0,035 longi. (v. s.)

III. *HOULTUYNIA* Thunb. flor. Jap. p. 12 (non Houtt. Schreb. gen. p. 814; Juss. gen. p. 25; E. Meyer de Houtt. p. 1. fig. 1-4; Wall. flor. Ind. p. 360; Lindl. veg. Kingd. p. 521; Endl. gen. n. 1825; Meissn. gen. p. 248; Schnitzl. icon. t. 82. — Polypara Lour. flor. Coch. v. 1. p. 77.

Bractea libera. Flos hermaphroditus. Stamina 3, libera. Ovarium unicum, emersum. Stigmata 3-4. Placentæ 3 raro 4, parietales, cum stigmatibus alternantes. Ovula cuicque placentæ circiter 6, horizontalia, biserialia. — Herba asiatica, habitu Polygonaceæ; amentum involuero patente cinctum.

H. *CORDATA* (Thunb. flor. Jap. p. 234. t. 26), foliis alternis petiolatis subrotundato-ovato-acuminatis, acumine acuto subulatoque, basi æquali cordatis supra subtusque ad nervos puberulis membranaceis subopacis subtilissime pellucido-punctulatis septuplinerviis, nervis centralibus 3 paulo supra basin interdum fere e basi solutis crebre reticulato-nervulosis, petiolo juniore parce adulto basi densius hirtello basi vaginante, stipulis quam petiolus quadruplo brevioribus connatis a petiolo supra vaginam liberis glabris ciliolatis pellucidis, amentis oppositifoliis pedunculatis quam folia multoties brevioribus, pedunculo glabro petiolum subæquante, involucri 4-phylli phyllis obovatis pellucidis membranaceis glabris ciliolatis, rhachi glabra, bractea minuta lineari glabra, staminibus 3, filamento antheram multoties ovarium multum superante, stigmatibus ovarium æquantibus, ovario glabro ovato-attenuato uniloculari. ① In Japonia ubique in fossis viarum et juxta villas vulgatissima inter Miaco et Jedo (Thunb l. c.), Nagasaki (Oldham, n. 667! in h. DC.), Japonia (Zolling. n. 145! in h. DC.), China Fortune, n. 106! in h. DC.); in hortis Cochinchinæ inter olera, in acetariis edulis (Lour. flor. Coch. v. 1. p. 78, sub Polypara), Napalia (Wall. in h. DC.), montib. Khasia et Himalaya alt. 4000 ped. (Hook. et Thoms. in h. DC.) Polypara cochinchinensis Lour. flor. Coch. v. 1. p. 78, Poir. Encycl. v. 4. Houttuynia cordata Willd. sp. v. 2. p. 290, Poir. Encycl. v. 2. t. 739, Bot. mag. t. 2731, Schnitzl. icon. t. 82. Houtt. foetida Loud. hort. Herba erecta, ramulosa, basi radicans. Rami glabri, ad nodos hirtelli. Limbi circ. 0,06 longi, 0,05 lati. Petioli 0,02 longi. (v. s.)

IV. *SAURURUS* L. gen. n. 464; Schkuhr, Handb. t. 103; Mirb. Ann. Mus. v. 16. t. 19; Jacq. eclog. p. 29. t. 18; Meyer de Houtt. p. 13; Rich. dict. class. 15. p. 195; Endl. gen. n. 1824; Meissn. gen. alt. pars. p. 248; Lindl. veg. Kingd. p. 521, Donn Hort. Cant. ed. 1845. p. 241. cum icon.; Schnitzl. icon. t. 82 — Mattuschka Gmel. Syst. p. 589 (fd. auct.).

Bractea cum flore concreta (raro libera). Flos hermaphroditus. Stamina 6. Ovaria 3-4, libera, biovulata. Ovula basi ovarii sessilia, adscendentia. — Herbæ americanæ et asiaticæ.

1. *S. CERNUUS* (L. sp. ed. 2. p. 489), foliis alternis petiolatis ovato-acuminatis, acumine acuto, basi æquali cordatis supra subtusque ad nervos breviter hirtellis membranaceis opacis crebre pellucido-punctulatis septemnerviis, vel nervis mediis supra basin solutis, septuplinerviis, omnibus crebre reticulato-nervu-

lohis, petiolo juniore parce hirtello adulto glabro ad limbum usque vaginante angustè alato, ardentis oppositifoliis maturis folia multum superantibus breviter pedunculatis, pedunculo quam petiolus brevior glabro, rhachi pubescente, bracteis lanceolata dorso pubescente cum floribus pedicello adnata (in monstros? libera) intus glabra margine ciliolata, pedicello ciliolato, filamentis antheras multoties ovaria multum superantibus, ovariis glabris glandulosis, stigmatibus recurvis supra bilineatim puberulis, uno ovulo sterili 4. In America boreali a Louisiana ad Canadā in pratis humidis ad stagna et rivulos (Tainturier! Lindley! in h. DC. Engelmann! Eatoh! in h. Petrop., Pursh flor., Ell. sketch., Mich. flor.; p. 218, Nutt. gen. v. 1. p. 240, Torrey flor. Un. st. v. 1. p. 381, Hook. flor., Darlingt. Flor. cestr., Chapm. flor. s. st.), et in China (Hook. et Arn. in Beech. voy. p. 216). *Serpentaria repens* Pluk. Alm. p. 343, exclus. synonym., Phyt. t. 117. f. 3. 4. S. foliis profunde cordatis, etc, L. Hort. Cliff. p. 139; Gron. flor. Virg. p. 56. *Mattuschkia aquatica* Gmel. Syst. p. 589 (fide auct.). S. cernuus L. l. c., Poir. Encycl. p. 668; Lam. ill. t. 276, Schk. Manih. t. 108, Diet. sc. nat. t. 295, Schinzl. icon. t. 82. S. lucidus Donn, Hort. Cant. ed. 1804. p. 66, Jacq. eclog. p. 29. t. 18. Herba erecta, ramulosa, basi radicans. Rami ramulique sparsim pilosuli, dein glabrati. Limbi efformati 0,09-0,13 longi, 0,07-1 lat! Petioli circ. 0,02 longi. Bacca circ. 0,001 longa. (v. s.)

2. S. LOURERII (Decaisne in Ann. sc. nat. ser. 3. v. 3. p. 102), caule valde angulato, spica folio æquilonga, staminibus ovaria subæquantibus filamentis brevissimis. In locis paludosis prope Cantonem Sinarum (Lour.), in Japonia (Thunb. Siebold in h. Mus. Par., Lugd. Bat., Deless.). *Spathium chinense* Lour. Coch. p. 217, Jussieu Ann. sc. nat. 1844. p. 107 notula. *Saururus cernuus* Thunb. flor. p. 154. (ex Decaisne, l. c.)

### Species exclusæ.

De Saururis in Plumiero sub phras. enūm. confer ad genera Piper et Peperomia.

V. SAURUROPSIS Turcz. in Bull. soc. mosc. v. 21.

Bractea cum flore coalita. Flos hermaphroditus. Stamina 6. Ovaria 4. — Frutices e China, Japonia et Philippinis.

\* Squamule hypogynæ duæ oppositæ stamina includentes (verisimiliter stamina exteriora abortiva).

1. S. CHINENSIS (Turcz. l. c.), foliis breviter petiolatis cordato-oblongis inæquilateris basi in petiolum brevem attenuatis apice sensim acuminatis quintuplinerviis, nervis in petiolum decurrentibus, amentis oppositifoliis, bracteis in pedicellum florigerum contracta, squamulis hypogynis oblongis interne concaviusculis, ovariis ovato-compressis utrinque marginatis apice in stylum incurvatum stigmatiferum attenuatis. 3 In China boreali (Fortunei n. 102, A, fide Turcz. l. c. et h. DC.). Herba glaberrima. Caulis simplex aut ramis unico auctus, angulatus. Petioli 2½-3 lin. longi. (ex Turcz. l. c.)

\*\* Squamule hypogynæ nullæ.

2. S. CUMINGII, foliis petiolatis ovatis apice attenuatis acutiusculis basi truncatis obtusis utrinque glabris coriaceis reticulato-nervulosis et quinque-septem-nerviis, nervis subtus prominulis, nervo centrali ad apicem usque ducto, lateralibus 2 ad apicem usque, 2 supra medium ductis externis subtilioribus, petiolo glabro anguste alato, amento oppositifolio, pedunculo pilosulo petiolum duplo superante, rhachi ochraceo-villosa, floribus oppositis, bractea spatulata apice obtusa basi in pedicellum florem æquantem ochraceo-villosam contracta, stam. 6, ovariis 4, apice in stylum incurvatum stigmatiferum attenuatis, stylo filiformi ovarium fere æquante. 3 In ins. Philippinis (Cuming n. 1259! in h. Brit. Mus.). Ramuli glabri subtrianguli, nodi

tumiduli. Limbi 0,09 longi, 0,055 lati. Petioli 0,015 longi. Pedunculi 0,03 longi. Amenta florentia 0,07 longa. (v. s.)

β, *Japonica*, foliis membranaceis basi magis cordatis. ♂ In Japonia ad Kanagavia (Fortune! in h. Brit. Mus.). Nagasaki (Oldham n. 668! in h. DC.). (v. s.)

Trib. II *PIPERÆ* Miq. *syst. Piper.* p. 45. — Systema fibrovasculare duplex. Ovarium unicum, uniovulatum. Anthera articulata vel non articulata.

VI. *PIPER* L. sp. pro parte (non L. gen., propter char.); Jacq. ic. rar. et coll. et Jacq. f. eclogæ (pro parte); Lam. ill. (pro parte); Gœrnter carp. v. 2. p. 67. t. 92; Willd. sp. (pro parte); Ruiz et Pav. flor. peruv. v. 1. p. 33 (mtas. char.); Vahl. enum. v. 1 (pro parte); Kunth. in H. et B. nov. gen. v. 1. Rasm. et Sch. syst. (pro parte); Spreng. syst. (pro parte); Opiz. Reliq. Hænk. fasc. 3; Bl. monogr. der oost Ind. Pepers in Verh. der Bat. Genoots. v. 11 et Enum. pl. Jav. (pro parte); Loud. Encycl. (pro parte); A. Dietr. sp.; Endlich. gen. n. 1820 (pro parte); Miq. syst. Pip. p. 305. — Zippelia Bl. fide Rœm. et Sch. syst. — Ottonia Spreng. neu. Entd. — Heckeria, Enckea, Steffensia, Schilleria, Ottonia Kunth. in Linn. v. 13. — Coccobryon, Caulobryon Klotzsch msc. in h. Berol. — Macropiper, Protomorphe, Cubeba, Muldera Miq. comm. phyt. — Carpunya Presl Epim. bot. — Schizonephos Griffith, notul. — Protomorphe, Macropiper, Rhyncholepsis, Muldera, Cubeba, Piper, Coccobryon, Callianira, Enckea, Peltobryon, Sphærostachys, Artantha, Ottonia, Zippelia, et pars Chavica Miq. syst. Piper. — Piper subgenus Eupiper Cas. DC. in Seem. journ. of bot. 1866. p. 161.

Bractea cum flore coalita vel libera. Flos unisexualis vel hermaphroditus. Stamina 1-10. Anthera matura quadrivalvata. Ovarium unicum. Stigmata 2-4-plura. — Frutices vel et arbores totius orbis calidioris.

#### Conspectus sectionum.

Sect. 1. *Schizonephos*. Flos e cupula, verisimiliter bracteis conatis formata, stamina vel ovarium includente, sistens. Amenta oppositifolia. Antheræ articulatae. — Species 1-9.

Sect. 2. *Enckea*. Bractea libera. Stamina 5-6, ovario circumposita. Amenta oppositifolia. Flores hermaphroditi. Antheræ articulatae. — Spec. 10-39.

Sect. 3. *Steffensia*. Bractea libera. Stamina 4, ovario circumposita. Amenta oppositifolia. Flores hermaphroditi vel unisexuales. Antheræ articulatae vel non articulatae. — Spec. 40-355.

Sect. 4. *Carpunya*. Bractea libera. Stamina 3, quorum duo lateralìa, unum posticum. Amenta oppositifolia. Flores hermaphroditi. Antheræ articulatae vel non articulatae. — Spec. 356-377.

Sect. 5. *Potomorphe*. Bractea libera. Stamina 2, lateralìa, vel 3, quorum unum posticum. Amenta axillaria vel apice ramuli axillaris umbellata. Flores hermaphroditi vel unisexuales. Antheræ articulatae. — Spec. 378-390.

Sect. 6. *Coccobryon*. Bractea libera. Stamina 2, lateralìa, vel 3, quorum unum posticum. Amenta oppositifolia. Flores hermaphroditi. Antheræ articulatae. — Spec. 391-405.

Sect. 7. *Eupiper*. Bractea libera. Stamina 2, lateralìa, vel 3, quorum unum posticum, vel rarissime 4, uno antico adjuncto. Amenta oppositifolia. Flores unisexuales vel basi amenti interdum hermaphroditi vel in rarissimis speciebus trioici, modo unisexuales modo hermaphroditi. Antheræ articulatae.

Sect. 8. *Apopiper*. Bractea libera. Stamina 2. Amenta oppositifolia. Antheræ non articulatae.

Sect. 9. *Nematanthera*. Bractea libera. Stamen unicum. Amenta oppositifolia. Flores hermaphroditi vel unisexuales. Plantæ monoicæ. Antheræ articulatae.

Sect. 1. SCHIZONEPHOS. — Genus *Schizonephos* Griff. not. 4. p. 383. — Gen. Muldera *Miq. comm. phyt.* p. 34. — Genus *Piper* Bl. in *Verh. der bat. genoots. t. 2. p. 172 (pro parte)*.

Flores e cupula verisimiliter bracteis connatis formata, stamina vel ovarium includente sistentes. Amenta oppositifolia. Antheræ articulatae.

#### α. Flores stipitati.

1. P. SCHIZONEPHOS, foliis petiolatis lanceolatis oblongo-lanceolatisve utrinque acutis et glabris rigide coriaceis 5-nerviis nervo centrali supra basin ad  $\frac{1}{2}$  alt. trifido nervulis subtransversim parallelis, petiolo glabro, pedunculo glabro petiolum paullo superante, stirpis masc. amento laxifloro, staminibus 8 biseriatim dispositis, cupula basi in stipitem cupulam æquantem contracta longitudinaliter dehiscente, stirpis fem. foliis conformibus bacca globosa crassit. *Piperis nigri*. ♀ In peninsula birmanica (Griffith n. 4400! in h. Kew. et Vindob.), et Java (h. Kew! ubi specim. fem. a Jungh. lect.). *Schizonephos glaucescens* Griff. ! l. c. Dioica. Ramuli glabri. Limbi 0,11 longi, 0,45 lati. Petioli 0,015 longi. Amenta matura 0,15 longa. (v. s.)

#### β. Flores sessiles.

2. P. RECURVUM (Bl. in *Verh. bat. Genoots. p. 176. t. 4*), foliis petiolatis inferioribus subrotundo-ovatis apice acute acuminatis basi æquali rotundatis superioribus elliptico-oblongis suboblique lanceolatis apice acute acuminatis, omnibus utrinque glabris coriaceis opacis septupli-quintupli-nerviis, centrali nervo ad  $\frac{1}{2}$  longitudinis usque nervos alternos adscendentes utrinque 2 et e basi utrinque unum supremos ante apicem evanidos mittente, petiolo glabro ad limbi basin usque vaginante, pedunculo glabro petiolum æquante, amento florente quam folium triplo brevior, stirpis masc. floribus remotis, stam. in quoque flore 10 biseriatim dispositis e cupula emergentibus, cupula intus pilosa, stirpis fem. rhachi glabra, cupula intus glabra, ovario glabro, stigmatibus plerumque 4. ♂ In sylvis humidis montium insulæ Java (h. DC. ! ex h. Lugd. bat., pl. fem.), sylvis montis Teapi ins. Java ad arbores scandens (Miq. pl. Jungh. fasc. 1. p. 18). — Bl. enum. pl. Jav. fasc. 1. p. 67. Muldera recurva *Miq. ! comm. phyt. p. 35, Syst. p. 342, illustr. p. 58. t. 60, flor. Ind. bat. v. 1. pars 2. p. 456*. Dioica, scandens. Truncus brachium haud raro crassior, lævis (Miq. syst.). Rami glabri. Limbi 0,08-0,12 longi, 0,25-0,055 lati. Petioli inferiores 0,015, superiores 0,01 longi. (v. s.)

3. P. BACCATUM (Bl. in *Verh. der bat. genoots. fasc. 11. p. 172. t. 3*), foliis petiolatis inferioribus subrotundo-ovatis apice acute acuminatis basi æquali rotundatis superioribus oblongo-ellipticis lanceolatisve utrinque glabris apice acute acuminatis basi subinæquali attenuatis acutis utrinque glabris coriaceis opacis septupli-quintuplinerviis, nervis supra subtusque subprominulis, centrali ad  $\frac{1}{2}$  longitud. usque nervum utrinque unum vel utrinque 2 alternos adscendentes et e basi sæpe utrinque unum mittente, petiolo glabro, pedunculo glabro petiolum æquante, stirpis fem. amentis multifloris sublaxifloris,

- cupula intus pilis albis pilosula, bacca globosa transversim dehiscente, stirpis masc. cupula ut in femina, staminibus 5 uniserialibus. ♂ In ins. Java (Zoll. n. 1525! in h. DC. stirpis fem. et h. DC. ex h. Lugd. bat. stirpis fem.), in Sadjira et Tjikandi dit. Bantam Javæ (Miq. in Linn. v. 21. p. 486), Bornœo (Korthals fide Miq. in Mus. Lugd. bat. v. 1. fasc. 5. p. 137). — Bl. enum. pl. Jav. fasc. 1. p. 66; Schnizl. iconog. t. 81. fig. 8-10. Muldera baccata Miq. comm. phyt. p. 34, Syst. p. 341, illustr. p. 58. t. 59, flor. ind. bat. v. 1. pars 2. p. 456, in Linn. v. 21. p. 486. Muld. recurva Zoll. cat. p. 84. Muld. Wightiana Miq. in Hook. Lond. Journ. bot. v. 5. p. 557; Wight icon. t. 1943. f. sinist. Dioica scandens. Ramuli glabri. Limbi 0,12 longi, 0,08 lati. Petioli 0,015 longi. Bacca 0,004 crassa. (v. s.)
4. *P. TRICHOSTACHYON*, foliis breviter petiolatis lanceolatis vel oblongo-lanceolatis apice acutis basi subattenuatis acutiusculis utrinque glabris subcoriaceis triplinerviis vel subtrinerviis, nervo centrali paulo supra basin trifido nervulis transversim parallelis, petiolo glabro, pedunculo glabro petiolum æquante, stirpis masc. cupula extus dense velutino-puberula intus ad basin staminum hirsuta, stam. 2. ♂ Prope urbem Mangalor in campis aridis et hortis terræ Canaræ (Hohenacker, n. 396! in h. Kew. et Boiss.), Malabar (Wight, fide Miq. in Hook. Lond. Journ. bot. v. 5. p. 556). Muldera trichostachya Miq. l. c.; Wight icon. t. 1944. Dioica, scandens. Ramuli glabri. Limbi 0,07-0,08 longi, 0,02-0,025 lati. Petioli 0,005 longi. Stirpis fem. ♀ baccæ polygono-globosæ 0,005 crassæ (Stocks! in h. Kew.). (v. s.)
5. *P. GALEATUM*, foliis petiolatis ovatis vel ovato-ellipticis oblique acuminatis, acumine acuto, basi subæquali rotundatis vel leviter emarginatis membranaceo-coriaceis pellucido-punctatis septuplinerviis, nervis 3 mediis ad apicem continuatis subtus rubentibus, amentis masc. longe pedunculatis filiformibus elongatis dissitifloris folium superantibus, cupulis reflexis clavato-galeatis transversim dehiscentibus intus hirtellis. ♂ In Courtallum (Wight, fide Miq. in Hook. Lond. Journ. bot. v. 5. p. 557). Muldera galeata Miq. l. c. et in Wight icon. t. 1943, fig. dextr. Dioica, scandens. Petioli 0,01-0,015 longi. Limbi 0,07-0,1 longi, 0,04-0,05 lati (ex Miq. l. c.).
6. *P. FIRNUM*, foliis breviuscule petiolatis superioribus e basi æquali vel leviter inæquali rotundata ovato-lanceolatis suboblongisve, superioribus angustioribus lanceolatis omnibus apice subacuminato-acutis crasse coriaceis sextupli-subquintupli-nerviis nervis tenuibus supremis fere ad apicem ductis, amento foemineo maturo longe pedunculato circ. 10-12-floro, pedunculo petiolum superante amentum subæquante, bacca globosa cupula orbiculari-explanata intus appresse pilosa. ♂ In Sumatra occidentali (Miq. in Ann. bot. Mus. Lugd. bat. v. 1. fasc. 5. p. 140). Muldera firma Miq. l. c. Dioica. Folia  $3\frac{1}{2}$ -2 poll. longa, 10 vulgo 8-4 lin. lata. Petioli semipollicares, sæpe breviores. Pedunculi tenues 6 lin. longi. Amenta baccifera fere pollicem longa. Bacca fusca vel nigro-rugulosa (ex Miq. l. c.).
7. *P. MALABARENSE*, foliis petiolatis superioribus ovatis vel elliptico-ovatis breviter acuminatis, acumine acuto, subæquilateralis utrinque glabris membranaceo-coriaceis pellucido-punctatis basi subæquali rotundatis septupli-vel subnovempli-nerviis, nervo centrali ad apicem usque lateralibus 2 fere ad apicem ductis, amento foemineo breviter pedunculato quam folium duplo brevior glabro subdensifloro, cupula transverse dehiscente, stigmatibus 3. ♂ In Malabar (Wight, fide Miq. in Hook. Lond. Journ. bot. v. 5, p. 558). Muldera multinervis Miq. l. c. Dioica. Limbi 0,12-0,14 longi, 0,005-0,07 lati. Petioli 0,01-0,02 longi. (ex Miq. l. c.)
8. *P. CEYLANICUM*, foliis breviter petiolatis oblongo-lanceolatis apice acuminatis, acumine acuto, utrinque glabris subcoriaceis opacis trinerviis, nervis subtus prominulis, centrali ad apicem ducto supra basin utrinque alternatim nervum fere ad apicem usque ductum mittente, lateralibus subtilioribus supra medium ductis, petiolo glabro, pedunculo glabro petiolum æquante, amento quam folium brevior subaxillifero. cupula glabra, bacca subglobosa. stigmatibus 3-4. ♂ In ins. Ceylon (Thwaites, n. 21751 in h. DC.). Dioica. Ramuli glabri. Limbi 0,08 longi, 0,025 lati. Petioli 0,005 longi. Bacca fere 0,005 crassa. (v. s.)



9. *P. UNGARAMENSE*, foliis petiolatis rotundato-ovatis apice breviter acuminatis, acumine acuto, subæquilateris basi cordatis utrinque glabris coriaceis novem-nerviis, nervis præter tres medios e basi ortis, amentis brevibus longiuscule pedunculatis nascentibus tenere puberulis, cupulæ labiis ore puberis intus pilosis, stigmatibus 4. ♀ In Javæ monte Ungaram alt. 3-4000 ped. (Miq. pl. Jungh. fasc. 1. p. 17), monte Salak (Miq. in Ann. Mus. Lugd. bat. v. 1. fasc. 5. p. 137), Sumatra occ. prope Lubu Sampit (Miq. in Sumatra, p. 186 et 473). Muldera cordata Miq. l. c. Dioica. Folia 8-5 poll. longa, 7-5 lata. Petioli 1-2 et superiores  $\frac{1}{2}$  poll. longi. Pedunculi 6-8 lin. longi. Amenta florentia  $1\frac{1}{2}$  poll. longa (ex Miq. l. c.).

Sect. II. ENCKEA. — Genus Enckea *Kunth in Linn. v. 13*, p. 590. — Gen. Callianira *Miq. Syst. p. 344*.

Bractea a flore distincta. Stamina 5-6, ovario circumposita. Amenta oppositifolia. Flos hermaphroditus. Antherae articulatae.

α. Flores stipitati.

10. *P. MUELLERI*, foliis breviter petiolatis ovato-acuminatis, acumine longiusculo acuto, basi æquali rotundatis supra glabris subtus ad nervos adpresse puberulis rigidis subopacis pellucido-punctulatis 7-nerviis, nervis subtus prominulis 3 centralibus ad apicem ductis, nervulis subtransversim parallelis, petiolo basi ima vaginante utrinque hirtello, pedunculo hirtello petiolum duplo superante, rhachi hirtella, amento folium æquante, bractea lanceolata subsessili dorso hirtella intus glabra, stipite hirtello florem æquante, stam. 6, antheris subglobosis connectivo supra loculos paulo producto, ovario ovato-oblongo glabro, stigmatibus 3 brevibus. ♀ In Orizaba (Ferd. Mueller, n. 180! in h. Mart.). Rami glabri. Ramuli juniores dense hirtelli. Limbi 0,125 longi, 0,045 lati. Petioli 0,01 longi. (v. s.)

β. Flores sessiles.

1° Folia penninervia.

11. *P. FLONGATUM*, foliis breviter petiolatis oblongis apice acuminatis, acumine acuto, basi leviter inæquali obtusis rotundatisve supra glabris subtus ad nervos parce hirtellis siccis rigidis opacis pellucido-punctulatis utrinque nitidulis supra areolatis subtus lacunosis albido-punctulatis, nervis subtus prominentibus, centrali ad apicem ducto nervos utrinque fere tota longitudine 11 nervulis fortioribus intermixtos alternos subadscendentes adscendentesve mittente, petiolo glabro ad limbum vaginante dorso albido-punctulato, alis cito deciduis, pedunculo petiolum subæquante glabro, amento quam folium multoties brevius, bractea ovarii proximæ cucullatæ glabræ vertice inflexo triangulari glanduloso-punctato, staminibus 6, ovario glabro, stigmatibus 3 abbreviatis. ♀ In Brasilia (Riedel n. 581! in h. Petrop.), Ramuli glabri albido-punctulati. Limbi 0,17 et inferiores 0,20 longi, 0,04-0,05 lati, petioli 0,005 et inferiores 0,015 longi. Amenta post anthesin 0,015 longa, 0,003 crassa (v. s.)
12. *P. HOHENACKERI*, foliis breviter petiolatis elliptico-oblongis apice breviter acuminatis, acumine acuto, basi æquali acutis utrinque glabris siccis membranaceis subopacis pellucido-punctulatis, nervis subtus subprominulis, centrali ad apicem ducto utrinque fere tota longitudine nervos circ. 13 alternos adscendentim patulos nervulis fortioribus sæpe intermixtos supremos ante apicem per anastomoses evanidos mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum æquante, amento quam folium multoties brevius cylindrico, rhachi foveolata glabra, bractea ovarii proximæ cucullatæ glabræ dorso carinata vertice inflexo angusto triangulari, stam. 5-6, bacca obovato-subquadrangula glabra basi subimmersa, pericarpio vertice pulposo, stigmatibus 3. ♀ In sylvis prope urbem Paramaribo (Kappler, n. 1668! in h. Boiss.). Ramuli glabri. Limbi 0,15 longi, 0,06 lati, petioli 0,006 longi. Amenta mat. 0,005 crassa. (v. s.)

2<sup>o</sup> Folia multinervia (1).

\* Nervuli primarii conspicue transverse paralleli.

13. *P. LÆVIGATUM* (Kunth in H. et B. nov. gen. v. 1. p. 56), foliis petiolatis apice longe acuminatis obtusiusculis infer. ovato-oblongis basi æquali rotundatis superioribus oblongo-lanceolatis basi æquali acutiusculis omnibus utrinque glabris rigidulo-membranaceis subpellucidis, foliis inf. 7- sup. 9-nerviis. nervis subtus prominulis 3 centralibus ad apicem ductis proximis fere marginalibus alte ductis infimis subtilioribus, petiolo glabro, pedunculo glabro petiolum subæquante, amento quam folium brevius, bractea sessili e basi cuneata spatulata vertice inflexa ad basin floris sita, staminibus 4-5, bacca emersa ovalo-elliptica apice in stylum brevem attenuata glandulosa, stigmatibus 3-5 sessilibus abbreviatis. ♀ In ripa flumin. Magdalene pr. San Bertholome et m. Barbacoa (H. B. l. c. et h. Willd. n. 690! ubi *P. attenuatum* ined. dicitur). *P. lævigatum* Kunth Syn. v. 1, p. 111. *Enckea lævigata* Kunth! in Linnæa v. 13. p. 602, exclus. num. 691 herb. Willd., Miq. Syst. p. 348. Rami et ramuli teretes, glabri, nodi tumidi, limbi infer. 0,2-0,22 longi, 0,09 lati, sup. 0,12-0,18 longi, 0,035-0,06 lati, petioli inf. 0,025, sup. 0,01 longi. (v. s.)
14. *P. TENUE* (Kunth! in H. et B. nov. gen. v. 1. p. 56), foliis petiolatis superioribus oblongo-lanceolatis apice longe subfalcatis acuminatis basi æquali attenuatis acutis utrinque glabris-rigido-membranaceis subpellucidis 5-nerviis, nervis utrinque prominulis 3 centralibus ad apicem ductis proximis subtilioribus marginalibus, petiolo glabro, amento filiformi subdensifloro, pedunculo petiolum subæquante glabro, bractea glabra sessili e basi cuneata spatulata apice conchæformi obtuso inflexo infra florem inserta, stam. 4-5 brevissimis, ovario ovato, stigmatibus 4 sessilibus, bacca glabra ovoidea emersa apice attenuata. ♀ In Peruvia (Poeppig, n. 311! in h. Cand. et Deless. et Mathews, n. 1708! in h. Kew.), Cuchero (Endl. in herb. Vind.), ripa fluminis Magdalene regione calidissima inter Penon et Cano de Batales (H. et B. l. c. et herb. Willd. n. 689! spec. Humb. sub nom. *P. leptostachyum* et *P. tenue*). — Kunth Syn. v. 1. p. 111. *Enckea tenuis* Kunth! in Linnæa, v. 13. p. 603. Miq.! Syst. p. 367. *Enckea Lessertiana* Miq.! Syst. p. 349. *Enck. lævigata* Miq. in Hook. Lond. journ. bot. v. 5. p. 440. *P. leptostachyum* Griseb. cat. pl. cub. p. 65. Ramuli angulato-sulcati, glabri. Limbi 0,17 circ. longi, 0,05 lati, petioli 0,015 longi. Pedunculi 0,01 longi. Amenta matura 0,06-0,07 longa. (v. s.)
15. *P. ENCKEA*, foliis brevissime petiolatis inferioribus ovato-oblongis super. lanceolato-oblongis apice longe acuminatis, acumine acutiusculo, utrinque glabris rigidulo-membranaceis subpellucidis subepunctulatis 5-nerviis, nervis utrinque prominulis 3 centralibus ad apicem et proximis vix ad medium ductis, petiolo juniore tenerrime puberulo limbum usque vaginante, amento quam folium paulo brevius maturo subdensifloro, rhachi tenere puberula, bractea sessili spatulata ad florem inserta extus pilosula, staminibus plerumque 6, bacca emersa ovato-subquadrangula puberula. ♀ In sylvis prope Rio Japure prov. Rio Negro (Mart. in h. reg. monac.). *Enckea Martiana* Miq.! in Linnæa, v. 20. p. 132 et in Mart. flor. bras. fasc. 11. p. 27. t. 3. f. 3. (*P. Martianum* adest, quod olim Schilleria Martiana). Ramuli juniores tenerrime puberuli. Limbi 0,12 longi, 0,047 lati, petioli 0,005 longi. (v. s.)
16. *P. HEPTANDRUM*, foliis breviter petiolatis oblongo-ovatis apice acuminatis acumine acuto basi æquali subattenuatis acutiusculis utrinque glabris, siccis firmis subopacis pellucido-punctulatis, 7-nerviis nervis utrinque prominulis 3 centralibus ad apicem proximis marginalibus ad medium ductis, petiolo glabro basi sub 1/2 longit. vaginante, pedunculo petiolum subæquante glabro, bractea glabra sessili concavo-oblonga apice obtusa, paulum infra florem inserta, staminibus 5 connectivo sub loculis cuneatis elongato quam filamentum longiori, ovario emerso glabro, stigmatibus 4 sessilibus. ♀ In Pe-

(1) Folium *multinervium* dico quando nervi omnes e basi ima limbi discedunt; *multiplinervium* quando partim e basi limbi, partim supra e nervo centrali abeunt.

ruvia (Pav. in h. Boiss. et h. reg. ber. ubi *P. heptandrum* dictum), Tocache Hualloga (Endlich. ! in herb. Vindob.). *Enckea heptandra* Miq. ! Syst. p. 350, ill. p. 63. t. 64. Ramuli glabri. Limbi 0,18 longi, 0,08 lati, petioli 0,01 longi. Amenta flor. 0,06 longa. (v. s.)

17. *P. SMILACIFOLIUM* (Kunth! in H. B. nov. gen. v. 1. p. 56), foliis breviter petiolatis amplis ovatis apice breviter acuminatis acumine obtusiusculo basi æquali foliorum infer. rotundato-cordata sup. obtusa acutiusculave utrinque glabris siccis rigidis epunctulatis 7-nerviis, nervis subtus prominulis 3 centralibus ad apicem proximis eorum alte ductis infimis subtilioribus, petiolo glabro ad  $\frac{1}{2}$  longit. vaginante alis cito deciduis, pedunculo glabro petiolum paulum superante, bractea glabra apice inflexo-dilatata, stam. 4, bacca emersa subglobosa subquadrangula glabra, stigmatibus 3 sessilibus. ♂ In sylvis prope urbem San Juan et Panama (Seemann, n. 418! in h. Kew. et 900 fide Miq. ! in Seem. bot. Herald.), in sylvis humidis Mirador et Zacuapan alt. 3000 ped. (Galeotti, n. 6010 fide Mart. et Gal. bull. ac. Brux. v. 10. n. 2. p. 22), locis umbrosis prope Caripe alt. 470 hexap. (h. Willd. n. 691. fol. 1 et 2!), in ins. Trinitatis (Schach! in h. Kew.) et Guadaloupa (h. Rich. ! in h. Francovil.). *Enckea smilacifolia* Kunth! in Linn. v. 13. p. 60 s., Miq. syst. p. 351. Ramuli glabri, limbi 0,14-0,19 longi, 0,11-0,16 lati, petioli 0,015 longi. Amenta mat. 0,11 longa. (v. s.)
18. *P. LATUM* (Kunth! in H. B. nov. gen. v. 1. p. 57), foliis petiolatis subrotundato-ovatis apice longiuscule acuminatis acumine acutiusculo basi æquali truncato-rotundatis utrinque glabris siccis rigido-membranaceis subpellucidis parce pellucido-punctulatis 9-nerviis, nervis subtus prominentibus 5 centralibus per anastomoses fere ad apicem ductis proximis fere marginalibus alte ductis infimis sub medio evanidis, petiolo glabro. ♂ In ripa flumin. Magdalenæ, juxta Guarumo (H. B. ! in nov. gen. l. c. et h. Willd. n. 692! spec. Humb. sub nom. *P. latum*). *P. latum* Kunth syn. v. 1. p. 112. Opiz reliq. Hænk. fasc. 3. p. 159. *Enckea lata* Kunth! in Linn. v. 13. p. 606, Miq. Syst. p. 351. Macropiper latum Presl in Epim. p. 223. Ramuli glabri, limbi 0,19 longi, 0,175 lati, petioli 0,025 longi. (v. s.)
19. *P. MELASTOMOIDES* (Schlecht. et Cham. in Linn. v. 5. p. 74), foliis petiolatis inferioribus oblongo-ovatis basi æquali rotundatis, superioribus lanceolato-ellipticis basi subinæquali acutis, omnibus apice longe lineari-acuminatis, acumine acuto, supra sparse pilosis subtus præsertim ad nervos et nervulos ochraceo-hirtello-villosis siccis rigido-membranaceis opacis crebre pellucido-punctulatis 7-nerviis, nervis subtus prominulis 3 centralibus ad apicem proximis fere ad medium ductis infimis subtilioribus, petiolo basi ima vaginante utrinque dense ochraceo-villoso, pedunculo ochraceo-villoso petiolum duplo superante, amento crasso cylindrico densifloro, rhachi foveolata, bractea spatulata extus villosa, stam. 4-5, filamentis antheram multum superantibus exsertis, bacca immersa glabra sub-4-angula apice stylum filiformem baccam fere æquantem gerente, stigmatibus 3 minutis imo apice. ♂ Prope Xalapa (Galeotti! in h. Kew. et Schiedel in h. Ber.), prope Miradores (Linden n. 115! in h. Boiss. et Galeotti n. 6011 fide Mart. et Gal. in Bull. ac. brux. v. 10. n. 2. p. 23). *Schilleria melastomoides* Kunth! in Linn. v. 13. p. 724. *Callianira melastomoides* Miq. Syst. p. 345. Ramuli ochraceo-villosi, limbi 0,17 longi, 0,075 lati, petioli 0,01. Pedunculi 0,025 longi. Amenta mat. 0,03 longa, fere 0,01 crassa. (v. s.)
20. *P. MARGINATUM* (Jacq. ic. rar. v. 2. p. 2. t. 215), foliis longe petiolatis e basi subæquali cordata ovato-acuminatis, acumine acuto, utrinque glabris vel supra pilosis membranaceis vel membranaceo-firmis pellucido-punctulatis 7-9-nerviis, nervis subtus prominulis 3 centralibus fere e basi solutis ad apicem ductis infimis sæpe obsolete, nervulis subtransversim parallelis, petiolo glabro ad  $\frac{2}{3}$  longitud. vel fere ad limbum vaginante, alis linearibus, pedunculo glabro quam petiolum dimidio brevior, amento densifloro folium subæquante, bractæ apice peltatæ pelta triangulari-semilunari margine ochraceo-hirsuta, stam. 4-5, stigmatibus 3-4. ♂ In Chincao Peruviae et Pernambuco (Miq. in Hook. Lond. Journ. bot. v. 4. p. 443), Pernambuco (Gardner, n. 1838! in h. Kew.), Brasilia prov. Santarem et Para (Spruce, n. 1! in h. DC. et 3658? et 723! in h. Francovil.), ins. Salutis (h. Lenormand!), Caracas (Humb.

in h. Willd. n. 697! 698! 699!), Novæ Granatæ prov. Barbacoas alt. 70 (Triana, n. 26! in h. Triana), Venezuela (Fendler n. 1142! in h. DC. et Moritz n. 1801! in h. brit. Mus.), Guyana gallica prope Karouany (Sagot n. 533! in h. Francovil.), Surinam (Hostman n. 629! in h. Francovil.), prope Poelebantje et Quarta (Kegel n. 85 et 135 fide Miq. in Linn. v. 22. p. 76), Panama (Seemann, n. 114!, Sutton et Hayes n. 741! in h. Kew.), Bahia Honda et communis circa Panama (Seemann n. 901! et 902! fide Miq. in Seem. bot. Herald. p. 198), Trinitate (Lockhart! in h. Kew.), Portorico (Wydlar n. 399! in h. DC.), Santo Domingo (Balbis! in h. DC.), Cuba (Wright n. 2267! in h. DC.). Nhandu Piso hist. nat. med. ed 1658. p. 197, cum ic. Saururus frutescens foliis plantagineis fructu brevior Plum. Am. ed. Par. t. 76. P. marginatum Jacq. collect. v. 4. p. 128, Vahl enum. v. 1. p. 331. P. caudatum Vahl eclog. v. 1. p. 3, enum. v. 1. p. 335. P. decumanum Aublet hist. pl. Guy. v. 1. p. 21? (exclus. syn. Marcg.), Willd. sp. v. 1. p. 162, Kunth! in H. et B. nov. gen. v. 1. p. 57, syn. v. 1. p. 112. P. catalpæfolium Kunth! nov. gen. v. 1. p. 58. P. anisatum Kunth! nov. gen. v. 1. p. 58, Opiz in Reliq. Hænk! fasc. 3. p. 161. Schilleria caudata Kunth! Linn. v. 13. p. 716. Sch. marginata Kunth! l. c. p. 718. Sch. catalpæfolia Kunth! l. c. Artanthe caudata Miq. Syst. p. 380, in Hook. Lond. journ. bot. v. 4. p. 443 et in Mart. flor. Bras. fasc. 11. p. 32. Schilleria caudata Grisebach flor. w. Ind. p. 170, Cat. pl. cub. p. 65. Ramuli glabri. Limbi 0,1-0,16 longi, 0,07-0,13 lati. Petioli 0,02-0,04 longi. Amenta florentia 0,002 crassa. (v. s.)

α. *Armum*, foliis firmis, amentis aromaticis. — In Santo Domingo (Balbis l. c.).

β. *catalpæfolium*, foliis supra pilosulis, amentis aromaticis. — In locis umbrosis prov. Cumanensis inter Cumanacoa et San Fernando alt. 100 hexap. (Kunth! l. c. et h. Willd. n. 698! et 699!). Artanthe catalpæfolia Miq. l. c.

\*\* Nervuli primarii non adeo conspique transversim paralleli.

21. P. TARAPOTIANUM, foliis petiolatis ovatis vel rotundato-ovatis apice acuminatis, acumine acuto, superioribus basi æquali truncato-rotundatis obtusiusve et infer. basi æquali rotundatis utrinque glabris supra nitidulis subtus pallidioribus siccis firmulis subpellucidis parce subtilissime pellucido-punctulatis, 7-nerviis, nervis subtus prominentibus, centralibus 3 ad apicem proximis alte supra medium ductis infimis subtilioribus, petiolo glabro basi ima vaginante, pedunculo glabro florente petiolum æquante, amento florente folium dimidium superante filiformi, rhachi pubescente, bractæ truncato-peltatæ pelta minuta semilunari margine inf. subtilissime ciliolata pedicello brevi ciliolato, staminibus 5, stigmatibus 3 sessilibus abbreviatis, ovario emerso glabro. ♀ In Peruvia orient. (Spruce n. 4157, in h. DC.). Ramuli glabri olivacei. Limbi 0,16 longi, 0,097 lati, petioli 0,012 longi. Amenta florentia 0,015 crassa. (v. s.)

22. P. VICTORIANUM (Cas. DC. in Seem. journ. of bot. 1866. p. 218), foliis breviter petiolatis ellipticis apice acuminatis, acumine obtusiusculo mucronulatoque, basi æquali subattenuato-acutiusculis supra glabris subtus ad nervos subtiliter hirtellis siccis firmule membranaceis subpellucidis subtiliter pellucido-punctulatis 5-7-nerviis, nervis subtus prominulis 3 centralibus per anastomoses fere ad apicem proximis 2 ad medium ductis infimis subtilioribus sæpe obsoletis, petiolo dorso subtiliter hirtello ad limbum vaginante, pedunculo glabro petiolum subsuperante, amento quam folium  $\frac{1}{4}$  brevior densifloro rhachi foveolata, foveolis margine hirta-puberulis, bractæ ovario proximæ obovato-cucullatæ dorso basi villosulæ vertice nudo inflexo peltam semilunarem margine inferiore ciliatam simulante, staminibus 5, hacca glabra emersa ovato-acuta basi rhachi subimmersa, stigmatibus 3 sessilibus. ♀ In Venezuela prope La Victoria (Fendler n. 1139! in h. DC.). Ramuli glabri, limbi 0,11 longi 0,06 lati, petioli 0,008 longi. (v. s.)

23. P. MEDIUM (Jacq. ic. rar. v. 1. p. 2. t. 8), foliis petiolatis elliptico-lanceolatis vel ovato-lanceolatis apice acuminatis, acumine obtusiusculo mucronulato, inferioribus basi æquali cordatis super. basi æquali rotundatis vel subattenuato-acutiusculis vel basi ima subcuneatis cuneatisve supra glabris subtus ad nervos puberulis vel glabris siccis membranaceis subpellucidis subtilissime

pellucido-punctatis 7-nerviis, nervis supra subprominulis subtus prominulis centrali ad apicem ducto proximis fere ad apicem sequentibus supra medium ductis infimis subtilioribus sæpe obsoletis, petiolo basi ima vaginante hirtello dein glabrato, pedunculo glabro petiolum æquante, amento filiformi quam folium  $\frac{1}{2}$  breviorē densifloro, rhachi puberula, bractæ ovario proximæ obovatæ dorso villosulæ vertice inflexo nudo, stam. 4, filamentis antheris æquantibus, ovario glabro, stigmatibus 3 sessilibus, bacca glabra emersa ovoidæ apice attenuata.  $\frac{3}{4}$  In Brasilia (Salzmann n. 623! in h. DC., Pohl n. 5020 in h. Vindob.), monte Corcovado (Langsdorff n. 84. 85. 98! in h. Petrop.), in sylvis Mirador Mexici alt. 3000 p. (Galeotti n. 6018 fide Martens et Gal. Bull. ac. Brux. v. 10. n. 2. p. 22), Porto-Rico (Wydler n. 344! in h. DC. et Deless.), Guadaloupa (Bertero in h. DC. et h. Balbis), Dominica (Mayerhoff n. 341, in h. Ber.), Jamaica (Murray n. 626! Swartz in H. J. Kew.), Martinica (Hahn n. 264! in h. Mus. brit.) P. medium Jacq. Coll. v. 1. p. 141, Lam. ill. p. 80. Vahl enum. v. 1. p. 330, Willd. spec. v. 1. p. 160, herb. ber. n. 681! P. discolor Sw. Prodr. p. 15, fl. ind. occ. p. 52. Enckea plantaginea Kunth! in Linn. v. 13. p. 599, Miq. Syst. p. 356 et Linn. v. 20. p. 133. Enck. Sieber Griseb. fl. w. Ind. p. 170. Enck. Swartzii Griseb. l. c. p. 169. Rami glabri, ramuli glabri. Limbi 0,11 longi, 0,05-0,06 lati, petioli 0,008-0,01 longi. Amenta flor. 0,06 longa, 0,0015 crassa (v. s.)

$\beta$ , *laxiflorum*, baccis remotioribus, foliis 5-nerviis. In Jamaicae memoribus Westmoreland (Purdie! in h. kew.).

24. *P. CEANOTHIFOLIUM* (Kunth! in H. et B. nov. gen. v. 1. p. 56), foliis breviter petiolatis ellipticis vel ovato-ellipticis apice acuminatis, acumine obtusiusculo mucronulato, basi æquali subrotundatis vel subattenuatis acutiusculisve supra glabris subtus ad nervos subtiliter hirtellis siccis membranaceis subpellucidis subtilissime pellucido-punctatis 7-nerviis, nervis subtus prominulis centrali ad apicem ducto proximis fere ad apicem sequentibus supra medium ductis infimis subtilioribus sæpe obsoletis, petiolo subtiliter hirtello basi vaginante, pedunculo glabro petiolum æquante, amento filiformi densifloro quam folium  $\frac{1}{2}$  breviori, rhachi puberula, bractæ ovario proximæ subrotundo-obovatæ extus puberulæ vertice nudo inflexo margine inferiore ciliolato, staminibus 4-6, ovario ovato emerso, stigmatibus 3-4, bacca ovoidæ apice attenuata.  $\frac{3}{4}$  In Brasilia (Mart. Herb. flor. Bras. n. 1208! in h. DC.), Rio Janeiro (Gaudichaud n. 81! 84! et 1108! in h. DC.), Brasilia (Riedel n. 295! in h. Petrop.), Bordones et Cumana (Humb. in h. Willd. n. 1108! et 688!), Ejutla prope Oaxaca (Galeotti n. 6014 fide Mart. et Gal. in bull. ac. Brux. v. 10. n. 2. p. 22), Porto Rico (Wydler n. 321! in h. DC.), St.-Thoma (Moritz! in h. Ber.), Guadaloupa (Lherminier! in h. DC.), Jamaica (Macfadayen in h. Hook. fide Miq. in Hook. Lond. Journ. bot. v. 4. p. 440, et Griseb. fl. w. Ind.) — h. Willd. l. c. Kunth syn. v. 1. p. 111. Enck. ceanothifolia Kunth! in Linn. v. 13. p. 597. P. reticulatum Vell. flor. flum. t. 61. Enckea ceanothifolia Miq. l. syst. p. 357, Linn. v. 20. p. 133, in Mart. flor. Bras. fasc. 11. p. 28. t. 2. f. 2. Enck. Amalago  $\beta$  hirtella Griseb. pl. W. Ind. p. 169. In h. Boiss. specim. extat ad *P. ceanothifolium* refer. quod in ins. Taïti (ubi cultum), ex Nova Hollandia ubi ab incolis in potionem inebriantem utitur, a Vieillard allatum dicitur. Rami glabri, ramuli subtilissime hirtelli, limbi 0,1-0,06 longi, 0,045-0,03 lati, petioli 0,003 longi. Amenta flor. 0,06 longa, 0,0015 crassa (v. s.)

25. *P. ORTHOSTACHYUM* (Kunth olim), foliis breviter petiolatis suboblique ovato-ellipticis ellipticisve apice acuminatis, acumine obtusiusculo mucronulato, basi æquali ima breviter cuneata supra glabris subtus ad nervos subtiliter puberulis siccis firmule membranaceis subopacis subtiliter pellucido-punctulatis 5-nerviis, nervis subtus prominulis 3 centralibus per anastomoses ad apicem fere laterales ad medium ductis, petiolo basi ima vaginante hirtello dein glabrato, pedunculo petiolum subsuperante glabro, amento quam folium  $\frac{1}{2}$  breviorē densifloro, rhachi subtiliter hirtella, bractæ ovario proximæ cucullato-obovatæ extus hirtellæ vertice nudo inflexo, staminibus 4, ovario emerso, stigmatibus 3-4.  $\frac{3}{4}$  In Brasilia (Sello! in h. Berol. et Mart. h. fl. Bras. n. 1208! Salzmann! in h. Deless. et Riedel n. 123! in h. Petrop.), prov. Rio Negro ad rio Parahiba (Mart. ! in h. Mart.), in sylvis prov. Bahia (Mart. !

- in h. Mart.), in Jamaica (Sloane h. ! cum ic. 79). *P. longum* Sloane hist. p. 13. n. 16. *Enckea orthostachya* Kunth ! in Linn. v. 13. p. 601, Miq. ! syst. p. 360, Linn. v. 20. p. 133 et in Mart. flor. Bras. fasc. 11. p. 28. *Enck. vernicea* Miq. syst. p. 359, in Linn. vol. 20. p. 132 et in Mart. fl. Bras. l. c. *Ramuli* glabri. Limbi 0,08-0,11 longi, 0,045-0,055 lati, petioli 0,008-0,01 longi. (v. s.)
26. *P. AMALAGO* (L. ! sp. p. 41, quoad syn. Sloane et Pluk.), foliis breviter petiolatis ovato-lanceolatis apice acuminatis, acumine obtusiusculo, basi æquali rotundatis utrinque glabris siccis membranaceis subopacis pellucido-punctatis 5-7-nerviis plerumque 5-nerviis, nervis subtus prominulis centrali ad apicem ducto proximis fere ad apicem sequentibus supra medium ultimis subtilioribus sæpe obsoletis, petiolo glabro, pedunculo petiolum æquante glabro, amento quam folium brevius, rhachi glabra, bractea obovata concava ovario proxima extus puberula, staminibus 4-6, ovario ovoideo, stigmatibus 3-4 sessilibus, bacca emersa ovoidea apice attenuata. ♀ In Jamaica (Swartz ! in h. DC., h. Lin. ! et h. hort. Cliff. ! h. Sloane p. 79. et 80 ! et h. Willd. n. 684 !), Barbadoes (Maycock pl. Barb. p. 39.), Antillis (Descourt. fl. Ant.). *P. frutex* Pluk. ! alm. p. 297, phyt. tab. 215. f. 2. prave delineata, h. Pluk. ! *P. longum* arboreum altius Sloane ! hist. p. 134. f. 87. f. 1. et h. Sloane l. c. *Saururus foliis lanceolato-ovatis* L. in H. Cliff. ! p. 140. *P. frutescens* diffusum ramis flexilibus geniculatis etc. Browne Jam. p. 121, et *P. arboreum erectum* etc. Browne l. c. p. 122, except. syn. L. *P. Amalago* Hill dict. v. 26. p. 17. f. 17. f. 4. Swartz ! Obs. p. 19. et h. Willd. l. c., Vahl emum. v. 1. p. 328., Willd. sp. v. 1. p. 160, Loudon encycl. n. 506. p. 28. *P. plantagineum* Lam. ill. p. 80. Rami et ramuli glabri. Limbi 0,09-0,1 longi, 0,04 circ. lati, petioli 0,008 longi. (v. s.)
27. *P. SIEBERI*, foliis breviter petiolatis ovato-ellipticis apice acuminatis, acumine acuto mucronulatoque, basi inæquali rotundatis vel attenuato-acutiusculis utrinque glaberrimis siccis membranaceis subpellucidis subtilissime pellucido-punctulatis 5-nerviis, nervis subtus prominulis 3 centralibus per anastomoses ad apicem aliis ad medium ductis, petiolo glabro, pedunculo glabro petiolum subsuperante, amento quam folium  $\frac{1}{2}$  brevius densifloro, rhachi glabra, bractæ ovario proximæ obovato-subrotundæ glabræ vertice inflexo, staminibus 5-6 (plerumque 6), bacca glabra, rhachi basi subimmersa ovato-acuta, stigmatibus 3-4 sessilibus. ♀ In ins. Trinitatis (Sieber, n. 224 ! in h. DC.), St. Crucis (h. Berol. !), Cuba (Rug n. 305. sec. Griseb. cat. pl. cub. p. 65). *Enckea Sieberi* Miq. ! syst. p. 358, illust. p. 65. f. 65. Griseb. l. c. *Ramuli* glabri. Limbi 0,1-0,12 longi, 0,05-0,065 lati, petioli 0,01 longi (v. s.)
28. *P. LINDENII*, foliis brevissime petiolatis elliptico-lanceolatis apice longiuscule acuminatis, acumine acuto mucronulatoque, basi inæquali dimidiato-sagittato-cordatis lobo majore rotundato auriculæformi petiolum velante supra glabris subtus ad nervos subtiliter puberulis siccis membranaceis subpellucidis sparse pellucido-punctulatis 4-5-nerviis, nervis 3 centralibus fortioribus subtus prominulis, nervo centrali ad apicem proximis supra medium ductis infimis subtilioribus brevibus, petiolo basi ima vaginante utrinque subtiliter puberulo petiolum superante, folio abortivo squamæformi opposito, rhachi subtiliter hirtella, bractea ovario proxima subrotundata cucullato-concava basi subtiliter hirtella, staminibus 4-5, ovario subgloboso emerso, stigmatibus 4 sessilibus abbreviatis 3. In sylvis densis ad Tabasco Mexici (Linden ! in h. Deless.). *Enckea Lindenii* Miq. ! syst. p. 318, ill. p. 68. f. 68. Rami ramulique subtiliter puberuli, limbi 0,07-0,1 longi, 0,02-0,04 lati, petioli 0,005 longi. Pedunculi 0,02-0,04 longi (v. s.)
29. *P. UHREI*, foliis breviter petiolatis elliptico-vel subovato-lanceolatis lanceolatis apice longiuscule acute acuminatis basi æquali acutis utrinque glabris siccis rigidulis opacis subtilissime pellucido-punctulatis 5-nerviis, nervis supra subprominulis subtus vix proximulis 3 centralibus ad apicem ductis infimis subtilioribus brevioribus sæpe deficientibus, petiolo basi ima vaginante glabro superne ad marginem hirtello, pedunculo glabro petiolum duplo superante, amento filiformi densifloro folium æquante, rhachi hirtella, bractæ ovario proximæ truncato-peltatæ breviter pedicellatæ pelta semilunari margine inferiori villosula pedicello dorso villosulo, staminibus 4, ovario emerso glabro

per anthesim secus rhachim elongato, stigm. 3 sessilibus abbreviatis planiusculis. ♀ In Mexico (Uhde n. 252 ! in h. Berol.) Ramuli glabri. Limbi 0,12 longi, 0,045 lati, petioli 0,007 longi. (v. s.)

30. *P. BERTEROANUM*, foliis breviter petiolatis in eodem specim. oblongo-ellipticis subrhombeisque utrinque acutis vel elliptico-lanceolatis apice longius acuminatis, acumine obtuso, basi subæquali vel parum inæquali subrotundatis utrinque glabris siccis membranaceis vel rigidulis subopacis subtiliter pellucido-punctulatis 5-nerviis, nervis 3 centralibus fere ad apicem lateralibus ad medium ductis, petiolo glabro, pedunculo glabro petiolum paulum superante, amento quam folium  $\frac{1}{2}$  breviori subaxillifero, rhachi puberula, bractæ ovario proximæ cucullato-concavæ basi extus puberulæ vertice nudo, staminibus 4-5, ovario cylindrico-oblongo glabro, stigmatibus 3 sessilibus, bacca oblongo-ovata apice attenuata. ♀ In Xalapa alt. 4000 ped. (Galeotti n. 6010 ! in h. Deless.), sylvis umbris prope Xalapam (Schiede ! in h. Berol.), Jamaica (Bertero ! in h. DC. spec. ster.) Enckea discolor Kunth ! in Linn. v. 13. p. 595., Miq. ! syst. p. 361. Ramuli glabri. Limbi 0,07-0,09 longi, 0,03-0,04 lati, petioli 0,005 longi. Amenta matura 0,06 longa (v. s.)
31. *P. WULFSCHLÆGELI*, foliis breviter petiolatis lanceolatis subovato-lanceolatis apice acuminatis, acumine acuto, basi æquali acutiusculis utrinque glabris siccis rigido-coriaceis subopacis pellucido-punctulatis 5-nerviis, centrali nervo ad apicem ducto proximis alte supra medium ultimis subtilioribus sub medium ductis, petiolo glabro basi vaginante, pedunculo glabro petiolum paulo superante, amento densifloro folium fere duplo superante, rhachi pubescente, bractea ovario proximæ ovato-subrotunda intus et dorso pubescente, staminibus 5-6, ovario glabro, stigmatibus 3. ♀ In Jamaica (Wulfschlagel ! in h. monac.) Rami ramulique glabri, parce lenticellosi. Limbi 0,06 longi, 0,02 lati, petioli 0,04 longi. (v. s.)
32. *P. RICHARDIANUM*, foliis breviter petiolatis lanceolato-ellipticis apice acuminatis, acumine obtusiusculo, basi æquali acutiusculis supra glabris subtus præsertim ad nervos puberulis siccis membranaceis subopacis subtiliter pellucido-punctulatis 5-nerviis, nervis subtus prominulis centrali ad apicem ducto proximis alte supra medium infimis subtilioribus sub medium per anastomoses evanidis, petiolo subtiliter hirtello basi ima vaginante, pedunculo subtiliter hirtello petiolum duplo superante, amento folium dimidium superante, rhachi dense pallide hirtello-pubescente, bractea ovario proximæ subrotundo-cucullata ciliata basi dorso hirtello-pubescente, staminibus 6, ovario basi immersulo, stigmatibus 4 abbreviatis. ♀ In Cuba orientali (Wright n. 514 ! in h. DC.) Rami glabri, ramuli subtiliter hirtelli. Limbi 0,06 longi, 0,033 lati, petioli 0,006 longi. (v. s.)
33. *P. CALIFORNICUM* (Opiz ! in Reliq. Hænk. fasc. 3. p. 160), foliis breviter petiolatis subrotundo-ovatis apice acuminatis, acumine obtuso, basi breviter cordatis utrinque glabris siccis firmulis subopacis subtilissime pellucido-punctulatis 7-nerviis, nervis subtus subprominulis 3 centralibus per anastomoses fere ad apicem proximis ad medium ultimis subtilioribus sub medium ductis, petiolo glabro, pedunculo quam petiolum paulum brevius, amento quam folium dimidio brevius subaxillifero, rhachi hirtella, bractea infra florem inserta obovata ciliata et basi dorso pilosa, staminibus 4-6, ovario emerso, stigmatibus 3 abbreviatis crassiusculis. ♀ In California ad montem Rey (Hænce ! in h. Berol.) Enckea californica Kunth ! in Linn. v. 13. p. 601, Miq. ! syst. p. 363. Ramuli glabri. Limbi 0,065 longi, 0,045-0,065 lati, petioli 0,01 longi. (v. s.)
34. *P. UNGUICULATUM* (Ruiz et Pav. flor. peruv. 1. p. 34. t. 37. fig. b), foliis breviter petiolatis oblongo-ovatis vel ovato-lanceolatis apice acuminatis, acumine acuto, basi æquali inferioribus cordulatis superioribus rotundatis utrinque glabris siccis membranaceis firmis sæpe glaucescentibus 3-5-nerviis, nervis subtus prominulis centrali ad apicem ducto proximis supra medium infimis subtilioribus, petiolo glabro ad  $\frac{1}{2}$  longit. fere vaginante alis anguste linearibus cito deciduis, pedunculo glabro petiolum æquante, amento densifloro quam folium brevius, rhachi hirt-puberula, bractæ ovario proximæ spatulata apice obtusæ basi dorso puberulæ vertice inflexo nudo, staminibus 4-5-6,

stigmatibus 3-4 sessilibus, bacca ovato-oblonga glabra apice attenuata.  $\beta$ . In Peruvia (herb. madrit. in h. DC.), in Peruviae nemoribus Chinca et Pozuzo (R. et Pav. l. c. et h. Pav. in h. Brit. Mus.), Chila dit. Puebla (Andrieux n. 96 in h. DC.), Mexico (Schiede in h. Brit. Mus., et h. Willd. n. 695! specim. Humb. et Schultz n. 512! in h. Vindob. et h. Pav. l. c.), Acapulco (Hænke! in h. Berol.), Vera Cruz (Schiede! in h. Berol.), temperatis prope Xalapa et montem Macultepec alt. 680 hexap. (H. et B. in nov. gen. v. 1. p. 57), Jamaica (Baibis! in h. Berol. et Griseb. fl. w. Ind), Cuba (Griseb. l. in Cat. pl. Cub. p. 65. num. 513 exclus.), Guyana anglica (Schomb. l. in h. Berol.). In caldariis vulgare. P. terminale Kunth, in H. et B. l. c., et syn. v. 1. p. 112. P. unguiculatum Vahl enum. v. 1. p. 331; Mart. syst. mat. med. p. 99. P. glaucescens Jacq. eclog. t. 76. P. pyrifolium Opiz! in Reliq. Hænk. fasc. 3. p. 158, non Vahl. P. celidifolium Desf. Cat. hort. Par. ed. 3. p. 414. Enckea unguiculata Kunth! in Linn. v. 18. p. 592; Miq. l. syst. p. 352. Enck. glaucescens Kunth in Linn. v. 13. p. 594; Miq. syst. p. 354, et in Hook. Lond. journ. bot. v. 4. p. 440. Enck. Amalago Griseb. fl. w. Ind. excl. syst. Sloane et Cat. pl. Cub. l. c. Rami glabri: ramuli glabri vel juniores subtiliter puberuli. Limbi 0,06-0,08 longi; 0,025-0,04 lati; petioli 0,008 longi. (v. s.)

$\alpha$ , minus, foliis firmioribus. In Mexico prope Orizaba (Botteri! et Sallé! in h. DC. et Boiss. et h. Willd. n. 684!) (v. s.)

$\beta$ , coriaceum, foliis coriaceis. In ins. Sacrificios (h. Le Normand!) (v. s.)

35. P. TIGERIANUM (Cas. DC. in Seem. journ. of bot., 1866. p. 300), foliis petiolatis ovato-lanceolatis apice acuminatis, acumine acuto, basi aequali rotundatis truncatisve vel inferioribus truncato-cordulatis supra glabris subtus ad nervos subtiliter hirtellis siccis membranaceis 7-nerviis, nervis subtus prominulis 3 centralibus ad apicem ductis, petiolo glabro juniore subtilissime hirtello, pedunculo glabro quam petiolus brevior, amento folium superante densifloro, rhachi foveolata foveolis margine hirtellis, bractea obovata ovario proximæ dorso basi hirtello, staminibus 5, bacca ovata basi rhachi subimmersa, stigmatibus 4-5 sessilibus abbreviatis.  $\gamma$  In insula Tiger (Barclay n. 2122! in h. Brit. Mus.), ad Realego (Barclay n. 2080! l. c.) et Honduras (Barclay n. 2633! l. c.), in Nova Hispania (Pav. l. in h. Boiss.), Cuba (Præpigg n. 1822! in h. Boiss.). Ramuli glabri. Limbi 0,085 longi, 0,05 lati. Petioli 0,02 longi. Pedunculi 0,01 longi. Amenta mat. 0,11 longa. Amenta cocta odorem foliorum Lauro-Cerasi præbent. (v. s.)

36. P. PANDURATUM, foliis breviter petiolatis e basi ovata irregulariter trilobis, lobis lateralibus obtusis brevissimis, lobo centrali longe lineari-acuminato acumine acuto, vel e basi irregulariter ovata subito lineari-acuminatis, omnibus utrinque glabris siccis rigidulis opacis epunctulatis 5-nerviis, centrali nervo ad apicem ducto proximis ad lobos laterales productis infimis subtilioribus, petiolo glabro basi ima vaginante, pedunculo petiolum paullo superante, amento quam folium paullo brevior subdensifloro, rhachi pubescente, bractea ovata ovario proxima dorso basi et intus pubescente, staminibus 4-5-6, stigmatibus 3, ovario glabro.  $\gamma$  In Jamaica (Wulfschlägel! in h. monac.) Enckea subpandureformis Miq. sylloge Pip. nov. in Linn. v. 26. p. 217. Rami ramulique glabri, sicci nigrescentes. Limbi 0,075 longi, sub lobis 0,01 lati; petioli 0,005 longi. (v. s.)

37. P. KUNTHII, foliis breviter petiolatis elliptico-lanceolatis apice longe acuminatis, acumine obtusiusculo submucronato supra appresse sparse pilosis subtus ad nervos præsertim pubescenti-hirtis siccis coriaceis subopacis parce pellucido-punctulatis 5-7-nerviis, nervis 3 centralibus fortioribus subtus prominulis centrali ad apicem ducto proximis supra medium sequentibus sub medium ductis infimis subtilioribus obsoletis, petiolo pubescenti-hirto, pedunculo petiolum subduplo superante pubescenti-hirto, amento folium subaequante, bractea ovario proxima subrotundato-spathulata extus villosa intus glabra, staminibus 5, stigmatibus tribus.  $\beta$  In montibus formationis gnéiss alt. 5500 ped. ad Oaxaca in Mexico (Galeotti n. 6014! in h. Deless.). Enckea Kunthii Miq. l. syst. p. 363. Rami glabri, ramuli hirtopubescentes. Limbi 0,05-0,075 longi, 0,015-0,04 lati; petioli 0,002-0,005 longi. (v. s.)



38. *P. MARTENSIANUM*, foliis breviter petiolatis inferioribus rotundato-ovatis apice breviter acutatis basi subinæquali rotundatis subcordatisque superioribus suboblique ovatis et ultimis suboblique ellipticis apice breviter acuminatis, acumine obtusiusculo mucronulato, basi subinæquali obtusis omnibus supra ad basin parce subtus ubique dense molliter pubescentibus siccis tenuiter membranaceis subopacis subtilissime pellucido-punctulatis 7-nerviis, nervis subtus vix prominulis, centrali ad apicem producto proximis fere ad apicem sequentibus ad medium ductis infimis brevioribus subtilioribus, nervulo obscuro juxta marginem, petiolo basi ima vaginante utrinque molliter dense hirtopubescente, pedunculo dense pubescente petiolum subæquante, amento folium æquante densifloro, rhachi hirtella, bractæ ovario proximæ subcullato-obovatæ intus extusque villosulæ vertice subinflexo semilunari nudo, staminibus 5-6, ovario ovato, stigmatibus plerumque 4 sæpe 3, bacca suboblongo-ovata apice subattenuata subtiliter pulverulento-puberula. ♀ In Nova Hispania (Pav. in h. Boiss.) Ramuli dense molliter subcinereo-pubescentes. Limbi infer. 0,09 longi, 0,07 lati, super. 0,07 longi, 0,046 lati, petioli 0,006 longi. Bacca matura 0,002 longa. (v. s.)

39. *P. DECRESCENS*, foliis petiolatis inferioribus lato-ovatis apice breviter acuminatis basi æquali profunde cordatis superioribus ovatis apice breviter acuminatis basi æquali vel inæquali rotundatis utrinque ad nervos puberulis siccis membranaceis subpellucidis 7-interdum 9-nerviis, nervis subtus prominulis centrali ad apicem ducto proximis fere ad apicem sequentibus ad medium ductis infimis subtilioribus, petiolo puberulo, pedunculo quam petiolum brevius puberulo, amento folium æquante, rhachi pubescente, bractæ ovario proximæ spatulatæ basi dorso pilosæ vertice inflexo nudo, staminibus 5, stigmatibus 3, bacca oblongo-ovata apice attenuata. ♀ In America centrali (Barclay! in h. Kew.), Mexico prope Hacienda de Los Marenjas (Seemann. n. 1455! in h. Kew.) Enckea decrescens Miq.! in Hook. Lond. journ. bot. v. 4. p. 440. Ramuli glabri. Limbi 0,08 longi, 0,05 lati, petioli 0,015-0,03 longi. Amenta mat. 0,08 longa. (v. s.)

♂, ovatum, foliis supra et subtus ad nervos puberulis, pedunculo petiolum æquante, bacca ovata, cæteris conformibus. — In Guatemala, Grenada (Friedrichsthal n. 905! in h. Vindob.).

Sect. III. STEFFENSIA. — Gen. Steffensia et Schilleria Kunth in Linn. v. 13. p. 609 et 676. — Enckea Kunth Kunth l. c. (pro parte). — Ottonia Sprengel in neue Endeck. v. 1. p. 225. — Serronia Gaudich. et Guillem. in Deless. icon. — Peltobryon Klotzsch mscr. in h. Berol. et Miq. Syst. p. 369. — Artanthe Miq. comm. phyt. p. 40. — Zippelia Blume in Rem. et Schult. v. 7. p. 1614 et 1651. — Piperis sect. Brachystachys et pro parte Macrostachys Cas. DC. in Seem. journ. of bot. 1866.

Bractea a flore distincta. Stamina 4. Amenta oppositifolia.

§ 1. *Stigmata 3-5 in eodem specimine. Flores hermaphroditi. Antheræ articulatæ.*

40. *P. DUCHASSAINGII*, foliis breviter petiolatis late ovatis apice acuminatis, acumine acuto, basi subinæquali æqualive subcordatis rotundatisve vel truncato-rotundatis utrinque glabris nitidulis subtus pallidioribus siccis rigidulis subopacis pellucido-punctulatis 7-9-nerviis, nervis subtus prominentibus 3 centralibus ad apicem ductis infimis valde subtilioribus sæpe deficientibus, petiolo glabro basi vel ad  $\frac{1}{2}$  longit. vaginante, pedunculo glabro petiolum æquante, amento quam folium brevius densifloro, rhachi hirtella glandulosa, bractæ pedicellatæ vertice truncato-peltatæ pelta minuta quadrangulati margine ciliolata, bacca obovato-sub-4-angula centro disci stigmata 3-5 abbreviata gerente. ♀ In Guadaloupe (Duchassaing! in h. Berol. et Bertero! in h. DC.). Rami ramulique glabri. Limbi usque ad 0.2 longi, 0,115 lati. Petioli 0,01 longi. (v. s.)

§ 2. *Stigmata* 4. *Flores hermaphroditi. Antheræ articulatae.*

## α. Flores stipitati.

## 1° Folia penninervia.

41. P. JABORANDI (Vell. flor. flum. tab. 54), foliis brevissime petiolatis subæqualiteris ovato-oblongis apice acuminatis, acumine acuto mucronulatoque, basi subinæquali rotundatis cordulatisve utrinque glabris subtus ad nervos dense hirtellis siccis subopacis epunctulatis, nervo centrali subtus prominulo ad apicem ducto utrinque tota longitudine nervos subtus prominulos alternos 6-8 patulo-subadscendentes mittente, petiolo basi ima vaginante utrinque dense hirtello, pedunculo hirtello, amento densifloro maturo folium subæquante, rhachi glabrata, bractea cucullato-unguiculata postice gibbosa puberula infra stipitem inserta, rhachi pilis subtilissimis punctiformibus obsita, stipite baccam duplo et plus superante hirtello, bacca ovato-globosa sulcata sub-4-angula foveolato-punctata. ♀ In Brasilia (Pohl! in h. Vindob.), prope Rio Janeiro (Wedd. n. 761 in h. DC. et h. Mus. Par. et Gaudichaud n. 1102 in h. DC.), monte Corcovado (Guillemin n. 728! in h. DC. et Deless. Vauthier n. 547! in h. DC. et Langsdorff n. 83 in h. Petrop.), prov. Sebastianopolitana (Mart. in h. Monac.), sylvis montium organensium (Gardner n. 5862! Miq. in Hook. Lond. journ. bot. v. 4. p. 470). Jaborandi sec. spec. frutescens Piso? Hist. nat. Med. ed. 1658. p. 215, cum icone. Serronia Jaborandi Gaudich. et Guillemin! in Deless. ic. p. 54. t. 90. Ottonia Anisum Sprengel neue Entdeck. v. 1. p. 255, Miq.! syst. p. 536, illustr. p. 82. t. 92. f. b., in Hook. Lond. journ. bot. v. 4. p. 470, Linn. v. 20. p. 175 et in Mart. fl. Bras. fasc. 11. p. 63. t. 32. f. 3; Mart. in Mat. med. p. 100. Ottonia Jaborandi Kunth in Linn. v. 13. p. 579. Rami glabri, ramuli subtiliter hirtelli. Limbi 0,08-0,14 longi, circ. 0,04 lati, petioli 0,003-0,004. Pedunculi 0,007-0,088; stipites mat. 0,005 longi. (v. s.)
42. P. SELLOI, foliis brevissime petiolatis oblongis lanceolatisve apice longe angustato-acuminatis basim versus subattenuatis ima basi æquali rotundatis supra glabris subtus ad nervos dense hirtellis siccis rigidulo-membranaceis subpellucidis obsolete pellucido-punctulatis epunctulatisve, centrali nervo ad apicem ducto subtus prominulo utrinque tota longitudine nervos circ. 10 alternos subpatulos nervulosque fortiores mittente, petiolo dense hirtello basi ima vaginante, pedunculo dense hirtello petiolum æquante, amento densifloro quam folium  $\frac{1}{3}$ - $\frac{1}{2}$  brevior, rhachi glabra, bractea pedicellata stipiti proxima apice saccato-galeata extus subtilissime hirtella, stipite subtilissime hirtello baccam superante, bacca patente sulcata tetragona. ♂ In Brasilia (Sello! in h. Berol.), Bahia (Blanchet, n. 1946! in h. DC.). Ottonia diversifolia Miq! Syst. p. 538 et in Mart. fl. Bras. fasc. 11. p. 63. Ramuli juniores subtiliter hirtelli. Limbi 0,12 longi, 0,03 lati, petioli 0,004 longi. Stipes 0,02-0,004 longus. (v. s.)
43. P. EUCALYPTOPHYLLUM, foliis petiolatis oblongo-lanceolatis apice longe angustato-acuminatis basi inæquali cuneato-angustatis utrinque glabris siccis subcoriaceis subopacis pellucido-punctulatis, centrali nervo subtus prominulo utrinque nervos 12-15 tenues alternos subpatulos mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum subsuperante, amento densifloro, bractea breviter pedicellata stipiti proxima saccato-galeata glabra, floris stipite glabro baccam paulo superante, bacca sulcata-4-angula glabra foveolato-punctulata. ♀ In Brasilia prope Rio de Janeiro (Gaudich. n. 1100! in h. DC. et Houlet! in h. Mart.), monte Corcovado (Guillemin. n. 728! in h. DC.). Ottonia eucalyptifolia Kunth in Linn. v. 13. p. 582. Miq.! Syst. p. 539 et in Linn. v. 20. p. 176 et in Mart. flor. Bras. fasc. 11. p. 66. Rami ramulique glabri. Limbi 0,12 longi, circ. 0,03 lati, petioli 0,005-0,008. Pedunculi 0,008-0,01. Stipes 0,0025 longus. (v. s.)
44. P. SEBASTIANOPOLITENSE, foliis breviter petiolatis ovato-lanceolatis vel ovato-oblongis apice acuminatis, acumine acuto, basim versus attenuatis subattenuatisve, basi ima æquali acutis acutiusculisve, utrinque glabris subtus non procul a margine serie una pilorum puberulis siccis membranaceis subpellucidis

ad nervulos subtiliter pellucide glanduloso-punctulatis, nervis subtus vix prominulis centrali utrinque fere tota longitudine nervos 11 circ. patulo-subadscendentes nervulosque aliquot fortiores mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum paulum superante, amento maturo folium æquante, rhachi fusco-punctulata, bractea stipiti proxima subsessili saccato-galeata glabra, stipite baccam multoties superante fusco-punctulata, bacca ovata sub-4-angula glabra. ♀ In sylvis primævis prov. Rio Janeiro (Mart. in h. Monac.). *Ottonia macrostachya* Miq. in Linn. v. 20. p. 176, et in Mart. flor. Bras. fasc. 11. p. 67. t. 22, f. 1. Rami ramulique glabri. Limbi 0,21 longi, 0,08 lati, petioli 0,008 longi. Bacca immat. 0,001 longa. (v. s.)

45. *P. HAYNEANUM*, foliis brevissime petiolatis oblongo-ovatis apice acuminatis, acumine acuto mucronulatoque, basi subinæquali rotundatis utrinque glabris siccis rigidis subpellucidis ad nervulos pellucido-punctulatis, centrali nervo subtus prominulo utrinque tota longitudine nervos circ. 12 alternos subpatulos nervulosque fortiores mittente, petiolo glabro, pedunculo glabro petiolum æquante, rhachi subtiliter puberula glanduloso-punctulata, bractea breviter pedicellata stipiti proxima apice saccato-galeata glabra, stipite ovarium multum superante glabro, ovario oblongo-tetragono. ♀ In Brasilia (Sello in h. Berol.). *Ottonia macrophylla* Kunth in Linn. v. 13. p. 583; Miq. Syst. p. 540. Ramuli glabri. Limbi 0,175 longi, 0,085 lati; petioli 0,01 circ. longi. Stipes 0,004 longus. (v. s.)

β, *macrophyllum*, foliis subtus haud procul a margine serie una pilorum puberulis siccis membranaceis. — In sylvis locis apicioribus prov. Minas-Geraes (Mart. in h. monac.). *Ottonia macrophylla* Miq. in Linn. v. 20. p. 176 et in Mart. flor. Bras. fasc. 11. p. 66. t. 22. f. 2.

46. *P. PADIFOLIUM*, foliis brevissime petiolatis subsessilibusve oblongis apice acuminatis, acumine acuto, basi æquali vel inæquali rotundatis cordulatisve utrinque glabris siccis rigidulo-membranaceis subpellucidis pellucido-punctulatis, nervo centrali tota longitudine nervos 10-13 alternos subpatulos mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum superante, amento densifloro quam folium brevius, rhachi glabra, bractea pedicellata glabra stipiti proxima saccato-galeata, stipite glabro baccam æquante, bacca ovata apice attenuata sulcato-subquadrangula glabra. ♀ In Brasilia (Sello in h. Berol.). *Ottonia padifolia* Kunth in Linn. v. 13. p. 580; Miq. Syst. p. 539. Rami ramulique glabri. Limbi 0,09 longi, 0,035 lati, petioli vix 0,002. Stipites vix 0,002 longi. (v. s.)

β, *Corcovadense*, foliis membranaceis parcius subtilissime pellucido-punctulatis apice longiuscule acuminatis, acumine acuto, basi inæquali rotundatis. — In monte Corcovado, prov. Sebastianopolitanæ (Mart. in h. Monac.). *Ottonia padifolia* Miq. in Linn. v. 20. p. 175 et in Mart. fl. Bras. l. c. p. 65.

47. *P. OVATUM* (Vahl! eclog. p. 3. t. 1), foliis brevissime petiolatis ovato-ellipticis apice acuto-subacuminatis versus basin subcuneato-angustatis basi ima subpellucidis obsolete pellucido-punctulatis epunctulatisve, centrali nervo utrinque tota longitudine nervos circ. 12 subpatulos mittente, petiolo glabro, pedunculo glabro petiolum æquante, amento densifloro quam folium  $\frac{1}{2}$  brevius, rhachi glabra, bractea pedicellata glabra stipiti proxima apice sulcato-galeata, stipite quam bacca paulum brevius, ovario ovato apice in stylum brevem attenuato, bacca elliptico-ovata sulcato-tetragona apice acuta. ♀ In ins. Trinitatis (Vahl in h. Willd. n. 651! et h. DC. et Schumacher! in h. Petrop.). *P. ovatum* h. Willd. Vahl enum. v. 1. p. 325. *Ottonia Vahlii* Kunth in Linn. v. 13. p. 585, Miq. Syst. p. 542; Griseb. fl. w. Ind. p. 172. Ramuli glabri. Limbi 0,14 longi, 0,065 lati, petioli 0,01 longi. Bacca fere 0,002 longa. (v. s.)

48. *P. BAPTISIANUM*, foliis brevissime petiolatis oblongis apice acuminatis, acumine acuto, basi leviter inæquali obtusis utrinque glabris siccis rigidomembranaceis subpellucidis parce pellucido-punctulatis, nervis subtus prominulis, centrali utrinque fere tota longitudine nervos circiter 10 patulo-subadscendentes nervulosque aliquot fortiores mittente, petiolo glabro basi ima

vaginant, amento quam folium brevior, rhachi glabra, bractea stipiti proxima pedicellata glabra apice saccato-galeata, bacca sulcato-subquadrangula glabra apice subattenuata, stipite glabro baccam æquante. ♀ In sylvis ad præs. S. John Baptiste prov. Minas-Geraes (Mart. l in h. monac. n. 434, sub nom. Ott. propinqua). Ottoniæ propinquæ forma Miq. l in Linn. v. 20. p. 176. Ott. propinqua Miq. l in Mart. fl. Bras. fasc. 11. p. 66 (non aliorum, nec P. propinquum C. DC.). Rami ramulique glabri. Limbi 0,17 longi, 0,06 lati, petioli 0,002 longi. Bacca mat. 0,003 longa. (v. s.)

49. *P. MACHADOENSE*, foliis brevissime petiolatis lanceolatis vel subovato-lanceolatis apice acuminatis, acumine acutiusculo, basi æquali acutiusculisve utrinque glabris siccis rigidulis subpellucidis pellucido-punctulatis, centrali nervo subtus prominulo utrinque fere tota longitudine nervos 8-10 subadscendenti-patulos mittente, petiolo glabro pulverulento basi ima vaginante, pedunculo petiolum æquante, amento quam folium paullo brevior, rhachi pulverulento-puberula, bractea stipiti proxima pedicellata saccato-galeata, stipite florum duplo superante, ovario ovato-attenuato. ♀ In sylvis primævis prov. Minas-Geraes, caldas Badista ad rio di Machado (G. A. Lindberg! in h. Mart.). Rami ramulique glabri. Limbi 0,14 longi, 0,045 lati, petioli 0,003 longi. (v. s.)

50. *P. BOUCHEANUM*, foliis breviter petiolatis ovato-oblongis apice acuminatis, acumine acutiusculo, basi rotundatis vel basim versus subattenuatis basi ima acutiusculis utrinque glabris siccis rigidulis subpellucidis epunctulatis, centrali nervo subtus prominulo utrinque tota longitudine nervos circ. 12 alternos patulo-adscendentes nervulosque fortiores mittente, petiolo glabro, pedunculo glabro petiolum æquante, amento quam folium  $\frac{1}{2}$  brevior densifloro, rhachi glabra, bractea stipiti proxima brevissime pedicellata glabra apice saccato-galeata, bacca subsessili oblongo-attenuata sulcato-tetragona glabra. ♀ In Brasilia (Sello l in h. Berol.) Ottonia leptostachya Miq. l in Mart. fl. Bras. fasc. 11. n. p. 68. Rami glabri. Limbi fere 0,14 longi, 0,052 lati, petioli fere 0,01 longi. Bacca 0,002 longa. (v. s.)

51. *P. GARDNERII*, foliis brevissime petiolatis oblongo-lanceolatis apice acuminatis, acumine acuto mucronulatoque, basi cuneato-attenuatis obtusiusculis utrinque glabris siccis firmo-membranaceis, centrali nervo subtus prominulo utrinque tota longitudine nervos 12-15 alternos nervulis fortioribus intermixtos patulo-subadscendentes mittente, petiolo puberulo, pedunculo petiolum æquante puberulo, amento densifloro quam folium multum brevior, rhachi glabra bractea infra stipitem inserta breviter pedicellata saccato-galeata glabra, stipite florem superante. ♀ In Brasilia (Gardner n. 5186! in h. Kew.). Ottonia Hookeriana Miq. l in Hook. Lond. journ. bot. v. 4. p. 470 et in Mart. fl. Bras. fasc. 11. p. 64. Ramuli glabri. Limbi 0,15 longi, 0,04 lati, petioli 0,005 longi. Amenta florent. 0,03 longa (v. s.)

52. *P. HIRTELLUM*, foliis brevissime petiolatis oblongis oblongo-lanceolatisve apice acuminatis, acumine acuto, basi rotundatis vel basim versus subattenuatis basi ima acutiusculis supra glabris subtus ad nervum centalem subtilissime puberulis siccis membranaceis subpellucidis subtilissime pellucidopunctulatis, nervo centrali subtus prominulo utrinque tota longitudine nervos 8-10 patulo-subadscendentes vel apicem versus nervulos fortiores mittente, petiolo glabro, pedunculo subtilissime puberulo petiolum æquante, amento quam folium triplo brevior subaxifloro, rhachi dense molliter hirtella, bractea breviter pedicellata apice saccato-galeata acuta dorso subtiliter hirtella infra stipitem inserta, floribus breviter stipitatis. ♀ In Brasilia (Sello l in h. Berol. n. 258 ubi cum Ott. Anisum mixta). Ramuli glabri vel juniores subtiliter puberuli. Limbi circ. 0,12 longi, 0,035 lati, petioli circ. 0,003 longi. Amenta flor. circ. 0,001 crassa. (v. s.)

53. *P. MIQUELIANUM*, foliis brevissime petiolatis ovato-vel elliptico-oblongis apice suboblique acuminatis, acumine acuto mucronulatoque, basi inæquali cordatis vel leviter inæquali excisis supra glabris subtus ad nervos tenerrime puberulis siccis rigido-membranaceis subopacis epunctulatis nervis subtus prominulis, nervo centrali utrinque tota longitudine nervos 8-9 alternos nervulosque fortiores patulo-subadscendentes subpatulosve mittente, petiolo glabro

basi ima vaginante, pedunculo petiolum æquante glabro, rhachi glabra, bractea brevissime pedicellata stipiti proxima saccato-galeata glabra, stipite florem paulum superante. § In Brasilia (herb. Mart.!) Ottonia Martiana Miq.! in Linn. v. 20. p. 178 et in Mart. fl. Bras. fasc. 12. p. 64. Rami ramulique glabri. Limbi 0,13 longi, fere 0,06 lati, petioli 0,005 longi. (v. s.)

54. *P. PUNCTATISSIMUM*, foliis petiolatis ovatis oblongisque apice acuminatis, acumine acutiusculo, basi æquali vel subæquali cordatis subtus ad nervum centralem pube densissima brevi subtomentosis siccis tenuiter coriaceis pellucido-punctatissimis, nervo centrali utrinque nervos 8-10 mittente, petiolo pilis densissime vestito, pedunculo petiolum paulo superante, amento folium æquante, bractea pedicellata cucullato-saccata ciliolato-albida, stipite pilis patentibus pubescente apice incrassato, ovario subgloboso obsolete anguloso, stigmatibus 3-4 magnis crassis patentissimis revolutis ad medium connatis et sic stigma 3-4 lobulatum peltatum efficientibus, bacca subglobosa obsolete angulata sub lente puberula. § In monte Corcovado ad Rio de Janeiro (Mikan fide Presl. in Epim. bot. p. 230). Ottonia consanguinea Presl. l. c.; Miq. in Mart. fl. Bras. fasc. 11. p. 65. Rami glabri, ramuli pilis densissimis flexuosis tomentosis. Folia fere quinquepollicaria, usque ad 2 poll. lata, petioli tri-biligneares (fide Presl. l. c.)

55. *P. CORCOVADENSE*, foliis breviter petiolatis elliptico-lanceolatis apice acuminatis, acumine acuto, basim versus attenuatis basi ima subæquali obtusis cordulatisve utrinque glabris siccis rigidulis subpellucidis obsolete pellucido-punctulatis, nervo subtus prominulo utrinque fere tota longitudine nervos 12-15 tenues alternos patulo-subadscendentes mittente, petiolo glabro basi ima vaginante, pedunculo petiolum æquante, amento densifloro maturo folium subæquante vel brevior glabro, bractea stipiti proxima glabra saccato-galeata breviter pedicellata, stipite glabro baccam subæquante, bacca sulcato-subquadrangula glabra. § In Brasiliæ prov. Bahia (Blanchet n. 2958! in h. DC.), prope Maritima (Blanchet n. 3519! in h. DC.), in sylvis prope Bahiam (Wawra bot. Ergebn. p. 139 et Maly n. 146 in Wawra l. c.), Brasilia (Sello et Kalkman! in h. Berol.), summitate montis Corcovado (Pohl 4776! et 5030! in h. Vindob.), sylvis insul. Sti Ludovici prov. Maragnanensis (Mart. n. 434! in h. monac.). Serronia acuminata Moric. pl. nouv. d'Am. t. 57. Ottonia Blancheti Moric. l. c. p. 87 et t. 57; Walpers ann. v. 1. p. 612; Miq.! in Mart. fl. Bras. fasc. 11. p. 68. Ottonia propinqua Kunth! in Linn. v. 13. p. 583; Miq.! syst. p. 541. Ott. Corcovadensis Miq.! in Linn. v. 20. p. 175 et in Mart. fl. Bras. fasc. 11. p. 67. t. 23. f. 1. Rami ramulique glabri. Limbi 0,16 longi, 0,065 lati, petioli 0,005 longi. Stipites fere 0,003 longi. (v. s.)

β, *sessilifolium*, foliis basi subattenuatis cordulatis, limbis 0,12 longis 0,04 latis, petiolis 0,002 longis. — In Bahia (Blanchet! n. 1094 in h. DC.).

γ, *petiolatum*, foliis minoribus longius petiolatis, limbis 0,12-0,14 longis, 0,04 latis, petiolis 0,005 longis. — In Brasilia (Gaudichaud n. 1101! in h. DC.).

56. *P. PTEROPODON*, foliis petiolatis oblongis apice longe acuminatis basi æquali vel subinæquali rotundatis supra glabris subtus ad nervos pulverulentis siccis membranaceis subtus pallidioribus, nervis subtus prominulis, centrali utrinque nervos 5 mittente, petiolo pulverulento ad  $\frac{1}{2}$  longit. vaginante, stipulis pulverulentis, stipite hirtello baccam superante, bacca ovoidea 4-sulcata pulverulenta. In prov. Bahia (Blanchet n. 1960 fide Moric. pl. nouv. d'Am. p. 88). Ottonia pteropoda Moric. l. c.; Walpers ann. v. 1. p. 612; Miq.! in Mart. flor. Bras. fasc. 11. p. 68. Petioli 4-6 lin. longi. Amenta mat. 4-3 poll. longa. (fide Moric. l. c.)

57. *P. CARPINIFOLIUM*, foliis breviter petiolatis oblongo-lanceolatis acuminatis basi æquali cordatis subtus pube densa velutinis siccis coriaceis impunctatis pinnatinerviis, nervis 5-7, petiolo marginato-alato pube minuta velutino, pedunculo petiolum æquante, rhachi glabra, bractea brevissime pedicellata cucullata ciliolata albida, stipite quam ovarium longiore, ovario ovato tetragono stigmatibus minutis, bacca ignota. § In Brasilia ad Rio de Janeiro (Bescke fide Presl. in Epim. bot. p. 229). Ottonia carpinifolia Presl. l. c.; Miq. in Mart. fl. Bras. fasc. 11. p. 64. Ramuli pilis minutis densis vestiti, folia usque semi-

quantum pollicem longa, in lat. ext. 6-10 in int. 5-8 lata, petioli bilineares. (Fide Preslii l. c.)

2° Folia multinervia.

58. *P. NEESIANUM*, foliis petiolatis lanceolatis vel elliptico-lanceolatis apice acuminatis, acumine acuto, basi æquali acutis utrinque glabris siccis rigidis subopacis epunctulatis nitidulis supra glaucescentibus 5-nerviis, nervis subtus subprominulis, centrali ad apicem ducto proximis alte supra medium ductis infimis subtilioribus brevioribus, petiolo glabro basi ima vaginante, amento florente folium subæquante sublaxifloro, rhachi glabra, bractea sessili subovato-rotunda stipiti proxima, stipite florem æquante. ♀ In Mexico (Karwinski n. 823! in h. Petrop.). Rami ramulique glabri, glaucescentes. Limbi 0,06-0,085 longi, 0,02-0,04 lati, petioli 0,005-0,01 longi. (v. s.)
59. *P. ZIPPÆIA*, foliis petiolatis ovato-lanceolatis apice acutis basi subinæquali cordatis utrinque glabris siccis membranaceis pellucidis 5-nerviis, nervo centrali ad apicem ducto proximis fere ad apicem infimis sub medium ductis, petiolo glabro ad  $\frac{1}{2}$  longit. vaginante, pedunculo glabro petiolum multum superante, amento laxifloro, bractea sessili ovato-lanceolata extus glandulosa stipiti proxima, stipite florem æquante, connectivo supra loculos producto, ovario oblongo apice tetragono glanduloso, bacca aculeis glochidiatis numerosissimis tecta. ♀ In ins. Java (Zollinger n. 2847 in h. DC. et h. Kew.). *Zippelia begoniæfolia* Bl. in Rœm. et Sch. syst. v. 7. p. 1614 et 1651; Miq. syst. p. 548, illustr. p. 82. t. 92. f. c, Fl. ind. bat. v. 1. pars 2. p. 456. *Zipp. lappacea* Bennett pl. pav. rar. p. 76. t. 16. Ramuli glabri. Limbi 0,065 longi, 0,045 lati, petioli 0,025 longi. Pedunculi 0,08 longi. Amenta mat. 0,03-0,04 longa, 7-8 flora. (v. s.)

β. Flores sessiles.

1° Folia penninervia.

60. *P. LETUM*, foliis breviter petiolatis oblongo-lanceolatis vel oblongis vel ovato-oblongis ovatisve apice angustato-acuminatis, acumine acuto, basi æquali vel subinæquali acutis acutiusculisve sæpe basi ima bicallosis utrinque glabris siccis coriaceo-membranaceis supra læte viridibus subopacis epunctulatis, nervo centrali subtus prominulo utrinque fere tota longitudine nervos 10-12 alternos venasque fortiores mittente, petiolo glabro basi ima vaginante, pedunculo quam petiolus paulum brevior. amento quam folium vix dimidio brevior densifloro, rhachi hirtella, bractea flori proxima pedicellata vertice saccato-galeata glabra pedicello basin versus ciliato, bacca sessili subgloboso-tetragona. ♂. Herb. Willd. n. 646 ! (specim. Hoffmannsegg). In Brasilia (Sello! in h. Berol. et Mart. n. 1212! in h. flor. Bras. in h. DC.) ad Rio de Janeiro, Wedd. n. 239 ! in h. DC., Mart. ! in h. monac), monte Corcovado prov. Rio Janeiro (Mart. in h. monac. et Casaretto n. 1701 ! in h. DC.), Brasilia (Riedel n. 293 ! n. 20 ! 64 ! 102 et 81 ! in h. Petrop.), Mandioca Brasiliæ (Langsdorff n. 110 ! et 107 ! in h. Petrop.), sylv. Bahiæ (Poeppig n. 3232 ! in h. DC.). *Ottonia læta* Kunth ! in Linn. v. 13. p. 587; Miq. ! syst. p. 544; in Hook. Lond. journ. of bot. v. 4. p. 470 et in Mart. Flor. Bras. fasc. 11. p. 70. Ramuli glabri, limbi 0,1-0,15 longi, 0,025-0,046 lati, petioli fere 0,004 crassi. Specimen Casaretti citatum formam foliis paulo minoribus basi vix glandulosis distinctam præbet. (v. s.)
61. *P. POHLIANUM*, foliis breviter petiolatis ovato-oblongis vel oblongo-ellipticis apice breviter acuminatis, acumine obtusiusculo mucronulatoque, utrinque glabris siccis membranaceis opacis pellucido-punctulatis, nervis subtus vix prominulis, centrali utrinque tota longitudine nervos circiter 15 subpatulos nervulosque fortiores mittente, petiolo glabro basi ima vaginante, amento folium æquante densifloro, pedunculo glabro petiolum æquante, bractea flori proxima brevissima pedicellata saccato-galeata superne glabra basi pilosula, rhachi pilosula, bacca sessili obovato-tetragona quadrisulcata vertice flavicante. ♀ In sylv. ad Coralfalga (Pohl n. 5033 ! in h. Vindob.). *Ottonia Pohlana* Miq. ! in Linn. v. 20. p. 179. et in Mart. fl. Bras. fasc. 11. p. 69. f. 23. f. 2. Rami ramulique glabri. Limbi 0,005 crassi. (v. s.)

62. *P. KLOTZSCHIANUM*, foliis brevissime petiolatis oblongo-vel-ovato-ellipticis apice acuminatis, acumine acuto, basi subinæquali obtusis acutiusculisve siccis rigidulo-membraneis obsolete pellucido-punctulatis subopacis, centrali nervo subtus prominulo utrinque ad apicem nervos circ. 12 alternos patulo-subadscendentes interdumque nervulos fortiores mittente, petiolo glabro, pedunculo petiolum subæquante glabro, amento quam folium duplo brevius densifloro, rhachi glabra, bractea breviter pedicellata infra florem inserta glabra vertice saccato-galeata, ovario subgloboso glabro. ♀ In Brasilia (Sello n. 205 ! in h. Berol.). *Ottonia Klotzschiana* Kunth ! in Linn. v. 13. p. 589, Miq. ! syst. p. 845 et in Mart. flor. Brasil. fasc. 11. p. 70. Rami ramulique glabri. Limbi circ. 0,11 longi, 0,045 lati, petioli 0,005-0,007 longi. Amenta florentia 0,06 longa, 0,003 crassa. (v. s.)
63. *P. COSMIANUM*, foliis brevissime petiolatis ellipticis apice longe acuminatis, acumine obtuso, basi æquali obtusis, supremis lanceolato-ellipticis apice conformibus basi æquali acutis utrinque glabris siccis firmulis subopacis pellucido-punctulatis, nervo centrali subtus prominente utrinque fere tota longitudine nervos alternos circ. 12 subadscendentes subpatulos subtus prominulos sæpeque nervulos fortiores mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum subæquante, amento densifloro florente folium dimidium superante, rhachi glabra vel subtilissime parce subpuberula, bractea breviter pedicellata valde infra florem inserta saccato-galeata glabra. ♀ In sylvis ad Bahia Cruz de Cosme (h. Fischeri n. 153 ! in h. Petrop.). Rami ramulique glabri. Limbi 0,16-0,185 longi, 0,057-0,084 lati; petioli 0,005 longi. Amenta florentia 0,002 crassa. (v. s.)
64. *P. WARAKABOURA*, foliis brevissime petiolatis oblongo-lanceolatis apice acuminatis, acumine acuto, basi attenuatis ima basi levissime inæquali acutis utrinque glabris siccis rigidis subtiliter pellucido-punctulatis, nervo centrali subtus prominulo utrinque tota longitudine nervos circ. 10 alternos patulo-subadscendentes nervulosque fortiores mittente, petiolo glabro basi ima vaginante, amento quam folium duplo brevius subdensifloro, pedunculo petiolum paulum superante glabro, rhachi foveolata foveolis hirtellis, bractea sessili flori proxima saccato-globosa dorso et margine villosa-hirtella, bacca sessili ovata 4-sulcata glabra basi immersa. ♀ In Surinam (Hostman pl. Hohenacker n. 687 ! a in h. Francovil. a Miq. ipso dicta) et Obidos (Spruce n. 524 in h. Francovil.), in Brasiliæ sylvis aboriginibus ad flum. Solimoes prope Coari et Ega prov. Rio Negro et in prov. paraensi (Mart. ! in h. monac.). *Ottonia Warakaboura* Miq. mant. ined., in Linn. v. 20. p. 180, in Mart. fl. Bras. fasc. 11. p. 69. f. 24. et in Hook. Lond. journ. bot. v. 4. p. 469. Ramuli glabri. Limbi 0,22 longi, 0,08 lati; petioli 0,003 longi. Amenta 0,001 crassa. (v. s.)
65. *P. BARTLINGIANUM*, foliis brevissime petiolatis lanceolato-ellipticis apice acuminatis, acumine obtuso, basi subæquali acutis utrinque glabris supra nitidis subtus nitidulis siccis rigidis subopacis crebre pellucido-punctulatis, nervo centrali subtus prominente tota longitudine nervos utrinque 8-12 patulo-subadscendentes subtus prominulos nervulosque fortiores mittente, petiolo glabro, pedunculo glabro petiolo paulum superante, amento quam folium quadruplo brevius, rhachi foveolata foveolis margine fusce puberulis, bractæ parvæ cucullato-calceolatæ puberulæ flori proximæ vertice nudo calloso, ovario glabro rhachi basi subimmerso. ♀ In Guyana gallica (Poiteau ! in h. Deless.) et vicinibus Obidos prov. Para (Spruce n. 5 ! in h. Kew.). *Artanthe Bartlingiana* Miq. ! syst. p. 510. Ramuli glabri. Limbi 0,14-0,2 longi, 0,06-0,09 lati, petioli 0,003 longi. Amenta florent. 0,0015 crassa. (v. s.)
66. *P. PRESLI*, foliis breviter petiolatis oblongis apice obtuse acuminatis, basi æquali attenuatis utrinque glabris siccis subcoriaceis nitidulis subopacis pellucido-punctulatis, nervis subtus prominulis, centrali utrinque fere tota longitudine nervos 7-8 alternos patulo-subadscendentes nervulis fortioribus aliquot intermixtos mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum æquante, amento quam folium multoties brevius, bractæ flori proximæ glabræ cucullatæ dorso basi puberulæ vertice glabro inflexo triangulari, ovario glabro. ♀ In America, monte Cassinisvord Ma-

ranyna (Wulfschlägel! in h. Mart. n. 1559? et DC. in h. Mart. ubi folia tantum duo). Rami ramulique glabri. Limbi 0,125-0,135 longi, 0,045 lati; petioli 0,01 longi. Amenta florent. 0,01 longa, 0,002 crassa. (v. s.)

67. *P. LESSERTIANUM*, foliis breviter petiolatis ovato-oblongis apice acutis basi parum inæquali cordulatis utrinque glabris siccis coriaceis obsolete punctulatis, centrali nervo subtus prominulo utrinque ad  $\frac{1}{2}$  longitud. nervos 5 alternos subadscendentes sursumque ad apicem nervulos fortiores mittente, petiolo glabro, amento breviter pedunculato quam folium duplo brevior, rhachi subtiliter hirtella, bacca subgloboso-trigona. ♀ In America meridion. (Mathews n. 1696. ! in h. Deless.). *Ottonia Lessertiana* Miq. ! syst. n. 546, et in Illustr. p. 82. t. 92. f. a. Rami ramulique glabri, hic illic veruculosi. Limbi 0,14-0,145 longi, 0,045 lati; petioli 0,01 longi. (v. s.)

68. *P. FRANCOVILLEANUM* (Cas. DC. in Seem. journ. of bot. 1866. p. 719), foliis brevissime petiolatis subovato-ellipticis apice protracto-acuminatis, acumine acuto, basi inæquali rotundatis, latere majore auriculatim producto, supra glabris subtus ad nervum centalem villosis margine ciliatis siccis rigidulo-membranaceis\* subtilissime pellucido-punctulatis, nervis nervulisque subtus prominentibus, centrali utrinque tota longitudine nervos circiter 20 alternos nervulosque fortiores margine anastomosantes patulo-subadscendentes mittente, petiolo villosio, pedunculo petiolum multoties superante parce villosio, amento quam folium triplo brevior densifloro, rhachi villosa foveolata, bractea minuta flori proxima dorso et margine villosula, ovario quadrisulcato, bacca quadrisulcata subtiliter glanduloso-puberula. ♀ In Brasilia prope Barra (Spruce n. 1784 ! in h. Francovil.) et prope San Gabriel da Cocheira ad Rio Negro (Spruce n. 2362 ! in h. DC.). Radix fibrosa. Caulis apicem versus fusce dense villosus. Limbi 0,2 longi, 0,085 lati; petioli 0,005 longi. Pedunculi 0,065 longi. Amenta matura 0,0025 crassa. (v. s.)

## 2° Folia multplinervia.

69. *P. VERBASCIFOLIUM*, foliis brevissime petiolatis oblongis vel oblongo-lanceolatis basi valde inæquali auriculatim cordatis coriaceis pellucido-punctulatis ciliolatis flavescenti-viridibus utrinque planis et tomentosis nec belatis nec scabris, nervo centrali subtus prominulo ad  $\frac{1}{2}$  longitudinis usque nervos utrinque 5-8 subtus prominulos subadscendentes mittente, petiolo tomentoso basi ima vaginante, pedunculo tomentoso petiolum æquante, amento obtusissimo, bractæ apice peltatæ pelta triangulari-semilunari convexa margine hirsuta pedicello hirtello, ovario glabro, stigmatibus linearibus. ♀ In Peruvia (Dombey n. 932 ! in h. DC. ex h. Mus. par.), prope Ouito (Jameison n. 61 ! in h. Kew.), Mexico prope La Puebla (Andrieux n. 94 ! in h. DC.). *Artanthe verbascifolia* Miq. ! syst. p. 431 et in Hook. Lond. journ. bot. v. 4. p. 455. *Artanthe leucophylla* Miq. ! in Hook. Lond. journ. bot. v. 4. p. 457. Ramuli tomentosi. Limbi 0,14-0,16 longi, 0,05-0,055 lati. Petioli 0,005-0,008 longi. Pedunculi 0,005 longi. Amenta immatura 0,005 crassa. (v. s.)

## § 3. Stigmata 3. Antheræ articulatæ. Flores, unica specie excepta, hermaphroditi.

### A. Ovarium apice styliiferum.

#### 1° Folia penninervia.

70. *P. MUNCHANUM*, foliis subsessilibus oblongo-obovato-lanceolatis apice breviter acuminatis, acumine acutiusculo, basi æquali cuneatis in petiolum brevissimum decurrentibus utrinque glabris siccis subrigidis, nervo centrali subtus prominulo utrinque tota longitudine nervos 15-20 alternos patulo-adscendentes subtus prominulos mittente, pedunculo glabro brevi, bractæ adultæ glabræ vertice triangulari subpeltato glabro, ovario obovato. ♀ In Peruvia orientali prope Tarapio (Spruce, n. 4341 ! in h. Kew.), sylvis humidis prope Muncha in Sancta Martha (Purdie ! in h. Kew.). *Peltobryon Hookeri* Miq. ! in Hook.



Lond. journ. bot. v. 4. p. 442. Ramuli glabri. Limbi 0,3 longi, 0,12 lati. Pedunculi 0,05-0,06 longi. Amenta florent. 0,08 longa 0,004 crassa. (v. s.)

β, *magnifolium*, foliis majoribus, limbo 0,4 longo, 0,19 lato. ♀ Ad rivulos in sylvaticis Cuchero (Herb. Endlich! in h. Vindob.) (v. s.)

71. *P. PELTOBRYON*, foliis breviter petiolatis oblongo-ellipticis apice breviter acuminatis, acumine acuto, basin versus attenuatis basi subæquali acutis vel acutato-contractis utrinque glabris subtus glandulose fusco-punctatis siccis membranaceis subtiliter pellucido-punctulatis, centrali nervo subtus prominulo utrinque tota longitudine nervos circ. 8 alternos patulo-ascendentes nervulosque fortiores mittente, petiolo glabro, pedunculo petiolum æquante, amento cylindrico densifloro quam folium multoties brevius, bractearum pedicellatæ vertice truncato-peltatæ pelta triangulari margine ochraceo-hirsuta, ovario ovato-globoso, stylo ovarium æquante. ♀ In Peruvia (Meyen! in h. Berol.), Venezuela ad ripas umbras fluminum prope Palmas, San Martheo et Valles de Aragoa (Ed. Otto in h. Berol.) et Biscaina alt. 2000-3000 p. (Fendler, n. 1136! in h. DC.), Venezuela (Funck et Schlim, n. 577! in h. Petrop. et Moritz, n. 1938! in h. Brit. Mus.), Nova Granata prov. Barbaçoas (Triana, n. 90! in h. Triana) et Bahia (Luschnatt, in h. Schlecht. fide Miq. in Linn. v. 18. p. 713). *Peltobryon attenuatum* Klotzsch! in h. Berol. Miq.! Syst. p. 371, illustr. p. 67. t. 67, in Mart. fl. Bras. fasc. 11. p. 29, t. 5. f. 3, et in Linn. l. c. Ramuli glabri, glandulis fuscis conspersi. Limbi 0,105 longi, 0,055 lati. Petioli 0,008 longi. (v. s.)

72. *P. LONGIFOLIUM* (Ruiz et Pav. fl. per. 1. p. 37. t. 61. f. a), foliis breviter petiolatis amplis oblongis apice subacuminatis, acumine acuto, basi subæquali attenuato-acutis utrinque glabris siccis subcoriaceis vel rigido-membranaceis subopacis pellucido-punctatis subtus glanduloso-punctatis, nervo centrali ad apicem ducto subtus prominulo fere tota longitudine nervos utrinque circ. 18 alternos patulo-subadscendentes mittente, petiolo glabro, pedunculo glabro petiolum æquante, amento quam folium triplo-quadruplo brevius cylindrico crasso densifloro bractearum truncato-peltatæ concavo-pedicellatæ pelta triangulari-semilunari margine pallide fusce et dense hirsuta, ovario ovato, stylo ovarium æquante. ♀ In Andium peruvianorum nemoribus (R. et P. h. Willd. n. 641! et h. Berol. l.). *P. longifolium* h. Willd.! Vahl enum. v. 1. p. 318. *Peltobryon longifolium* Miq.! Syst. p. 373. Ramuli fusco-punctulati. Limbi 0,4 longi, 0,1-0,175 lati, petioli fere 0,025 longi. Amenta circ. 0,12 longa, 0,005-0,006 crassa. (v. s.)

73. *P. BRASILIENSE*, foliis breviter petiolatis lanceolato-oblongis apice acuminatis, acumine acuto, basi ima obtusiusculis supra glabris subtus pubescentibus siccis rigidulo-membranaceis opacis subtilissime pellucido-punctulatis, nervis subtus prominulis, centrali utrinque tota longitudine nervos circ. 12 subpatulos mittente, petiolo fusco-hirsuto ad limbum vaginante, pedunculo fusco-pubescente petiolum æquante, amento quam folium multoties brevius cylindrico, bractearum cucullato-spathulatæ vertice inflexo, hacca obovata puberula, stylo quam bacca brevius, rhachi puberula. ♀ In sylvis Japurensibus prov. Rio Negro (Mart. l. in h. Monac.). *Peltobryon pubescens* Miq.! in Linn. v. 20. p. 134, et in Mart. fl. Bras. fasc. 11. p. 30. t. 4. Rami subglabrati. Ramuli fusce pubescentes Limbi 0,11 longi 0,04 lati. Petioli 0,006 longi, Amenta mat. 0,015 longa, 0,006 crassa. (v. s.)!

#### 2° Folia multiplinervia.

74. *P. CARABOENSE*, foliis breviter petiolatis oblongo-lanceolatis acuminatis, acumine acuto mucronulatoque, basi leviter inæquali acutis utrinque glabris siccis membranaceis subpellucidis pellucido-punctulatis nervis subtus haud prominulis, centrali utrinque ad  $\frac{2}{3}$  longit. nervos 4 patulo-subadscendentes supremos ante apicem per anastomoses evanidos sursumque nervulos fortiores mittente, petiolo glabro basi ima vaginante, amento quam folium multoties brevius mucronato cylindrico densifloro, pedunculo petiolum æquante, bractearum vertice truncato-peltatæ pelta triangulari margine hirsuta pedicello subconcavo margine et dorso hirtello, stigmatibus abbreviatis.

- ♂ In Venezuelæ prov. Carabobo (Funck et Schlim, n. 577! in h. Boiss.). Ramuli glabri. Limbi 0,12 longi, fere 0,04 lati; petioli 0,006 longi. Amenta florentia 0,01 longa, 0,003 crassa. (v. s.)
75. *P. TEAPENSE*, foliis brevissime petiolatis subobovato-elliptico-oblongis apice acuminatis, acumine acuto mucronulatoque, basi inæquali cordatis, lobo majore auriculiformi producto petiolum velante, supra glabris subtus ad nervos nervulosque pubescentibus siccis membranaceis subopacis subtilissime pellucido-punctulatis, nervis subtus prominulis, centrali utrinque ad  $\frac{1}{2}$  altitudinis nervos supra basin 3 alternos supremos per anastomoses ante apicem evanidos et 3 e basi subtiliores divaricantes mittente, petiolo ad  $\frac{1}{2}$  longit. vaginante basi dorso et sursum utrinque pubescente, pedunculo puberulo petiolum fere duplo superante, bractæa vertice truncato-peltatæ pelta triangulari margine villosa pedicello angusto sulcato margine et dorso villosulo, ovario villosulo, stigmatibus filiformibus. ♂ In Teapa (Linden, n. 121! in h. Boiss.). Ramuli juniores pubescentes. Limbi 0,13 longi, 0,058 lati; petioli 0,005 longi. Amenta florent. 0,002 crassa. (v. s.)
76. *P. TENUISTYLUM*, foliis petiolatis ovato-oblongis vel subovatis apice acuminatis, acumine lineari-lanceolato subfalcato acutissimo, basi late rotundata subæquali in petiolum protractis siccis membranaceis subepunctatis subtus glandulosis supra glabris subtus tenerrime ad nervos puberulis cito glabratis, nervo centrali tenui ad  $\frac{3}{4}$  longit. nervos utrinque 6-8 patulos versus marginem adscendentes supremos per anastomoses fere ad apicem ductos nervulosque fortiores mittente, petiolo cito glabrato, pedunculum superante, amento quam folium  $\frac{1}{2}$  brevius cylindrico recto, bractæa conchæformi peltata hirta decidua, stylo filiformi ovarium superante, stigmatibus linearibus patulis. ♂ In Peruvia (Mathews, n. 1710! in h. Kew.). Peltobryon Mathewsii Miq! in Hook. Lond. journ. bot. v. 4. p. 441. Ramuli glabrati. Limbi 0,09-0,15 longi, 0,045-0,06 lati petioli 0,008-0,015 longi. Amenta florentia 0,055 longa, 0,003 crassa. (v. s.)
77. *P. CALLOSUM* (Ruiz et Pav.! fl. per. 1. p. 34. t. 53. f. a.), foliis breviter petiolatis oblique subrotundato-ovatis ovatisve apice suboblique acuminatis, acumine acuto, basi subinæquali acutiusculis rotundatisve siccis rigidulis utrinque glabris subtilissime pellucido-punctulatis, nervo centrali utrinque ad  $\frac{3}{4}$  longit. nervos 5 alternos subadscendentes mittente, petiolo glabro, pedunculo glabro petiolum æquante, amento quam folium quadruplo-septuplo brevius densifloro, bractæa cucullatæ vertice inflexo subpeltato, ovario ovato. ♂ In Peruvia (R. et P. in h. Willd. n. 653! et in h. Berol.!). *P. callosum* h. Willd.! l. c. Vahl enum. v. 1. p. 316. Schilleria callosa Kunth! in Linn. v. 13, p. 689. Peltobryon callosum Miq. Syst.! p. 369. Ramuli glabri. Limbi 0,13 longi, 0,072 lati, petioli 0,007 longi. (v. s.)
78. *P. PÆPPIGII*, foliis breviter petiolatis rhombeo-ellipticis ellipticisve vel subovatis apice longe lineari-acuminatis, acumine acutiusculo, basi subæquali acutiusculis utrinque glabris siccis rigido-membranaceis obsolete siccis pellucido-punctulatis subtus sparse punctatis pallidis, nervo centrali subtus prominulo utrinque ad  $\frac{3}{4}$  alt. nervos circ. 5 alternos subadscendentes mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum æquante, amento quam folium multoties brevius cylindrico densifloro, bractæa glabræ vertice truncato triangulari pedicello dorso carinato margine extenuato, ovario ovato, stylo longo. ♂ Ad flumen Amazonum (Pæppig! n. 1738 in h. Berol. et DC.). Peltobryon Pæppigii Klotzsch! mss. in h. Berol. et Miq.! Syst. p. 370. Ramuli glabri. Limbi 0,11 longi, 0,05 lati, petioli 0,01 circ. longi. Amenta post anthesin 0,015 circ. longa, 0,005 circ. crassa. (v. s.)
79. *P. DICHOTOMUM* (Ruiz et Pav.! fl. peruv. 1. p. 25. t. 60. f. b), foliis breviter petiolatis oblongis apice subito acuminatis, acumine acuto mucronulatoque, basi subæquali attenuatis acutis supra glabris subtus ad nervos subtiliter puberulis glanduloso-punctulatis siccis membranaceis subpellucidis subtiliter pellucido-punctulatis, nervo centrali utrinque ad  $\frac{1}{2}$  longit. nervos 7 alternos patulo-adscendentes nervulosque fortiores mittente, petiolo puberulo, pedunculo petiolum subæquante, amento quam folium multoties brevius cylindrico densifloro, bractæa spatulatæ glabræ vertice inflexo subpeltato triangulari, ovario ovato-globoso glanduloso-punctato, stylo quam ovarium bre-

viora, stigmatibus brevibus. ♀ In Peruvia et Chili (Meyen! et Ruiz! in h. berol.), in sylvis Chincac ad Macora et Messapata prædia (R. et P. l. c.). — Vahl en. v. 1. p. 316. Schilleria! longifolia Kunth! in Linn. v. 13. p. 725. Peltobryon dichotomum Klotzsch! mss. in h. berol. et Miq.! Syst. p. 372. Ramuli subtiliter puberuli. Limbi 0,16 longi, 0,08 lati, petioli 0,012 longi. (v. s.)

80. *P. OVALE* (Cas. DC. in Seem. journ. of bot. 1866. p. 166), foliis petiolatis ovato-ellipticis apice acuminatis basi subæquali obtusis utrinque glabris siccis firmulo-membranaceis subtiliter pellucido-punctulatis, nervis subtus prominulis, centrali utrinque ad  $\frac{3}{4}$  longitudinis nervos 6 alternos subadscendentes supremos ad apicem ductos mittente, petiolo glabro ad  $\frac{3}{4}$  longit. vaginante, pedunculo petiolum subæquante subtiliter apicem versus aureo-puberulo, amento cylindrico densifloro quam folium dimidio brevior, bractæ cucullatæ vertice inflexo-truncato subpeltato triangulari margine et postice longius aureo-ochraceo-hirsuto pedicello aureo-ochraceo, ovario ovato-globoso, stylo ovarium subæquante, stigmatibus recurvis lanceolatis, bacca globosa apice mucronata nigra. ♀ In Venezuela prope Tovar alt. 4000 p. (Fendler n. 2398! in h. DC.) et Nova Granata prov. Barbacoas (Triana n. 19! in suo herb.). Ramuli glabri. Limbi 0,17 longi, 0,09 lati, petioli 0,015 longi. Amenta 0,09 circ. longa, mat. 0,005 crassa. (v. s.)

81. *P. PROPINQUUM* (Cas. DC. in Seem. journ. of bot. 1866. p. 167), foliis longiuscule petiolatis suboblongo-ellipticis apice acuminatis, acumine acuto mucronulatoque, basin versus subattenuatis basi ima æquali acutis utrinque glabris siccis firmule membranaceis subpellucidis glanduloso-punctulatis nec pellucido-punctulatis subtus pallidioribus, nervis subtus prominulis, centrali utrinque ad  $\frac{1}{2}$  longitudinis nervos 5 subalternos subadscendentes supremos ad apicem ductos mittente, petiolo glabro ad limbum vaginante, pedunculo quam petiolus paulum brevior apicem versus aureo-puberulo, amento quam folium fere  $\frac{1}{2}$  brevior cylindrico densifloro, bractæ cucullatæ extus aureo-ochraceo-hirsutæ vertice inflexo truncato semilunari subpeltato margine aureo-ochraceo hirtello, ovario ovato-globoso, stylo ovarium fere æquante, stigmatibus lineari-lanceolatis, bacca globosa mucronata nigra. ♀ In Peruvia orientali prope Tarapoto (Spruce n. 4032! in h. DC.). Ramuli glabri. Limbi 0,135 longi, 0,065 lati, petioli 0,022 longi. Amenta mat. 0,045 longa, fere 0,004 crassa. (v. s.)

82. *P. EXSERENS*, foliis petiolatis inferioribus ovatis superioribus elliptico-ovatis apice acuminatis, acumine obtusiusculo, superioribus basi ima inæquali subattenuatis acutiusculis, omnibus utrinque glabris siccis subcoriaceis opacis subtilissime pellucido-punctulatis, nervo centrali subtus prominente ad  $\frac{1}{2}$  longit. nervos utrinque 5 alternos subadscendentes et supremos adscendentes ad apicem ductos infimos e basi solutos mittente, petiolo glabro ad  $\frac{3}{4}$  longit. vaginante, pedunculo sursum puberulo quam petiolus paulo brevior, amento quam folium haud multum brevior mucronato, bractæ apice truncato-peltatæ vertice subrotundo margine fusce hirtio pedicello subcucullato fusce hirtio, ovario ovato glabro, bacca glabra sub-4-angula. ♀ In Minas-Geraës (Ackermann! in h. Mart. ex h. Vindob.). Peltobryon exserens Miq.! in Mart. flor. Bras. fasc. 11. p. 31. t. 5. f. 2. Rami ramulique glabri. Limbi infer. 0,2 longi, 0,11 lati, super. 0,14 longi, 0,05 lati, petioli 0,015 longi. Amenta mat. 0,005 crassa. (v. s.)

B. *Sylus nullus*.

+ Bacca fere tetragona vel tetragona.

a. *Bractea apice peltata, truncato-peltata vel truncata*.

1° Folia penninervia.

\* Centralis nervus tota vel fere tota longitudine nervos mittens.

83. *P. MATHEWSII*, foliis breviter petiolatis apice attenuatis subacuminatis basi æquali subattenuatis acutis supra scabris subtus hirtio-pubescentibus coriaceis opacis, nervo centrali subtus prominulo utrinque fere tota longitudine nervos 11-13 alternos subadscendentes subtus vix prominulos mittente, petiolo hirtio, pedunculo petiolum æquante hirtello, bractæ pelta triangulari margine

hirsuta. ♂ In Peruvia prope Chachapoyos (Mathews n. 3224! in h. Kew.). Ramuli molliter hirtopubescentes. Limbi 0,12-0,14 longi, 0,03-0,035 lati. Petioli 0,005 longi. Amenta ante anthesim 0,04 longa, 0,003 crassa. (v. s.)

84. *P. STELLIPILUM*, foliis breviter petiolatis ovato-oblongis apice anguste acuminatis, acumine acuto, basi subinæquali rotundato-obtusis supra sparse et subtus densius pilis stellatis pubescentibus ciliolatis, junioribus fusce tomentosis firmo-membranaceis opacis pellucido-punctulatis, nervis subtus prominulis, centrali utrinque fere tota longitudine nervos circ. 13 alternos patulo-subadscendentes mittente, petiolo ad  $\frac{1}{2}$  longit. vaginante basi dorso sursum utrinque dense hirtello-tomentoso, pedunculo hirtello tomentoso petiolum æquante, amento florente quam folium multoties brevior, bractæ pelta semilunari margine villosa, filamentis exsertis. ♂ In sylvis ad flum. Solimoës prov. Rio Negro (Mart. in h. monac.). Artanthe stellipila Miq. in Mart flor. Bras. fasc. 11. p. 58. Rami fusce hirtello-tomentosi. Limbi 0,28 longi, 0,115 lati. Petioli 0,006 longi. Amenta florent. 0,004 crassa. (v. s.)
85. *P. MOLLIPILOSUM*, foliis breviter petiolatis elliptico-lanceolatis apice longiuscule acuminatis, acumine acuto mucronulatoque, basi subæquali acutis supra sparse pilis brevissimis pilosis subtus præsertim ad nervos sparse molliter pubescentibus firmule membranaceis subopacis pellucido-punctulatis, centrali nervo utrinque fere tota longitudine nervos circ. 8 alternos patulo-adscendentes mittente, petiolo basi ima vaginante utrinque pubescente, pedunculo hirsuto-pubescente petiolum æquante, amento quam folium multoties brevior, bractæ pelta triangulari margine canescenti-hirsuta, ovario puberulo, bacca lateraliter compressa. ♂ In sylvis ad Ceari prov. Rio Negro (Mart. in h. monac.). Artanthe mollis Miq. in Mart. fl. Bras. fasc. 11. p. 58. Rami pilis recurvulis puberuli, ramuli densius hirtello-pubescentes. Limbi 0,15 longi, 0,036 lati. Petioli 0,005 circ. longi. (v. s.)
86. *P. MAYNASIANUM*, foliis breviter petiolatis oblongo-ellipticis apice subprotracto-acuminatis, acumine acuto, basi leviter inæquali acutis supra glabris subtus villosulis rigidulo-membranaceis subopacis pellucido-punctulatis, nervis subtus prominulis, centrali utrinque fere tota longitudine nervos circ. 15 alternos subpatulo-adscendentes supremos ad apicem ductos mittente, petiolo villosulo basi vaginante, pedunculo parce villosulo petiolum æquante, amento florente folium subæquante, bractæ vertice truncato-peltatæ pelta triangulari margine ciliolata pedicello ciliolato dorso subproducto. ♂ In Peruvia ad Maynas et Cuchero (Poeppig? n. 2553! et 1463! in h. Vindob.). Rami glabri, ramuli juniores ad nodos villosi. Limbi 0,19 longi, 0,075 lati. Petioli 0,007 longi. (v. s.)
87. *P. CUMARALENSE*, foliis brevissime petiolatis suboblique lanceolatis apice acuminatis, acumine acuto mucronulatoque, basi leviter inæquali cuneatis utrinque glabris rigidulis opacis subtus crebre nigro-punctulatis supra nitidulis, nervo centrali subtus prominulo utrinque nervos circ. 9 alternos superiores breviores subadscendentes patulos inferiores utrinque 4 longiores patulo-subadscendentes mittente, petiolo glabro basi ima vaginante, amento mat. quam folium fere quadruplo brevior cylindrico obtuso, pedunculo glabro petiolum duplo superante, bractæ truncato-peltatæ pelta triangulari margine griseo-ciliata pedicello nigro-punctulato dorso ciliato, bacca nigra glabra. ♂ In nova Granata ad Cumaral (h. Vindob. l.) Rami glabri. Limbi 0,012 longi 0,04 lati. Petioli 0,001 longi. Amenta matura 0,006 crassa. Bacca 0,001 crassa. (v. s.)
- 88 (1). *P. AMPLIFOLIUM*, foliis amplis petiolatis ovato-oblongis apice breviter acute acuminatis basi leviter inæquali obtusis utrinque glabris coriaceis opacis parce pellucido-punctulatis nitidulis, nervis subtus prominulis, centrali utrinque fere tota longitudine nervos circiter 13 alternos nervulis fortioribus sæpe intermixtos patulo-subadscendentes mittente, petiolo glabro pedunculum glabrum fere duplo superante ad limbi basim et paulo ultra vaginante, amento florente quam folium multoties brevior, bractæ apice truncato-peltatæ vertice triangulari nudo pedicello cucullato-ciliolato, bacca lateraliter compressa

(1). Vide sp. sub n. 293.

- glabra. ♀ In parte meridionali prov. Bahia (Blanchet n. 3160! A. in h. DC.), Rio Janeiro (Gaudich. n. 87! in h. DC.). Ramuli glabri. Limbi 0,31 longi, 0,09 lati. Petioli 0,023 longi. (v. s.)
89. *P. MACROPHYNCHON*, foliis petiolatis sublanceolato-oblongis apice apiculatis basi leviter inæquali acutis obtusius utrinque glabris subcoriaceis opacis parcellissime pellucido-punctatis utrinque et præsertim subtus glanduloso-punctulatis, nervis subtus prominulis, centrali fere tota longitudine nervos circiter 8 patulo-subadscendentes alternos mittente, petiolo glabro ad limbum usque vaginante, pedunculo petiolum superante, amento apice longiuscule mucronato quam folium multoties brevius, bractæa pelta triangulari carnosula margine subaureo-ciliata, stigmatibus abbreviatis. ♀ In sylvis primævis prov. Paraensis (Mart. in h. monac.). *Artanthe macrorhyncha* Miq. in Mart. flor. Bras. fasc. 11. p. 61. Rami ramulique glabri. Limbi 0,12 longi, 0,045 lati. Petioli 0,018 longi. Amenta matura 0,02 longa, 0,004 crassa (v. s.)
90. *P. JAPURENSE*, foliis breviter petiolatis subelliptico-oblongis vel lanceolato-oblongis apice subacuminatis basi leviter inæquali acutis supra glabris subtus ad nervos parce hirtellis firmulis opacis parce pellucido-punctulatis, nervis subtus prominulis centrali utrinque tota longitudine nervos circ. 10 nervulis fortioribus aliquando intermixtos alternos subalternosve patulo-subadscendentes mittente, petiolo ad limbum usque vaginante dorso glabro, pedunculo glabro petiolum paulo superante, amento maturo quam folium multoties brevius, bractæa pelta semilunari margine subaureo lacero-hirtella, bacca glabra. ♀ In Brasiliæ prov. Rio Negro (Mart. n. 309! in h. monac., specimine n. 310 excluso verisimiliter speciem novam præbente). *Artanthe japurensis* Miq. in Linn. v. 20. p. 168 et in Mart. flor. Bras. fasc. 11. p. 59. Rami glabri. Limbi 0,2 circ. longi, 0,085 lati. Petioli 0,006 longi. Amenta matura fere 0,005 crassa. (v. s.)
91. *P. INDECORUM* (Kunth! olim), foliis petiolatis oblongo-ellipticis apice anguste acuminatis, acumine acuto, basi æquali attenuatis acutis supra glabris subtus ad nervos villosos-barbatis membranaceis, nervo centrali subtus prominulo utrinque fere tota longitudine nervos 13 alternos arcuatim adscendentes mittente, petiolo villosos-barbato, pedunculo petiolum æquante, bractæa apice inflexo-peltatæ vertice triangulari glabro. ♂ In Peruvia nemoribus Pangoa Mathews n. 1144! in h. Kew. et Poeppig! in h. Berol.). *Schilleria indecora* Kunth! in Linn. v. 13. p. 722. *Artanthe polyneura* Miq.! syst. p. 504 et in Hook. Lond. journ. bot. v. 4. p. 467. Ramuli juniores villosos-barbati vel pilosi. Limbi 0,17 longi, 0,055 lati. Petioli 0,01 longi. Amenta florent. 0,06 longa, 0,002 crassa. (v. s.)
92. *P. LANCEOLATUM* (Ruiz et Pav. flor. peruv. 1. p. 36. t. 61. f. b), foliis breviter petiolatis oblongo-lanceolatis apice acuminatis, acumine subobliquo acuto mucronulatoque, basi æquali attenuato-acutis vel cuneatis utrinque glabris fusco-punctulatis opacis membranaceis, nervo centrali subtus prominulo nervos utrinque tota longitudine 8-10 alternos subpatulos subtus subprominulos nervulis fortioribus intermixtos mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum æquante vel paulo superante, bractæa pelta triangulari vel semilunari margine luteo-hirsuta. ♀ In Peruvia (h. Pav. in h. DC. et Boiss.), prope Maynas ad flumen Amazonum (Poeppig n. 1434! in h. DC. et Berol.), nemoribus Chacahuassi (R. et P. l. c.), et Caracas (Bredemeyer! in h. Vindob. ex h. Jacquin.).—Vahl enum. p. 313. *P. evonymifolium* Kunth! olim. *Steffensia evonymifolia* Kunth! in Linn. v. 13. p. 624 (fide syn. Poeppig.). *Artanthe Brunoniana* Miq.! syn. p. 509. Ramuli glabri, fusco-punctati. Limbi 0,12-0,14 longi, 0,03-0,04 lati. Petioli 0,003 longi. Amenta florent. 0,003 crassa. (v. s.)
- β, *latifolium*, foliis multo latioribus (0,17 longis, 0,05 latis) oblongo-lanceolatis. — In Brasilia prope San Carlos ad Rio Negro (Spruce n. 3090! in h. Kew.).
93. *P. TRICUSPE*, foliis breviter petiolatis e basi cuneata trilobis, lobis cuspidatis cuspidate mucronulato, utrinque glabris nigro-punctatis membranaceis, centrali nervo subtus subprominulo utrinque tota longitudine nervos 12-15 alternos infimos adscendentes cæteris patulos mittente, petiolo glabro, pedunculo

- petiolum superante, bractæ pelta triangulari margine hirsuta. ♀ In Nova Granata (Seeman n. 1068! in h. Kew.) et prov. Barbacoas in planitie (Triana n. 54! in h. suo), Darien merid. (Miq. in Seeman bot. Herald p. 200). *Artanthe tricuspis* Miq. ! in Seem. l. c. t. 42. Ramuli glabri. Limbi 0,1-0,12 longi, 0,05-0,06 lati. Petioli 0,008 longi. Pedunculi 0,015 longi. Amenta florent. 0,01 longa, vix 0,003 crassa. Verisimiliter *P. lanceolati* R. et P. forma. (v. s.)
94. *P. EGENSE*, foliis brevissime petiolatis amplis subelliptico-oblongis apice brevissime acuminatis, acumine acutiusculo, basi ima æquali acutis adultis utrinque glabris (junioribus subtus pubescentibus) rigido-subcoriaceis opacis, centrali nervo subtus prominulo utrinque tota longitudine nervos 14 alternos patulo-subadscendentes mittente, petiolo glabro ad limbum usque vaginante, pedunculo glabro petiolum multoties superante, amento maturo quam folium quadruplo breviori crasso cylindrico recto, bractæ pelta semilunari margine fusco-villosa, bacca glabra. ♀ In sylvis ad Ega et Maribi juxta prov. Rio-Negro et in prov. Paraensi (Mart. ! in h. monac.), Ilheos (Wawra et Maly n. 28! fide Wawra in bot. Ergeb. p. 138). *Peltobryon Martianum* Miq. ! in Linn. v. 20. p. 133 et in Mart. flor. Bras. fasc. 11. p. 29. t. 5. f. 1. Rami glabri, ramuli juniores pubescentes. Limbi 0,21 longi, 0,08 lati. Petioli 0,003 longi. Amenta matura 0,015 crassa. (v. s.)
95. *P. BENTHAMIANUM*, foliis breviter petiolatis oblongo-ovatis apice longe acuminatis, acumine acuto, basi subattenuatis obtusiusculis utrinque glabris membranaceo-firmis, centrali nervo subtus prominulo utrinque nervos 12-13 alternos nervulis fortioribus intermixtos patulo-adscendentes subtus prominulos mittente, petiolo glabro, pedunculo petiolum æquante glabro, amento quam folium multum breviori recto, bractæ subconchæformis glabræ vertice tuberculiformi triangulari. ♀ Prope Caracas (Linden n. 216! in h. Kew.). *Artanthe Benthamiana* Miq. ! in Hook. Lond. journ. bot. v. 4. p. 447. Ramuli glabri. Limbi 0,22 longi, 0,11 lati. Petioli 0,015 longi. Amenta florent. 0,003 crassa. (v. s.)
96. *P. CALOCOMA*, foliis breviter petiolatis ellipticis apice breviter attenuatis acutis basi æquali subattenuatis acutiusculis utrinque glabris nigro-punctulatis membranaceis, nervo centrali utrinque tota longitudine nervos 15 subalternos patulo-adscendentes mittente, petiolo glabro nudo, pedunculo petiolum subæquante glabro, bractæ pelta triangulari margine hirsuta. ♀ In Nova Granata prope La Paila (Holton n. 29! in h. Kew. et Seeman n. 1067! in h. Kew.), Darien meridion. (Miq. in Seem. bot. Herald p. 199). *Artanthe calcoma* Miq. ! in bot. Herald l. c. Ramuli glabri. Limbi 0,19 longi, 0,11 lati. Petioli 0,015 longi (1). (v. s.)
97. *P. TOCACHEANUM*, foliis brevissime petiolatis elliptico-oblongis apice breviter acuminatis, acumine acuto, basi æquali acutis utrinque glabris coriaceis opacis, nervo centrali utrinque tota longitudine nervos 12-15 patulo-subadscendentes mittente, nervis subtus prominulis, petiolo glabro, amento florente quam folium triplo breviori cylindrico crasso, bractæ pelta semilunari margine fusce hirsuta, stigmatibus abbreviatis. ♀ In Tocache Huallaga (Endlicher! in h. Vindob.) et Peruvia (Mathews n. 1707! in h. Vindob.). Rami ramulique glabri. Limbi 0,25-0,3 longi, 0,07-0,12 lati. Petioli 0,005 longi. Amenta flor. 0,005 crassa. (v. s.)
98. *P. CORDATUM* (Cas. DC. in Seem. journ. of. bot. 1866. p. 213), foliis subsessilibus oblongo-ovatis apice attenuatis acutis basi æquali cordatis utrinque glabris margine ciliolatis membranaceis, nervo centrali subtus prominulo utrinque tota longitudine nervos 8-10 alternos patulo-adscendentes subtus prominulos nervulis fortioribus intermixtos mittente, petiolo brevissimo glabro subnullo, pedunculo brevissimo, bractæ pelta triangulari-subquadragulari margine hirsuta. ♀ In Peruvia orientali prope Tarapoto (Spruce n. 4847! in h. Kew. et DC.). Ramuli glabri. Limbi 0,23 longi, 0,11 lati. Amenta nondum efformata. (v. s.)
99. *P. RIGIDUM*, foliis brevissime petiolatis anguste oblongis apice acutatis basi

---

(1) Præsens sp. ex observationibus recentioribus speciminis Rudgeani authenticici in h. Brit. Mus. forsan ad *P. augustum* Rudge pertinet.

satis inæquali acutiusculis supra glabris nitidulis subtus dense subcinereo-pubescentibus rigide subcoriaceis opacis punctulatis, nervo centrali subtus prominente utrinque nervos tota longitudine 9-10 infimos subadscendentes sub medio per anastomoses evanidos superiores adscendentium subpatulos nervulosque fortiores mittente, petiolo dorso dense pubescente ad limbi lobum minorem usque vaginante, alis diutius persistentibus, pedunculo dense puberulo petiolum triplo superante, amento post anthesin folium dimidium superante, rhachi hirtella, bractæ apice truncato-peltatæ pelta semilunari margine subcinereo-hirsuta, ovario glabro, stigmatibus brevissimis. ♀ In Cuba (Wright n. 2271! in h. DC.). *Artanthe geniculata* Griseb. cat. pl. cub. p. 65. Rami ramulique dense subcinereo-pubescentes. Limbi 0,095 longi, 0,022 lati. Petioli 0,004 longi. Amenta post anthesin 0,0015 crassa. (v. s.)

100. *P. SWARTZIANUM*, foliis brevissime petiolatis oblongo-lanceolatis apice acutatis submucronulatis basi valde inæquali obtusis supra glabris subtus pubescenti-hirtellis coriaceis opacis, nervo centrali subtus prominulo utrinque tota longitudine nervos 10-12 subpatulos nervulosque fortiores mittente, petiolo ad limbum usque vaginante dorso pubescenti-hirtello, alis linearibus cito deciduis, pedunculo petiolum æquante puberulo, amento folium æquante, bractæ pelta triangulari margine aureo-hirsuta, bacca glabra lateralter subcompressa. ♀ In Santo Domingo (Poiteau! in h. Deless.) et Cuba (Wright n. 2269. 2270! fide Griseb. in Cat. pl. cub. p. 65). *Artanthe Swartziana* Miq.! syst. p. 490, Griseb. l. c. Ramuli puberuli. Limbi 0,13 longi, vix 0,04 lati. Petioli vix 0,002 longi. Amenta matura 0,004 crassa. (v. s.)

101 (1).

102. *P. OBUMBRATUM*, foliis brevissime petiolatis late oblongis apice breviter acuminatis, acumine acuto, basi valde inæquali rotundatis, lobo longiore cum petiolo adnato, supra glabris subtus præsertim ad nervos pubescenti-hirtellis coriaceis opacis, nervo centrali subtus prominente utrinque tota longitudine nervos 10-12 subpatulos subtus prominulos nervulosque fortiores mittente, petiolo glabro ad limbi lobum minorem usque vaginante, alis cito deciduis, pedunculo glabro petiolum subæquante, amento folium subæquante, bractæ pelta triangulari margine fusco-hirta, bacca glabra lateralter compressa. ♀ In sylvis ad cataractas Cupatenses in flumin. Japura prov. Rio Negro (Mart.! in h. monac.). *Artanthe obumbrata* Miq.! in Linn. v. 20. p. 142 et in Mart. flor. Bras. fasc. 11. p. 52. t. 19. f. 1. Rami cinerascetes. Ramuli nascentes subvillosi. Limbi 0,27 longi, 0,11 lati. Petioli pars libera vix 0,003 longa. Amenta matura 0,003 crassa. (v. s.)

103. *P. TUBERCULATUM* (Jacq. ic. rar. v. 2. p. 2. t. 210), foliis petiolatis oblique ovatis ovato-oblongis oblongisque apice acutis vel subacuminatis basi valde inæquali rotundatis rarissime subcordatis, latere majore cum petiolo fere omnino vel dimidio adnato, utrinque glabris vel subtus ad nervos subtiliter puberulis sæpe subtus fusce vel nigro-punctulatis membranaceis subopacis parcissime pellucido-punctulatis, nervo centrali subtus prominulo utrinque tota longitud. nervos 8-10 subtus subprominulos subpatulos nervulis fortioribus intermixtos mittente, petiolo glabro vel dorso subtilissime hirtello ad limbi latus minus usque vaginante, alis linearibus angustis, pedunculo petiolum subæquante glabro vel subtiliter puberulo, amento folium æquante vel superante, bractæ pelta triangulari subrhomboidali margine hirsuta pedicello lato subcucullato, bacca glabra lateralter compressa. ♂ Late per novum orbem sparsa. In Peruvia (Poeppig n. 3114! in h. DC. et 1036! in h. Vindob.), Maynas, ad rivulos umbrosos (Poeppig n. 2123! in h. Boiss. et 3114! in h. DC.), Brasiliæ prov. Para prope Santarem (Spruce n. 31! in h. DC.) et Rio Janeiro (Weddell n. 596! in h. DC.), in sylvis primævis prope Rio Janeiro et in prov. Maragnan et sylvis udis sec. flumen Itapicura prope S. Miguel prov. Maragnanensis (Mart.! in h. monac.), prov. Ceara (Gardner n. 1846! in h. Boiss.), ad rio Maranhao Brasiliæ (Pohl n. 2195! et 1558! in h. Vindob.), ad ripas locosque humidus ad Maiquetia (Wagner n. 128 in Linn. v. 26. p. 632),

(1) Species hic erat propter observationem speciminis authentici nuperrime factam nunc omissa.

Ruiz in h. Willd.<sup>1</sup> n. 635). — Vahl enum. v. 1. p. 320. Schilleria obovata Kunth! in Linn. v. 13. p. 686. Artanthe obovata Miq.<sup>1</sup> syst. p. 508. Ramuli glabri. Limbi 0,12-0,13 longi, 0,04-0,045 lati, petioli 0,01-0,015 longi. Amenta matura 0,055-0,07 longa, 0,005-0,007 crassa. Pedunculi 0,002 longi. (v. s.)<sup>1</sup>

\*\* Centralis nervus e parte longitudinis nervos mittens.

107. *P. MAROWYNENSE*, foliis brevissime petiolatis elliptico-oblongis apice acuminatis, acumine acuto acutiusculove mucronulato, basi leviter inæquali acutis supra glabris junioribus puberulis cito basi excepta ad nervum centalem appresse puberulis subtus ad nervos nervulosque appresse puberulis margine ciliolatis membranaceis opacis pellucido-punctulatis, nervis subtus subprominulis, centrali utrinque ad  $\frac{1}{2}$  longitudinis nervos circiter 5 subpatulo-subadscendentes alternos sursumque tota longitudine nervulos fortiores aliquot mittente, petiolo basi ima vaginante basi dorso sursum utrinque pubescente, pedunculo pubescente petiolum æquante, amento florente quam folium quadruplo brevior, bractæ pelta semilunari margine puberula. ♂ In monte Cassinisvard Marowyna Guyanæ (Wulschlaegel n. 1562! in h. Mart.). Ramuli pilis retrorsis pubescentes. Limbi 0, 125 longi, 0,05 lati. Petioli 0,005 longi. Amenta florentia 0,002 crassa. (v. s.)
108. *P. GABRIELANUM*, foliis brevissime petiolatis elliptico-oblongis apice longe acuminatis, acumine acuto, basi leviter inæquali acutiusculis acutisve supra glabris lævibus subtus dense molliter pubescentibus rigidulo-membranaceis subopacis creberrime pellucido-punctulatis, centrali nervo utrinque ad  $\frac{1}{2}$  longitud. nervos 6 alternos patulo-subadscendentes nervulosque fortiores mittente, petiolo hirtello-pubescente, pedunculo pubescente petiolum æquante, amento mucronato florente quam folium multoties brevior, bractæ brevissime pedicellatæ pelta triangulari carnosula margine hirtella, ovario.... ♂ In Guyana gallica (Gabriel n. 1802! in h. Deless. Arthante Gabriellana Miq.<sup>1</sup> syst. p. 476. Ramuli pubescentes. Limbi 0,18 longi, 0,08 lati. Petioli 0,005 longi. (v. s.)
109. *P. CUMBOTIANUM*, foliis brevissime petiolatis amplis elliptico-oblongis apice breviter acuminatis, acumine obtusiusculo, basi inæquali obtusis latere majore rotundato minore acuto utrinque dense scabrido-pubescentibus subtus ad nervos pilis longioribus villosis margine ciliolatis firmule membranaceis opacis pellucido-punctulatis, nervis subtus prominulis centrali utrinque ad  $\frac{1}{2}$  longitud. nervos circ. 9 alternos subadscendentes supremos ante apicem per anastomoses evanidos infimos subtiliores magis patulos sursumque nervulos fortiores mittente, petiolo fere ad limbum usque vaginante dorso villosa, pedunculo villosa petiolum multoties superante, amento florente folium dimidium vix æquante, bractæ truncato-peltatæ vertice semilunari antice calloso-subtuberculato margine exteriori ciliolato pedicello lato cucullato intus juniiori hirtello margine et dorso basi hirtello, ovario.... ♂ In Venezuela inter Blanca et Cumboti alt. 5000 p. (Fendler n. 2571! in h. DC.). Rami ramulique dense villosi. Limbi 0,2 longi, 0,11 lati. Petioli 0,01 longi. Pedunculi 0,04 longi. (v. s.)
110. *P. NERVULOSUM*, foliis breviter petiolatis oblongo-lanceolatis apice acuminatis, acumine acuto, basi æquali cuneatis, supra glabris subtus ad nervos appresse puberulis rigidis opacis subtiliter pellucido-punctulatis, centrali nervo subtus prominente nervos utrinque ad  $\frac{1}{2}$  longitud. circ. 9 alternos subadscendentes nervulis fortioribus sæpe intermixtos supremos ad apicem ductos mittente, petiolo ad  $\frac{1}{2}$  longitud. vaginante, pedunculo glabro petiolum æquante, amento quam folium brevior, bractæ margine et dorso sublaevide hirtellæ vertice truncato triangulari nudo, bacca lateraliter compressa glabra. ♂ In Brasilia (Riedel n. 2881 in h. Petrop.). Rami ramulique glabri. Limbi 0,24 longi, 0,055-0,065 lati. Petioli 0,007 longi. (v. s.)
111. *P. DUMOSUM* (Rudge! pl. Guy. p. 13. t. 14), foliis breviter petiolatis suboblongo-lanceolatis apice acuminatis, acumine acuto, basi subinæquali acutis utrinque pilis parvis hirsutis supra brevioribus subtus ad basin pilorum quasi punctatis, centrali nervo utrinque ad  $\frac{1}{2}$  longitud. nervos circ. 6 alternos supremos ad apicem ductos mittente, pedunculo petiolum parum superante quam folium triplo brevior, bractea vertice truncato semilunari-triangulari



ciliato, bacca obovato-trigona glabra. ♀ In Guyana (Rudge in h. Mus. brit.!). Rami ramulique glabri. Limbi 0,17 longi, 0,068 lati. Petioli 0,006 longi. (Ex Rudge l. c.)

112. *P. FRASERI*, foliis subsessilibus obovato-oblongis apice longiuscule suboblique protracto-acuminatis, acumine acuto mucronulatoque, basi breviter inæquali cordulatis supra valde bullatis scabrisque subtus lacunosis ad nervos et nervulos pilis longis villosis et inter nervulos pilis brevibus scabriusculis margine ciliolatis, nervis subtus prominulis centrali utrinque ad  $\frac{1}{2}$  longitud. nervos 10-11 alternos adscendentium subadscendentes nervulis fortioribus intermixtos supremos ante apicem per anastomoses evanidos mittente, petiolo brevissimo longe villoso, amento adhuc juniori quam folium multoties brevius mucronato, bractæ truncato-peltatæ pelta elliptica margine dense villosa pedicello villoso angusto. ♀ In Ecuador (Fraser! in h. DC.). Rami a-mulique villosi. Limbi 0,2 longi, 0,065 lati. Petioli vix 0,002 longi. (v. s.)
113. *P. DASYOURA*, foliis petiolatis oblongo-ellipticis apice acuminatis, acumine obtusiusculo, basi æquali acutiusculis coriaceis opacis supra scaberrimis verrucosis subtus scabre pubescentibus, nervo centrali utrinque ad  $\frac{1}{2}$  longitud. nervos 8 alternos adscendentes supremos fere ad apicem ductos mittente, petiolo hirsuto, pedunculo hirsuto petiolum bis terve superante, bractæ subpeltatæ pelta anguste triangulari margine longe villosa pedicello longe villoso, ovario pubescente, stigmatibus linearibus. ♀ In Peruvia (Matthews n. 1716 in h. Kew.). *Artanthe dasyoura* Miq.! in Hook. Lond. Journ. bot. v. 4. p. 452. Ramuli hirsuti. Limbi 0,19 longi, 0,07 lati. Petioli 0,015 longi. Pedunculi 0,035 longi. Amenta florentia 0,19 longa, 0,006 crassa. (v. s.)
114. *P. ACUTIFOLIUM* (Ruiz et Pav. pl. peruv. p. 38. t. 64. f. a), foliis brevissime petiolatis ovatis apice longe acuminatis, acumine acuto mucronulatoque, basi valde inæquali profunde auriculatim cordatis, lobis rotundatis lobo majore petiolum velante, supra scabriusculo-hirtis subtus lacunosis et molliter hirsutopubescentibus rigido-coriaceis opacis, nervis subtus prominulis centrali utrinque ad  $\frac{1}{2}$  longitudinis nervos subadscendentes latere minore 8 majore 10 supremos fere ad apicem ductos mittente, petiolo hirsuto, pedunculo molliter hirsuto petiolum superante, amento cylindrico quam folium multum brevius recto, bractæ pelta triangulari-orbiculari margine hirsuta. ♀ In Cuchero et Pozuzo sylvis (Ruiz et Pav. l. c.), Andium peruvianorum montibus (Ruiz in h. Willd. n. 678! et in h. Berol.), provinc. Chachapoyas (Matthews n. 1715! in h. Kew.). — Vahl enum. v. 1. p. 325; Opiz in Reliq. Haenk. fasc. 3. p. 156. *Steffensia acutifolia* Kunth! in Linn. v. 13. p. 650. *Artanthe acutifolia* Miq.! syst. p. 428, illust. p. 76. t. 78 et in Hook. Lond. Journ. bot. v. 4. p. 455. Ramuli pilis longiusculis crispulis villosis. Limbi 0,26 longi, 0,11 lati. Petioli 0,05 longi. Amenta florentia 0,04 longa, 0,004 crassa. (v. s.)
115. *P. LECHLERIANUM*, foliis petiolatis oblongo-ellipticis apice acuminatis, acumine acuto, basi parum inæquali cordulatis supra scabriusculis subtus molliter pubescentibus membranaceo-firmis margine ciliolatis, centrali nervo subtus prominulo utrinque ad  $\frac{1}{2}$  longitud. nervos 8-6 alternos subadscendentes mittente, petiolo dense molliter villoso, pedunculo villoso petiolum æquante, amento elongato recto apice mucronulato, bractæ pelta triangulari margine hirsuta. ♀ In Peruvia (Lechler, n. 2438! in h. Kew.). Ramuli dense molliter villosi. Limbi 0,2 longi, 0,08 lati. Petioli centim. vix longi. Amenta florent. 0,12 longa, 0,002 crassa. (v. s.)
116. *P. CROCATUM* (Ruiz et Pav.! fl. peruv. 1. p. 35. t. 55. f. a), foliis breviter petiolatis oblongo-ellipticis apice acuminatis basi æquali vel subinæquali acutis acutiusculisve supra scabre hirtis sæpe bullatis subtus molliter dense pubescentibus coriaceis opacis, nervo centrali subtus prominulo utrinque ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitudinis nervos 7-9 nervulis fortioribus intermixtos subadscendentes mittente, petiolo pubescente, pedunculo pubescente petiolum paulo superante, amento maturo quam folium paulo brevius, bractæ pelta angusta elliptica margine longe hirsuta, bacca glabra lateraliter subcompressa. ♀ In Pillao et Chacahuassi nemoribus (R. et P. l. c.), Peruvia (Matthews n. 1711! et 1712! in h. Kew., Ruiz! in h. Willd. n. 657 et in h. Berol.), monte Pichincha alt. 1700 hexap. (Francis Half! in h. Berol. cum nomine *P. nubige-*

125. *P. BARYANUM*, foliis longiuscule petiolatis inferioribus subrotundis apice brevissime acuminatis, acumine acuto, basi æquali cordatis superioribus ovatis apice breviter acuminatis basi truncato-rotundatis omnibus supra glabris lævibus albedo-punctulatis subtus ad nervos subtiliter hirtellis membranaceis subopacis subtilissime pellucido-punctulatis, nervis subtus prominulis, centrali utrinque ad  $\frac{1}{2}$  longit. nervos 7 subalternos ante apicem per anastomoses evanidos sursumque nervulos fortiores (inferis nervis in foliis inferioribus utrinque 2-3 e basi vel fere e basi ortis) mittente, petiolo glabro ad  $\frac{1}{2}$  longit. vaginante, pedunculo glabro quam petiolus multum brevior, amento florente folium dimidium æquante, bracteæ apice truncato-peltatæ pelta triangulari margine subaureo-ciliolata, ovario glabro. ♀ In Brasilia (Riedel n. 49! in h. Petrop. et Riedel et Langsdorff n. 103 ibid.). Ramuli glabri. Limbi superiores 0,2 longi, 0,14 lati, inferiores 0,18 longi latique. Petioli 0,04 longi. Amenta florentia 0,005 crassa. (v. s.)

126. *P. AMAZONICUM*, foliis longiuscule petiolatis late ovato-acuminatis, acumine acuto, inferioribus basi æquali cordatis superioribus rotundato-truncatis utrinque glabris subtus ad nervos pubescenti-tomentellis rigidulo-flaccidis subopacis subtiliter pellucido-punctulatis, nervis subtus prominentibus, centrali nervos utrinque ad  $\frac{1}{2}$  longit. in foliis inferioribus 6-7 in superioribus 5-6 subadscendentes oppositos inferiores magis patulos supremos ad apicem ductos mittente, petiolo ad limbum usque vaginante dorso subtiliter fuscescenti-tomentello, amento quam folium dimidio brevior, pedunculo quam petiolus brevior, bracteæ apice truncato-peltatæ pelta triangulari margine aureo-pubescente pedicello angusto, bacca glabra vertice ciliolata. ♀ In Costa de Matary in ripa flumin. Solimoes, sylvis prope Apapuris flumen in prov. Rio Negro et sylvis ad flumen Amazonum prov. Paraensis (Mart.! in h. monac.). Artanthe amazonica Miq. in Linn. v. 20, p. 146 et in Mart. flor. Bras. fasc. 11. p. 39. t. 11. Rami fuscescentes; ramuli glabri. Limbi 0,25 longi, 0,15 lati. Petioli 0,04 longi. Amenta matura vix 0,003 crassa. (v. s.)

127. *P. ESTRELLENSE*, foliis petiolatis ovato-acuminatis, acumine acuto, basi subinæquali rotundato-cordulatis utrinque fusco-pubescentibus rigidis opacis pellucido-punctulatis, nervis subtus prominulis, centrali utrinque ad  $\frac{1}{2}$  longit. nervos 5-6 alternos subadscendentes supremos ad apicem ductos infimos magis patulos mittente, petiolo limbum usque vaginante fusco-pubescente, amento quam folium dimidio brevior, pedunculo puberulo petiolum æquante, bracteæ apice truncato-peltatæ vertice triangulari margine subaureo-pubescente pedicello latiusculo fusco subaureo-pubescente. ♀ In sylvis ad Regiato Jacaremi prov. San Paoli (Mart. in h. monac.), Brasilia in Serra d'Estrella (Pohl n. 4770! et 3465! in h. Vindob.). Artanthe fuscescens Miq. in Linn. v. 20. p. 144 et in Mart. flor. Bras. fasc. 11. p. 38. t. 10. Ramuli subtiliter fusco-puberuli. Limbi 0,17 longi, 0,105 lati. Petioli 0,015 longi. Amenta florentia 0,003 crassa. (v. s.)

β, *pilosius*, pilis densioribus rigidioribus, petiolis dense pilosis, amentis brevius pedunculatis. — In Santa Cruz do Terza (Pohl n. 3465! in h. Vindob.). (v. s.)

128. *P. MACROPHYLLUM* (Kunth! in H. et B. nov. gen. 1. p. 46), foliis petiolatis elliptico-oblongis apice acute acuminatis basi subinæquali rotundatis cordulatisve vel subattenuato-obtusis supra glabris subtus ad nervos subtiliter pilosis carnosulo-membranaceis, nervo centrali subtus prominulo utrinque nervos 5 subadscendentes alternos subtus prominulos supremos ad apicem ductos mittente, petiolo glabro limbum usque vaginante, pedunculo glabro petiolum subæquante. amento cylindrico obtuso mucronatove, bracteæ subcucullatæ apice truncato-peltatæ vertice rhomboidali glabro pedicello dorso hirsuto lateraliter producto, stigmatibus linearibus. ♀ In insulis Portorico (Wyder n. 260! in h. DC.), Martinica (Isert in h. Willd. n. 640! et Fouché! in h. Boiss.), Saint-Vincent (Goulding! in h. Hook.), ad Miradores (Linden n. 120! in h. Boiss.), in Peruvia (Mathews n. 1145 bis! in h. Kew.). *P. macrophyllum* Kunth synopsis. v. 1. p. 103. *Schilleria macrophylla* Kunth! in Linn. v. 18. p. 715. *Artanthe macrophylla* Miq. syst. p. 423 et in Hook. Lond. journ. bot. v. 4. p. 452. Ramuli glabri. Limbi 0,17-0,19 longi, 0,8-0,103-

0,125 lati. Petioli 0,015 longi. Amenta matura 0,05 circ. longa, 0,003 circ. crassa. (v. s.)

129. *P. PANAMENSE* (Cas. DC. in Seem. Journ. of bot. 1866. p. 216), foliis brevissime petiolatis elliptico-lanceolatis apice breviter acuminatis, acumine acutiusculo, supra glabris lævibus subtus ad nervos puberulis subcoriaceis nitidis centrali nervo subtus prominulo utrinque ad  $\frac{1}{2}$ - $\frac{2}{3}$  longit. nervos 4-5 alternos subadscendentes subtus prominulos sursumque sæpe nervulos fortiores mittente, petiolo appresse retro-puberulo, pedunculo puberulo petiolum æquante, bractæ apice truncato-peltatæ vertice angusto semilunari-triangulari margine ciliolato bractearum pedicellis latis inter se dorso productis connatis foveolas margine ciliolatas sic formantibus, bacca immersa vertice hirsuto-pubescente lateraliter compressa. ♀ In isthmo panamensi prope Chagres (Fendler n. 2701 in h. Kew.). Ramuli subglaucescentes. Limbi 0,1 longi, 0,055 lati. Petioli 0,005 longi. Amenta matura 0,03 longa, 0,004 crassa. (v. s.)

130. *P. ADENANDRUM*, foliis brevissime petiolatis subovato-oblongo-lanceolatis apice suboblique longe acuminatis, acumine acuto, basi subæquali acutis supra glabris subtus imprimis ad nervos hirtis margine subtiliter ciliolatis rigidulo-membranaceis creberrime pellucido-punctulatis subobscuris, centrali nervo subtus prominulo ad  $\frac{1}{2}$ - $\frac{2}{3}$  longit. utrinque nervos 6-7 subadscendentes supremos per anastomoses ante apicem evanidos mittente, petiolo basi vaginante dorso et sursum utrinque hirtio, pedunculo hirtello petiolum quadruplo superante, amento maturo quam folium multoties brevius mucronato, bractæ apice truncato-peltatæ vertice triangulari carnosulo margine ciliolato angulo anteriore carnosulo, bacca glabra. ♀ In Guyana gallica (Leprieur n. 1521 in h. Deless. et DC.). Artanthe adenandra Miq.! syst. p. 515. Ramuli hirti. Limbi 0,155 longi, 0,045 lati. Petioli 0,003 longi. Amenta matura 0,015-0,02 longa, 0,003 crassa. (v. s.)

131. *P. OBLONGIFOLIUM*, foliis brevissime petiolatis oblongo-ellipticis apice attenuatis acutis acutiusculis vel obtusiusculis basi subinæquali attenuato-obtusiusculis supra glabris subtus ad nervos appresse hirtellis membranaceis subtilissime pellucido-punctulatis, nervo centrali subtus prominulo utrinque ad  $\frac{2}{3}$  longitud. nervos 4-5 alternos patulo-subadscendentes mittente, petiolo ad  $\frac{2}{3}$  longit. vaginante dorso puberulo, pedunculo hirtello petiolum æquante, amento maturo quam folium dimidio vel duplo brevius mucronato, bractæ pelta triangulari-semilunari margine hirsuta, bacca glabra. ♀ In Guyana gallica (Leprieur! in h. DC.). Artanthe oblongifolia Klotzsch in h. Berol., Miq. Syst. p. 530 et in Hook. Lond. Journ. bot. v. 4. p. 322. Ramuli hirtelli. Limbi 0,07-0,11 longi, 0,03-0,04 lati. Petioli 0,003-0,005 longi. Amenta florentia 0,015 crassa. (v. s.)

β, *glabrum*, foliis et ramuli glabris. — In Guyana anglica (Schomb. in h. DC.)

132. *P. DECIPIENS*, foliis brevissime petiolatis oblongis vel subovato-oblongis apice acuminatis, acumine acuto mucronulatoque, basi inæquali auriculatim semicordatis, folio majore petiolum velante libero, minore rotundato, supra glabris læte viridibus subtus subtiliter puberulis ad nervos hirtellis membranaceis subopacis obsolete pellucido-punctulatis, nervo centrali utrinque ad  $\frac{2}{3}$  longit. nervos alternos latere minore 4 majore 5 subadscendentes supremos ante apicem per anastomoses evanidos mittente, petiolo hirtello, amento maturo quam folium multoties brevius, bractæ pelta semilunari margine villosa-hirsuta, bacca subglobosa glabra. ♀ In Mexici cordilleris alt. 1200 ped. (Galeotti n. 6016! in h. Deless.). Artanthe decipiens Miq.! syst. p. 462. Ramuli pubescentes. Limbi 0,14 longi, 0,065 lati. Petioli 0,005 longi. (v. s.)

133. *P. APICULATUM*, foliis breviter petiolatis subobovato-oblongis apice brevissime apiculatis acutis basi æquali acutis utrinque glabris rigidis opacis parce pellucido-punctulatis supra albido-punctulatis, nervis subtus prominulis, centrali utrinque fere ad  $\frac{2}{3}$  longitudinis nervos alternos 8 nervulis fortioribus sæpe intermixtos patulo-subadscendentes supremos ante apicem per anastomoses evanidos mittente, petiolo glabro, pedunculo glabro petiolum æquante,

amento maturo quam folium paullo brevior, rhachi hirtella, bractæ apice truncato-peltatæ vertice triangulari margine hirsuta, bacca parce hirtella, stigmatibus abbreviatis. ♀ In Brasilia (Riedel n. 435! in h. Petrop.). Rami ramulique glabri, abido-verruculosi. Limbi 0,12 longi, 0,04 lati. Petioli 0,012 longi. Amenta matura 0,003 crassa. (v. s.)

134. *P. ADENOPHORUM*, foliis brevissime petiolatis sublanceolato-oblongis apice breviter acuminatis, acumine acutiusculo mucronulato, basi inæquali acutis utrinque imprimis subtus præcipue ad nervos parce hirtellis margine subtiliter ciliolatis membranaceis subpellucidis utrinque glandulis fuscis conspersis, nervo centrali subtus prominulo utrinque ad  $\frac{1}{2}$  longitud. nervos 6-7 adscendentes nervulosque fortiores supremos ante apicem per anastomoses evanidos mittente, petiolo basi ima vaginante patentim hirtello dein glabrato glandulis fuscis utrinque consperso, pedunculo petiolum duplo superante glandulis fuscis consperso, amento quam folium multoties brevior, bractæ glandulis fuscis conspersæ pelta triangulari margine ciliolata pedicello margine ciliato. ♀ In Guyana gallica (Leprieur n. 153! in h. Cand. et Deless.), Cayenne (Martin! in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 468.) *Artanthe adenophora* Miq.! syst. p. 514 et in Hook. Lond. journ. bot. l. c. Ramuli glabri, glandulis fuscis conspersi. Limbi 0,18 longi, 0,075 lati. Petioli 0,004 longi. Amenta florentia 0,002 crassa. (v. s.)
135. *P. SCHLIMII*, foliis brevissime petiolatis lanceolato-oblongis apice acuminatis, acumine acutiusculo mucronulato, basi æquali acutis, supra glabris subtus præsertim ad nervos appresse puberulis membranaceis subopacis pellucido-punctulatis, centrali nervo subtus prominulo utrinque ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. nervos 4 alternos subtus prominulos patulo-subadscendentes margine per anastomoses adscendentes supremos per anastomoses ad apicem ductos nervulosque fortiores sæpe multos patulos mittente, petiolo pubescente basi ima vaginante, pedunculo puberulo petiolum æquante, amento quam folium multoties brevior, bractæ glabræ apice truncato-peltatæ vertice rhomboidali nudo, connectivo supra loculos producto. ♀ In Venezuela (Funck et Schlim n. 591! in h. Petrop. et Boiss.). Ramuli appresse pubescentes. Limbi 0,11 longi, 0,22 lati. Petioli 0,005 longi. (v. s.)
136. *P. OAXACANUM*, foliis breviter petiolatis oblongo-ovato-ellipticis acuminatis, acumine obtuso, basi inæquali acutis, utrinque glabris rigidulis opacis pellucido-punctulatis, nervis subtus prominulis, centrali utrinque ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitudinis nervos 6 alternos subadscendentes supremos ad apicem ductos infimos subtiliores mittente, petiolo glabro basi ima vaginante, amento folium æquante, bractæ apice truncato-peltatæ pelta semilunari margine exteriori ciliata. ♀ In Oaxaca (h. Vindob. n. 51!). Ramuli glabri, verruculis albidis minutis conspersi. Limbi 0,165 longi, 0,065 lati. Petioli 0,008 longi. Amenta florentia 0,003 crassa. (v. s.)
137. *P. MELANOSTICTUM*, foliis petiolatis subovato-oblongis apice paullo protracto-obtusiusculis basi subinæquali gibbose obtusis, latere majore rotundato minore acuto, utrinque glabris nitidulis supra scabriusculis rigido-coriaceis opacis utrinque glandulis fuscis conspersis, nervo centrali subtus prominulo ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. nervos 5 subadscendentes subtus prominulos supremos per anastomoses ante apicem evanidos mittente, bractæ pelta subrotunda margine aures-pubescente. ♀ In Mexico ad Tabasco (Linden! in h. Deless.). *Artanthe melanosticta* Miq.! syst. p. 404. Ramuli glabri, glandulis fuscis conspersi. Limbi 0,19 longi, 0,075 lati. Petioli 0,03 longi. (v. s.)
138. *P. NITIDUM* (Vahl enum. v. 1. p. 314, exclus. syn. Swartz), foliis breviter petiolatis oblongo-ovato-lanceolatis apicem versus sensim acuminatis, acumine acuto, basi inæquali rotundatis utrinque glabris supra nitidis subtus pallidioribus utrinque nigro-punctulatis subcoriaceis subopacis, nervis subtus prominulis, centrali utrinque ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitudinis nervos 4 alternos patulo-adscendentes supremos ad apicem fere ductos infimos subtiliores mittente, petiolo glabro, pedunculo petiolum æquante glabro, amento quam folium brevior, bractæ pelta triangulari-semilunari margine hirsuta, stigmatibus filiformibus. ♀ In Surinam (Hostman n. 709! in h. Berol. et Boiss.), Demerara (Parker in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 464),

*Paramaribe* (Kegel n. 170 et 465 fide Miq. in Linn. v. 22. p. 77), in regionibus inter. ad Savanhe (Kappler n. 1668! in h. Boiss.). P. præmorsum Rothb. fide Vahl l. c. *Artanthe nitida* Miq. syst. p. 495, in Hook. Lond. Journ. l. c., in Linn. l. c. et in Comment. phyt. p. 47. t. 7. Rami glabri, sicci fuscescentes nigro-punctulati. Limbi 0,1 longi, 0,045 lati. Petioli 0,007 longi. Amenta florentia 0,04 longa, 0,002 crassa. (v. s.)

139. P. *SANTAREMENSE*, foliis breviter petiolatis oblongo-ovatis apice acuminatis, acumine acuto, superioribus basi subinæquali obtusis inferioribus basi æquali rotundatis cordatisve utrinque glabris nitentibus membranaceis opacis subtiliter pellucido-punctulatis, nervo centrali subtus prominulo utrinque ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. nervos 3 subtus prominulos nervulis fortioribus intermixtos infimos fere e basi ortos subadscendentes cæteros patulo-subadscendentes mittente, petiolo glabro ad limbum usque vaginante, pedunculo glabro petiolum æquante, bracteæ pelta triangulari margine hirsuta, stigmatibus brevibus. ♀ In vicinibus Santarem prov. Para (Spruce! n. 7 et h. DC. in h. Kew.). Ramuli glabri. Limbi 0,13 longi, 0,06 lati. Petioli 0,015 longi (v. s.)

140. P. *MODESTUM*, foliis brevissime petiolatis oblongo-lanceolatis apice acutiusculis mucronulatisque basi subæquali acutis utrinque glabris membranaceis subtilissime pellucido-punctulatis subopacis, nervo centrali subtus prominulo utrinque ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. nervos 4 alternos adscendentes supremos ante apicem per anastomoses evanidos infimos subtiles mittente, petiolo glabro glandulis fuscis punctulato, pedunculo glabro petiolum triplo superante, amento quam folium multoties brevius, bracteæ pelta semilunari margine villosula. ♀ In Guyana gallica (Leprieur n. 1839! in h. Deless.). *Artanthe modesta* Miq. syst. p. 517 et? in Hook. Lond. Journ. bot. v. 4. p. 468. Ramuli glabri, glandulis fuscis conspersi. Limbi 0,19 longi, 0,03 lati. Petioli 0,003 longi. Amenta florentia 0,002 crassa. (v. s.)

141. P. *SUBANDINUM*, foliis brevissime petiolatis suboblique elliptico-oblongis apice protracto-acuminatis, acumine acuto, basi breviter inæquali obtusis, lobo minore basi subacuto, majore rotundato, utrinque glabris membranaceis subopacis subtiliter pellucido-punctulatis, nervis subtus subprominulis, centrali utrinque ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitudinis nervos alternos subadscendentes supremos ad apicem ductos mittente, petiolo dorso pilis longis hirsuto, pedunculo petiolum æquante puberulo, amento florente quam folium duplo brevius mucronato, bracteæ apice truncato-peltatæ pelta semilunari margine fusce ciliolata pedicello cucullato. ♀ In Peruvia subandina ad Cuchero (h. Vindob.). Ramuli dense fusco-pubescentes, pilis ad nodos longioribus. Limbi 0,21 longi, 0,09 lati. Petioli 0,002 longi. (v. s.)

142. P. *MANDONI*, foliis breviter petiolatis ovatis apice longiuscule acuminatis, acumine acuto mucronulatoque, basi valde inæquali cordatis, lobo majore auriculæformi petiolum velante libero, supra bullatis scabris subtus molliter pallidius dense tomentosis lacunosus coriaceis opacis olivaceis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis usque nervos latere majore 9 minore 8 subadscendentes alternos supremos ad apicem ductos inferos magis patule arcuatos mittente, petiolo ad limbi lobum majorem usque vaginante dorso molliter dense hirsuto, pedunculo dense molliter hirsuto petiolum æquante, amento per anthesim folium dimidium æquante obtuso, bracteæ apice truncato-peltatæ pelta semilunari margine hirsuta, pilis marginis anterioris longioribus, pedicello basi hirsuto, ovario glabro, stigmatibus linearibus acutis. ♀ In Bolivia prov. Larecaja vicinis Sorata ad rivum Challasuyo et in sepibus alti. reg. temp. alt. 2600 m. (Mandon n. 1113! in h. DC.). Rami dein glabrati. Ramuli densius molliter hirsuti. Limbi 0,16-0,17 longi, 0,08-0,09 lati. Petioli 0,01 longi. (v. s.)

143. P. *CLAUSSENIANUM*, foliis breviter petiolatis oblongo-ovatis apice acuminatis, acumine acuto, basi subinæquali obtusis supra bullatis scabris subtus ad nervos nervulosque molliter hirtellis glanduloso-punctatis rigidis subopacis pellucido-punctatis, nervis subtus prominulis centrali utrinque ad  $\frac{1}{2}$  longitudinis usque nervos 5 alternos subadscendentes mittente, petiolo molliter hirtello basi ima vaginante, pedunculo molliter hirtello petiolum triplo superante, amento folium subsuperante, bracteæ pelta semilunari margine dense

griseo-pubescente, bacca glabra. ♀ In Brasilia ad Minas-Geraes (Clausen, n. 155! in h. Mart.). Artanthe Clauseniana Miq.! in Mart. flor. Bras. fasc. 11, p. 43, et in Linn. v. 20. p. 153. Rami serius fere glabrati, asperuli. Ramuli juniores molliter hirti. Limbi 0,09 longi, 0,03 lati. Petioli vix 0,005 longi. (v. s.)

144. P. MEXICANUM, foliis brevissime petiolatis oblongo-lanceolatis apice acuminatis basi inæquali rotundatis vel subcordatis supra bullatis scabris in nervo subpuberulis nitidulis subtus lacunosis ad nervos præsertim appresse longiuscule pubescenti-hirtis subcoriaceis opacis, centrali ad  $\frac{1}{4}$  longitudinis usque nervos utrinque 5-6 subadscendentes subtus prominulos supremos ad apicem fere ductos sursumque aliquando nervulos fortiores mittente, petiolo hirtio, amento quam folium brevior, bractæ pelta subrotunda margine cinereo-hirta, bacca vertice subprotracto puberulo. ♀ In corjilleris Mexici ad Oaxaca (Galeotti, n. 6015! in h. Deless.), ad rivulos planitie Eintla, præcipue ad Hacienda de la Compagnia alt. 4400-5500 ped. (Martens et Gal. l. c.). P. hispidum Martens et Gal. in bull. ac. Brux. v. 10. p. 21 (non Kunth). B. Artanthe mexicana Miq.! in Linn. v. 20. p. 153. Ramuli pubescenti-hirti. Limbi 0,14 longi, 0,045 lati. Petioli fere 0,004 longi. Amenta matura 0,003 crassa. (v. s.)

145. P. HIRSUTUM (Swartz! prodr. p. 15), foliis breviter petiolatis inferioribus elliptico-lanceolatis oblongis superioribus oblique elliptico-lanceolatis omnibus apice acuminatis, acumine acuto, basi inæquali rotundatis vel obtusis vel obtusiusculis vel et attenuatis lobo longiore cum petiolo adnato membranaceo-rigidis pellucido-punctulatis supra verruculoso-scabris subtus ad nervos appresse hirtellis, nervo centrali ad apicem ducto subtus prominulo ad  $\frac{1}{4}$  longitudinis usque nervos utrinque 5-6 alternos subadscendentes subtus prominulos supremos ad apicem ductos mittente, petiolo hirtio basi ima vaginante, pedunculo hirtello petiolum æquante vel parum superante, amento maturo quam folium paulo brevior, bractæ apice peltatæ pelta triangulari-semilunari margine hirsuta, bacca apice glanduloso-puberula. ♀ In Tocache et Huallaga (Endlich! in h. Vindob.), Peruvia (Mathews n. 1703! in h. Kew.), Brasilia (Riedel n. 440! in h. Petrop.), Bahía (Lhotzky! in h. DC.), Nova Granata (Linden n. 121! in h. Kew. et Boiss.), Surinam (Buchinger n. 312! in h. Lenormand), Costa-Rica (Hoffman n. 348! in h. Berol.), sylvis Jalapæ (Miq. in Linn. v. 18. p. 715), Santo Domingo (Mayerhoff, n. 342! et Ehrenberg! in h. Berol.), Jamaica (March n. 1902! in h. Kew. et Swartz! in h. DC. et in h. Willd. n. 666!), Sanct. Vincent (L. G.! in h. Kew.), Cuba (Ossa! in h. DC.), in ins. Java culta (Zoll. n. 1471! in h. Francovil.). — Swartz! fl. Ind. occ. v. 1. p. 60, Vahl enum. v. 1. p. 323, Willd.! sp. v. 1. p. 163. Steffensia hirsuta Kunth! in Linn. v. 13. p. 641. P. scabrum Swartz, fl. Ind. occ. v. 1. p. 59. Steff. scabra Kunth! l. c. p. 640. Artanthe hirsuta Miq.! Syst. p. 446 et in Hook. Lond. journ. bot. v. 4. p. 456. Art. scabra Miq.! Syst. p. 447 et in Hook. Lond. journ. bot. l. c. Grisebach, fl. w. Ind. p. 171, cat. fl. Cub. p. 65. Art. Kunthiana Miq. in Hook. Lond. journ. bot. l. c. Art. asperifolia Miq.! in Linn. v. 20. p. 155 156, et in Mart. flor. Bras. fasc. 11. p. 45. Art. Ollersiana Miq.! in Linn. v. 20. p. 156 et in Mart. flor. Bras. l. c. p. 46. Art. aspera Miq. in Hook. Lond. journ. bot. l. c. Ramuli verruculosi scabri hirtelli. Limbi 0,15 longi, 0,06 lati. Petioli et pedunculi 0,005 longi. Amenta florentia 0,1-0,11 longa, vix 0,002 crassa. (v. s.)

α, *fusco-punctulatum*, foliis subtus fusco-punctulatis. — In San Gabriel (Spruce, n. 2138! in h. Francov.). (v. s.)

β, *Jamaicense*, foliis basi inæquali subattenuatis attenuatisve obtusis obtusiusculisve. — In Jamaica (March n. 1379! in h. Kew.), Novæ Granatæ prov. Barbacoas (Triana, n. 38! in h. suo), Venezuela (Fendler, n. 1134! in h. DC.). (v. s.)

γ, *magnifolium*, foliis majoribus (limbis 0,2-0,25 longis, 0,08-0,1 latis) ovato-ellipticis apice recte longe acuminatis, acumine acuto, basi subattenuatis, basi ima inæquali acutis. — In Brasiliæ prov. Para (Mart.! in h. Monac.), Jamaica (Wilson! et Alexander! in h. Kew.), Cuba (Wright, n. 493! in h. DC.), Portorico (Bertero! in h. DC.). (v. s.)

- δ, *Olfersianum*, limbis 0,14 longis, 0,07 latis, pedunculo petiolum fere duplo superante. — In Brasilia (Sello! in h. Berol.), ad Eugenio do Bernardino (Pohl n. 1274! in Vindob.), Guadaloupa (Krauss! in h. DC.), Cuba (Ossa! in h. DC.). P. Olfersianum Kunth! olim. Steffensia Olfersiana Kunth! in Linn. v. 13. p. 645. Artanthe Olfersiana Miq.! syst. p. 445. (v. s.)
- ε, *Barclayanum*, foliis oblique obovato-ellipticis apice oblique acuminatis, acumine acuto, basin versus subattenuatis basi ima inæquali obtusis. — In Peruvia (Mathews, n. 1145 et 1146! in h. Kew.) et Nova Granata (Barclay! in h. Kew.). Artanthe Barclayana Miq.! in Hook. Lond. Journ. bot. v. 4. p. 460. (v. s.)
- η, *asperiusculum*, foliis apice longe protracto-acuminatis, acumine acuto, pedunculo petiolum duplo superante, limbis 0,15 longis, 0,055 latis, ramulis reflexe hirtellis, bractææ pelta marginæ ciliolata. — In Nova Granata (Holton, n. 238! in h. DC.), convalle umbrosa flumin. Magdalenzæ prope Santo Bartholome et Boca de Opon (Humb. et Bonpl. nov. gen. v. 1. p. 51 et in h. Willd. n. 665!), locis humidis prope Xalapa alt. 4-5000 ped. (Galeotti, n. 6019 fide Martens et Galeotti in Bull. ac. Brux. v. 10. p. 20). P. asperiusculum Kunth! l. c. et synopsis, v. 1. p. 107. Steffensia asperiuscula Kunth! in Linn. v. 13. p. 631. Artanthe asperiuscula Miq.! syst. p. 464. (v. s.)
146. P. JALAPENSE, foliis petiolatis subobovato-elliptico-oblongis apice suboblique acuminatis, acumine acuto, basin versus attenuatis basi inæquali-obtusiusculis supra verruculis albis obsitis scabris junioribus parce pilosis subtus pallidioribus ad nervos pilis mollibus dense pubescentibus, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis usque nervos utrinque 6 rarissime 7 subadscendentes alternos supremos ad apicem ductos mittente, petiolo hispidulo-hirto basi ima vaginante, pedunculo hirtello petiolum paulo superante, amento maturo quam folium breviori cylindrico, bractææ apice truncato-peltatæ pelta suborbiculari fere tota griseo-hirsuta, bacca glabra. ♀ In Cordilleris Mexici alt. 4000 ped. (Galeotti, n. 6009! in h. Deless. et DC. Coulter! h. Kew.), prope Xalapa (Schiede et Deppel in h. Petrop.). Artanthe jalapensis Miq.! syst. p. 444. Ramuli dense hispido-hirti. Limbi 0,13-0,19 longi, 0,04-0,07 lati. Petioli 0,01-0,15 longi. Amenta matura 0,1 longa, 0,004 crassa. (v. s.)
147. P. GAUDICHAUDIANUM (Kunth! olim), foliis brevissime petiolatis oblongo-lanceolatis apice acuminatis, acumine acuto, basi inæquali acutis supra scabris subtus adpresse canescenti-pubescentibus margine ciliolatis membranaceis subopacis epunctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis usque nervos utrinque 5 alternos adscendentes supremos ad apicem ductos mittente, petiolo basi dorso sursum utrinque hirtello-pubescente basi ima vaginante, pedunculo puberulo petiolum multoties superante, amento maturo folium subæquante, bractææ apice truncato-peltatæ pelta semilunari vel suborbiculari margine griseo-luteo-ciliata, bacca glabra. ♀ In Brasiliæ prov. Cearnsi (Mart. in h. Monac.), prope Bahia (Blanchet, n. 408! in h. DC., Gaudich. n. 1116! in h. DC.), et monte Corcovado (Kunth, l. c.), Rio Janeiro (Gardner n. 118! in h. Deless. et Weddell! in h. DC.), in Brasilia (Riedel n. 19! et 286! in h. Petrop.). Steffensia Gaudichaudiana Kunth! in Linn. v. 13. p. 636. Artanthe Gaudichaudiana Miq.! Syst. p. 469. Ram. puberulo-hirtelli. Limbi 0,155 longi, 0,04 lati. Petioli 0,005 longi. Amenta matura 0,004 crassa. (v. s.)
148. P. FISCHERIANUM, foliis petiolatis lanceolato-ellipticis apice oblique longiuscule acuminatis, acumine acuto mucronulatoque, basi inæquali acutis supra verruculis albis obsitis junioribus scabriusculis subtus ad nervos adpresse hirtellis membranaceis subpellucidis pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis usque nervos utrinque 6 alternos subadscendentes supremos ad apicem ductos mittente, petiolo basi ima vaginante utrinque juniore hirtello dein glabrato, pedunculo florente petiolum æquante glabro, amento florente folium dimidium æquante, bractææ apice truncato-peltatæ pelta subrotunda margine dense ciliata. ♀ In Mexico (Kar-

winski n. 826! in h. Petrop.). *Ramuli glabri, adulti breves. Limbi 0,195 longi, 0,08 lati. Petioli 0,01 longi.* (v. s.)

149. P. *PLEXUOSUM* (Jacq. eclog. p. 139. t. 93), foliis brevissime petiolatis oblique oblongo-ellipticis apice acuminatis, acumine acuto mucronulatoque, basi inæquali cordatis supra pilosulis subtus ad nervos et nervulos pilis adpressis conspersis membranaceis, nervo centrali ad  $\frac{1}{2}$  longitud. usque nervos utrinque 5-6 subadscendentes mittente, petiolo pilosulo. ♀ Ad Caracas (h. Willd. n. 650!). P. Jacquinianum Rœm. et Schult. mant. v. 1. p. 234. Steffensia? flexuosa Kunth! in Linn. v. 13. p. 644. *Artanthe flexuosa* Miq.! syst. p. 454. *Ramuli flexuosi, glabri, fusco-virentes. Limbi 0,12 longi, 0,05 lati. Petioli, 0,005 longi.* (v. s.)
150. P. *MIERSINUM*, foliis brevissime petiolatis suboblique ovato-vel obovato-ellipticis apice acuminatis, acumine acuto mucronulatoque, basi inæquali obtusis cordulatis supra ad nervos pubescentibus areolato-bullatis brevibus subtus molliter dense hirsuto-pubescentibus margine ciliatis coriaceis opacis, nervo centrali subtus prominulo utrinque nervos 4-5 alternos subadscendentes subtus prominulos nervulis fortioribus intermixtos mittente, petiolo hirsuto ad  $\frac{1}{2}$  longitud. usque yaginate, pedunculo hirsuto petiolum superante, amento crassoquam folium brevior, bractæ apice peltatæ pelta triangulari-semilunari margine hirsuta, filamentis exsertis, ovario pubescente. ♀ In Audibus Quitensibus alt. 10000 ped. (Jameson n. 665! et 666! in h. Kew. et 297! in h. DC.) et prov. Chachapoyas (Mathews n. 186! in h. Kew.). *Artanthe Miersina* Miq.! in Hook. Lond. journ. bot. v. 4. p. 449. *Rami puberuli. Ramuli hirsuto-puberuli. Limbi 0,1 circ. longi, 0,04-0,05 lati. Petioli 0,005 longi. Pedunculi 0,007-0,008 longi. Amenta florentia 0,05 longa, 0,005 crassa.* (v. s.)
151. P. *LEUCOPHYLLUM*, foliis brevissime petiolatis oblongo-lanceolatis apicem versus attenuatis subacuminatis, acumine acuto mucronulatoque, basin versus paulo angustatis basi valde inæquali subdimidiato-cordatis supra verrucis numerosissimis albicantibus et asperimis ad nervos hirtis subtus pubescenti-tomentosis rigidis obscuris subtiliter pellucido-punctulatis, nervo centrali ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. usque nervos utriusque 6-7 subadscendentes subtus prominulos supremos ad apicem ductos mittente, petiolo dense pubescenti-tomentoso, pedunculo petiolum parum superante pubescenti-tomentoso, amento folium subæquante cylindrico, bractæ apice peltatæ pelta subrotunda margine villosa pedicello angusto. ♀ In Mexico (Andrieux n. 134! in h. Deless.), in angustis Guadalupe in Mexico alt. 3000 ped. (Galeotti n. 6024! in h. Deless.), Atlacomulco (Schiede n. 1393! fide. Miq. in Linn. v. 18. p. 715), Acapulco (Benth. in bot. of Sulph. p. 167). *Artanthe leucophylla* Miq.! syst. p. 460. *Ramuli pubescenti-tomentosi. Limbi 0,14 longi, 0,04 lati. Petioli 0,004 longi. Amenta ferè 0,005 crassa.* (v. s.)
152. P. *PISOENSE*, foliis breviter petiolatis suboblique oblongo-ellipticis apice longiuscule acuminatis, acumine acuto mucronulatoque, basi inæquali rotundatis supra subtusque dense molliter pubescentibus supra nigro-punctulatis subtus griseo-pallidioribus margine ciliatis firmule membranaceis opacis crebre pellucido-punctulatis, nervis subtus prominulis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos latere majore 7 minore 6 supremos ad apicem ductos superos subadscendentes inferos magis arcuatim patulos infimos subtiles mittente. ♀ In Brasilia (Riedel n. 2! et Langdorff! 94! in h. Petrop.). *Rami ramulique dense molliter griseo-hirsuti. Limbi 0,14 longi, 0,06 lati. Petioli 0,007 longi.* (v. s.)
153. P. *PSEUDO-MOLLIOMA*, foliis breviter petiolatis inferioribus ovato-ellipticis superioribus elliptico-lanceolatis omnibus apice modice acuminatis acutis basi subæquali subrotundatis subattenuatisve supra scabris et minute pilosis subtus molliter pubescentibus membranaceis subopacis pellucido-punctatis, centrali nervo subtus prominente ad  $\frac{3}{4}$  longitud. nervos utrinque 6 subadscendentes subtus prominulos alternos supremos ad apicem ductos infimos subtiliores mittente, petiolo hirsuto, amento densifloro, pedunculo petiolum superante hirtello, bractæ apice peltatæ pelta triangulari margine hirsuta. ♀ In Peruvia (Poeppig n. 2290! in h. DC. et Mathews n. 1705! in h. Kew.). *Artanthe Poeppigii* Miq.! syst. p. 437 et in Hook. Lond. Journ. bot. v. 4. p. 456.



Ramuli adspectu ochraceo-grisei, pilis retrorsis adpresse pubescentes. Limbi 0,16-0,17 longi, 0,06-0,07 lati. Petioli 0,008 longi. Pedunculi 0,01 longi. Amenta 0,12-0,15 longa, 0,005 crassa. (v. s.)

♂, *dichotomum*, foliis minoribus magis acuminatis oblongo-ovatis glabrioribus. — In Peruvia (Poeppig! in h. Berol. sub nomine *P. dichotomum*. (v. s.)

154. *P. SUBFLAVUM*, foliis brevissime petiolatis ovato-lanceolatis apice acuminatis, acumine acuto, basi inæquali cordulatis utrinque et subtus densius molli-ter flavicanti-pubescentibus subcoriaceis opacis, nervo centrali ad  $\frac{1}{2}$  longitudi-  
nervos utrinque 4-5 alternos subadscendentes supremos ad apicem ductos mittente, petiolo dense flavicanti-hirsuto basi ima vaginante, pedunculo petiolum æquante dense flavicanti-hirsuto, bractæ apice peltatæ pelta triangulari margine flavicanti-hirsuta, ovario puberulo. ♂ In Novæ Granatæ prov. Pasto Ortega alt. 1500 m. (Triana n. 23 in h. DC.). Ramuli retrorsum flavicanti-hirsuti. Limbi 0,145 longi, 0,055 lati. Petioli 0,004 longi. (v. s.)
155. *P. LEPTOPHYLLUM*, foliis brevissime petiolatis oblongo-ellipticis apice longiuscule acuminatis, acumine subulato, basi inæquali acutiusculis supra intense viridibus scabridis planis subtus ad nervos molliter pilis longiusculis villosulis membranaceis subpellucidis subtiliter pellucido-punctulatis, nervis subtus prominulis, centrali nervo ad  $\frac{1}{2}$  longitudinis nervos utrinque alternos subadscendentes supremos ad apicem ductos mittente, petiolo utrinque villosulo basi ima vaginante, pedunculo pubescente petiolum æquante, bractæ apice truncato-peltatæ pelta semilunari-subrotunda margine subtiliter aureo-puberulo, amento mucronulato. ♂ In Venezuela (Funck et Schlim n. 408! in h. Petrop.). Ramuli molliter villosuli. Limbi 0,2 longi, 0,078 lati. Petioli 0,005 longi. (v. s.)
156. *P. CHIMONANTHIFOLIUM* (Kunth! olim), foliis breviter petiolatis oblique oblongo-lanceolatis apice acuminatis, acumine acuto mucronulatoque, basi inæquali subattenuatis vel acutiusculis obtusiusculisve supra punctulato-scabris subtus ad nervos sparse adpresse hirtellis membranaceis subpellucidis, nervis subtus prominulis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos utrinque 4-5 alternos patulo-adscendentes sursumque ad apicem usque nervulos fortiores mittente, petiolo hirtello, pedunculo hirtello dein glabrato petiolum fere triplo superante, amento quam folium multum brevius, bractæ apice peltatæ pelta triangulari margine hirsuta. ♂ In Brasilia meridionali ad Rio Janeiro et Bahiam (Sello! in h. Berol.). *Steffensia chimonanthifolia* Kunth in Linn. v. 13. p. 628. *Artanthe chimonanthifolia* Miq.! syst. p. 466 et in Mart. Flor. Bras. fasc. 11. p. 48. Ramuli teretiusculi, sparsissime puberuli et scabriusculi, juniores dense hirtelli. Limbi 0,13 longi, 0,03 lati. Petioli 0,005 longi. Amenta florentia 0,002 crassa. (v. s.)
157. *P. BACCANS*, foliis brevissime petiolatis suboblongo-ovato-acuminatis, acumine acutiusculo mucronulatoque, basi breviter inæquali rotundatis supra præsertim ad nervum centalem puberulis subtus dense molliter pubescentibus firmis obsolete pellucido-punctulatis, nervis subtus prominulis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos utrinque 4 alternos subalternosve patulo-subadscendentes mittente, petiolo utrinque dense molliter pubescente basi ima vaginante, amento maturo folium dimidium æquante, pedunculo petiolum æquante molliter pubescente, bractæ apice peltatæ pelta semilunari margine canescenti-hirsuta, bacca glabra glandulosa. ♂ In Brasiliæ prov. Paraensi (Mart.! in h. monac.). *Artanthe baccans* Miq.! in Linn. v. 20. p. 166 et in Mart. flor. bras. fasc. 11. p. 58. Rami pilis recurvulis molliter pubescentes, ramuli densius molliter canescenti-pubescentes. Limbi 0,08 longi, fere 0,035 lati. Petioli 0,005 longi. Amenta matura 0,004 crassa. (v. s.)
158. *P. PRUNIFOLIUM* (Jacq. eclog. v. 1. p. 140. t. 94), foliis breviter petiolatis ovatis apice acuminatis, acumine acuto, basi inæquali cordatis utrinque glabris subtus ad nervos sub lente villosulis supra rugosis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos utrinque 4 alternos subadscendentes supremos ad apicem ductos mittente, petiolo pubescente. ♂ In horto Schoenbr. culta ex Caracæ a Bredemeyero allata (Jacq. l. c.). — Rœm. et

Schult. mant. v. 1. p. 235. Rami ramulique glabri, immaculati, virides. Limbi 0,1 longi, 0,04 lati. Petioli 0,008 longi.

159. P. BOLIVIANUM, foliis longiuscule petiolatis late ovato-acuminatis, acumine longiusculo acuto, basi parum inæquali haud profunde cordatis lobis rotundatis supra excepta basi ad nervos glabris subtus ad nervos fusce adpresse hirtellis rigidis opacis pellucido-punctulatis, centrali nervo subtus prominente ad  $\frac{1}{2}$  longitud. nervos alternos patulo-subadscendentes supremos ante apicem per anastomoses evanidos infimos subtiles mittente, petiolo ad  $\frac{1}{2}$  longitud. vaginante basi dorso et sursum utrinque hirsuto, pedunculo fusce hirtello maturo quam petiolus paullo brevior, amento maturo limbum æquante, bractæ apice truncato-peltatæ vertice semilunari margine superiore ciliolato inferiore ciliato pedicello subcucullato dorso hirsuto, bacca vertice dense pubescente. ♀ In Bolivia (Kelly, Pentland ! in h. Petrop.). Rami subtiliter hirtelli. Limbi 0,245 longi, 0,18 lati. Petioli 0,03 longi. Amenta matura 0,004 crassa. (v. s.)
160. P. PETIOLARE, foliis longe petiolatis subrotundato-ovatis apice attenuatis acutiusculis basi rotundato-truncatis supra glabris subtus ad nervos nervulosque pubescentibus rigidulis opacis, nervo centrali subtus prominente ad  $\frac{1}{2}$  longitud. nervos 7 alternos subtus prominulos subadscendentes inferos subpatulo-adscendentes et gradatim subpatulos; supremos ad apicem ductos mittente, petiolo hirtello postea glabrato ad  $\frac{1}{2}$  longitudinis vaginante, pedunculo quam petiolus brevior, bractæ apice peltatæ pelta semilunari margine dense et margine inferiore densius hirsuta pedicello margine hirtello. ♀ In Novæ Granatæ prov. Pasto alt. 1600 m. (Triana n. 53 ! in h. suo). Ramuli glabri. Limbi 0,16 longi, 0,11 lati. Petioli 0,055 longi. (v. s.)
161. P. JACQUINI, foliis petiolatis ovato-oblongis apice acuminatis, acumine acuto, basi inæquali obtusis supra glabris subtus pallidioribus ad nervos adpresse puberulis membranaceis subpellucidis subtilissime pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos 5 subadscendentes alternos supremos ad apicem ductos infimos subtiliores arcuatos mittente, petiolo subtiliter hirtello basi vaginante, pedunculo subtiliter hirtello petiolum æquante, amento florente quam folium duplo brevior. ♀ Patria ignota, an Caracas? (Jacquin ! in h. Vindob.). Rami fuscescentes glabri. Ramuli juniores subtiliter hirtelli. Limbi 0,095 longi, 0,04 lati. Petioli 0,006 longi. Amenta florentia 0,002 crassa. (v. s.)
162. P. WYDLERIANUM, foliis brevissime petiolatis ovato-lanceolatis apice longe acuminatis, acumine acuto subulatoque, basi parum inæquali obtusis acutiusculisve supra atro-viridibus nitidis glabris subtus fuscescenti-glaucis ad nervos pilosulis membranaceis subopacis pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos utrinque 5 subalternos patulo-adscendentes supremos ad apicem ductos ab inferis parum remotiores mittente, petiolo pilosulo, pedunculo pilosulo petiolum æquante, amento quam folium multoties brevior, bractæ apice inflexo-peltatæ dorso villosulæ vertice semilunari margine hirsuto, stigmatibus linearibus, ovario pubescente. ♀ In Porto Rico (Wydler n. 321 ! in h. Deless.), St. Domingo (Ehrenberg ! in h. Berol. et Mayerhoff n. 343 ! in h. Berol.), locis humidis ad Maiquetia (Wagner n. 127. in Linn. v. 26. p. 632.) Ramuli pilosuli. Limbi 0,17-0,23 longi, 0,07 lati. Petioli 0,005 longi. Amenta florentia 0,002 crassa. (v. s.)
163. P. MURRAYANUM, foliis brevissime petiolatis elliptico-oblongis apice acuminatis, acumine acuto mucronulato, basi inæquali latere majore rotundatis minore acutis supra glabris subtus ad nervos parce hirtellis membranaceis subopacis pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos 5 alternos subadscendentes supremos per anastomoses ad apicem ductos infimos subtiliores mittente, petiolo hirtello basi ima vaginante, pedunculo petiolum triplo superante, amento folium dimidium superante, bractæ apice truncato-peltatæ pelta semilunari-subrotunda margine ciliolata pedicello brevi, bacca glabra lateraliter compressa. ♀ In Jamaica (Murray n. 497 ! in h. DC.). Ramuli glabri. Limbi 0,13 longi, 0,055 lati. Petioli 0,003 longi. Amenta florentia 0,002 crassa. (v. s.)

164. *P. FISTULOSUM*, foliis breviter petiolatis oblique ovato-ellipticis apice attenuato-acutis basi subæquali subrotundatis supra glabris subtus ad nervos pubescentibus membranaceis subopacis subtiliter pellucido-punctulatis, centrali nervo subtus prominulo ad  $\frac{1}{2}$  longitud. nervos utrinque 4 alternos subtus prominulos subadscendentes supremos supra medium ductos infimos subtiliores mittente, petiolo puberulo, pedunculo glabro petiolum æquante, amento folium æquante, bractæ apice peltatæ pelta triangulari margine aureo-hirsuta, bacca glabra. ♀ In Novæ Granatæ prov. Pasto (Triana n. 25 ! in h. DC.). Rami glabri, intus fistulosi, nigri. Ramuli glabri. Limbi 0,1 longi, 0,055 lati. Petioli 0,01 longi. (v. s.)
165. *P. TRICHONEURON*, foliis breviter petiolatis sublanceolato-oblongis apice longe acuminatis basi inæquali obtusis supra glabris nitidulis subtus ad nervos nervulosque hirtis membranaceo-rigidis subopacis epunctulatis, centrali nervo subtus prominente ad  $\frac{1}{2}$  longitud. nervos utrinque 6 alternos subadscendentes subtus prominulos supremos fere ad apicem ductos mittente, petiolo ad limbum vaginante dorso dense hirtio, amento quam folium triplo brevior filiformi, bractæ apice peltatæ pelta triangulari fusca margine dense longe ciliata, stigmatibus obtusis. ♀ In Guyana gallica (Leprieur n. 143 ! in h. Deless.). Ramuli hirtio-tomentosi. Limbi 0,25 longi, 0,085 lati. Petioli 0,007 longi. Amenta florentia 0,001 crassa. (v. s.)
166. *P. LEDEBOURII*, foliis breviter petiolatis elliptico-oblongis apice acuminatis, acumine acuto mucronulatoque, basi breviter inæquali acutis supra glabris breviter bullatis subtus ad nervos subtiliter velutino-puberulis rigidulis opacis pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos utrinque 6 alternos inferos subadscendentes supremos ad apicem ductos infimos subtiliores mittente, petiolo glabro basi ima vaginante, pedunculo glabro quam petiolum dimidio brevior, amento maturo folium paullo superante, bractæ apice truncato-peltatæ vertice semilunari margine exteriori triangulariter ciliato, bacca glabra lateraliter compressa apice olivaceo-glandulosa. ♀ In Brasilia (Riedel n. 803 ! in h. Petrop. et DC.), in Mexico (herb. Ledebour ! in h. Petrop. et h. DC.) Ramuli verrucis albidis dense conspersi, glabri. Limbi 0,12 longi, 0,045 lati. Petioli 0,007 longi. Amenta matura 0,004 crassa. (v. s.)
167. *P. DESCOITTSIANUM*, foliis brevissime petiolatis oblongo-ellipticis apice protracto-acuminatis, acumine acuto, basi inæquali acutis acutiusculisve supra glabris lævibus nitidulisque subtus ad nervos puberulis supra subbullatis firmulo-membranaceis subopacis crebre pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos utrinque 5 subadscendentes supremos per anastomoses fere ad apicem ductos infimos subtiliores mittente, petiolo glabro ad  $\frac{1}{2}$  longitudinis vaginante, pedunculo glabro petiolum æquante, amento maturo folium æquante, bractæ apice truncatæ vertice triangulari margine ciliolato pedicello dorso producto et margine ciliolato, bacca vertice fusco-hirsuta et basi glandulis flavicantibus conspersa, stigmatibus abbreviatis. ♀ In sylvis prope Xalapa (Fischer ! in h. Petrop.) et Miradores ad Puente Nacional (Linden n. 116 ! in h. Boiss.). Rami ramulique glabri, adulti verruculis albidis sub lente obsiti. Limbi 0,155 longi, 0,07 lati. Petioli 0,005 longi. Amenta matura fere 0,003 crassa. (v. s.)
168. *P. RUGULOSUM* (Cas. DC. in Seem. journ. of bot. 1866. p. 211), foliis brevissime petiolatis ellipticis apice breviter attenuatis vel acutiusculis basi inæquali obtusis junioribus supra subtiliter pubescentibus subtus ad nervos præsertim adpresse pubescentibus subcoriaceis opacis parce pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos latere majore 4 minore 3 subadscendentes mittente, petiolo dorso pilis recurvulis adpresse pubescente basi ima vaginante, pedunculo puberulo petiolum duplo superante, amento maturo quam folium duplo brevior, bractæ apice subtruncato-peltatæ pelta triangulari margine pubescente, bacca glabra. ♀ In Novæ Granatæ prov. Pasto (Triana n. 46 ! in h. DC.). Rami glabri, plicato-rugulosi. Ramuli juniores pilis recurvulis pubescentes. Limbi 0,05 longi, 0,025 lati. Petioli 0,003 longi. Amenta matura 0,004 crassa. (v. s.)
169. *P. GRISNUM*, foliis brevissime petiolatis suboblique oblongo-lanceolatis

apice longiuscule acuminatis, acumine acuto, basi leviter inæquali obtusis basi ima cordulatis supra adpresse puberulis subtus pallidioribus et præsertim ad nervos adpresse pubescentibus membranaceis subopacis subtilissime pellucido-punctulatis, petiolo dense cinereo adpresse pubescente, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos utrinque 6 alternos subadscedentes supremos ad apicem ductos mittente, pedunculo petiolum triplum superante adpresse et dense cinereo-pubescente, amento cinereo florentia folium dimidium æquante filiformi, bractæ apice truncato-peltatæ vertice triangulari margine cinereo-hirsuto. ♀ In Peruvia ad Cuchero (Poeppig? n. 1737! in h. Vindob.). Rami ramulique pilis retrorsis cinereo-pubescentibus. Limbi 0,12 longi, 0,04 lati. Petioli 0,001 longi. Pedunculi 0,017 longi. Amenta florentia 0,0015 crassa. (v. s.)

170. *P. MASTERSIANUM*, foliis breviter petiolatis suboblongo-ovatis apice acute acuminatis vel attenuato-acutis basi subæquali cordulatis vel obtusis supra glabris nitidis subtus pallidioribus ad nervos nervulosque adpresse hirtellis firme membranaceis creberrime subtilissime pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos utrinque 5 subadscedentes supremos ad apicem ductos infimos subtiliores brevissimos subpatulos nervulosque fortiores aliquot fere tota longitudine mittente, petiolo ad  $\frac{1}{2}$  longitudinis vaginante basi dorso et sursum utrinque fusce hirsuto-pubescente, pedunculo hirsuto-pubescente quam petiolum paulo breviori, bractæ apice peltatæ pelta triangulari margine ciliata, bacca vertice glanduloso-hirtella. ♀ In Brasiliæ prov. Para (Mart. in h. monac.). *Artanthe pyrifolia* Miq. in Mart. flor. Bras. fasc. 11. p. 50. t. 16. f. 1 (non *P. pyrifolium* Vahl). Rami vetustiores glabrati. Ramuli juniores hirsuti. Limbi 0,14 longi, 0,07 lati. Petioli 0,007 longi. (v. s.)

171. *P. CABRALIANUM*, foliis breviter petiolatis ovato-ellipticis apice acuminatis, acumine acuto, basi inæquali acutiusculisve supra glabris subulter pubescentibus rigidulis opacis crebre pellucido-punctulatis, nervo centrali subtus prominente ad  $\frac{1}{2}$  longitud. nervos utrinque 5 alternos subadscedentes supremos per anastomoses ad apicem ductos infimos subtiliores mittente, petiolo utrinque hirtello-pubescente basi ima vaginante, pedunculo hirtello-pubescente petiolum paulo superante, amento post anthesin quam folium paulo breviori, bractæ apice truncato-peltatæ pelta semilunari margine subaureo hirsuta pedicello brevi, rhachi hirtella, ovario lateraliter compresso sursum pubescente, ovariis arcte approximatis, stigmatibus abbreviatis. ♀ In Brasilia (Riedel n. 65! in h. Petrop., Riedel et Langsdorff ibid. n. 106!). Rami glabri, fusciscentes. Ramuli hirtello-pubescentes. Limbi 0,11-0,13 longi, 0,055-0,086 lati. Petioli 0,01 longi. Amenta post anthesin 0,002 crassa. Sp. ill. navigatori ad Brasiliam dicata: (v. s.)

172. *P. PSEUDOVELUTINUM*, foliis brevissime petiolatis ovato-oblongis apice acuminatis, acumine acuto mucronulatoque, basi inæquali subcordatis vel subattenuatis membranaceis pellucido-punctatis margine ciliatis supra puberulis, pilis adpressis haud rigidis brevibus, subtus pubescentibus pilis adpressis haud rigidis, nervo centrali subtus prominulo nervos ad  $\frac{1}{2}$  longitud. utrinque 5-6 subtus prominulos subadscedentes supremos ad apicem ductos mittente, petiolo pubescente basi ima vaginante pedunculo petiolum triplo superante, bractæ apice peltatæ pelta triangulari margine hirsuta, bacca apice glandulosa glabra. ♀ In Brasilia meridionali (Lhotsky! in h. DC.); in America merid. (Bertero! in h. DC.). Ramuli pubescentes. Limbi 0,14-0,16 longi. Petioli 0,005 longi. Amenta 0,002-0,003 crassa. (v. s.)

173. *P. PSEUDO-PERUVIANUM*, foliis breviter petiolatis ovato-vel elliptico-lanceolatis apice linear-acuminatis, acumine obtusiusculo, basi æquali acutis utrinque glabris membranaceis subtiliter pellucido-punctatis, nervo centrali ad  $\frac{1}{2}$  longitud. nervos utrinque 4-5 alternos subadscedentes subtus prominulos supremos ad apicem ductos infimos et basi fere solutos mittente, petiolo glabro basi ima vaginante, pedunculo petiolum æquante glabro vel subtilissime puberulo, bractæ apice truncato-peltatæ vertice triangulari margine subtiliter ciliolato et glanduloso. ♀ In Brasilia (Poeppig n. 1037! in h. DC. et Berol.), in Peruvia (Mathews n. 1718! in h. Kew.). *Artanthe peruviana forma levigata*

- Miq. syst. p. 483. Art. cuspidata Miq. ! in Hook. Lond. Journ. bot. v. 4. p. 468. Ramuli glabri, rugulosi. Limbi 0,13-0,17 longi; 0,06-0,075 lati. Petioli 0,015 longi. Pedunculi circiter 0,01 longi. Amenta florentia 0,04 longa, 0,0015 crassa. (v. s.)
174. *P. CHAMISSONIS* (Steud. nom.), foliis breviter petiolatis oblongis apice breviter acuminatis, acumine acuto, basi subinæquali attenuatis acutis utrinque glabris subtus ad nervos subtiliter puberulis rigido-membranaceis opacis; nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos laterē majore 6 minore 5 alternos subadscendentes supremos ad apicem ductos mittente, petiolo glabro pedunculum glabrum duplo superante, amento cylindrico folium æquante, bracteæ apice peltatæ pelta triangulari margine exteriorē ciliolatâ interiorē subnuda, stigmatibus exiguis, bacca glabra. ♀ In Mexico propè Hacienda de la Laguna (Schiede! in h. Berol.) et ad Tlacolula (Ehrenb. in h. Schlecht. fide Miq. ! in Linn. v. 18. p. 715.). *Steffensia oblonga* Kunth! in Linn. v. 13. p. 625. *P. oblongum* Cham. et Schlecht. ! in Linn. v. 5. p. 73. *Artanthe Chamissonis* Miq. ! syst. p. 457. Rami albedo-punctulati. Limbi 0,15 longi, 0,07 lati. Petioli 0,01 longi. Amenta 0,004 crassa. (v. s.)
175. *P. PERUVIANUM*, foliis breviter petiolatis oblongo-elliptico-lanceolatis apice longe acuminatis, acumine acuto mucronulatoque, basi æquali subattenuato-acutiusculis subrotundatisve utrinque glabris et glanduloso-punctulatis membranaceis, nervis subtus prominulis, centrali utrinque ad  $\frac{1}{2}$  longitudinis nervos 4-5 alternos subadscendentes supremos ad apicem ductos mittente, petiolo glabro pedunculo petiolum subæquante glabro, amento cylindrico quam folium multum brevius mucronato, bracteæ apice inflexo-peltatæ dorso ciliolatæ vertice inflexo triangulari margine hirtello. ♀ In Peruvia subandina ad Maynas (Poeppig! in h. Berol.), in montibus ad Pangoa (Mathews n. 1145! fide Miq. in Hook. Lond. Journ. bot. v. 4. p. 461), Guyaquil (Haenke in h. Berol.). Rami subtilissime verruculosi vel subpulverulenti. Ramuli glabri. Limbi 0,19-0,23 longi, 0,075-0,08 lati. Petioli 0,01-0,015 longi. Amenta florentia 0,1 longa, 0,0025 crassa. (v. s.)
176. *P. SPRENGELIANUM*, foliis breviter petiolatis elliptico-lanceolatis lanceolatisve apice acuminatis, acumine acuto mucronulatoque, basi æquali acutis utrinque glabris membranaceis rigidulisve subopacis subtilissime pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos utrinque 4 alternos subadscendentes supremos ante apicem per anastomoses evanidos infimos subtiliores mittente, petiolo basi ima vaginante glabro, pedunculo glabro petiolum æquante, amento florente folium æquante, bracteæ apice truncato peltatæ pelta rhomboidali glabra pedicello dorso producto hirsuta, ovario glabro. ♀ In Brasilia (Riedel n. 1093! et 296! in h. Petrop.) Rami ramulique glabri, fusciscentes. Limbi 0,12 longi, 0,06 lati. Petioli 0,01 longi. Amenta florentia 0,003 crassa. (v. s.)
177. *P. PALUDOSUM*, foliis longiuscule petiolatis oblongo-ovatis apice acuminatis, acumine acutiusculo, basi æquali cordatis utrinque glabris subtus ad nervum centramem subtilissimè puberulis membranaceis subopacis pellucido-punctulatis utrinque nigro-punctulatis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos utrinque 4 suboppositos subadscendentes sursumque nervulos fortiores raros mittente, petiolo dorso subtiliter hirtello, pedunculo subtiliter puberulo petiolum æquante, amento quam folium multo brevius, bracteæ apice peltatæ pelta triangulari margine albo-cinereo ciliata. ♀ In Maynas sylvis paludosis (h. Vindob.). Rami ramulique nigro-punctulati, juniores hirtelli. Limbi 0,115 longi, 0,045 lati. Petioli 0,015-0,02 longi. Amenta florentia 0,015 longa. (v. s.)
178. *P. RUDE* (Kunth! in H. et B. nov. gen. 1. p. 49), foliis petiolatis oblongo-lanceolatis apice acute acuminatis, basi inæquali subattenuatis obtusiusculis supra primum planis remotiuscule scabro-pilosulis demum bullatis scabris subtus lacunosis et molliter hirtopubescentibus coriaceis opacis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos subalternos subadscendentes latere majore 5 minore 4 mittente, petiolo et pedunculo pilis rectivulis hirtellis, pedunculo petiolum æquante, amento quam folium multum brevius; bracteæ apice peltatæ pelta triangulari-semilunari margine albo-hirtello.

- suta. ♂ In temperatis umbris montis Quindiu juxta La Palmilla et El Moral alt. 800-1100 hexap. in Andibus Novæ Granatæ (Humb. et Bonpl. l. c. et h. Willd. n. 660!). *Schilleria rudis* Kunth! in Linn. v. 13. p. 703. P. rude Kunth synopsis v. 1. p. 105. *Artanthe rudis* Miq.! syst. p. 467. Ramuli pilis recurvulis hirtopubescentes. Limbi 0,09-0,15 longi, 0,025-0,04 lati. Petioli 0,008-0,014 longi. Amenta florentia 0,003 crassa. (v. s.)
179. P. ALNOIDES (Kunth olim), foliis breviter petiolatis ovato-oblongis oblongisve apice acuminatis, acumine acuto mucronulatoque, basi subinæquali rotundatis supra papillis conicis pilisque sparsis scabris subtus præsertim ad nervos piloso-pubescentibus rigido-membranaceis subopacis, nervo centrali subtus prominulo nervos utrinque 4-5 adscendentes alternos subtus prominulos supremos ad apicem ductos mittente, petiolo hirtello basi ima vaginante, pedunculo petiolum paullo superante hirtello, bractæ apice peltatæ pelta elliptica vel elliptico-rotunda margine dense hirtovillosa. ♂ In Brasilia (Riedel et Langsdorff n. 704! in h. Petrop.), ins. S<sup>m</sup> Catharinæ (Gaudich. n. 276! in h. DC.; Humb. in h. Berol.). *Steffensia alnoides* Kunth! in Linn. v. 13. p. 643. *Artanthe alnoides* Miq.! syst. p. 462, et in Mart. flor. Bras. fasc. 11 p. 48. Rami ramulique scabriusculi. Limbi 0,14 longi, 0,045 lati. Petioli 0,007 longi. Pedunculi 0,01 longi. Amenta florentia 0,03 longa, vix 0,002 crassa. (v. s.)
180. P. RARUM, foliis breviter petiolatis elliptico-lanceolatis apice longiuscule acuminatis, acumine acuto mucronulatoque, basi breviter inæquali acutis supra verruculis pilisque scabris subtus glabris ad nervos adpresse puberulis rigidis opacis pellucido-punctulatis, nervis subtus prominulis, centrali utrinque nervos alternos adscendentes supremos ad apicem ductos latere majore 6 minore 5 mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum æquante, amento maturo folium dimidium æquante, bractæ apice truncato-peltatæ pelta semilunari margine ciliolata, rhachi foveolata, bacca basi subimmersa glabra apice glandulis flavicantibus conspersa interdum trigona. ♂ In Brasilia (Riedel n. 469! in h. Petrop.). Ab uno tantum viatore allatum, ideo rarum. Rami ramulique glabri, fusciscentes. Limbi 0,2 longi, 0,065 lati. Petioli 0,007 longi. Amenta matura 0,003 crassa. (v. s.)
181. P. FADYENII, foliis brevissime petiolatis oblique oblongo-ovato-ellipticis apice breviter acuminatis, acumine acuto, basi subæquali vel inæquali rotundatis supra subtusque molliter pubescentibus firme membranaceis subopacis subtilissime pellucido-punctatis, nervo centrali subtus prominulo utrinque nervos 5 alternos adscendentes subtus prominulos mittente, petiolo fusce villosa, pedunculo petiolum multum superante glabro, bractæ apice peltatæ pelta triangulari vel semilunari margine hirtella, bacca lateraliter compressa vertice glanduloso-pubescente. ♂ In Jamaica (Fadyen n. 82! in h. Kew.). *Artanthe scabra* v. foliis minoribus Miq.! in h. Kew. Ramuli fusce remote villosi. Limbi 0,09 longi, 0,04 lati. Petioli 0,002-0,003 longi. Pedunculi 0,012 longi. Amenta florentia 0,07 longa, 0,003 crassa. (v. s.)
182. P. GRACILE (Ruiz et Pav. flor. peruv. 1. p. 35. t. 59. f. b), foliis brevissime petiolatis subovato-subelliptico-lanceolatis apice acutis basi inæquali vel subæquali subattenuatis supra rugosulis non scabris pubescentibus subtus molliter dense pubescentibus subcoriaceis opacis, nervo centrali subtus prominulo nervos utrinque 4-5 alternos adscendentes subtus prominulos nervulis fortioribus intermixtos mittente, petiolo hirsuto, pedunculo hirsuto petiolum superante, amento mucronulato, bractæ apice peltatæ pelta triangulari margine hirsuta, stigmatibus linearibus recurvis, bacca obconica glabra. In Chinchao nemoribus (R. et P. l. c.), San Carlos (Mathews n. 1695! in h. Kew.). — Vahl enum. v. 1. p. 316. *Artanthe gracilis* Miq. syst. p. 448 et in Hook. Lond. journ. bot. v. 4. p. 456. Ramuli molliter hirsuti. Limbi 0,08-0,085 longi, 0,03-0,035 lati. Petioli 0,006 longi. Pedunculi 0,012 longi. Amenta matura 0,003 crassa. (v. s.)
183. P. ERIOPODON, foliis breviter petiolatis elliptico-lanceolatis apice acuminatis, acumine acuto, basi subæquali attenuatis acutis supra verruculoso-scabris subtus ad nervos molliter hirtellis membranaceo-rigidis, nervo centrali subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes subtus prominulos utrinque plerumque 7 mittente, petiolo hirtello, pedunculo glabro petiolum

paulo superante, amento mucronulato, bractæ apice peltatæ pelta elliptico-semilunari margine brevius hirsuta, bacca glabra. ♂ In Nova Granata (Linden n. 840! in h. DC.). Artanthe eriopoda Miq.! in Hook. Lond. journ. bot. v. 4. p. 459. Ramuli breves remote pilosuli. Limbi 0,15 longi, 0,05 lati. Petioli 0,01 longi. Amenta ante anthesin 0,09-0,1 longa, 0,002 crassa. (v. s.)

184. *P. aduncum* (L. sp. p. 41), foliis breviter petiolatis oblongo-ellipticis apice acuminatis, acumine acuto, basi plus minusve inæquali rotundatis cordulatisve vel latere minore acutis majore rotundatis rigido-membranaceis subopacis pellucido-punctatis supra scabris pilosulisque subtus præsertim ad nervos subpubescentibus pubescentibusve sæpe margine ciliolatis, centrali nervo subtus prominente ad  $\frac{1}{2}$  longitud. nervos utrinque 6-7 alternos ascendentes subtus prominentes supremos ad apicem ductos mittente, petiolo basi ima vaginante utrinque hirtello, pedunculo petiolum superante puberulo, amento plus minusve uncinato mucronato maturo quam limbus parum brevior, bractæ apice peltatæ pelta triangulari margine fusce vel aureo-hirsuta, bacca glabra. ♀ In Brasilia (L.! herb. excl. syn. Piso; Luschnath n. 72! et Riedel n. 439! in h. Petrop.; Sello! in h. Berol.; Martius n. 1204! in h. DC.), ad oram septentrion. Amazonum ad ostium Rio Negro (Spruce n. 1763! in h. DC.), prope Bahiã (Salzmänn n. 594! in h. DC.), in Nova Granata ad Antioquia (Triana n. 33! et 338! in h. Triana), in declivitate montis Impossibile inter Caripe et Cumanacoa (Humb. et Bonpl.! nov. gen. v. 1. p. 50 et h. Willd. n. 664!), republica æquatoriali (Fraser! in h. DC.), in Guyana anglica (Schomburgk n. 1! in h. DC.), Surinam (Weigelt! in h. DC.), Guyana gallica (Aublet hist. pl. Guy. v. 1. p. 21), Panama (Sutton et Hayes n. 740! 742! 743! 746! 747! in h. Kew.), Mexico prope Xalapa (Galeotti! in h. Kew. et Galeotti n. 6008! in h. Deless.), prope Hacienda de la Laguna (Schiede! in h. Berol.), in Trinitate (Lockhart! Crueger! in h. Kew. et Sieber n. 259! in h. DC.), Portorico (Wydyler n. 261! et Bertero! in h. DC.), S<sup>o</sup> Domingo (Mayerhoff n. 340! in h. Berol.), Jacquin in h. Willd. n. 663!), Jamaica (Swartz! in h. DC. et Murray n. 603!, 604! et 367! in h. DC.; March n. 1900! et Alexander! in h. Kew.), Cuba (Grisebach Cat. pl. cub. p. 65), Barbades (Maycock flor. Barb. p. 39). P. longum folio nervoso pallide viridi humilior Sloane! hist. p. 137. t. 87. f. 2 et collect. orig. icon. cum planta origin. p. 135. t. 81 in Brit. Mus. Stururus arborescens fructu aduncum Plum. Am. ed. Par. p. 58. t. 77; Linn. hort. C. iff. p. 140. P. aduncum Hill dict. v. 26. p. 18. t. 18. f. 2; Aublet. l. c.; Jacq. ic. rar. v. 2. p. 2. t. 210; Willd. spec. v. 1. p. 163; Vahl enum. v. 1. p. 319; Opiz in Reliq. Haenk. fasc. 3. p. 154; Vell. flor. flum. t. 60; Mart. syst. mat. med. p. 99; Loudon Encycl. p. 28. n. 502. P. scabrum Lam. ill. p. 80. P. cettidifolium Kunth! in H. et B. nov. gen. v. 1. p. 50, synopsis. v. 1. p. 106. Steffensia adunca Kunth! in Linn. v. 13. p. 633. Artanthe adunca Miq.! in Comm. phyt. p. 49. t. 4. f. f. syst. p. 449, in Mart. flor. Bras. fasc. 11. p. 46. t. 14. f. 2 et in Nederl. Kruidk. v. 1. p. 222; Grisebach l. c. Artanthe Velloziana Miq.! in Linn. v. 20. p. 155, in Mart. flor. Bras. l. c. p. 44 et in Hook. Lond. journ. bot. v. 4. p. 456. Poivrier à fruit crochu Descourtils flor. Ant. v. 3. p. 355. t. 230. Rami ramulique hispidulo-scabri vel pubescentes, adulti minute tuberculati. Limbi 0,17-0,23 longi, 0,07-0,08 lati. Petioli 0,01 longi. Amenta matura 0,14 longa, 0,004 crassa. (v. s.)

185. *P. angustifolium* (Ruiz et Pav.) flor. peruv. l. p. 38. f. 57. f. a), foliis breviter petiolatis suboblique lanceolato-elliptico-oblongis longe acuminatis, acumine acuto mucronulatoque, basi inæquali rotundatis semicordatisve supra verruculis et pilis asperis subtus molliter hirtis-pubescentibus rigido-subcoriaceis subopacis pellucido-punctatis, nervo centrali subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos ascendentes subtus prominulos utrinque 7-8 supremos ad apicem ductos mittente, petiolo hirsuto basi ima vaginante, pedunculo hirtis petiolum vix duplo superante, bractæ apice peltatæ pelta triangulari margine hirsuta, stigmatibus filiformibus, bacca glabra. ♀ In Peruvia (Pav. l in h. Boiss.), Andibus peruvianis in nemoribus Huanuco (R. et P. in h. Willd. n. 659! et Pœppig n. 1526! in h. Berol. et DC.), Brasilia (Riedel n. 439! in h. Petrop.), prov. Bahia (Blanchet, n. 3148! et 33! in h. DC. et Luschnath! in h. Boiss.), Panama prope Chagres (Fendler, n. 269! in h. Kew.). P. granulosum R. et P. flor. peruv. v. 1. p. 38. t. 64. f. b, Vahl enum. v. 1. p. 313,

*Opiz* in Reliq. Haenk. fasc. 3. p. 157. *P. elongatum* Vahl enum. v. 1. p. 312. *P. purpurascens* A. Dietr. syn. v. 1. *Steffensia elongata* Kunth! in Linn. v. 13. p. 637, Gaudich. iter Bonite, t. 95. *Artanthe elongata* Miq.! Syst. p. 434 et in Hook. Lond. journ. bot. v. 4. p. 455. *Artanthe granulosa* Miq. Syst. p. 435. Benth. bot. of Sulph. p. 167. Rami glabri, ramuli molliter hirtelli vix aliquid asperuli. Limbi 0,12-0,2 longi, 0,055-0,06 lati. Petioli 0,005-0,006 longi. Pedunculi 0,01-0,015 longi. Amenta matura 0,12-0,13 longa, 0,004 crassa. Confer ad num. 119. (v. s.).

α, *cordulatum*, limbo basi cordulato, lobo majore plus minusve libero rotundato-subauriculæformi, petiolo brevissimo hirsuto. Variat: 1° ramulis dense molliter hirsutis. In Cuba prope Havanam (Sagra, n. 71 in h. DC.), Venezuela (Funcke n. 1301 in h. DC.). — 2° Ramulis dense appresse puberulis scabridisque, foliis rigidioribus. In Brasiliæ prov. Minas-Geraes (Clausen, n. 5001 et 3001 in h. Deless. et DC.), Santa Luzia (Pohl, n. 746! in h. Vindob.), Caracas (Jose de Vargas, n. 2781 in h. DC.), prope urbem Panama (Seemann! in h. Kew.), Mexico (Sallé et Botteri! in h. DC., Pav. 1 in h. Boiss.). *Artanthe Kunthiana* Miq.! Syst. p. 453, in Linn. v. 20. p. 156 et in Mart. flor. Bras. fasc. 11. p. 47. — 3° Ramulis dense molliter olivaceo-pubescentibus, bractææ pelta margine flavicanti-hirsuta. In Peruvia prope Biscaina alt. 3000 ped. (Fendler, 11301 in h. DC.). — 4° Ramulis hirsutis adultis verruculoso-scabris, bractææ pelta margine aureo-hirsuta. In Peruvia (Mathews, n. 17061 et 32251 in h. Kew.), Brasiliæ prov. Ceara (Gardner, n. 18471 in h. Deless.), et Goyaz (Gardner, n. 43551 in h. Kew.), Caracas (Jose de Vargas! in h. DC.), Nova Granata (Holton, n. 2371 in h. DC.), Colombia (Hartweg n. 13961 in h. Kew., Linden n. 338 in h. Kew. et Triana n. 421). *Artanthe Cearensis* Miq.! Syst. p. 456 et in Mart. flor. Bras. fasc. 11. p. 47. — 5° Ramulis fusciscentibus junioribus pulverulento-puberulis, foliis oblongis apice acuminatis, acumine acuto mucronulatoque, amento ante florat. quam folium multo brevius aureo. In Peruvia (Poeppig n. 541 vel 1526 β1 in h. Boiss.).

β, *Ossanum*, limbis angustioribus 0,035-0,045 latiss, limbi latere majore libero rotundato auriculæformi petiolum velante, foliis supra scabris bullatis subtus dense molliter tomentosis. In Bolivia (h. Petrop. n. 1461) et Cuba (Sagra, in 288! Ossa! n. h. DC.). (v. s.)

186. *P. MISANTLENSIS*, foliis brevissime petiolatis oblongis, apice acuminatis, acumine obtuso, basi inæquali acutis supra glabris subbullatis subtus ad nervos velutino-puberulis rigidis subopacis pellucido-punctulatis, nervis subtypis prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos adscendentes utrinque 5 supremos ad apicem ductos mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum æquante, amento maturo folium æquante obtuso, bractææ apice truncato-peltatæ pelta triangulari margine exteriori ciliolata, rhachi foveolata, bacca glabra vertice olivacea basi subimmersa. ♀ In Misantla (Fischer, n. 761 in h. Petrop.). Rami verruculis albidis subasperuli. Ramuli glabri. Limbi 0,14 longi, 0,04 lati. Petioli 0,005 longi. Amenta matura 0,004 crassa. (v. s.)

187. *P. FORMOSUM*, foliis petiolatis ample ovatis apice longe acuminatis, acumine acuto, basi æquali cordatis supra glabris lævibus subnitidulis sublus sericeo-pubescentibus membranaceis opacis sub lente fortiori subtiliter pellucido-punctulatis, nervis subtus prominentibus, nervo centrali ad  $\frac{1}{2}$  longitud. nervos adscendentes utrinque 4-5 quorum 3-4 fere e basi ortos mittente, nervulis primariis fere transversis, petiolo ad  $\frac{1}{2}$  longitud. vaginante pubescente, bractææ apice peltatæ pelta subrotunda margine dense hirsuta. ♀ In Peruvia? (Mathews, n. 16971 in h. Kew.), prope Casapi (Mathews, n. 16981 in h. Kew.). *Artanthe formosa* Miq.! Syst. p. 386, illustr. p. 71. t. 69. Art. *Endlicheriana* Miq.! in Hook. Lond. journ. bot. v. 4. p. 441. Ramuli puberuli. Limbi 0,36 longi 0,12 lati. Petioli 0,04 longi. Amenta nascentia. (v. s.)

188. *P. AFFINE*, foliis brevissime petiolatis oblongis apice breviter attenuato-acuminatis basi inæquilateraliter leviter inæquali acutiusculis vel obtusis supra lævissimis glabris subnitidulis subtus in nervis canescenti-hirtellis pallidis



tenuiter membrananeis crebre pellucido-punctulatis, nervo centrali subtus prominulo nervos utrinque circ. 6 inferiores tenuiores approximatos supremos ad apicem ductos mittente, petiolo præsertim postice dense hirtello, pedunculo quam petiolus parum brevior hirtus, amento quam folium multoties brevior breviter mucronato, bractæ conchæformi-peltatæ pelta subrotunda margine præsertim interiore villosa-ciliata, stigmatibus crassiusculis. ♀ Prope Paramaribo (Kegel, n. 821, fide Miq. in Linn. v. 22. p. 77) et Surinam (Miq. in Linn. v. 18. p. 228). *Artanthe affinis* Miq. in Linn. v. 18. l. c. Piper affine Martens et Gal. est *Peperomia quadrifolia* Kunth. Ramuli flexuosi, hirtuli. Limbi 0,13-0,15 longi, 0,05-0,07 lati. Petioli 0,005 longi. (Miq. l. c.)

189. *P. BENNETTIANUM*, foliis petiolatis superioribus elliptico-lanceolatis inferioribus lanceolato-ellipticis omnibus apice longiuscule acuminatis, acumine acuto subulatoque, basi æquali vel breviter inæquali acutis utrinque glabris membranaceis vel firmule membranaceis supra subfusciscentibus creberrime pellucido-punctulatis opacis, nervis supra vix prominulis subtus prominulis, centrali ad  $\frac{1}{2}$  -  $\frac{1}{4}$  longitudinis nervos alternos adscendentes utrinque 5 supremos ad apicem ductos infimos subtiles mittente, petiolo glabro ad  $\frac{1}{2}$  longitud. vaginante, pedunculo florente petiolum æquante glabro, amento florente quam folium quadruplo brevior, bractæ apice truncato-peltatæ pelta subrotunda margine dense hirsuta, stigmatibus linearibus. ♀ In Brasilia (Langsdorff n. 99! in h. Petrop.). Ramuli glabri, fusciscentes. Limbi 0,14-0,15 longi, 0,07-0,062 lati. Petioli 0,012 longi. Amenta florentia 0,004 crassa. (v. s.)

190. *P. TEJUCENSE*, foliis petiolatis ellipticis vel subovato-ellipticis apice brevissime acuminatis, acumine acutiusculo, basi breviter inæquali acutis vel obtusis utrinque glabris supra saturate viridibus verruculis albidis et glandulis atris adpersis hinc tactu asperulis subtus pallidioribus fusciscenti-nitidulis rigidiuscule coriaceis sparse subpellucido-punctulatis, centrali nervo subtus prominente ad  $\frac{1}{2}$  -  $\frac{1}{4}$  longitud. nervos adscendentes utrinque 5-6 supremos ad apicem ductos infimos multum subtiliores mittente, bractæ apice peltatæ pelta exigua rotundata glabra cæterum pilis longis luteo-griseis fere tota hirtotomentosa, stamp., stigmatibus 3 abbreviatis crassis puberulis. ♀ In Brasilia ad Tejuco (Vauthier, n. 481! in h. Deless.). *Artanthe Tejucensis* Miq. Syst. p. 496. Ramuli glabri, læves. Limbi 0,12-0,14 longi, 0,06-0,07 lati. Petioli 0,01 longi. (ex Miq. l. c.)

191. *P. HOSTMANNIANUM*, foliis brevissime petiolatis ovato-oblongis apice acuminatis, acumine acuto mucronulato, basi subinæquali latere uno longiore rotundatis altero attenuato-acutis supra glabris atro-viridibus nitidis subtus ad nervos nervulosque subtiliter adpresse puberulis coriaceo-membranaceis obsolete pellucido-punctulatis, nervo centrali subtus prominulo ad  $\frac{1}{2}$  longitud. nervos adscendentes nervulis fortioribus intermixtos utrinque 4-6-8 supremos per anastomoses ferè ad apicem ductos mittente, petiolo ad  $\frac{1}{2}$  longitud. vaginante basi dorso sursum utrinque hirtello, pedunculo petiolum æquante, bractæ apice peltatæ pelta triangulari vel semilunari margine hirsuta, bacca lateraliter compressa vertice glanduloso-puberula vel glabra. ♀ In Brasilia boreali prope San Gabriel da Cocheira, ad Rio Negro (Spruce, n. 2213! in h. DC.) et in prov. Barra (Spruce, n. 11! in h. DC. et 1348! in h. Francovil.), montib. Orgaos (Lhotszky! in h. DC.), Guyana anglica (Appun! n. 398! in h. Kew.) et Guyana gallica (Sagot, n. 536! in h. Kew.), ad Surinam (Hostmann, n. 116! in h. Kew.), Belize (March n. 1899! in h. Kew.). P. prægnans Rich.! in h. Francovil. *Artanthe Hostmanniana* Miq.! in Hook. Lond. journ. bot. v. 4. p. 465. Ramuli glabri læves, juniores hirtelli. Limbi 0,16 longi, 0,065 lati. Petioli pedunculique 0,005-0,007 longi. Amenta matura 0,09 circ. longa, 0,003 crassa. (v. s.)

α, *Berbicense*, foliis basi inæquali acutiusculis vel obtusis nervo centrali nervos utrinque 3-4 mittente, bacca vertice hirtella. ♀ In Surinam (Forbes n. 1234! et 943! in h. Kew.) et Guyana anglica (Schomb. n. 23! aut 53! in h. DC. et Kew.). *Artanthe berbicensis* Miq. syst. p. 500.

β, *ramiflorum*, ramulis pubescentibus. ♀ In Brasiliæ prov. Rio Negro in vicinibus Barra (Spruce in h. DC.), sylvis Guyanæ (Rich.! in h. Francovil.)

Surinam (Kappler n. 1397 in h. Francovil.). *Artanthe ramiflora* Miq. in Kappler exsicc. 1. c.

192. P. PSEUDO-CHURUMAYU, foliis breviter petiolatis amplis subrotundato-ovatis apice breviter acuminatis, acumine acuto, basi æquali lato-rotundatis supra glabris subtus ad nervos puberulis membranaceo-coriaceis opacis pellucido-punctatis, nervo centrali subtus prominente ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes supremos ad apicem ductos utrinque 5-6 infimos subtiliores mittente, petiolo glabro ad limbum usque vaginante, pedunculo glabro petiolum superante, bractæ apice peltatæ pelta triangulari semilunari vel subrotundata margine late griseo-ochraceo-villosa. ♀ In Brasilia ad flumen Amazonum (Poeppig n. 3059! in h. DC. et Berol.). Steffensia pseudo-Churumayu Kunth! in Linn. v. 13. p. 662. *Artanthe pseudo-Churumayu* Miq. syst. p. 407, illustr. p. 74. t. 75. Ramuli glabri. Limbi 0,23 longi, 0,18 lati. Pedunculi 0,025 longi. Amenta florentia 0,16 longa, 0,004 crassa (v. s.).
- β, *membranaceum*, foliis membranaceis magis pellucidis pellucido-punctatis, Ad flumina Cassiquari, Vasiva et Pacimoni (Spruce n. 3433! in h. DC.) (v. s.)
193. P. LHOTZKYANUM, foliis petiolatis oblongo-ellipticis apice acuminatis, acumine acuto, basi subinæquali acutis subrotundatisve utrinque glabris membranaceis pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos subadscendentes utrinque 4 supremos ad apicem ductos mittente, petiolo glabro pedunculum glabrum subsuperante, amento cylindrico quam folium dimidio brevius, bractæ apice peltatæ pelta semilunari margine hirsuta, bacca glabra. ♀ In Brasilia prope Rio Janeiro (Lhotzky! in h. Berol.). *Artanthe Lhotzkyana* Miq. syst. p. 417 et in Mart. flor. Bras. fasc. 11. p. 40. Rami glabri. Ramuli glabri. Limbi 0,14 longi, 0,045 lati. Petioli 0,015 longi. Amenta matura 0,004 crassa (v. s.).
194. P. SCHACHII, foliis petiolatis inferioribus late ovatis apice oblique acuminatis, acumine acuto, basi lato-rotundatis superioribus ellipticis basi obtusis vel acutiusculis omnibus utrinque glabris membranaceis pellucido-punctatis, centrali nervo subtus prominulo ad  $\frac{1}{2}$  longitud. usque nervos alternos patulo-adscendentes utrinque 3-4 nervulosque mittente, petiolo glabro, pedunculo quam petiolus duplo brevius, bractæ pelta triangulari-semilunari margine ciliolata, staminibus 4, ovario basi villosulo. ♀ In ins. Trinitatis (Schach! in h. Kew., Lockhart ibid.). *Artanthe Schachii* Miq. in Hook. Lond. journ. of. bot. v. 4. p. 448. Ramuli glabri. Limbi 0,17-0,18 longi, 0,1-0,11 lati; superiores 0,15-0,17 longi, 0,06-0,085 lati. Petioli 0,035-0,01 longi (v. s.).
195. P. COCCOLOBOIDES (Kunth! olim), foliis longiuscule petiolatis elliptico-ovatis apice vix acuminatis basi valde inæquali profunde cordatis, lobis incumbentibus lobo majore auriculatim extenso, supra pilis sparsis adpresse scabridohirtellis subtus densissime tomentosis coriaceis, nervis subtus prominentibus, centrali ad  $\frac{3}{4}$  longitudinis nervos alternos subadscendentes latere majore 8 minore 5 infimos magis approximatos fere e basi solutos mittente, petiolo ad limbum vaginante dorso fusce villosulo, alis linearibus persistentibus, pedunculo fusce villosulo quam petiolus duplo brevius, amento cylindrico folium superante, bractæ subcucullato-truncato-peltatæ pelta semilunari margine fusce hirsuta pedicello brevi lato subcucullato villosulo, stigmatibus subulatis. ♀ In Brasilia meridionali (Seil! in h. Berol.), Serra de Mendanha Brasiliæ (Gardner n. 5185, Miq. in Hook. Lond. journ. bot. v. 4. p. 446). Steffensia coccoloboides Kunth! in Linn. v. 13. p. 666 et in h. Berol. *Artanthe coccoloboides* Miq. syst. p. 397 et in Hook. Lond. journ. bot. 1. c. Ramuli villosi. Limbi 0,22-0,24 longi, 0,11-0,14 lati. Petioli 0,04 longi. Amenta florentia 0,003 crassa. (v. s.)
196. P. MORITZIANUM, foliis longiuscule petiolatis ovato-oblongis subæquilatis breviter acuminatis basi inæquali cordatis, lobo majore perpendiculari minore horizontaliter truncato coriaceo-membranaceis subepunctulatis supra glabris subtus molliter pubescentibus, centrali nervo ad  $\frac{1}{2}$  longitud. nervos 8-9 supremos fere ad apicem ductos omnes crebre anastomosantes mittente, petiolo ad  $\frac{1}{2}$  longitud. vaginante, alis linearibus, pedunculum multum superante, amento folium superante, bractæ cucullato-peltatæ vertice triangulari margine villosulo

stigmatibus linearibus filamentis longiuscule exsertis. ♀ In Colombia (Moritz in h. Schlecht. fide Miq. in Linn. v. 18. p. 714). *Artanthe Moritziana* Miq. in Linn. l. c. Limbi 0,37 longi, 0,205 lati. Petioli 0,08 longi. Amenta 0,42 longa, 0,004 crassa. (v. s.)

197. *P. NOBILE*, foliis longiuscule petiolatis oblongo-ovato-lanceolatis apice acuminatis acutiusculis basi valde inæquali profunde cordatis supra glabris subtus ad nervos tenuissime puberulis firmis opacis, nervis subtus prominentibus, centrali nervo ad  $\frac{3}{4}$  longitud. nervos alternos subadscendentes latere majore 7 minore 6 mittente, infimis duobus magis approximatis subpatulis petiolo glabro ad limbum vaginante, alis linearibus persistentibus, pedunculo glabro quam petiolus brevior, amento folium æquante cylindraceo crasso, bracteæ apice truncato-peltatæ vertice subrotundo margine pedicelloque dorso hirsutis, pedicelli lati subcucullati pilis longioribus, bacca glabra. ♀ In Venezuela prope Tovar (Fendler n. 1144! in h. DC.). *Artanthe flagellaris* Miq. syst. p. 393? Ramuli glabri. Limbi 0,25-0,30 longi, media parte 0,14-0,20 lati. Petioli 0,07 longi. Amenta matura 0,008 crassa. Bacca 0,0025 longa. (v. s.)

2° Folia multiplinervia.

198. *P. INSIGNE* (Kunth olim), foliis longiuscule petiolatis oblongo-ellipticis apice subito breviter acuminatis, basi valde inæquali dimidiato-cordatis supra glabris subtus nitidulis ad nervos hirtellis rigidis opacis subtiliter pellucidopunctulatis, centrali nervo subtus prominente ad  $\frac{3}{4}$  longitud. nervos alternos latere majore 8 minore 7 mittente, inferioribus magis patulis fere e basi solutis infimis subtilioribus; petiolo ad limbi basin usque vaginante, alis linearibus, dorso hirtopubescente, pedunculo quam petiolus brevior, amento cylindrico quam folium longiore, bracteæ apice truncato-peltatæ pelta semilunari margine ciliolata pedicello cucullato margine ciliolato, bacca glabra lateraliter compressa. ♀ In Guyana gallica (Poiteau! in h. Deless.), prope Cajennam (Kunth in Linn. v. 13. p. 667). *Steffensia insignis* Kunth l. c. *Artanthe insignis* Miq. l. syst. p. 394. Ramuli hirtopubescentes. Limbi 0,26 longi, 0,14 lati. Petioli 0,06-0,07 longi. Amenta matura 0,4 longa, 0,004 crassa. (v. s.)

199. *P. PSEUDO-POTHIFOLIUM*, foliis longiuscule petiolatis subovato-lanceolato-oblongis apice angustato-acuminatis, acumine acuto, inferioribus basi æquali profunde cordatis, lobis discretis, superioribus basi valde inæquali profunde cordatis, lobo majore subauriculæformi petiolum velante supra glabris subtus ad nervos dense hirtellis, centrali nervo subtus prominulo ad  $\frac{3}{4}$  longitud. nervos alternos subadscendentes utrinque 6-7 supremos longe ab apice per anastomoses evanidos inferos utrinque 2-3 e basi divaricantes sursumque nervulos aliquot fortiores mittente, petiolo dorso molliter villosa ad limbum vaginante, alis linearibus persistentibus, pedunculo dense hirtello quam petiolus paulo brevior, amento maturo quam folium paulo brevior, bracteæ apice truncato-peltatæ pelta triangulari margine hirsuta, stigmatibus abbreviatis. ♀ In Brasilia (Riedel n. 513! in h. Petrop.). Rami glabri. Ramuli dense molliter villosi. Limbi 0,21 longi, basi 0,075 lati. Petioli 0,026-0,04 longi. Amenta matura 0,004 crassa. (v. s.)

200. *P. POTHIFOLIUM* (Kunth! olim), foliis longiuscule petiolatis oblique ovato-lanceolatis angustato-acuminatis basi valde inæquali profunde cordatis utrinque glabris subcoriaceis opacis, centrali nervo subtus prominente nervos utrinque 4 alternos adscendentes nervulis fortioribus intermixtos infimos utrinque 1-2 e basi solutos per lobos divaricantes omnes subtus prominulos mittente, petiolo ad limbum usque vaginante glabro, alis angustis linearibus coriaceis persistentibus glabris, pedunculo quam petiolus multum brevior glabro, bracteæ apice truncato-peltatæ vertice subrotundo semilunari margine ciliato pedicello cucullato dorso et margine hirsuto, bacca vertice fusce puberula lateraliter compressa. ♀ In Brasilia prope Rio de Janeiro (Gaudich. n. 1105! in h. DC. et Riedel n. 284! 17! 88! 92! 93! in h. Petrop.), Petropolim (Wawra et Maly n. 472 fide Wawra bot. Ergeb. p. 138), in sylvis umbrosis ad Coralfaga Brasiliæ (Pohl n. 22! et 5024! in h. Vindob.). *P. truncatum* Vell. flor. flum. v. 1. t. 57. *Steffensia pothifolia* Kunth! in Linn. v. 13. p. 670. *Artanthe pothifolia* Miq. l. syst. p. 899, illustr. p. 73. t. 74 et in Mart.

flor. Bras. fasc. 11. p. 38. Rami ramulique glabri. Limbi 0,32-0,25 longi, sub medio 0,08-0,07 lati. Petioli 0,05-0,06 longi. Pedunculi 0,01 longi. Amenta ante florationem 0,1 longa, 0,0015 crassa. (v. s.)

201. *P. LANGSDORFFIANUM*, foliis longe petiolatis lanceolato-oblongis apice angustato-acuminatis basi valde inæquali cordatis, lobo minore rotundato majore subauriculæformi petiolum velante partim cum petiolo adnato, utrinque glabris subtus fusco-punctulatis firmis opacis parce pellucido-punctulatis, centrali nervo subtus prominente ad  $\frac{1}{2}$  longitud. nervos alternos latere minore 7 majore 9 mittente, superis adscendentibus nervulis fortioribus sæpe intermixtis subtus vix prominulis supremis ante apicem per anastomoses evanidis inferis latere minore 2 majore 4 e basi solutis divaricantibus subtilioribus, petiolo glabro ad limbi latius majus vaginante, alis linearibus persistentibus glabris, pedunculo quam petiolus multum brevior glabro, amento maturo quam folium parum brevior, bractæ apice truncato-peltatæ pelta triangulari margine dense hirsuta pedicello lato subcucullato dorso producto hirsutoque, rhachi foveolata, bacca fere tota immersa vertice fusce pubescente, stigmatibus abbreviatis. ♀ In Brasilia (Riedel et Langsdorff n. 95! in h. Petrop.). Ramuli glabri. Limbi 0,2 longi, 0,037 media parte lati. Petioli 0,035 longi. Amenta matura 0,004 crassa. (v. s.)
202. *P. HILLIANUM*, foliis longe petiolatis lanceolato-oblongis apice acuminatis, acumine acutiusculo, basi profunde inæquali cordatis, lobo majore auriculæformi petiolum velante, utrinque glabris subtus nigro-punctulatis pallidioribus membranaceis opacis parce pellucido-punctulatis, centrali nervo ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. nervos alternos subadscendentes latere majore 7 minore 6 sursumque nervulos fortiores mittente, supremis ante apicem per anastomoses evanidis, petiolo glabro ad limbum vaginante, alis linearibus persistentibus, pedunculo glabro quam petiolus multoties brevior, amento florente quam folium multoties brevior, bractæ apice truncato-peltatæ vertice semilunari margine fusce ciliato pedicello subcucullato. ♀ In Brasilia (Riedel n. 481 et Langsdorff n. 101! in h. Petrop.). Ramuli glabri. Limbi 0,26 longi, 0,12 lati. Petioli 0,055 longi. (v. s.)
203. *P. PARAENSE*, foliis breviter petiolatis oblongo-lanceolatis lanceolatisve apice acuminatis, acumine acuto mucronulatoque, basi leviter inæquali obtusis acutiusculisve supra glabris nitidis subtus pallidioribus ad nervos nervulosque hirtello-puberulis membranaceo-rigidulis opacis pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$ - $\frac{3}{4}$  altitud. nervos alternos suboppositosve subadscendentes utrinque 4 supremos ad apicem ductos magis patulos infimum utrinque unum e basi solutum sursumque tota longitudine nervulos fortiores subpatulos numerosos mittente, petiolo ad  $\frac{1}{2}$  longitud. vaginante dorso hirtello, pedunculo puberulo petiolum æquante, amento florente quam folium triplo brevior, bractæ apice peltatæ pelta triangulari margine fusce hirsuta. ♀ In sylvis cæduis prov. Paraensis passim (Mart! in h. monac.). *Artanthe paraensis* Miq! in Mart. flor. Bras. fasc. 11. p. 60. Ramuli hirtelli. Limbi 0,135 longi, 0,043 lati. Petioli 0,006 longi. Amenta florentia 0,0015 crassa. (v. s.)
204. *P. MAREQUITENSE*, foliis breviter petiolatis ovato-acuminatis apice acutis basi æquali obtusis supra glabris subtus dense molliter pubescentibus membranaceis subopacis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos suboppositos adscendentes utrinque 4-5 inferiores utrinque 2 fere e basi ortos infimos subtiliores supremos alternos fere ad apicem ductos mittente, petiolo ad  $\frac{3}{4}$  longitud. vaginante basi dorso et sursum utrinque dense fusce tomentoso pedunculum fusce tomentosum superante, amento quam folium multum brevior, bractæ apice peltatæ pelta triangulari margine aureo-hirsuta pedicello brevissimo dorso inter flores producto. ♀ In Novæ Granatæ prov. Marequita alt. 800 hexap. (Linden n. 1696! in h. DC.) et prov. Antioquia alt. 1700 m. (Triana n. 24! in h. Triana). Rami fusce hirtelli. Ramuli fusce tomentosi. Limbi 0,14 longi, 0,08 lati. Petioli 0,015 longi. (v. s.)
205. *P. MONTANUM* (Cas. DC. in Seemann journ. of bot. 1866 p. 214), foliis longe petiolatis subrotundo-ovatis apice subattenuatis acutiusculis basi cordatis utrinque glabris rigidulis subopacis epunctulatis 13-plinerviis, medianis ner-

vis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos suboppositos subadscendentes utrinque 6 supremos ad apicem ductos inferiores utrinque 3 fere e basi solutos patulo-subadscendentes mittente, petiolo glabro ad  $\frac{1}{2}$  longitud. vaginante, pedunculo glabro quam petiolus brevior, amento quam folium paullo brevior, bracteæ apice peltatæ pelta triangulari margine aureo-villosa pedicello lato subcucullato, bacca glabra secundum rhachin elongata. ♀ In Novæ Granatæ prov. Marequita monte Quindiu alt. 2200 m. (Triana n. 14! in h. DC.). Ramuli glabri. Limbi 0,21 longi, 0,17 lati. Petioli 0,05 longi. Amenta matura 0,003 crassa. (v. s.)

206. *P. OTTONLEFOLIUM* (Cas. DC. in Seem. journ. of bot. 1866 p. 213), foliis brevissime petiolatis elliptico-oblongis apice brevissime acuminatis, acumine acuto, basi subinæquali cordulatis utrinque glabris opacis, nervis subtus prominentibus, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos adscendentes supremos magis discretos ad apicem ductos mittente, nervulis primariis subtransversim parallelis, petiolo glabro, amento quam folium duplo brevior, pedunculo petiolum fere duplo superante, bracteæ apice peltatæ pelta triangulari margine dense pubescente. ♀ In Novæ Granatæ prov. Choco alt. 170 m. (Triana n. 32! in h. DC.) et Barbacoas alt. 200 (Triana n. 31, in h. suo). Ramuli glabri. Limbi 0,15 longi, 0,06 lati. Petioli 0,003 longi. (v. s.)

207. *P. MARTIANUM* (Kunth! olim), foliis petiolatis elliptico-ovatis apice acuminatis, acumine acutiusculo, basi æquali obtusis vel obtusiusculis utrinque glabris coriaceo-membranaceis obsolete pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis trifido nervos utrinque 2 fere e basi solutos sursumque interdum nervulos fortiores subpatulos mittente, petiolo glabro pedunculum glabrum superante, amento cylindrico quam folium quadruplo brevior, bracteæ apice truncato-peltatæ pelta triangulari subquadrangulari margine ciliolata, stigmatibus filiformibus. ♂ In Bræsilia (Sello! in h. Berol.), in Antillis (Grisebach flor. w. Ind. p. 172). Schilleria Martiana Kunth in Linn. v. 13. p. 690. Artanthe Martiana Miq.! syst. p. 422, illustr. p. 74. t. 77 et in Mart. flor. Bras. fasc. 11. p. 41; Grisebach l. c. (except. synonym. A. Schachii Miq.) Ramuli glabri. Limbi 0,1 longi, 0,045 lati. Petioli 0,01 longi. Amenta florentia 0,003 crassa. (v. s.)

208. *P. FUSCESCENS*, foliis petiolatis ovatis apice acuminatis, acumine acuto basi æquali cordatis breviter cordulatis supra præsertim ad nervos adpresse puberulis subtus dense molliter pubescentibus rigidis opacis fuscis, nervis subtus vix prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos subadscendentes utrinque 3 supremos ad apicem ductos infimos e basi medios paullo supra basin solutos mittente, petiolo basi ima vaginante dense molliter pubescente, pedunculo puberulo petiolum æquante, amento crasso maturo folium subsuperante obtuso, bracteæ apice peltatæ pelta triangulari margine dense fusce hirsuta, bacca nigra. ♂ In Bogota (h. Vindob.). Rami glabri, nigrescentes. Ramuli fusce pubescentes. Limbi 0,09 longi, 0,05 lati. Petioli 0,01 longi. Amenta matura 0,006 crassa. (v. s.)

209. *P. PEDUNCULARE*, foliis petiolatis ovato-vel elliptico-lanceolatis apice acutis basi æquali rotundatis cordulatisve utrinque glabris membranaceis subpellucidis pellucido-punctatis utrinque glandulosis, nervo centrali subtus prominulo nervos utrinque 3-4 ad  $\frac{1}{2}$  longitud. usque inferos utrinque 2 e basi solutos subadscendentes infimos subtiliores et 1-2 supremos longe ab apice per anastomoses evanidos sursumque tota longitudine nervulos fortiores magis patulos mittente, petiolo fere ad limbum usque vaginante, alis persistentibus linearibus, pedunculo hirtello petiolum superante, amento ramulum lateralem unifolium terminante, bracteæ apice truncato-peltatæ pelta semilunari carnosula margine exteriore pilosula, filamentis brevibus. ♂ In Guyana anglica (Schomburgk n. 696! in h. DC. et Berol.). Artanthe Schomburgkii Klotzsch! mss. in h. Berol. Artanthe peduncularis Miq.! syst. p. 531, illustr. p. 81. t. 91 et in Hook. Lond. journ. bot. v. 4. p. 322. Ramuli glabri. Limbi 0,06-0,07 longi, 0,025-0,03 lati. Petioli 0,008-0,012 longi. Pedunculi 0,025 longi. (v. s.)

210. *P. SOLMSIANUM*, foliis longiuscule petiolatis ovatis vel subrotundo-ovatis apice breviter attenuato-acutatis basi æquali rotundatis utrinque glabris fir-

mule membranaceis opacis subtilissime pellucide-punctulatis, nervis supra vix prominulis subtus prominentibus, centrali ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitudinis nervos alternos subadscendentes utrinque circ. 6 supremos ante apicem per anastomoses evanidos sursumque sæpe nervulos fortiores et inferos utrinque 2 magis divaricantes e basi solutos mittente, petiolo glabro basi vaginante, pedunculo per floracionem quam petiolus multum brevior glabro, amento florente folium subæquante, bractæ apice truncato-peltatæ vertice triangulari glabro pedicello subcucullato dorso subproducto carinato hirtello, ovario glabro. ♀ In Brasilia (Riedel n. 1171! in h. Petrop.). Rami ramulique glabri, fuscесcentes. Limbi 0,125-0,18 longi, 0,075-0,125 lati. Petioli 0,025-0,04 longi. Amenta florentia 0,003 crassa. (v. s.)

211. *P. CORUSCANS* (Kunth! in H. et B. nov. gen. 1. p. 53), foliis brevissime petiolatis ovato-acuminatis vel oblongo-ovato-acuminatis, acumine acuto, basi subinæquali subattenuato-obtusis utrinque glabris rigidis opacis, nervis subtus prominulis centrali ad  $\frac{1}{2}$ - $\frac{1}{4}$  longitud. nervos alternos subadscendentes utrinque 5-7 inferos utrinque 2 e basi solutos mittente, petiolo glabro, pedunculo glabro petiolum subæquante, bractæ apice peltatæ pelta triangulari parva margine villosa. ♀ Ad ripas fluminis Magdalenæ inter S. Bartholome et Carapatas (Humb. in h. Willd. n. 672!), Colombia (Hartweg n. 1398! in h. Boiss. et 1400! in h. Kew.), Trinitate (Grisebach in flor. w. Ind. p. 171). — Kunth synopsis v. 1. p. 108. Schilleria coruscans Kunth! in Linn. v. 13. p. 708. Artanthe coruscans Miq.! syst. p. 411; Grisebach l. c. Ramuli glabri. Limbi 0,1 longi, 0,07 lati. Petioli vix 0,005 longi. (v. s.)

β, *aureum*, bractæ pedicello subcucullato margine hirtello pelta magis aureo-hirsuta. In Novæ Granatæ prov. Bogota alt. 2700 m. et prov. Mariquita alt. 2200 (Triana n. 20 et 50 in h. suo). (v. s.)

212. *P. SAGRANUM* foliis breviter petiolatis ovatis apice acuminatis, acumine acuto, basi cordatis vel truncato-rotundatis utrinque glabris rigidulis opacis subtiliter pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos subadscendentes supremos ad apicem ductos inferos utrinque 2 e basi solutos magis arcuatos breviores mittente, petiolo glabro ad limbum usque vaginante, pedunculo glabro petiolum æquante, amento folium æquante, bractæ apice truncato-peltatæ pelta semilunari margine (præsertim postico) hirsuta, stigmatibus abbreviatis, bacca vertice subflavescente. ♀ In Cuba (R. de la Sagra! in h. Vindob.). Rami ramulique glabri. Limbi 0,13 longi, 0,07 lati. Petioli 0,008 longi. Amenta matura 0,003 crassa. (v. s.)
213. *P. HOOKERIANUM*, foliis petiolatis late ovato-ellipticis apice acuminatis, acumine acuto basi truncato-rotundatis utrinque glabris firmis fuscесcentibus, nervo centrali subtus prominulo ad  $\frac{1}{2}$  longitud. nervos subadscendentes subtus prominulos utrinque 4-5 inferos utrinque 3 e basi solutos mittente, petiolo glabro basi vaginante, pedunculo petiolum subæquante glabro, bractæ apice peltatæ pelta triangulari margine aureo-fusco-hirsuta, ovario glabro. ♀ In Peruvia ad Chachapoyas (Mathews n. 3222! in h. Kew. et Lechler n. 2081! in h. Kew., DC. et Boiss.). Artanthe Hookeriana. Miq.! in Hook. Lond. journ. bot. v. 4. p. 446. Ramuli glabri. Limbi 0,2 longi, 0,1-0,12 lati. Petioli 0,015 longi. Amenta florentia 0,04 longa. (v. s.)
214. *P. TRINERVE*, foliis breviter petiolatis oblongis apice acuminatis, acumine acuto, basi æquali cuneatis acutis utrinque glabris rigido-membranaceis opacis subtilissime pellucido-punctatis, nervo centrali subtus prominulo paulo supra basin nervos suboppositos utrinque 2 adscendentes supremos ad apicem ductos et nervulos permultos tota longitudine transversim parallelos mittente, petiolo glabro, pedunculo glabro petiolum paulo superante, amento cylindrico recto, bractæ apice peltatæ pelta semilunari margine hirtella bacca glabra. ♀ In isthmo panamensi prope St-Lorenzo Veraguas (Seemann n. 1217! in h. Kew., Sutton et Hayes n. 739! in h. Kew.). Artanthe trinervis Miq.! in Seem. Bot. Herald. p. 200. t. 41 (non Piper trinerve Vahl). Ramuli glabri. Limbi 0,12-0,15 longi, 0,035-0,045 lati. Petioli 0,008 longi. Amenta matura 0,05-0,06 longa, 0,003-0,0055 crassa. (v. s.)

215. *P. AMPLECTENS*, foliis breviter petiolatis ovatis apice attenuato-acutis obtusisve basi inæquali cordatis supra junioribus adpresse piloso-pubescentibus adultis scaberrimis subtus pubescentibus adultisque scabris rigidis opacis obsoleto pellucido-punctatis nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos suboppositos adscendentes utrinque 6 inferos 4 e basi solutos magis patulos mittente, petiolo basi ima vaginante basi dorso et sursum utrinque pubescente, pedunculo dense pubescente petiolum triplo superante, amento folium æquante, bractæ apice peltatæ pelta subrotunda margine fusce hirsuta, bacca glabra. ♂ In prov. Minas Geraës versus Salgado et Contendas inter arbusta (Mart.† in h. monac.). Artanthe amplectens forma longe petiolata Miq.† in Linn. v. 20. p. 154, in Mart. flor. Bras. fasc. 11. p. 44. Rami asperuli. Ramuli dense pubescentes. Limbi 0,08 longi, 0,035 lati. Petioli 0,005 longi. Amenta florentia 0,003 crassa. Pedunculi 0,015 longi. (v. s.)
216. *P. FLAVICANS*, foliis breviter petiolatis oblongo-ovatis apice acuminatis, acumine acuto, basi inæquali cordatis supra bullatis verruculis scabris et ad nervos puberulis subtus verruculis scabridis et ad nervos hirtellis rigidis opacis pellucido-punctulatis, nervis subtus prominentibus, centrali ad  $\frac{1}{2}$  longitud. nervos alternos latere majore 7 minore 6 mittente, superis adscendentibus inferis utrinque 3-4 e basi solutis divaricantibus supremis ad apicem ductis, petiolo ad  $\frac{1}{2}$  longitud. vaginante basi dorso sursumque utrinque dense pubescente, pedunculo petiolum quadruplo superante juniore dense pubescente, amento folium æquante, bractæ apice truncato-peltatæ pelta triangulari margine dense aureo-villosa, bacca glabra glandulis flavicantibus conspersa. ♂ In Brasilia (Riedel n. 289† in h. Petrop.). Rami verruculis albis obsiti. Ramuli pilis retrorsis dense hirsuti. Limbi 0,1 longi, 0,033 lati. Petioli 0,005 longi. Amenta matura 0,003 crassa. (v. s.)
217. *P. PALUSTRE*, foliis brevissimis petiolatis ovatis apice attenuato-acutiusculis inferioribus basi æquali superioribus basi inæquali cordatis supra pilis verruculisque scabris subtus hirtellis asperis coriaceis opacis epunctulatis utrinque flavicantibus, nervis subtus prominulis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos alternos superiores utrinque 5 adscendentes supremos ad apicem ductos inferos utrinque 4 e basi solutos divaricantes infimos subtiliores mittente, petiolo basi ima vaginante utrinque dense flavide hirsuto, pedunculo florente flavide dense hirsuto, amento florente quam folium brevius, bractæ apice truncato-peltatæ pelta semilunari subrotunda margine dense aureo-hirsuta. ♂ In Brasilia ad Barra do Tiquiliba (Riedel n. 114† et 125† in h. Petrop. et DC.). Rami flavicantes, pilis et verruculis asperati. Ramuli dense flavide hirsuti. Limbi inferiores 0,08 longi, 0,046 lati, superiores 0,06 longi, 0,035 lati. Petioli 0,003 longi. Amenta florentia 0,002 crassa. Pedunculi 0,015 longi. (v. s.)
218. *P. MULTIPLINERVIUM* (Cas. DC. in Seemann journ. of bot., 1866. p. 214), foliis petiolatis ovatis apice breviter acuminatis, acumine acuto, basi subtruncato-rotundatis ima parte ad petiolum acutiusculis utrinque glabris membranaceis opacis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervum utrinque alternatim fere ad apicem ductum adscendentem et 4 e basi solutos adscendentes mittente, petiolo glabro, pedunculo glabro quam petiolum paullo brevius, amento quam folium paullo brevius filiformi, bractæ apice peltatæ pelta semilunari margine pubescente. ♂ In Novæ Granatæ prov. Barbacoas (Triana n. 27† in h. DC.). Ramuli glabri. Limbi 0,095 longi, 0,065 lati. Petioli 0,012 longi. (v. s.)
219. *P. XYLOSTEIOIDES* (Steud. nom. p. 344), foliis breviter petiolatis elliptico-lanceolatis apice longe acuminatis, acumine acuto, basi æquali acutiusculis acutisve utrinque glabris rigidule membranaceis pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos subadscendentes utrinque 3 inferum nervum utrinque 1 paullo supra basin solum infimumque e basi solum mittente, petiolo glabro basi ima vaginante, pedunculo petiolum paullo superante, amento folium æquante filiformi, bractæ apice truncato-peltatæ pelta semilunari glabra pedicello cucullato intus ad apicem villosulo, staminibus 4?, rhachi pubescente, stigmatibus filiformibus. ♂ In Brasilia meridionali (Sello† in h. Berol. et Riedel n. 37† in h. Petrop.), Trinitate (Grisebach flor. w. Ind. p. 172). Schilleria xylosteoides Kunth† in Linn. v. 13.

- p. 677 et Humb. in h. Berol. *Artanthe xylosteoides* Miq.† syst. p. 422 et in Mart. flor. Bras. fasc. 11. p. 42. t. 13. f. 1, Grisebach† l. c. Ramuli glabri. Limbi 0,075 longi, 0,028 lati. Petioli 0,005 longi. (v. s.)
220. P. PAVONI, foliis petiolatis ovato-acuminatis, acumine acuto, basi truncato-obtusis supra ad nervos tenuissime puberulis subtus molliter pubescentibus membranaceo-firmis, nervo centrali ad  $\frac{1}{2}$  longitud. nervos utrinque 5-6 quorum 5 fere e basi soluti mittente, petiolo pubescenti-villoso, pedunculo pubescenti-villoso quam petiolus multum brevior, bractæ apice truncato-peltatæ vertice triangulari nudo vix reflexo margine interiore aureo-villoso unde bractæ primo aspectu oblonga. § In Peruvia (Mathews in h. Kew.). *Artanthe Pavoni* Miq.† in Hook. Lond. journ. bot. v. 4. p. 450. Ramuli pubescentes. Limbi 0,12 longi, 0,09 lati. Petioli 0,02 longi. (v. s.)
221. P. BOGOTENSE (Cas. DC. in Seem. journ. of bot., 1866. p. 214), foliis petiolatis ovatis apice acuminatis, acumine acuto, basi subæquali subrotundatis supra glabris subtus ad nervos hirtellis rigido-membranaceis subopacis subtilissimè pellucido-punctulatis, nervis adscendentibus subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos utrinque 2 supremos ad apicem ductos et utrinque 3 e basi solutos infimos subtiliores mittente, petiolo dorso hirsuto, bractæ apice peltatæ pelta triangulari margine aureo-hirsuta pedicello subcucullato margine aureo-hirtello. § In Novæ Granatæ prov. Bogota (Triana n. 501 in h. DC.) et prov. Mariquita Quidio Paras (Triana n. 201 in h. DC.). Ramuli juniores hirsuti, dein glabrati. Limbi 0,16 longi, 0,095 lati. Petioli 0,01 longi. (v. s.)
222. P. PAULIANUM, foliis petiolatis ovatis apice acuminatis basi subæquali rotundatis utrinque glabris rigidulis subtilissimè pellucido-punctulatis opacis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos utrinque 6-7 supremos ad apicem ductos inferos utrinque 3 e basi ortos subtiliores mittente, petiolo glabro ad  $\frac{1}{2}$  longitud. vaginante, pedunculo quam petiolus brevior, amento quam folium dimidio brevior, bractæ apice peltatæ pelta triangulari subrotunda margine aureo-villosa, pilis marginis exterioris longioribus, bacca glabra. § In Serra d'Estrella (Pohl n. 4770! in h. Vindob.) et in sylvis Jacarens. prov. St. Pauli (Mart.† in h. monac.). *Artanthe Haenkeana* Miq.† in Linn. v. 20. p. 145 (non Piper Haenkeanum Opiz). Ramuli glabri. Limbi 0,17-0,23 longi, 0,85-0,125 lati. Petioli 0,02 longi. (v. s.)
223. P. INCURVUM (Sieber† pl. Mart. n. 254 in h. Berol.), foliis petiolatis subrotundo-ovatis apice breviter acuminatis, acumine acuto mucronulatoque, basi cordulatis utrinque glabris nigro-punctulatis membranaceis, nervis subtus prominulis 3 centralibus supra basin solutis fere ad apicem ductis 3 utrinque fere e basi solutis, petiolo glabro pedunculum glabrum æquante, amento folium æquante, bractæ apice inflexo-peltatæ vertice inflexo semilunari subrotundo margine exteriori ciliolato, stigmatibus 3-4. § In Martinica (Sieber n. 254! in h. Berol.). Ramuli glabri. Limbi 0,08 longi, 0,06 lati. Petioli 0,025 longi. (v. s.)
- α. *majus*. Limbi 0,09 longi, 0,08 lati. Petioli 0,08 longi. — In sylvis Guyanæ (Rich.† in h. Francovil.).
224. P. VERAGUENSE, foliis longe petiolatis amplis ad  $\frac{1}{2}$  longitud. limbi peltatis ovatis apice acuminatis, acumine acuto, basi subcordato-rotundatis utrinque glabris membranaceis subopacis pellucido-punctulatis, nervis subtus subprominulis centrali ad  $\frac{1}{2}$  longitudinis a petiolo nervos utrinque 2 alternos adscendentes supremos ad apicem ductos inferos utrinque 5 e petiolo basi solutos divaricantes (inimum e basi solum adscendentem ad  $\frac{1}{2}$  longitud. a petiolo bifidum) mittente, petiolo glabro basi ima vaginante, pedunculo glabro quam petiolus multoties brevior, amento florente quam folium multum brevior densifloro, bractæ apice peltatæ pelta subrotundo-subquadrangulari coriacea glabra margine attenuata subflavicante pedicello clavato crasso, rhachi glabra, stigmatibus linearibus. § In Costa Rica et Veragua (Warszewicz! in h. Boiss.). Limbus (folium in spec. unicum) 0,28 longus, 0,14 latus. Petiolus fere 0,12 longus. Pedunculus 0,015 longus. Amentum florens 0,003 crassum (v. s.)



## 3° Folia multinervia.

225. *P. SORORIUM*, foliis longe petiolatis late ovatis apice abrupte acuminatis, acumine acuto, basi concavo-truncatis vel repando-cordatis utrinque præter margines dense adpresse hirtellos imamque basin juniorum glabris tenuiter membranaceis pellucido-punctatis 9-nerviis, nervis 3 centralibus ad apicem ductis subtus prominentibus, petiolo glabro ad  $\frac{3}{4}$  longitud. vaginante, alis angustis, amento breviter pedunculato folium subæquante, bractæ apice peltatæ pelta triangulari margine villosa. ♀ In Peruvia ad Chinchao (R. et Pav. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 443), America centrali (Barclay fide Miq. l. c.), sinum Fonseca (Benth. bot. Sulph. p. 167). *Artanthe sororia* Miq. l. c. Ramuli glabri. Limbi 0,14-0,17 longi, 0,11-0,13 lati. Petioli 0,03-0,05 longi. Pedunculi 0,005 longi. (fide Miq. l. c.)
226. *P. UNDENINERVIVM*, foliis longiuscule petiolatis e basi cordata vel repando-cordata ovato-longiuscule-acuminatis, acumine acuto, supra glabris utrinque basi ad nervos subtiliter puberulis margine ciliolatis membranaceis opacis pellucido-punctulatis 11-nerviis, nervis subtus prominulis 3 centralibus ad apicem ductis infimis brevioribus subtilioribus, petiolo glabro fere ad limbum vaginante, alis linearibus, pedunculo glabro quam petiolus multum brevior, amento florente folium æquante, bractæ apice truncato-peltatæ pelta subrotunda margine ciliata. ♀ In Maynas (Pöppig! in h. Petrop.). Ramuli glabri. Limbi 0,135 longi, 0,09 lati. Petioli 0,035 longi. Pedunculi 0,005 longi. (v. s.)
227. *P. RETICULATUM* (L. spec. p. 41, exclus. syn. Piso et Marcg.), foliis petiolatis ovatis apice acuminatis, acumine acuto, basi rotundatis cordatisve utrinque glabris firmis 7-9-nerviis, nervis subtus prominulis 3 centralibus ad apicem ductis infimis sæpe deficientibus nervulis primariis subtransversim parallelis, petiolo puberulo ad  $\frac{1}{2}$  longitud. vaginante, pedunculo glabro petiolum æquante, amento filiformi, bractæ apice peltatæ paullo infra florem insertæ pelta angusta margine anteriore hirtella pedicello brevi. ♀ In Mexico (Schiede, in h. Berol. ubi A. Schlechtendahl dicta), Jamaica (Swartz obs. p. 19). *Saururus Botrytis* major foliis plantagineis Plum. Am. ed. Par. p. 57. t. 75. *P. reticulatum* Hill dict. v. 26. p. 18. f. 18. f. 1, Swartz l. c., Lam. illustr. pl. 80. Willd. sp. v. 1. p. 162, Vahl enum. v. 1. p. 330. *Enckea reticulata* Miq. syst. p. 365. *Artanthe Schlechtendahlia* Miq. ! syst. p. 386. Ramuli glabri, læves. Limbi 0,19 longi, 0,12 lati. Petioli 0,015 longi. (v. s.)
228. *P. CINEREUM*, foliis longiuscule petiolatis e basi parum inæquali profunde cordata suboblique lanceolatis acutis supra pulverulento-puberulis subtus subtiliter præsertim ad nervos et nervulos hirtellis rigidulo-membranaceis epunctulatis 11-nerviis, nervis subtus prominulis, centrali ad apicem ducto laterali-bus latere minore 4 majore 5 mediis utrinque 2 fere ad apicem ductis sequentibus ad medium ductis infimis subtilioribus nervulis subtransversim parallelis, petiolo ad  $\frac{3}{4}$  longitudinis vaginante, alis linearibus persistentibus, pilis recurvulis dense pulverulento-hirtello, pedunculo quam petiolus subdimidio brevior pilis recurvulis dense pulverulento-hirtello, amento folium parum superante, bractæ apice truncato-peltatæ pelta semi-lunari subrotunda margine glanduloso-puberta pedicello longo, rhachi villosula ovario villosulo, bacca vertice flavide villosula. ♀ In Novæ Granatæ prov. Choco (Triana! in h. DC.). Ramuli pilis recurvulis dense pulverulento-hirtelli. Limbi 0,08 longi, 0,05 lati. Petioli 0,025 longi. Amenta matura 0,002 crassa. (v. s.)
229. *P. BERLANDIERI*, foliis petiolatis subobovato-oblongis apice breviter acuminatis, acumine acuto, basi leviter inæquali acutiusculis utrinque glabris membranaceis subopacis subtilissime pellucido-punctulatis 5-nerviis, nervis subtus prominulis, centrali ad apicem ducto sequentibus sub apicem evanidis infimis subtilioribus, petiolo basi ima vaginante glabro, amento maturo folium subæquante, bractæ apice truncato-peltatæ pelta triangulari-subrotunda margine subtiliter ciliolata, rhachi foveolata bacca ovato-subquadrangulati rhachi basi immersa, stigmatibus abbreviatis. ♀ Inter Tampico et Real del Monte (Berlandier n. 383! in h. DC. et Vindob.). Rami glabri. Ramuli glabri. Limbi 0,095 longi, 0,045 lati. Petioli 0,015 longi. (v. s.)

## b. Bractea cucullata apice inflexa.

## 1° Folia penninervia.

\* Nervo centrali tota vel fere tota longitudine nervos mittente.

230. *P. CONSANGUINEUM* (Kunth! olim), foliis brevissime petiolatis anguste oblongo-lanceolatis apice acuminatis, acumine acutiusculo, basi subæquali acutis supra glabris subtus ad nervos puberulis rigidulo-membranaceis, centrali nervo subtus prominulo utrinque tota longitudine nervos 12-15 alternos subpatulos margine anastomosantes mittente, petiolo puberulo, pedunculo petiolum paullo superante, amento quam folium multo breviori mucronato cylindrico, bractea rotundato-subcucullata glabra, stigmatibus 3-4 linearibus. ♀ In Guyana gallica (Lam.! in h. Willd. n. 632! Leprieur! in h. Deless. et DC., Poiteau! in h. Berol. ubi *P. consanguineum* Kunth. dictum), Maynas (Endl! n. 2289! in h. Vindob.). *P. angustifolium* Lam.! ill. p. 81, Vahl enum. v. 1. p. 312 (non Ruiz et Pav.). *Steffensia?* *angustifolia* Kunth! in Linn. v. 13. p. 622. *P. consanguineum* Kunth! l. c. et in Linn. l. c. *Artanthe angustifolia* Miq.! syst. p. 528. *Art. consanguinea* Miq.! syst. p. 529. Ramuli pilis recurvulis adpressis hirtelli. Limbi 0,07-0,11 longi, 0,012-0,023 lati. Petioli 0,002-0,003 longi. (v. s.)
231. *P. BRACHYSTACHYUM*, foliis subsessilibus lanceolatis apice acutis basi æquali acutis utrinque glabris rigidis opacis, centrali nervo utrinque tota longitudine nervos nervulosque fortiores patulos mittente, pedunculo glabro, amento recto cylindrico brevi mucronato, bractæ oblongo-cucullatæ apice inflexo triangulari angusto glabro, baccis glabris maturitate coalitis. ♀ In Guyana gallica prope Karouany (Sagot n. 542! in h. Kew.), Novæ Granatæ prov. Cumaral (Triana! n. 48 in h. suo). *Artanthe brachystachya* Miq.! mss. in h. Kew. Ramuli glabri. Limbi 0,06 longi, 0,022 lati. Petioli vix 0,002 longi. Pedunculi 0,003-0,004 longi. Amenta matura 0,01 longa, 0,005 crassa. (v. s.)
232. *P. ROHRII*, foliis petiolatis lato-ellipticis apice breviter acuminatis, acumine acuto, basi subæquali subattenuatis utrinque glabris opacis rigidulo-membranaceis, nervo centrali subtus prominente utrinque tota longitudine nervos circ. 15 suboppositos subtus prominulos subpatulos mittente, petiolo glabro paullo ultra limbi basin vaginante, alis linearibus, pedunculo glabro quam petiolus breviori, amento maturo quam folium breviori, bractæ cucullatæ basi ciliatæ vertice triangulari inflexo peltam ciliolatam simulante, bacca glabra. ♀ In Brasilia prope Rio Janeiro (Gaudich. n. 87! in h. DC., Sello! in h. Berol.), sylvis montosis Gabrielæ (Rich.! in h. Francovil.), sylvis Mandioca (Pohl n. 5036! in h. Vindob.), sylvis primævis prope Sebastianopolin (Mart. in h. monac.), Corcovado (Riedel et Langsdorff n. 86! et 91! in h. Petrop.), Guyana gallica (Poiteau! et Von Rohr! ex h. Schumacher in h. Petrop.). *P. sylvestre* Vell. flor. flum. t. 56. *P. macrophyllum* Loud. encycl. n. 503. p. 28. *Steffensia ampla* Kunth! in Linn. v. 13. p. 618. *Artanthe ampla* Miq.! syst. p. 501, in Linn. v. 20. p. 163 et in Mart. flor. Bras. fasc. 11. p. 55. t. 20. *Art. rugosa* Miq. syst. p. 489 (exclus. syn. Lam.). Ramuli glabri. Limbi 0,21-0,23 longi, 0,11 circ. lati. Petioli 0,025 longi. Pedunculi 0,012 longi. Amenta mat. 0,09 circ. longa, 0,0035 crassa. (v. s.)
- β, *crassum*, amentis crassioribus, maturis 0,007 crassis, 0,09 longis. — In Oaxaca (Buchinger! in h. Lenormand). (v. s.)
- γ, *grandifolium* (Miq.! Syst. p. 502). Limbi 0,25 longi, 0,12-0,125 lati. — In Brasilia prope Rio Janeiro (Gaudich. n. 87! et 1104! in h. Deless.) (v. s.)
233. *P. SURINAMENSE*, foliis breviter petiolatis oblongo-subelliptico-lanceolatis apice attenuatis acutiusculis utrinque glabris, basi subinæquali cordulatis membranaceis, nervo centrali subtus prominulo utrinque tota longitudine nervos alternos 7-12 nervulosque fortiores patulos mittente, petiolo puberulo, amento brevi, bractea rotundato-subcucullata glabra. ♀ In Caripi juxta Para (Spruce, n. 121! in h. Kew.), Barra (Spruce, 574! et 1346! in h. Francovil.),

Guyana gallica (Sagot, n. 537! in h. Kew.), Surinam (Hostmann, n. 1275! in h. Kew. et Boiss.). Ramuli pilis recurvulis adpresse puberuli. Limbi 0,055 longi, 0,012 lati. Petioli 0,003-0,004 longi. (v. s.)

234. *P. LINDBERGHII*, foliis breviter petiolatis elliptico-oblongis apicem versus attenuatis apice acutiusculis obtusiusculisve, basin versus attenuatis basi subæquali acutis utrinque glabris rigidis opacis pellucido-punctulatis, nervis subtus prominulis, centrali utrinque tota longitudine nervos circiter 12 alternos nervulosque fortiores subpatulos mittente, petiolo glabro, pedunculo glabro petiolum æquante, bractæ obovato-cucullatæ glabræ vertice inflexo semilunari. ♀ In Caldas prov. Minarum (S. A. Lindberg! in h. Mart.). Rami ramulique glabri. Limbi 0,17 longi, 0,065 lati. Petioli 0,006 longi. Amenta florent. 0,002 crassa. (v. s.)
235. *P. NEGROENSE*, foliis brevissime petiolatis oblongis apice acuminatis, acumine acuto, basi inæquali rotundatis, latere majore auriculæformi producto petiolum velante supra glabris subtus ad nervos nervulosque adpresse pubescentibus membranaceis subpellucidis creberrime pellucido-punctulatis, nervis subtus prominulis, centrali utrinque tota longitudine nervos circiter 12 alternos subalternosve subpatulo-subadscendentes mittente, petiolo utrinque hirsuto basi ima vaginante, pedunculo hirtello petiolum paulo superante, amento maturo quam folium multoties brevior, bractæ cucullatæ glabræ vertice inflexo glanduloso-punctato petalum triangularem simulante. ♀ In Brasiliæ prov. Rio Negro (Mart. n. 340! in h. Monac.). Ramuli dense adpresse pubescentes. Limbi 0,2 longi, 0,063 lati. Petioli vix 0,003 longi. Amenta florentia 0,004 crassa. (v. s.)
236. *P. PARAMARIBENSE*, foliis brevissime petiolatis subobovato-oblongo-ellipticis apice attenuato-acutatis, acumine obtusiusculo mucronulatoque, basin versus subattenuatis basi ima leviter inæquali obtusis supra glabris subtus ad nervos adpresse hirtellis rigidis opacis subtilissime pellucido-punctulatis, nervis subtus prominulis, centrali utrinque tota longitudine nervos circiter 12 patulo-subadscendentes mittente, petiolo dorso hirsuto, pedunculo glabro petiolum superante, amento florente quam folium multoties brevior mucronato, bractæ ovato-cucullatæ dorso hirtellæ vertice inflexo semilunari, rhachi hirtella, stigmatibus linearibus. ♀ In Paramaribo Surinam (Wulfschlägel, n. 1563! in h. Mart.). Rami ramulique sordide hirsuti. Limbi 0,13 longi, 0,057 lati. Petioli vix 0,004 longi. Amenta florentia 0,007 longa, vix 0,002 crassa. (v. s.)
237. *P. JULIFLORUM* (Nees et Mart. in nov. act. Ac. Leop. 11. p. 9), foliis brevissime petiolatis ellipticis vel suboblongo-ellipticis apice attenuato-obtusis basi parumper inæquali obtusis subexcisis utrinque glabris rigido-membranaceis subopacis pellucido-punctulatis, nervis subtus prominulis, centrali fere tota longitudine nervo utrinque circiter 10 alternos subalternosve subpatulo-subadscendentes mittente, petiolo ad limbum vaginante dorso hirtello, pedunculo glabro petiolum fere duplo superante, amento brevi quam folium multoties brevior, bractæ ovato-subcucullatæ glabræ vertice subinflexo nigro. ♀ In Brasiliæ sylvis (Mart. in Monac.). *P. juliflorum* Miq. Syst. p. 375. *Artanthe Maximiliani* Miq. in Linn. v. 20. p. 169, et in Mart. flor. Bras. fasc. 11. p. 59. Rami glabri. Ramuli una facie crispulo-hirtelli. Limbi 0,1 longi, 0,045 lati. Petioli vix 0,002 longi. Amenta florentia 0,003 crassa, vix 0,01 longa. (v. s.)
238. *P. LEIOPHYLLUM*, foliis petiolatis elliptico-oblongis apice breviter acuminatis, acumine acuto, basi inæquali cordulatis, lobo majore cum petiolo adnato, utrinque glabris subcoriaceis opacis pellucido-punctulatis, nervis subtus prominulis, centrali tota longitudine nervos utrinque 12-13 patulo-adscendentes mittente, petiolo ad limbi lobum minorem usque vaginante glabro, pedunculo glabro quam petiolum multum brevior, bractæ calceolatæ, basi hirtellæ vertice glabro triangulari. ♀ In Brasiliâ prope Bahia (Blanchet, n. 3151! in h. Vindob.). Ramuli glabri. Limbi 0,1, longi, 0,08 lati. Petioli 0,016 longi. Amenta florentia 0,003 crassa. (v. s.)
239. *P. RUMICIFOLIUM*, foliis petiolatis lanceolato-acuminatis, acumine acutiusculo, basi valde inæquali profunde cordatis, lobo majore petiolum velante, supra glabris subtus ad nervos hirtellis rigidulo-membranaceis opacis epunc-

- tatis, nervis subtus prominulis, centrali tota longitudine nervos circ. 12 patulo-subadscendentes supremos subpatulos nervulosque fortiores mittente, petiolo ad limbi basin usque vaginante dorso hirt, amento maturo quam folium triplo brevior, pedunculo hirtello petiolum æquante, bractæ subalceolatæ dorso et margine hirtellæ apice inflexo nudo carnosulo peltam triangularem simulante, bacca lateraliter compressiuscula fuligineo-tomentella. ♀ In Brasilia (Mollian! in h. Mart.). *Artanthe rumicifolia* Miq.! in Linn. v. 20. p. 141 et in Mart. flor. Bras. fasc. 11. p. 37. Rami vetustiores subglaberrati. Ramuli hirtelli. Limbi 0,17-0,18 longi, 0,045-0,055 lati. Petioli 0,03 longi. Amenta matura 0,003 crassa. (v. s.)
240. *P. GLANDULIGERUM* (Cas. DC. in Seem. *journal of bot.*, 1866. p. 215), foliis breviter petiolatis oblongo-ovato-ellipticis apice longe acuminatis, acumine acuto, basi æquali attenuatis acutiusculis utrinque glabris rigidis opacis, nervo centrali subtus prominulo tota longitudine nervos utrinque 12-14 alternos subpatulo-subadscendentes nervis fortioribus intermixtos mittente, petiolo glabro ad limbi basin usque vaginante, pedunculo glabro petiolum circ. æquante, amento cylindrico recto mucronulato vel nudo quam folium dimidio brevior, bractæ cucullatæ vertice inflexo peltam triangularem margine interiore ciliolatam angulo exteriori carnosulo-tuberculatam simulante. ♀ In Venezuela prope Tovar (Fendler, n. 1137! et 2400! in h. DC. Moritz n. 284! in h. Brit. Mus.), Caracas (Bushell! in h. Kew.), Colombia (Hartweg n. 1400! in h. Kew.). Ramuli glabri. Limbi 0,15 longi, 0,055-0,06 lati. Petioli 0,01 longi. Amenta matura 0,05-0,06 longa, 0,004 crassa. (v. s.)
- β, *subcoriaceum*, foliis subcoriaceis. — Hartweg. (n. 1400! in h. Kew.)
241. *P. CORDULATUM* (Cas. DC. in Seem. *journal of bot.*, 1866. p. 217), foliis petiolatis oblongo-lanceolatis apice acuminatis, acumine acuto, basi subæquali cordulatis utrinque glabris rigidis, nervo centrali subtus prominulo tota longitudine nervos utrinque 11-13 alternos margine anastomosantes mittente, petiolo ad limbi basin usque vaginante, pedunculo glabro quam petiolum brevior, bractæ cucullatæ vertice inflexo peltam triangularem fingente. ♀ In isthmo Panama prope Chagres (Fendler, n. 267! in h. Kew.). Ramuli glabri. Limbi 0,14 longi, 0,045 lati. Petioli 0,02 longi. (v. s.)
242. *P. PHTHINOTRICHON*, foliis breviter petiolatis elliptico-vel subovato-lanceolatis apice acuminatis, acumine acuto, basin versus cuneatis basi ima æquali acutis obtusive vel foliis ovato-lanceolatis basi tantum ima acutis utrinque glabris glanduloso-punctulatis membranaceis, nervo centrali subtus prominulo tota longitudine nervos utrinque circ. 10 versus marginem anastomosantes subalternos subpatulos nervulis fortioribus intermixtos mittente, petiolo foliorum inferiorum ad limbi basin usque vaginante, pedunculo glabro petiolum multum superante, amento cylindrico brevi quam folium multoties brevior, bractæ obovatæ glabræ carnosulæ basin versus pilosulæ vertice subinflexo, stigmatibus linearibus. ♀ In Peruvia ad flumen Amazonum (Poeppig n. 2283! in h. Berol. et 2289! in h. DC. et Vindob.), Maynas (Poeppig n. 2289! in h. Boiss.), in vicinibus Barra ad Rio Negro (Spruce, n. 10! in h. DC.). *Artanthe phtthinotricha* Miq.! *Syst.* p. 527, illustr. p. 80. t. 90. Rami glabri. Limbi 0,07 longi, 0,025 lati. Petioli 0,005 longi. (v. s.)
243. *P. DEMERARANUM*, foliis brevissime petiolatis vel subsessilibus oblongis vel oblongo-ovato-ellipticis apice acuminatis, acumine acuto mucronulatoque, basi parum inæquali cordulatis, lobo longiore auriculæformi petiolum velante, supra glabris subtus ad nervos et nervulos hirtellis, nervo centrali tota longitudine nervos alternos subadscendentes utrinque 10-12 nervulis fortioribus intermixtos mittente, petiolo glabro, pedunculo glabro petiolum superante, bractæ cucullatæ dorso aureo-puberulæ vertice inflexo glabro peltam simulante, haccis glabris lateraliter compressis ad aspectum subcoarctatis vertice subaureo-puberulis. ♀ In Surinam (Hostmann n. 312! in h. Kew.), regione Para (Wulfschlagel! in h. Mart.). *Artanthe Demerana* Miq.! in Hook. *lond. journal of bot.* v. 4. p. 464. Ramuli pilosuli. Limbi 0,2 longi, 0,08 lati. Petioli 0,003 longi. Pedunculi 0,005 longi. Amenta florentia 0,02 longa. (v. s.)
244. *P. ANONÆFOLIUM* (Kunth olim), foliis breviter petiolatis oblongo-ellipticis

apice breviter acuminatis, acumine acuto, basi æquali attenuatis acutis obtusiusculis supra nitidulis glabris subtus splendentibus glabris vel ad nervos subtilissime hirtellis rigidis vel subcoriaceis opacis, nervo centrali fere tota longitudine nervos patulo-subadscendentes utrinque 8-10 nervulis fortioribus intermixtos inferiores utrinque 2 subadscendentes mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum paulo superante, amento quam folium multoties brevius recto cylindrico mucronulato, bractearum obovato-cucullatæ extus glabræ vertice inflexo peltam triangularem fingente, bacca glabra. ♀ Prope Bahiam (Lhotzky apud Kunth sub nomine *P. rhododendrifolium*), in Guyana gallica prope Poura et Harouang (Sagot n. 844! in h. Kew. et 543! in h. Vindob.). Surinam (Hostmann n. 313! in h. Kew. et Boiss.), sylvis montosis ad Mariepatan (Kegel fide Miq. in Linn. v. 22. p. 77), Guyana anglica (Parker! in h. Kew.), Essequibo (Schomburgk n. 53! in h. Berol.), Demerara (Parker! in h. Kew. ubi *Artanthe rhododendrifolia* a Miq. ipso dictum), monte Cassinivord, Maranyana (Wulfschlaegel n. 1560! in h. Mart.). *Steffensia anonæfolia* Kunth in Linn. v. 13. p. 619. *P. rhododendrifolium* Kunth olim. *Steffensia rhododendrifolia* Kunth in Linn. l. c. p. 621. *Artanthe anonæfolia* Miq.! syst. p. 504, et in flor. Bras. fasc. 11. p. 56. Miq.! syst. p. 506 et in Hook. Lond. journ. bot. v. 4. p. 468. *Art. apiculata* Klotzsch! in Hook. Lond. journ. bot. v. 4. p. 321. Ramuli glabri vel uno latere hirtelli. Limbi 0.13 circ. longi, 0.055-0.06 lati. Petioli 0.003-0.01 longi. Pedunculi 0.005-0.015 longi. Amenta matura 0.035 longa, 0.004 crassa. (v. s.)

α, *rigidulum*, foliis rigidulo-membranaceis. — In Peruvia (Poeppig n. 3113! in h. DC. et Berol.). *Artanthe Durvilleana* Miq.! syst. p. 507. (v. s.)

β, *latifolium*, limbis 0.18-0.2 longis, 0.08-0.085 latis. — Prope Paramaribo (Miq. in Linn. v. 18. p. 227). Statura vegetior, fruter elatior (ex Miq. l. c.)

245. *P. SPRUCEANUM* (Cas. DC. in Seem. journ. of bot., 1866. p. 216), foliis brevissime petiolatis oblongo-lanceolatis apice acuminatis, acumine obtusiusculo, basi æquali cuneatis supra subtusque pilosis membranaceis, nervo centrali subtus prominulo nervos tota longitudine alternos subtus prominulos subadscendentes utrinque 8-10 mittente, petiolo villosopiloso, pedunculo piloso petiolum duplo superante, bractearum cucullatæ glabræ vertice inflexo peltam fingente, bacca glabra. ♀ In Peruvia orientali prope Tarapoto (Spruce n. 4072! in h. Kew.). Ramuli villosopilosi. Limbi 0.17 longi, 0.05 lati. Petioli 0.01 longi. Amenta matura 0.055 longa, 0.002-0.003 crassa. (v. s.)

246. *P. LENORMANDIANUM*, foliis brevissime petiolatis oblongo-ovato-ellipticis apice acuminatis, acumine acutiusculo, basi æquali cordulatis vel subattenuato-obtusis supra subtusque glabris vel subtus remote pilosis coriaceis opacis, nervo centrali subtus prominente nervos fere tota longitudine alternos subtus prominentes subadscendentes utrinque 5-9 nervulis fortioribus subtus prominulis intermixtos mittente, petiolo pilosulo alte vaginante, alis diutius persistentibus, pedunculo petiolum duplo superante, amento recto cylindrico non admodum densifloro quam folium multoties brevius, bractearum cucullatæ glabra, bacca subquadrangulo-obovata glabra. ♀ In Guyana gallica prope Akarouang (Sagot n. 540! in h. Kew. Lenormand et Berol.). Ramuli glabri. Limbi 0.16 longi, 0.06 lati. Petioli 0.005-0.006 longi. Pedunculi 0.015 longi. Amenta matura 0.025-0.03 longa, 0.004 crassa. (v. s.)

247. *P. PURDIEI*, foliis petiolatis e basi subcuneata oblongo-obovatis apice attenuatis obtusis basi acutis utrinque glabris coriaceis, nervo centrali subtus prominente fere tota longitudine nervos alternos patulo-adscendentes utrinque nervulis 5-6 fortioribus intermixtos subtus prominulos mittente, petiolo glabro plicato-rugoso, pedunculo glabro petiolum bis terve superante, amento recto cylindrico quam folium brevius, bractearum cucullatæ intus pilosæ vertice triangulari nudo peltam simulante, ovario puberulo. ♀ In Nova Granata prope S. Sebastian, S<sup>ta</sup> Martha (Purdie! in h. Kew.). Ramuli plicato-rugulosi. Limbi 0.05 longi, 0.02 lati. Petioli 0.01 longi. Pedunculi 0.02 longi. Amenta florentia 0.03 longa, 0.004 crassa. (v. s.)

\*\* Centrali nervo e parte tantum longitudinis nervos mittente.

248. *P. TEQUENDANENSE*, foliis petiolatis elliptico-lanceolatis apice acuminatis, acumine acuto, basi inæquali attenuatis acutiusculis obtusisve, latere longiore omnino cum petiolo adnato, supra glabris subtus ad nervos molliter pubescentibus membranaceis subopacis pellucido-punctulatis, nervo centrali subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes subtus prominulos utrinque 4-5 supremos ab aliis remotiores fere ad apicem ductos nervulosque fortiores mittente, petiolo juniore dorso molliter pubescente ad limbum vaginante, alis cito deciduis linearibus sursum liberis ultra limbi basin productis, pedunculo quam petiolus paullo brevior juniore pubescente, amento folium subæquante, bractæe obovato-cucullatæ junioris dorso villosæ postea pubescentis vertice inflexo, bacca secundum rhachin elongata glabra. ♀ In Nova Granatæ prov. Tequendana (Triana n. 351 et 391 in h. DC.). Ramuli glabri. Limbi 0,12 longi, 0,045 lati. Petioli fere 0,02 longi. Amenta matura 0,003 crassa. (v. s.)
249. *P. HOLTONII*, foliis petiolatis ovato-ellipticis apice acuminatis, acumine obtusiusculo mucronulato, basi inæquali rotundatis obtusisve utrinque glabris membranaceis subopacis subtilissime pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos subadscendentes utrinque circ. 6 nervulis fortioribus aliquando intermixtos supremos ante apicem per anastomoses evanidos mittente, petiolo glabro ad limbi latus minus et paullo ultra vaginante, pedunculo glabro per anthesin quam petiolus brevior, amento per anthesin folium dimidium vix æquante, bractæe suboblongo-obovato-cucullatæ dorso hirtellæ intus glabræ vertice semilunari inflexo margine inferiore ciliato. ♀ In Nova Granata prope La Paila (Holton n. 236! in h. DC.). Ramuli glabri. Limbi 0,12 longi, 0,06 lati. Petioli 0,02 longi. Amenta per anthesin 0,002 crassa. (v. s.)
250. *P. SECUNDUM* (Ruiz et Pav.! flor. Per. 1. p. 36. t. 62. f. b), foliis breviter petiolatis oblongo-ellipticis apice acuminatis, acumine acuto mucronulatoque, basi subæquali attenuatis acutis acutiusculisve utrinque glabris membranaceis subopacis pellucido-punctulatis, nervis subtus vix prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos subalternos subadscendentes utrinque 6 supremos fere ad apicem ductos mittente, petiolo glabro, pedunculo glabro petiolum superante, amento quam folium paullo brevior cylindrico, bractæe cucullatæ margine ciliatæ vertice inflexo peltam triangularem simulante, ovario glanduloso-punctulato glabro obovato. ♂ In nemoribus Andium peruvianorum (Ruiz! in h. Berol.), nemoribus Pillao, Muna, Cuchero, Chinchao (R. et P. l. c.). — Vahl enum. v. 1. p. 317. Steffensia secunda Kunth! in Linn. v. 13. p. 626. Artanthe secunda Miq.! syst. p. 516. Ramuli glabri, fusco-punctulati vel verruculoso-scabriusculi. Limbi 0,14-0,185 longi, 0,05-0,07 lati. Petioli 0,007 longi. Amenta florentia 0,1-0,12 longa, 0,003-0,004 crassa. (v. s.)
251. *P. RUBESCENS*, foliis brevissime petiolatis oblongo-ellipticis apice breviter acuminatis, acumine acuto mucronulatoque, basi subinæquali breviter cordatis supra glabris subtus ad nervos nervulosque pilis rubellis rubello-hirtellis rigidulis subopacis subtilissime pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos subadscendentes utrinque 6 supremos fere ad apicem ductos mittente, petiolo fere ad limbi basin usque vaginante dorso et apice utrinque rubello-hirto, amento quam folium fere dimidio brevior, pedunculo petiolum æquante rubello-hirtello, bractæe obovato-cucullatæ dorso hirtellæ vertice inflexo, rhachi inter baccas hirtella, bacca rhachi immersa subtiliter hirtella, stigmatibus linearibus. ♀ In Guyana gallica ad Karouany (Sagot n. 844! in h. Mart. et Berol.). Rami ramulique pilis rubellis hirtis. Limbi 0,095 longi, 0,04 lati. Petioli 0,003 longi. Amenta matura fere 0,004 crassa. (v. s.)
252. *P. DIOSPYRIFOLIUM* (Kunth! olim), foliis breviter petiolatis oblongo-ellipticis apice breviter attenuatis acutis mucronulatisque basi æquali attenuatis acutiusculis acutisve supra glabris subtus ad nervos sparsissime pilosulis rigi-

- dulo-membranaceis opacis, centrali nervo subtus prominulo ad  $\frac{3}{4}$  longitud. nervos alternos subadscendentes subtus prominulos utrinque 6-7 mittente, petiolo ad limbi basin usque vaginante, alis cito deciduis angustissime linearibus, pedunculo glabro petiolum paullo superante, bractæ cucullatæ extus glabræ intus pilosulæ vertice inflexo triangulari, bacca glabra. ♀ In Brasilia (Sello! in h. Berol. et Kew.), sylvis prov. Sebastianopolitanæ (Mart.! in h. monac.). *Steffensia diospyrifolia* Kunth! in Linn. v. 13. p. 677. *Artanthe diospyrifolia* Miq.! syst. p. 477, ill. p. 76. t. 82 et in Mart. flor. Bras. fasc. 11. p. 50. t. 14. Ramuli glabri. Limbi 0,13-0,18 longi, 0,06-0,08 lati. Petioli 0,01-0,015 longi. Amenta matura 0,08 longa, 0,004 crassa. (v. s.)
253. *P. GUIANENSE*, foliis breviter petiolatis elliptico-vel subovato-lanceolatis apice breviter acuminatis, acumine acuto, basi subinæquali obtusis supra glabris subtus margine et ad nervos ciliolatis membranaceis subopacis pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{3}{4}$  longitudinis nervos subalternos patulo-adscendentes utrinque 3-4 nervulosque fortiores interdum mittente, petiolo glabro, pedunculo glabro petiolum superante, amento cylindrico quam folium brevior, bractæ obovatæ subcucullatæ basi margine ciliolatæ vertice dein inflexo, stigmatibus linearibus recurvis. ♀ In crepidinibus ad ostia fluv. Atapuris in flum. Japura prov. Rio Negro (Spruce n. 1875! in h. DC. et 1194! aut 10 in h. Francovil.), Guyana anglica (Schomburgk n. 901! in h. Berol.). *Artanthe guianensis* Klotzsch! in Hook. Lond. journ. bot. v. 4. p. 322; Miq.! syst. p. 530 et in Mart. flor. Bras. fasc. 11. p. 62. Caulis glaber basi lignescens radicans. Limbi 0,06-0,09 longi, 0,02-0,035 lati. Petioli 0,01 longi. (v. s.)
254. *P. ACULEATUM* (Cas. DC. in Seem. journ. of bot., 1866, p. 216), foliis breviter petiolatis oblongo-subovato-lanceolatis apice acuminatis, acumine subulato, basi æquali acutiusculis obtusiusculisve supra glabris subtus ad nervos sparsissime hirtellis membranaceis subopacis creberrime subtiliter pellucido-punctulatis, centrali nervo subtus prominulo ad  $\frac{3}{4}$  longitud. nervos alternos subadscendentes utrinque 4 supremos ad apicem per anastomoses ductos nervulosque crassiores mittente, petiolo glabro, pedunculo glabro petiolum superante, amento pedunculum vix superante quam folium multoties brevior, bractæ obovatæ cucullatæ vertice inflexo margine ciliolato peltam triangularem fingente, stigmatibus linearibus recurvis. ♀ In regione Amazonica prope Panure ad Rio Uapes (Spruce n. 2887! in h. Francovil. et Mart.). Ramuli glabri. Limbi 0,085 longi, 0,025 lati. Petioli 0,003-0,005 longi. Pedunculi 0,01 longi. Amenta florentia 0,0025 crassa. (v. s.)
255. *P. CRASSIFOLIUM*, foliis brevissime petiolatis elliptico-oblongis apice attenuatis acutis basi subæquali cordatis utrinque glabris coriaceis opacis subtus nitidis, nervis subtus prominulis, centrali ad  $\frac{3}{4}$  longitudinis nervos alternos subadscendentes utrinque 7-8 sursumque nervulos crassiores mittente, petiolo glabro petiolum paullo superante, bractæ cucullatæ glabræ vertice inflexo carnosulo quadrangulari. ♀ In Brasilia (Pohl n. 53! in h. Vindob.). *Artanthe colubrina* var. *crassifolia* Miq.! in Linn. v. 20. p. 170. Rami ramulique glabri. Limbi 0,13 longi, 0,045 lati. Petioli 0,004 longi. (v. s.)
256. *P. INÆQUALE* (Cas. DC. in Seem. journ. of bot., 1866, p. 215), foliis petiolatis oblongo-lanceolatis apice acuminatis, acumine acuto basi inæquali rotundatis subcordatis utrinque glabris subcoriaceis opacis pellucido-punctatis, nervis subtus prominentibus, centrali nervo ad  $\frac{3}{4}$  longitud. nervos alternos subadscendentes latere majore 6 minore 5 supremos fere ad apicem ductos mittente, petiolo glabro, alis cito deciduis, pedunculo glabro quam petiolum brevior, amento cylindrico, bractæ cucullatæ vertice inflexo margine ciliolato peltam subsimulante. ♀ In Guyana (Rich.! in h. Francovil. ubi *P. recurvum* dictum). Ramuli glabri. Limbi 0,18 longi, 0,075 lati. Petioli 0,03 longi. (v. s.)
257. *P. NOVÆ-HISPANIÆ*, foliis petiolatis late suboblongo-ellipticis apice breviter acuminatis, acumine acuto, basi inæquali cordatis supra glabris basi ad nervum centralem adpresse hirtis subtus ad nervos nervulosque hirsutis membranaceis subopacis subtilissime pellucido-punctulatis, nervis subtus subprominulis, centrali ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitudinis nervos alternos subadscendentes utrinque 7 supremos ad apicem ductos mittente, petiolo dorso hirtello ad

limbi basin usque vaginante, pedunculo glabro quam petiolus triplo brevior, amento florente quam folium multoties brevior cylindrico mucronato, bractæ cucullatæ basi dorso hirsutæ vertice inflexo peltam triangularem simulante. ♀ In Nova Hispania (Pav. in h. Boiss.). Ramuli glabri. Limbi 0,16 longi, 0,09 lati. Petioli 0,025 longi. (v. s.)

258. *P. AVELLANUM*, foliis brevissime petiolatis ellipticis apice breviter et latiuscule acuminatis, acumine obtuso, basi inæquali gibbosis, latere majore rotundato minore acuto, supra glabris subtus ad nervos præsertim adpressè hirtellis rigidulo-membranaceis opacis pellucido-punctulatis, centrali nervo subtus prominulo ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. nervos subpatulo-adscendentes utrinque 5-6 sursumque nervulos aliquot mittente, petiolo dense adpressè hirtello ad  $\frac{1}{2}$  longitudinis vaginante, pedunculo dense hirtello puberulo petiolum paululum superante, amento quam folium duplo brevior mucronulato, bractæ cucullato-obovatæ vertice inflexo peltam triangularem margine ciliolatam puberulento-glandulosam simulante, stigmatibus truncato-abbræviatis. ♀ In Guyana gallica (Leprieur! in h. Deless.), prope Paramaribo (fide Miq. in Linn. v. 18. p. 227). *Artanthe avellana* Miq. syst. p. 478. Ramuli juniores dense retrorsum adpressè hirti. Limbi 0,1 longi, 0,055 lati. Petioli 0,003 longi. Amenta florentia 0,0015 crassa. (v. s.)

259. *P. WAGNERI*, foliis petiolatis ovato-elliptico-sublanceolatis apice acuminatis, acumine obtusiusculo, basi inæquali acutis utrinque glabris rigidulis opacis pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos subalternos subadscendentes utrinque 5 supremos ad apicem ductos infimos subtiles mittente, petiolo glabro fere ad limbi basin usque vaginante, alis cito deciduis, pedunculo glabro quam petiolus paullo brevior, amento brevissimo quam folium multoties brevior mucronato, bractæ cucullato-obovatæ glabræ vertice inflexo, filamentis exsertis. ♀ In Panama prov. Chiriqui (Wagner! in h. monac.). Rami ramulique glabri. Limbi 0,13 longi, 0,06 lati. Petioli 0,015 longi. Amenta florentia 0,005 longa, 0,003 crassa. (v. s.)

260. *P. TECTONIAEFOLIUM* (Kunth! olim), foliis petiolatis ovatis apice acuminatis, acumine acuto, basi inæquali rotundatis supra subtiliter hirtellis subtus mollior tomentoso-hirtis nascentibus utrinque sericeo-villosis, nervis subtus prominentibus, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos patulo-subadscendentes utrinque 5-6 mittente, petiolo villosio-hirto basi vaginante, pedunculo hirtello petiolum superante, amento cylindrico folium æquante, bractæ cucullatæ extus villosulæ vertice inflexo peltam margine hirtellam simulante, bacca vertice glanduloso puberula. ♀ In Brasilia meridionali ad Serra di Renaldo (Sello! in h. Berol.). *Steffensia tectoniaefolia* Kunth! in Linn. v. 13. p. 660. *Artanthe tectoniaefolia* Miq. syst. p. 396; illustr. p. 73. t. 73 et in Mart. flor. Bras. fasc. 11. p. 36. Ramuli fusce villosio-hirti. Limbi 0,25 longi, 0,15 lati. Petioli 0,02-0,03 longi. Amenta matura 0,004 crassa. (v. s.)

261. *P. DECURRENS* (Cas. DC. in Seem. journ. of bot. 1866. p. 2. p. 215), foliis breviter petiolatis lanceolatis apice acuminatis, acumine acuto basi æquali cuneato-acutis in petiolum subdecurrentibus utrinque glabris subcoriaceis opacis crebre subtiliter subpellucido-punctulatis, nervo centrali subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes subtus subprominulos utrinque 4 supremos ad apicem ductos infimos subtiliores mittente, petiolo glabro basi ima vaginante, amento cylindrico crasso quam folium duplo brevior, pedunculo glabro petiolum fere duplo superante, bractæ oblongo-subobovato-cucullatæ basi dorso subaureo-villosæ vertice carnosulo subinflexo margine ciliolato peltam simulante, bacca glabra. ♀ In Costæ-Ricæ monte Candalaria (Hoffmann n. 8531 in h. Berol.). Ramuli glabri. Limbi 0,12 longi, 0,045 lati. Petioli 0,008 longi. Amenta matura 0,006 crassa, ebullitione aromatica. (v. s.)

262. *P. VENEZUELENSE* (Cas. DC. in Seem. journ. 1866. p. 216), foliis breviter petiolatis amplis oblongo-ellipticis apice acuminatis, acumine acuto, basi subæquali obtusiusculis supra glabris subtus ad nervos hirtellis firmulis subopacis subtilissime pellucido-punctulatis, centrali nervo subtus prominente ad  $\frac{1}{2}$  longitud. nervos subtus prominentes subadscendentes oppositos utrinque 6 oppositos supremos ad apicem ductos mittente, petiolo glabro ad limbi



basin usque vaginante, alis cito deciduis, pedunculo glabro petiolum æquante, amento crasso quam folium duplo brevius, bracteis cucullatis glabris rhachis extenuatæ ope junctis inter flores productis, bracteæ vertice inflexo peltam simulante, baccae glabræ inter bracteas immersæ lateraliter compressæ vertice carnosulo margine extenuato. ♀ In Venezuela inter Agua Blanca et Cumbote alt. 5000 p. (Fendler n. 2572! in h. DC.). Ramuli glabri. Limbi 0,2 longi, 0,15 lati. Petioli 0,012 longi. Amenta matura 0,006 crassa. (v. s.)

263. *P. LACUNOSUM* (Kunth! in H. et B. nov. gen. 1. p. 51), foliis petiolatis oblique rotundato-ovatis apice acutis basi subinæquali rotundatis supra glabris bullatis subtus lacunosis ad nervos nervulosque adpresse hirtis, nervo centrali subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos subpatulo-adscedentes subtus prominulos mittente, petiolo basi vaginante cum pedunculo pilis recurvulis adpresse hirtello, amento crasso cylindrico folium subæquante, bracteæ ovato-lanceolatæ utrinque dense hirtæ vertice nudo recurvo obtusiusculo, ovario oblongo apice subattenuato, stigmatibus linearibus recurvis. ♂ In montanis temperatis prov. Jaen de Bracamoros inter Passo de Matara et Cavico alt. 400 hexap. (Humb. in h. Berol.).—Kunth! synops. v. 1. p. 107. *Steffensia lacunosa* Kunth! in Linn. v. 13. p. 651. *Artanthe lacunosa* Miq.! syst. p. 427. Ramuli pilis recurvulis adpresse hirtelli. Limbi 0,16 longi, 0,12 lati. Petioli 0,025 longi. Amenta post anthesin 0,01 crassa. (v. s.)

264. *P. VERRUCULOSUM*, foliis brevissime petiolatis oblongo-subovato-lanceolatis apice longiuscule acuminatis, acumine obtusiusculo mucronulato, basi inæquali subrotundatis subsemicordatis latere minore basi acutiusculo utrinque glabris rigidulo-membranaceis supra albo-punctulatis glaucescentibus, centrali nervo subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos subadscedentes subtus prominulos utrinque 4-5 supremos ad apicem ductos mittente, petiolo glabro basi ima vaginante subtus verruculis albidis punctulato, pedunculo petiolum quater superante, amento cylindrico folium subæquante, bracteæ oblongo-cucullatæ vertice inflexo triangulari peltam margine ciliolatam simulante, pedicello glabro, bacca lateraliter compressa vertice subtiliter puberula. ♀ In Costæ-Ricæ monte Candelaria (Hoffmann n. 8! in h. Berol.). Ramuli glabri, verruculis albidis scabridi. Limbi 0,1 longi, 0,045 lati. Petioli 0,003 longi. Amenta matura 0,002 crassa. (v. s.)

265. *P. CHURUMAYU* (Ruiz et Pav. flor. Per. 1. p. 35. t. 58. f. b), foliis petiolatis ovatis acuminatis, acumine acuto, basi subæquali obtusis utrinque glabris, centrali nervo ad  $\frac{1}{2}$  longitud. nervos alternos subadscedentes utrinque 5 supremos ad apicem ductos mittente, pedunculo petiolum æquante, amento folium æquante obtuso, bracteæ obovato-cucullatæ intus et extus sub pubescentibus vertice triangulari inflexo nudo. ♀ In nemoribus Huacahuassi ad fluvii margines (Dombey! in h. Berol.), in Peruvia (R. et P. l. c.). — Vahl enum. v. 1. p. 325. *Artanthe Churumayu* Miq.! syst. p. 481. Limbi 0,135 longi, 0,055 lati. Petioli 0,005 longi. (v. s.)

266. *P. GRANATENSE*, foliis breviter petiolatis lanceolatis apice acuminatis, acumine acuto, basi æquali cordatis supra glabris subtus ad nervos adpresse puberulis membranaceis, nervo centrali subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos subadscedentes subtus prominulos utrinque 3-4 supremos ad apicem ductos mittente, petiolo adpresse puberulo, pedunculo petiolum multum superante, amento brevi gracili, bracteæ obovato-cucullatæ vertice inflexo triangulari margine hirtello peltam simulante. ♀ In Nova Granata (Purdie! in h. Kew.). Ramuli pilis retrorsis adpresse puberuli. Limbi 0,07-0,08 longi, 0,025-0,035 lati. Petioli 0,005 longi. Amenta florentia 0,02 longa, 0,0015 crassa. Pedunculi 0,03 longi. (v. s.)

267. *P. AREOLATUM*, foliis longiuscule petiolatis ovato-acuminatis vel ovato-lanceolatis apice acutis basi æquali subattenuatis supra glabris areolatis subtus molliter pilosis margine ciliolatis subcoriaceis opacis, nervo centrali subtus prominulo nervos alternos subadscedentes utrinque 3-4 subtus prominulos mittente, petiolo ad limbi basin usque vaginante dorso ciliato, alia angustis, pedunculo petiolum superante piloso, bracteæ conchæformis intus pubescentis dorso extus pubescentis vertice nudo peltam subfingente, stig-

- matibus lanceolatis, ovario glabro. ♀ In Peruviae Serra Chachapoyas (Mathews n. 3227! in h. Kew.). *Artanthe areolata* Miq.! in Hook. Lond. journ. bot. v. 4. p. 454. Ramuli glabri. Limbi 0,13 longi, 0,08 lati. Petioli 0,03 longi. Pedunculi 0,055 longi. Amenta florentia 0,1-0,12 longa, 0,003 crassa. (v. s.)
268. *P. PACHYSTACHYON* (Cas. DC in Seem. journ. 1866, p. 116), foliis longiuscule petiolatis ovatis vel elliptico-ovatis supra bullatis lævigatis (an ætate?) subtus villosis lacunosulis coriaceis opacis, nervo centrali subtus prominulo utrinque nervos suboppositos 5-6 subadscendentes subtus subprominulos mittente, petiolo utrinque dense fusco-pubescente basi ima vaginante pedunculo pubescente quam petiolus brevior, amento cylindrico crasso, bractæe oblongo-subcucullatæ utrinque glanduloso-puberulæ vertice nudo obtuso, ovario glabro, stigmatibus linearibus. ♀ In Colombia (Linden n. 162! in h. Kew.), prov. Bogota (Triana n. 52! in h. suo), Costa-Rica (Hoffmann n. 856! in h. Berol.). Ramuli adpresse puberuli. Limbi 0,1-0,12 longi, 0,065-0,08 lati. Petioli 0,02 longi. Pedunculi 0,012 longi. Amenta florentia 0,12 longa, 0,006 crassa. (v. s.)
269. *P. MINARUM*, foliis petiolatis lanceolato-ovatis apice acuminatis, acumine acuto, basin versus paullo attenuatis basi ima acutiusculis supra glabris subtus ad nervos parvis pilis brevissimis hirtellis firmo-membranaceis subopacis pellucido-punctulatis lævibus, nervis subtus prominulis ad  $\frac{1}{2}$  longitudinis nervos alternos adscendentes utrinque 4 (et aliquando infimum utrinque quintum) supremos ad apicem ductos mittente, petiolo glabro ad  $\frac{1}{2}$  longitud. vaginante, amento quam folium duplo brevior, bractæe subcalceatæ peltatæ vertice carnosulo semilunari margine aureo-hirtello pedicello superne concavo dorso aureo-hirto, pedunculo glabro petiolum subæquante, bacca obovata vel obovato-tetragona glabra, stigmatibus linearibus. ♀ In prov. Minas-Geraes (Lindberg n. 640! in h. Mart.). Rami ramulique glabri. Limbi 0,125 longi, 0,045 lati. Petioli 0,01 longi. Amenta matura 0,005 crassa. (v. s.)
270. *P. DASYPODUM*, foliis breviter petiolatis oblongo-ovatis apice oblique acuminatis, acumine acuto, basi æquali læviter cordatis utrinque (præsertim ad nervos longius) molliter pubescentibus margine ciliatis membranaceis opacis, nervo centrali vix ad  $\frac{1}{2}$  longitud. nervos alternos adscendentes utrinque 5-6 supremos ad apicem ductos mittente, petiolo valde hirsuto, pedunculo valde hirsuto petiolum æquante, bractea lanceolata carnosula margine longe hirsuta. ♀ In Peruvia (Mathews n. 1702! in h. Kew.). *Artanthe dasypoda* Miq.! in Hook. Lond. journ. bot. v. 4. p. 458. Ramuli dense molliter fusce hirsuti. Limbi 0,17 longi, 0,07 lati. Petioli 0,01 longi. Amenta ante evolutionem 0,01 longa, 0,006 crassa. (v. s.)
271. *P. TENUICUSPE*, foliis breviter petiolatis late oblongo-ellipticis apice acuminatis, acumine acutissimo, basi inæquali rotundatis, latere longiore basi ima a petiolo libero auriculatim petiolum velante, utrinque glabris membranaceis subpellucidis, nervo centrali ad  $\frac{1}{2}$  longitud. nervos alternos adscendentes utrinque 5-6 mittente, petiolo hirsuto, pedunculo glabro petiolum æquante, amento mucronulato, bractea gibboso-calceolata glabra intus pilosula. ♀ In Peruvia (Mathews n. 1704! in h. Kew.). *Artanthe tenuicuspis* Miq.! in Hook. Lond. journ. bot. v. 4. p. 457. Ramuli hirti. Limbi 0,23 longi, 0,09 lati. Petioli 0,01 longi. Amenta ante evolutionem 0,1 longa, 0,002 crassa. (v. s.)
- 2° Folia multiplinervia.
272. *P. RICHARDIÆFOLIUM* (Kunth! olim), foliis longe petiolatis oblongo-ellipticis apice acuminatis basi valde inæquali profunde cordatis supra glabris subtus ad nervos subtilissime puberulis subcoriaceo-membranaceis, nervis subtus prominentibus, centrali ad  $\frac{2}{3}$  longitudinis nervos subadscendentes latere majore 8 minore 6 inferos latere minore 3 majore 5 fere e basi solutos mittente, petiolo ad limbi basin usque vaginante, alis linearibus, pedunculo quam petiolum multum brevior, amento folium superante cylindrico, bractæe cucullatæ extus margine et intus hirtellæ vertice inflexo peltam simulante. ♀ In Brasilia (Sello! in h. Berol.), in sylvis umbrosis mont. Organensium (Miq.! in Hook. Lond. journ. bot. v. 4. p. 445). *Steffensia richardiæfolia* Kunth! in Linn.

- v. 13. p. 668. *Artanthe richardiæfolia* Miq. l. syst. p. 395, illustr. p. 72. t. 72, et in Mart. flor. Bras. fasc. 11. p. 36. t. 9. f. 2. Ramuli glabri. Limbi 0,19-0,27 longi, 0,1-0,13 lati. Petioli 0,06 longi. Amenta florentia 0,25 longa, 0,003 crassa. (v. s.)
273. *P. subfuscum* (Cas. DC. in Seem. Journ. of bot. 1866, p. 217), foliis longiuscule petiolatis rotundato-ovatis apice subattenuatis acutis basi subæquali profunde cordatis supra parce pilosulis demum glabris subtus dense fusce pubescentibus rigidulis opacis epunctulatis, nervis subtus prominentibus, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos patulo-adscendentes utrinque 4 supremos ad apicem ductos mittente, lateralibus utrinque 4 e basi divaricantibus, petiolo ad limbi basin usque vaginante dorso fusce hirtello demum glabrato, alis linearibus glabris cum petiolo omnino adnatis, pedunculo glabro petiolum æquante, amento cylindrico crasso folium superante, bractæ calceolatæ vertice carnosulo margine ciliolato, bacca glabra. ♀ In Costæ-Ricæ regione sylvatica ad Alto de la Cruz (Hoffmann n. 544! in h. Berol.). Ramuli juniores dense fusce pubescentes. Limbi 0,32 longi, 0,24 lati. Petioli 0,04 longi. Amenta matura 0,01 crassa. (v. s.)
274. *P. ovalifolium*, foliis longe petiolatis ample ovatis apice acuminatis inferioribus basi æquali profunde cordatis superioribus basi valde inæquali profunde cordatis lobis discretis supra glabris subtus fusce pubescentibus, centrali nervo ad  $\frac{1}{2}$  longitudinis nervos alternos latere majore 8 minore 7 superiores utrinque subadscendentes inferiores magis patulos supremos ad apicem ductos inferos latere majore 3 minore 2 e basi divaricantes mittente, petiolo ad  $\frac{1}{2}$  longitudinis vaginante basi dorso et apice utrinque fusce veluti-pubescente, alis linearibus glabris cum petiolo omnino adnatis, pedunculo quam petiolus brevior post anthesin glabrato, amento post anthesin folium æquante, bractæ cucullatæ dorso et intus fusce hirsutæ vertice inflexo triangulari margine fusce hirsuto, ovario apice villosulo, stigmatibus linearibus. ♀ In Brasilia (Riedel n. 285! in h. Petrop.). Rami et ramuli densius fusce velutino-puberuli. Limbi 0,21 longi, 0,16 lati. Petioli 0,06 longi. Amenta post anthesin 0,004 crassa. (v. s.)
275. *P. pulchrum*, foliis longiuscule petiolatis ample ovato-oblongis apice acuminatis basi inæquali profunde cordatis supra glabris obsolete bullatis subtus ad nervos et nervulos dense fusce-hirtis rigidis opacis, centrali nervo subtus prominente ad  $\frac{1}{2}$  longitudinis usque nervos supra basin utrinque 4 subadscendentes subtus prominulos inferos magis divaricantes mittente, lateralibus basi utrinque 4 divaricantibus, petiolo ad limbi basin usque vaginante dorso dense fusce hirtello, alis linearibus glabris cum petiolo omnino adnatis, pedunculo dense fusce hirtello quam petiolus paullo brevior, amento cylindrico folium æquante, bractæ spatulatæ utrinque puberulæ vertice inflexo semilunari peltam simulante. ♀ In Novæ Granatæ prov. Pasto (Triana n. 12! in h. DC.) et prov. Marequita (Triana n. 13! in h. suo). *Artanthe Moritziana* Miq. in Linn. v. 18. p. 710? Ramuli dense fusce hirtelli. Limbi 0,3 longi, 0,16 lati. Petioli 0,04 longi. Amenta matura 0,005 crassa. (v. s.)
276. *P. Boissierianum*, foliis longiuscule petiolatis oblongo-ovatis apice attenuatis acutis? basi valde inæquali cordatis, lobis discretis a petiolo liberis, supra glabris subtus dense fusce pubescentibus coriaceis opacis subtiliter pelliculo-punctulatis, nervis subtus prominentibus, centrali utrinque supra basin nervos 4 alternos subadscendentes supremos ad apicem ductos mittente, lateralibus e basi utrinque 3-4 magis divaricantibus brevioribus, petiolo ad limbi lobum majorem vaginante dorso dense fusco-hirsuto, alis linearibus glabris apice acutis breviter liberis, pedunculo dense pubescente quam petiolus circ.  $\frac{1}{2}$  brevior, amento maturo folium fere duplo superante, bractæ oblongo-cucullatæ dorso et intus fusce villosulæ vertice inflexo semilunari nudo, bacca glabra. ♀ In Nova Granata (Linden n. 1100! in h. Boiss.). Ramuli dense fusce pubescentes. Limbi circ. 0,205 longi, 0,185 lati. Petioli 0,025 longi. Amenta fere mat. 0,004 crassa. (v. s.)
277. *P. Cabellense*, foliis longe petiolatis ample elliptico-ovatis apice attenuato-acutatis basi valde inæquali profunde cordatis, lobis conniventibus, supra glabris subtus subfusce molliter pubescentibus membranaceo-rigidis

opacis subtiliter pellucido-punctulatis, nervis subtus prominentibus, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos utrinque 8 superiores utrinque 4 subadscendentes inferiores utrinque 4 e basi divaricantes mittente, petiolo ad limbi lobum majorem usque vaginante dorso fusce hirsuto, alis linearibus glabris cum petiolo omnino adnatis, bractæ oblongo-cucullatæ dorso extus fusce villosæ vertice truncato semilunari margine fusce villosa, ovario sursum fusce villosulo, stigmatibus linearibus. ♀ In Venezuela prov. Carabobo ad P<sup>er</sup> Cabello (Funck n. 784) in h. Petrop.). Ramuli fusce hirsuti. Limbi 0,31 longi, 0,24 lati. Petioli 0,06 longi. Amenta florentia 0,006 crassa. (v. s.)

278. *P. GIGANTIFOLIUM*, (Bredem.!) foliis longe petiolatis amplissime oblongo-ovatis apice subito breviuscule acuminatis, acumine acuto mucronulato, basi valde inæquali cordatis, lobis conniventibus liberis lobo majore auriculæformi, supra glabris subtus dense fusce pubescentibus rigidis opacis parce pellucido-punctulatis, nervis subtus prominentibus, centrali ad  $\frac{1}{2}$  longitud. nervos latere majore 9 minore 7 superiores utrinque subadscendentes supremos ad apicem ductos inferiores latere majore 4 minore 2 e basi divaricantes mittente, petiolo fere ad limbi basin vaginante dorso dense fusce hirsuto, alis linearibus cum petiolo omnino adnatis glabris, pedunculo fusce pubescente petiolum æquante, amento folium æquante, bractæ cucullatæ dorso hirsutæ margine ciliolatæ vertice inflexo triangulari nudo margine subtiliter ciliolato, bacca glabra. ♀ In Brasilia (Riedel n. 77! in h. Petrop.). Limbi a petiolo 0,35 longi, 0,026 lati, lobus major a petiolo 0,12 longus. Petioli fere 0,07 longi. Amenta matura 0,005 crassa. (v. s.)

279. *P. CARACASANUM* (Bredem.!) in h. Willd. n. 679!), foliis longe petiolatis oblongo-ovatis apice obtusis? basi inæquali profunde cordatis supra glabris subtus fusce tomentosis firmo-membranaceis, nervis subtus prominentibus, centrali nervos alternos adscendentes latere majore 8 minore 7 inferiores magis approximatos infimos e basi solutos magis patulos mittente, petiolo ad limbi basin vaginante dorso fusce tomentoso. ♂ In Caracas (Bredem. l. c.). — Link Jahrb. 1. 3. p. 61 (fide R. et Schult. mant.), Opiz in Reliq. Haenk. fasc. 3. p. 135. Steffensia? Caracasana Kunth! in Linn. v. 13. p. 664. Artanthe Caracasana Miq. l. syst. p. 398. Limbi 0,3 longi, 0,2 lati. (v. s.)

β, *firmius*, foliis apice acute acuminatis firmioribus subtus magis tomentosis, bractæ calceolatæ dorso villosulæ vertice semilunari carnosio hirsuto, pilis dorsi longioribus, stigmatibus linearibus; bacca glabra lateraliter compressa. — In Caracas (Bredem. in h. Willd. n. 679!), Venezuela prope coloniam Tovar (Fendler n. 1143! in h. DC.). Ramuli tomentosi. Limbi 0,36 longi, 0,25 lati. Petioli 0,08 longi. Amenta 0,006 crassa. (v. s.)

280. *P. CERNUUM* (Vell. flor. Num. t. 58), foliis longe petiolatis oblongo-ellipticis apice attenuatis breviter acuminatis, acumine acuto, basi valde inæquali profunde cordatis, lobo majore auriculæformi petiolum velante, supra glabris subtus ad nervos tomentellis rigido-membranaceis epunctatis, nervis subtus prominentibus, centrali ad  $\frac{1}{2}$  longitudinis usque nervos latere majore 8 minore 7 subadscendentes alternos inferiores latere majore 4 minore 3 e basi solutos divaricantes mittente, petiolo fere ad limbi basin usque vaginante dorso fuligineo-tomentello, amento maturo cylindrico folium æquante, bractæ subcalceolatæ dorso pubescentis vertice nudo triangulari margine ciliolato petalum simulante. ♂ In Serra d'Estrella in via a Praja Alta ad Margaritiva (Pohl! n. 281, 50231; 4771! in h. Vindob.), in udis sylvis primævis ad Sebastianopolin prov. Rio Negro (Mart. l. in h. monac.). Artanthe spectabilis Miq. l. in Lina. v. 20. p. 138 et forma glabrior l. c. p. 139, in Mart. flor. Bras. fasc. 11. p. 35. t. 8. f. 1 et 2. Art. cernua Presl! in Epim. bot. p. 225; Miq. in Mart. flor. Bras. fasc. 11. p. 72. Ramuli fuligineo-tomentelli. Limbi 0,32-0,34 longi, 0,23-0,24 lati. Petioli 0,06-0,08 longi. (v. s.)

281. *P. OBLIQUUM* (Ruiz et Pav. flor. peruv. 1. p. 37. t. 63. f. a), foliis longe petiolatis oblongo-ovatis apice attenuatis obtusiusculis basi valde inæquali profunde cordatis supra glabris subtus ad nervos puberulis subcoriaceis, nervis subtus prominentibus, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos latere majore 9 minore 7 superiores utrinque subadscendentes inferiores latere majore 5-6 minore 3 tenuiores magis approximatos e basi solutos divaricantes mittente,

petiolo ad limbi basin usque vaginante puberulo pedunculum superante, amento pendulo cylindraceo folium superante, bractæ calceolatæ seu cucullato-obovatæ basi extus et apice utrinque villosulæ vertice inflexo peltam subsimulante, bacca glabra lateraliter compressa. § In Cuchero sylvis ad Cayumba terminum (R. et P. l. c.), Peruvia subandina (Poeppig! in h. Berol.), Casapi (Mathews n. 1699! in h. Kew.), Brasiliâ (Riedel n. 285! in h. Petrop. et Blanchet n. 286! in h. DC.), Mandiocca (Langsdorff n. 92! in h. Petrop.), Ilheos sylvis (Wawra et Maly n. 380! fide Wawra bot. Ergeb. p. 128), sylvis terræ firmæ ad Japura prope Maribi prov. Rio Negro (Mart. in h. monac.) et Canta de Gallo prov. Rio Negro (Peckott! in h. Mart.), Venezuela (Fendler n. 1145! in h. DC. et Moritz n. 286! in h. Brit. Mus.), locis umbrosis juxta Buenavista inter Valle del Tuy et Caracas alt. 835 hexap. (Humb. et Bonpl. in nov. gen.). P. obliquum Vahl enum. v. 1. p. 324. P. flagellare Kunth in Nov. gen. v. 1. p. 54, synopsis v. 1. p. 109. Steffensia obliqua Kunth! in Linn. v. 13. p. 663. Artanthe magnifica Miq. syst. p. 391, illustr. p. 72. t. 71, in Hook. Lond. journ. bot. v. 4. p. 445. Art. flagellaris Miq. syst. p. 393. Ari. superba Miq. in Linn. v. 20. p. 137 et in Mart. flor. Bras. fasc. 11. p. 34. t. 7. Ramuli glabriusculi vel pubes tenuissima pulveraceo-adpressi. Limbi usque ad 0,39 longi, 0,26 lati. Petioli 0,07 longi. Amenta matura usque ad 0,6 longa, 0,006 crassa. Bacca ferè 0,002 longa. (v. s.)

β, *eximium*, foliis subtus ad nervos magis puberulis apice breviter acuminatis, lobis magis convergentibus, ramulis hirtellis. — In Brasilia meridionali (Sello! in h. Berol.), prov. Minarum prope Xipoto (Miq. in Mart. flor. Bras.), prope San Gabriel da Cocheira (Spruce, n. 2001! in h. Kew. et DC.), ins. Cattivo (Friedrichsthal, n. 608! in h. Vindob.). P. eximium Kunth! olim. Steffensia eximia Kunth! in Linn. v. 13. p. 665. Artanthe eximia Miq. Syst. 393 et in Mart. flor. Bras. fasc. 11. p. 35. t. 9. v. 1.

282. P. CHAVICOIDE, foliis longiuscule petiolatis inferioribus ovatis superioribus oblongo-ovatis omnibus apice acuminatis, acumine acuto, basi profunde gordatis utrinque glabris atque crebre nigro-punctulatis membranaceis subpellucidis, centrali nervo utrinque ad  $\frac{1}{2}$  longitudinis nervos supra basin 2 oppositos patulo-subadscendentes et e basi utrinque 3 subpatulos infimos subtiliores mittente, petiolo glabro crebre nigro-punctulato ad limbi basin vaginante, bractæ cucullato-spathulatæ vertice inflexo triangulari margine ciliolato peltam simulante. § In sylvis ad flum. Solimoes prope Ega, prov. Rio Negro (Mart. in h. Monac.). Artanthe chavicoide Miq. in Linn. v. 20. p. 135 et in Mart. flor. Bras. fasc. 11. p. 33. Ramuli glabri, crebre nigro-punctulati. Limbi 0,065–0,09 longi, 0,03–0,055 lati. Petioli 0,025 longi. (v. s.)

283. P. REGNELLI, foliis petiolatis ovato-subrotundis apice obtusiuscule apiculatis basi breviter cordatis vel truncatis coriaceis pellucido-punctulatis supra ad nervos puberulis, nervo centrali ad  $\frac{1}{2}$  longitud. trifido et e basi nervos utrinque 4–5 mittente, petiolo ad  $\frac{1}{2}$  longitud. vaginante juniore puberulo, pedunculo juniore puberulo quam petiolus multum brevior, amento florente recto cylindrico quam folium triplo brevior, bractæ spathulato-calceolatæ pedicellatæ sub vertice extus villosulæ vertice planiusculo. § In prov. Minarum prope Caldas (Regnell fide Miq. in Mart. flor. Bras. fasc. 11. p. 34). Artanthe Regnellii Miq. l. c. t. 6. f. 2. Limbi circ. 0,14 longi latique. Petioli 0,03 longi. Amenta florentia 0,005 crassa (fid. Miq. l. c.).

284. P. CALCEOLATUM (Cas. DC. in Seem. journ. of bot. 1866. p. 217), foliis petiolatis e basi cordata ovato-lanceolatis supra crebre bullatis pilosisque subtus molliter pubescentibus lacunosis membranaceo-coriaceis, opacis nervis subius prominulis, centrali supra basin nervos utrinque 2 subalternos adscendentes et e basi utrinque 1–2 subadscendentes divaricantes mittente, petiolo utrinque villosulo ad  $\frac{1}{2}$  longitud. vaginante, pedunculo villosulo quam petiolus brevior, amento cylindrico mucronato quam folium brevior, bractæ calceiformi subtus et margine subvillosa, ovario oblongo apice attenuato glabro stigmatibus linearibus. § In Colombia (Linde n. 339! in h. Kew.). Ram. villosi. Limbi 0,07 longi, 0,05 lati. Petioli 0,015 longi. Amenta florentia 0,05 longa, 0,004 crassa. (v. s.)

β, *magnifolium*, foliis inferioribus 0,13 longis, 0,06 latis, petiolis 0,025

longis, ramis retrorsum villosis. — In Novæ Granatæ prov. Antioquia (Triana, n. 211 in h. DC.)

3° Folia multinervia.

285. *P. FLUMINENSE*, foliis longiuscule petiolatis subrotundato-ovatis apice subattenuatis obtusis basi rotundatis cordatisve utrinque glabris nigro-punctulatis membranaceis novemnerviis, nervis omnibus fere e basi solutis subtus prominulis, centrali ad apicem ducto aliis fere ad apicem ductis, petiolo glabro fere ad limbi basin usque vaginante, pedunculo glabro quam petiolus brevior, amento cylindrico folium superante, bractea obovata subcucullata basi dorso et margine pubescente, stigmatibus 3-4. ♂ In Brasilia (Sello in h. Berol.), Tejuco prope Rio Janeiro (Pohl n. 4794 in h. Vindob.). Artanthe obtusa Miq. Syst. p. 416, in Linn. v. 20. p. 149 et in Mart. flor. Bras. fasc. 11. p. 39. t. 12. Ramuli glabri. Limbi 0,08 longi, 0,06 lati. Petioli 0,03 longi. Amenta matura 0,005 crassa. (v. s.)

286. *P. POTOMORPHE*, foliis longe petiolatis ovato-rotundatis brevissime acuminatis, acumine obtusiusculo, basi subæquali profunde cordatis, lobis rotundatis parum divergentibus, subcoriaceis vel potius membranaceis obsolete pellucido-punctulatis supra atro-viridibus lævibus glabris vix nitidulis sub lente atomulis albidis conspersis subtus pallidis ad nervulos puberulis cæterum utrinque glabris 13-nerviis, nervis 3 mediis paulo supra basin solutis, centrali patulo-costato, amentis breviter pedunculatis apice obtusiusculis breviter mucronulatis, bractæ inflexo-subconchæformis vertice semilunari margine ciliato, stam., bacca glabra vertice glandulosa. ♂ In sylvis humidis montium Mexici alt. 3500 p. ad Vera Cruz (Galeotti, n. 6117 in h. Deless. fide Miq. Syst. p. 403). Artanthe potomorphe Miq. l. c. Limbi 0,2-0,24 longi, 0,145-0,19 lati. Petioli 0,07-0,08 longi. (ex Miq. l. c.)

287. *P. RUIZIANUM*, foliis petiolatis inferioribus cordiformibus superioribus basi truncato-rotundatis omnibus apice subprotracto-acuminatis, acumine acutiusculo, utrinque glabris tenuiter membranaceis subopacis parca pellucido-punctulatis 7-nerviis, nervis omnibus haud prominulis, centrali ad apicem ducto sequentibus sub apice evanidis inferioribus brevioribus infimis subtilioribus, petiolo glabro basi ima vaginante, pedunculo glabro petiolum æquante, amento sublaxifloro folium superante gracili, rhachi albo-villosula, bractea ovata apice acutata subtus et supra basi ima margineque villosula subpeltata subsessili, ovario oblongo discreto, stigmatibus abbreviatis. ♂ In Nova Hispania (Pav. in h. Boiss.). Ramuli graciles, glabri. Limbi inferiores 0,09 longi, 0,07 lati, super. 0,08 longi, 0,043 lati. Petioli fere 0,01 longi. (v. s.)

288. *P. VARIIFOLIUM*, foliis breviter petiolatis lanceolatis vel ovato-ellipticovel lanceolatis vel ellipticis vel subovatis apice acuminatis vel acutis, acumine acute obtusove vel rotundato, basi subæquali rotundatis siccis membranaceis subtilissime pellucido-punctulatis supra saturatius subtus pallidius glaucescentibus opacis utrinque glabris præter imam basin paginæ superioris puberulam, 5-vel sub-7-nerviis, petiolo subciliato-puberulo, amento densifloro, bractæ pedicellatæ apice subspathulato-dilatata concavo-conchæformi vertice glabriusculo dorso pedicelloque floccoso-hirtello, staminibus 4 (an plura?), ovario.... ♂ In Mexico (Andrieux, n. 133 in h. Deless. fide Miq. Syst. p. 355). Enckea variifolia Miq. l. c. Rami ramulique glabri, petioli 0,006-0,008 longi, limbi 0,075-0,055, longi 0,025 lati.

++ Bacca trigona vel obovata.

a. *Bractea apice peltata vel truncato-peltata.*

1° Folia penninervia.

\* Nervus centralis nervos tota vel fere tota longitudine mittens.

289. *P. FLEXICAULE*, foliis brevissime petiolatis oblongo-ellipticis apice acuminatis, acumine acutiusculo, basi subinæquali obtusis, latere minore basi ima acutis utrinque dense pubescentibus, centrali nervo nervos fere tota longitudine utrinque 6 subadscendentes mittente, pedunculo petiolum duplo

superante, amento maturo folium dimidium vix æquante, bractæ apice peltatæ pelta triangulari (in icone subrotunda) margine ciliata, pedicello ciliato, bacca glabra obpyramidato-trigona. ♂ In Guyana (Rudge). *P. flexuosum* Rudge, pl. Guy. p. 13. t. 13 (non Jacq.). *Artanthe flexicaulis* Miq. Syst. p. 533. Ramuli dense pubescentes. Limbi 0,145 longi, 0,043 lati. Petioli vix 0,001 longi (ex Rudge, l. c.)

290. *P. HALESIAEFOLIUM* (Kunth! olim), foliis petiolatis subovato-ellipticis apice acuminatis, acumine acuto mucronulatoque, basi subinæquali vel inæquali acutiusculis vel subrotundatis supra glabris subtus ad nervos nervulosque molliter hirtellis supra bullulis albidis notatis, centrali nervo subtus prominulo tota longitudine nervos alternos utrinque 12 nervulosque mittente, petiolo subtiliter puberulo, amento quam folium  $\frac{1}{2}$  brevior, bractæ apice truncato-peltatæ dorso medio basi hirtellæ et acute productæ pelta triangulari lateraliter anguste producta pedicello lato, bacca obovato-trigona vertice glanduloso-hirtella. ♀ In Rio Janeiro (Gaudich. n. 1114! in h. Berol.). *Artanthe halesiaefolia* Miq. in Mart. flor. Bras. fasc. 11. p. 40. Ramuli glabri, juniores hirtelli. Limbi 0,115 longi, 0,045 lati. Petioli 0,01 longi. (v. s.)
291. *P. REGELII*, foliis inferioribus longiuscule superioribus breviter petiolatis ovato-oblongis apice acuminatis, acumine acuto, basi leviter inæquali rotundatis utrinque glabris supra subbullatis subtus creberrime fusce-punctulatis, nervis subtus subprominulis, centrali nervo patulo-subadscendentes-utrinque 8-9 nervulis validioribus sæpe intermixtos supremos ante apicem per anastomoses evanidos mittente, petiolo glabro ad limbi latum minus usque vaginante, alis cito deciduis, pedunculo glabro petiolum superiorem æquante amento maturo quam folium multoties brevior, bractæ truncato-peltatæ vertice triangulari glabro pedicello cucullato lato dorso carinato carina hirsuta, stigmatibus linearibus, bacca obovata glabra. ♀ In Brasilia (Riedel n. 791 in h. Petrop.). Rami ramulique glabri. Limbi 0,155 longi, 0,06 lati. Petioli inferiores 0,03, superiores 0,007 longi. Amenta matura 0,025 longa, 0,046 crassa. (v. s.)
292. *P. JUSSIÆANUM* (Kunth olim), foliis petiolatis oblique ovato-lanceolatis rarius ovato-oblongis angustato-acuminatis, acumine acutiusculo, basi subæquali rotundatis membranaceis subtilissime pellucido-punctulatis supra glabris junioribus pilis raris conspersis nitidis subtus pallidioribus et præsertim ad nervos pilosis, centrali nervo nervos utrinque plures subtus prominentes mittente, petiolo hirsuto, amento obtuso leviter curvato pedunculato, bractea cuneata apice truncato-peltata glabra, ovario glabro, stigmatibus filiformibus subulatis, baccis densis obovato-turbinatis trigonis. ♂ In Serra d'Estrella Brasiliæ meridionalis (Luschnath apud Kunth.). *Schilleria Jussæana* Kunth, in Linn. v. 13. p. 680. *Artanthe Jussæana* Miq. Syst. p. 420 et in Mart. flor. Bras. fasc. 11. p. 41. Ramuli hirsuti. Limbi 0,14-0,16 longi, 0,055-0,06, rarius 0,08 lati. Petioli 0,015-0,025 longi. (v. s.)
293. (1) *P. AUGUSTUM* (Rudge, ic. p. 10. t. 7), foliis brevissime petiolatis grandibus ovato-oblongis apice leviter acutis basi leviter inequali acutis utrinque præsertim subtus punctulatis, centrali nervo tota longitudine nervos subadscendentim-patulos utrinque circ. 13 nervulis validioribus intermixtos mittente, pedunculo scabriusculo petiolum paulo superante, amento quam folium triplo brevior, bractæ apice peltatæ pelta triangulari margine hirsuta, stam. 4, bacca obovato-tetragona. In Guyana anglica (Rudge! l. c. et in h. Rudge et DC.). *Artanthe augusta* Miq. Syst. p. 398. Ramuli scabriusculi. Limbi 0,2 longi, 0,09 lati. Petioli 0,004 longi. (v. s.)
294. *P. PEDICELLARE* (Cas. DC. in Seem. journ. 1866. p. 211), foliis brevissime petiolatis elliptico-oblongis apice attenuato-acuminatis, acumine acuto, basi attenuatis acutis utrinque glabris membranaceis subopacis subtiliter pellucido-punctulatis, nervis subtus prominulis, centrali tota longitudine nervos patulo-subadscendentes suboppositos utrinque circ. 13 nervulis validioribus intermixtos mittente, petiolo glabro pedunculo petiolum æquante glabro, amento florente quam folium multoties brevior bractæ apice peltatæ pelta

- triangulari margine dense aureo-pubescente, bacca obpyramidato-trigona glabra. ♀ In Novæ Granatæ prov. Barbacoas (Triana, n. 161 in h. suo). Ramuli glabri. Limbi 0,2 longi, 0,09 lati. Petioli 0,008 longi. (v. s.)
295. *P. PUNCTATUM* (Ruiz et Pav. flor. Per. 1. p. 36. t. 62. f. a), foliis breviter petiolatis elliptico-oblongis apice breviter acuminatis, acumine acuto, basi leviter inæquali acutis utrinque glabris punctatisque, centrali nervo fere tota longitudine nervos patulo-subadscendentes utrinque circiter 10 nervulis validioribus intermixtos mittente, pedunculo petiolum æquante, amento cylindrico maturo folium subæquante. ♂ In nemoribus Pillao ad Iscutuniam (R. et P. l. c.) — Vahl, enum. v. 1. p. 317, Opiz reliq. Haenk. fasc. 3. p. 150. *Artanthe punctata* Miq. Syst. p. 522. Frutex glaber. Limbi 0,14 longi, 0,06 lati. Petioli 0,01 longi. (ex R. et P. l. c.)
296. *P. GRANDIFOLIUM* (Kunth! in H. et B. nov. gen. v. 1. p. 46), foliis petiolatis amplis ovatis apice acuminatis, acumine acuto, basi truncato-rotundatis utrinque glabris firmule membranaceis, nervo centrali subius prominulo nervos alternos subius prominulos subadscendentes utrinque 10-12 mittente nervulis subtransversim parallelis, petiolo glabro, bractæ apice truncato-peltatæ pelta triangulari antice callosa nuda marginis villosula pedicello longiusculo villosa, bacca obovato-trigona vertice glandulosa puberula. ♂ In monte prope villam Arranjon (Seemann, n. 417! in h. Kew.), temperatis montis Ouladiu juxta Quebrada de Bogueia (Humb. l. c. et n. 639! et 656! in h. Willd.), Oaxaca (h. Vindob.). — Kunth synops. v. 1. p. 103. *Schilleria grandifolia* Kunth! in Linn. v. 13. p. 709. Schill. riparia Kunth! l. c. p. 707. *P. riparium* Kunth! in H. et B. nov. gen. v. 1. p. 48, Kunth synops. v. 1. p. 105. *Artanthe grandifolia* Miq. Syst. p. 408. Art. riparia Miq. l. c. p. 409. Ramuli glabri. Limbi 0,25-0,28 longi, 0,16 lati. Petioli 0,035 longi. Amenta florentia 0,002 crassa. (v. s.)
- \*\* Nervus centralis e parte longitudinis nervos mittens.
297. *P. FUNCKII*, foliis breviter petiolatis subelliptico-oblongis apice acuminatis, acumine obtusiusculo, basin versus attenuatis basi ima æquali acutis utrinque glabris firmulo-membranaceis subpellucidis epunctulatis, nervis subius subprominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos subalternosve subpatulo-subadscendentes utrinque 6 nervulis validioribus intermixtos supremos ad apicem ductos mittente, petiolo glabro ad limbi basin usque vaginante, pedunculo glabro petiolum æquante, amento maturo folium dimidium æquante cylindrico recto, bractæ apice peltatæ pelta triangulari glabra glandulosa, bacca subobovato-trigona glabra vertice flavicante. ♀ In Venezuelæ prov. Caracas alt. 4000 ped. (Funck et Schlim in Linden exsicc. n. 134! in h. Petrop. et DC.). Rami ramulique glabri nitiduli. Limbi 0,16 longi, 0,066 lati. Petioli 0,007 fere longi. Amenta matura 0,004 crassa. (v. s.)
298. *P. TRIGONUM* (Cas. DC. in Seem. journ. 1866. p. 212), foliis breviter petiolatis oblongis apice acuminatis, acumine acuto, basi æquali attenuatis acutis supra glabris subius præsertim ad nervos adpresse puberulis margine tenerrime ciliolatis membranaceis opacis pellucido-punctulatis, centrali nervo subius prominulo ad  $\frac{1}{2}$  longitud. nervos subalternos subadscendentes utrinque 5 nervulis validioribus intermixtos supremos ad apicem fere ductos mittente, petiolo utrinque pubescente basi ima vaginante, pedunculo pubescente petiolum fere duplo superante, amento folium dimidium æquante, bractæ apice peltatæ pelta triangulari margine ciliata pedicello angusto subcucullato, bacca obovato-trigona vertice glanduloso-puberula. ♀ In Novæ Granatæ prov. Barbacoas (Triana n. 44! in h. DC.). Ramuli pilis subrecurvulis adpresse pubescentes. Limbi 0,13 longi, 0,04 lati. Petioli 0,006 longi. Amenta matura 0,002 crassa. (v. s.)
299. *P. ÆQUALE* (Vahl! eclog. p. 4. t. 3), foliis breviter petiolatis oblongo-ellipticis apice acuminatis, acumine obtuso, basi subæquali acutis acutiusculatis utrinque glabris rigidulis subpellucidis subtilissime pellucido-punctulatis, nervis subius vix prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos subalternos patulo-subadscendentes utrinque 6-7 nervulis validioribus intermixtos mit-



tenis, petiolo glabro, pedunculo glabro petiolum paullo superante, amento cylindrico folium dimidium æquante, bractæ apice peltatæ pelta triangulari glabra vel minute ciliolata, pedicello glabro clavato, bacca subobovato-trigona glabra. ♀ In Brasilia (Riedel n. 287! in h. Petrop.), umbrosis ad Bihai prope Sapativa (Pohl n. 5035! in h. Vindob.), ins. Trinitatis (Sieber n. 192! in h. Vindob.), Montserrat (Vahl! in h. Willd. n. 649!), S<sup>a</sup> Luzia (Crudy! in h. monac.), S<sup>o</sup> Domingo (Griseb. flor. w. Ind. p. 172), Guadalupe (Funcke et Schlim n. 29! in h. Petrop., Murray n. 329! in h. Hook. fide Miq. in Hook. Lond. journ. bot. y. 4. p. 468). P. æquale Vahl enum. v. 1. p. 315. P. punctatum R. et P. ♀ flor. peruv. v. 1. p. 36. t. 62. f. a; Opiz? in Reliq. Haenk. fasc. 3. 151. Rami ramulique glabri, Limbi 0,145 longi, 0,065 lati. Petioli 0,008-0,01 longi. Amenta matura 0,06 longa, 0,003 crassa. (v. s.)

β, *ovalifolium*, foliis ovato-ellipticis apice acuminatis, acuminis acutis, basi æquali rotundatis, bractæ vertice magis carnosulo. — In ins. Trinitatis (Sieber n. 192! in h. DC.). (v. s.)

300. P. MORISONIANUM, foliis breviter petiolatis lanceolatis apice acuminatis, acuminis acuto mucronulatoque, basi æquali acutis utrinque glabris rigidulo-membranaceis subopacis subtiliter pellucido-punctulatis, centrali nervo subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos patulo-subadscendentes utrinque 7 nervulis validioribus sæpe intermixtos supremos ante apicem evanidos infimos subadscendentes mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum æquante, amento folium dimidium superante, rhachi hirtella, bractæ dorso hirtella apice truncato-subinflexa, bacca obovato-trigona glabra. ♀ In Brasilia (Riedel n. 191! in h. Petrop.). Rami ramulique glabri. Limbi 0,095 longi, 0,025 lati. Petioli 0,006 longi. Amenta matura 0,003 crassa. (v. s.)

301. P. LINEATUM (R. et Pav.! flor. peruv. v. 1. p. 35. t. 60. f. a), foliis breviter petiolatis elliptico-lanceolatis apice longe acuminatis, acuminis acuto mucronulatoque, basi subacutis supra nitidis glaberrimis verruculoso-scarbis subtus ad nervos pilosulis coriaceis opacis obsolete pellucido-punctulatis, nervo centrali subtus prominente ad  $\frac{1}{2}$  longitud. nervos alternos adscendentes subtus prominentes utrinque 5-7 supremos ad apicem ductos mittente, petiolo glabro, pedunculo glabro petiolum circ. æquante, amento folium subsuperante, bractæ apice peltatæ pelta triangulari-semilunari margine hirtella, bacca obovata apice truncato-rotundata glabra. ♀ Ad margines fluv. Huanuci et Panao et in Chincho (R. et P. l. c. et in h. Willd. v. 662!), Guatemala (Friedrichshall! in h. Lenormand). — Vahl v. 1. p. 315. Schilleria lineata Kunth! in Linn. v. 13. p. 704. Artanthe lineata Miq.! syst. p. 459. Ramuli hirtopubescentes. Limbi 0,09-0,1 longi, 0,03-0,04 lati. Petioli 0,01 longi. (v. s.)

β, *magnifolium*, foliis rigido-membranaceis, limbis 0,15 longis, 0,055 latis, ramulis glabris. — In sylvis umbrosis et temperatis alt. 6000 ped. reipublicæ æquatorialis (Jameson n. 362! in h. DC. et Kew.). (v. s.)

γ, *cordulatum*, foliis basi rotundato-cordulatis. — In Peruvia (Mathews n. 1691! in h. Kew.). (v. s.)

302. P. KAPPLERI (Cas. DC. in Seem. journ. 1866. p. 213), foliis brevissime petiolatis oblongo-elliptico-lanceolatis apice acuminatis, acuminis acuto, basi æquali rotundatis supra glabris subtus præsertim ad nervos pilis recurvulis sparse hirtellis membranaceis opacis pellucido-punctulatis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitudinis nervos. alternos utrinque 6 inferiores subadscendentes supremos patulo-adscendentes nervulosos validiores tota longitudine mittente, petiolo pilis recurvulis hirtello, pedunculo hirtello petiolum duplo superante, amento cylindrico quam folium multoties brevius, bractæ apice peltatæ pelta angusta triangulari margine villosa, pedicello lato dorso villosa, stigmatibus brevibus, baccis discretis obovatis nigris vix angulosis. ♀ In Surinam (Kappler n. 1885! in h. Francov.). Ramuli nigri, pilis recurvulis hirtelli. Limbi 0,13 longi, 0,035 lati. Petioli 0,005 longi. Amenta 0,025 longa, 0,005 crassa. (v. s.)

303. P. ACAROUANYANUM, foliis brevissime petiolatis oblongo-ellipticis apice

attenuatis acutiusculis basi inæquali attenuatis acutis supra glabris subtus præsertim ad nervos subtiliter adpresse puberulis membranaceis, nervo centrali subtus prominulo nervos ad  $\frac{1}{2}$  longitud. alternos subpatulo-subadscendentes utrinque 5-6 nervulis validioribus intermixtos subtus prominulos supremos per anastomoses fere ad apicem ductos mittente, petiolo puberulo-hirsuto, pedunculo puberulo petiolum duplo superante, amento brevi quam folia quadruplo brevioræ sæpe androgyno et tum floribus masc. basi amenti vel inter feminas mixtis, florum inferiorum masc. bractea rotundato-cucullata dorso ciliata, stam. ...., flor. femin. bractea conformi, bacca plerumque trigono-obpyramidata rhachi immersa glabra. ♀ In Guyana gallica prope Acaouany (Sagot n. 5411 in h. Kew. et Berol.). Ramuli puberuli. Limbi 0,115 longi, 0,035 lati. Petioli 0,002-0,004 longi. Amenta matura 0,02 longa, 0,003 crassa. (v. s.)

304. *P. OBLONGUM* (Kunth! in H. et B. nov. gen. v. 1. p. 52), foliis brevissime petiolatis oblongo-ellipticis apice acuminatis, acumine acuto mucronulatoque, basi subæquali acutiusculis utrinque glabris subtus ad nervos obsolete puberulis membranaceis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos patulo-adscendentes utrinque 4-5 mittente, petiolo glabro, pedunculo glabro petiolum æquante, amento cylindrico quam folium brevioræ, bractæ apice truncato-peltatæ vertice triangulari margine hirsuto, bacca obovato-trigona flavo-glandulosa. ♀ In convalle umbrosa juxta Cumanacoa prov. Cumanæ alt. 140 hexap. (Humb. l. c. et in h. Willd. n. 6711 et pro parte 647 vel 644!), ad rivulos Xalapa et Miradores alt. 3000-4000 ped. (Martens et Gal. in bull. ac. Brux. v. 10. p. 20). — Kunth synops. v. 1. p. 108. Schilleria oblonga Kunth! in Linn. v. 1, 3. p. 696. Artanthe oblonga Miq.! syst. p. 480. Ramuli glabri hic illic verruculosi. Limbi 0,11-0,12 longi, 0,055 circ. lati. Petioli 0,005-0,008 longi. Amenta matura 0,08 circ. longa, 0,003 crassa. (v. s.)

305. *P. GLABELLUM*, foliis breviter petiolatis ellipticis apice breviter acuminatis, acumine acutiusculo, basi subæquali acutis vel attenuato-acutis basi ima in petiolo decurrentibus, utrinque glabris glandulis nigris atomulisque albis conspersis, centrali nervo ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. nervos adscendentes utrinque 5 supremos ad apicem ductos infimos et basi solutos subtilissimos mittente, bractæ apice peltatæ pelta triangulari margine flavo-ciliata, stam. ...., bacca obovato-triangulari, stigmatibus abbreviatis. ♀ In Guyana gallica (Poi-teau in h. Deless.). Artanthe glabella Miq. syst. p. 518. (non *P. glabellum* Sw.) Ramulus fistulosus, glaber, sparsissime et minute verruculosus. Limbi 0,13-0,17 longi, 0,09-0,075 lati. Amenta matura 0,09-0,1 longa, 0,003 crassa. (v. s.)

306. *P. RIVINOIDES* (Kunth! olim), foliis breviter petiolatis oblique ellipticis vel subovato-ellipticis apice subito suboblique acuminatis, acumine acuto vel acutiusculo, basi subæquali acutis utrinque glabris membranaceis subpellucidis subtilissime glanduloso-pellucidis, nervo centrali subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos adscendentes utrinque 4-5 subtus vix prominulos nervulis validioribus intermixtos supremos ad apicem ductos mittente, petiolo glabro ad limbi basin usque vaginante, pedunculo petiolum subæquante glabro, rhachi foveolata, foveolis parce ciliolatis, bractæ apice peltatæ pelta triangulari glabra glandulosa, pedicello villosulo, bacca obovato-trigona subquadrangulave glabra. ♀ In Brasilia prope Rio de Janeiro (Gaudich. n. 1109! in h. DC.; Riedel n. 526! in h. Petrop.), monte Corcovado (Luschnath! in h. Mart.), inter Vittoria et Bahia (Sello! in h. Berol.). *P. lentaginoides* Kunth! olim. Schilleria rivinoides Kunth! in Linn. v. 13. p. 682. Schill. lentaginoides Kunth! in Linn. l. c. p. 683. Artanthe rivinoides Miq.! syst. p. 519, in Linn. v. 20. p. 171 et in Mart. flor. Bras. fasc. 11. p. 57. Art. lentaginoides Miq.! syst. p. 520, in Mart. flor. Bras. l. c. Ramuli glabri. Limbi 0,11-0,135 longi, 0,055-0,06 lati. Petioli 0,012 longi. Pedunculi 0,01 longi. Amenta matura 0,07-0,08 longa, 0,002-0,004 crassa. (v. s.)

307. *P. HETEROPHYLLUM* (R. et Pav.! flor. Peruv. v. 1. p. 34. t. 56. f. a), foliis breviter petiolatis ellipticis vel subovato-ellipticis apice breviter subacuminato-acutiusculis basi subæquali acutis utrinque glabris glandulis fuscis

obsitis membranaceo-subcoriaceis subpellucidis pellucido-punctulatis, centrali nervo subtus subprominulo ad  $\frac{3}{4}$  longitud. nervos alternos patulo-adscentes subtus vix prominulos utrinque 7-8 nervulis validioribus intermixtos supremos fere ad apicem ductos mittente, petiolo glabro, pedunculo glabro petiolum æquante, bractæ apice peltatæ pelta triangulari margine hirsuta, bacca glabra obovato-trigona.  $\frac{3}{4}$  In Peruvia (Dombey n. 930! in h. DC. ex h. Mus. par.; Pav. in h. Boiss. et Berol.; Ruiz! in h. Willd. n. 642! et h. Berol.), nemoribus Pillao, Chacahuassi, Chinchao et Muna (R. et P. l. c.). P. heterophyllum Vahl enum. v. 1. p. 318. Schilleria heterophylla Kunth! in Linn. v. 13. p. 688. Artanthe heterophylla Miq.! syst. p. 502. Ramuli glabri fusco-punctulati. Limbi 0,12-0,16 longi, 0,07-0,08 lati. Petioli circ. 0,01 longi. (v. s.)

$\beta$ ?, *angustifolia* (Miq. in Hook. Lond. journ. bot. v. 4. p. 467). In Peruvia (Mathews n. 1711 fide Miq. l. c.)

308. P. SEPTUPLINEAVIUM, foliis brevissime petiolatis ovato-lanceolatis apice acuminatis, acumine acuto mucronulato, utrinque glabris subcoriaceis opacis, centrali nervo subtus prominulo ad  $\frac{3}{4}$  longitud. nervos alternos subadscentes subtus prominulos utrinque 4-3 supremos fere ad apicem ductos mittente, nervulis subtransversalibus, petiolo glabro, pedunculo glabro petiolum 2-3° superante, amento mucronulato maturo quam folium brevior, bractæ apice peltatæ pelta semilunari angusta ciliata, pedicello cucullato ciliato, bacca obpyramidato-trigona vertice flavo-glandulosa.  $\frac{3}{4}$  In isthmo Panama (Seemann, n. 1066! in h. Kew.). Artanthe septuplinervia Miq.! in Bot. Herald. p. 199. Ramuli glabri. Limbi 0,1 longi, 0,04 lati. Petioli 0,003-0,004 longi. Amenta matura 0,03-0,04 longa, 0,003-0,004 crassa. (v. s.)

309. P. TENUINERVE, foliis brevissime petiolatis lanceolato-oblongis apice acuminatis, acumine acuto mucronulatoque, basi inæquali cordatis supra glabris subtus villosis pubescentibus pallidioribus membranaceis subopacis pellucido-punctulatis, nervis subtus vix prominulis, centrali ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. nervos alternos subadscentes utrinque 4-5 supremos ad apicem ductos infimos subtiles interdumque nervulos fortiores mittente, nervulo obscuro juxta marginem percurrente, petiolo basi vaginante basi dorso et sursum utrinque pubescenti-villosa, pedunculo villosulo petiolum æquante, amento post anthesim quam folium multoties brevior, bractæ apice truncato-peltatæ pelta triangulari ciliolata pedicello concavo, bacca obpyramidato-trigona glabra.  $\frac{3}{4}$  In Brasilia (Blanchet! in h. Boiss.). Ramuli fuscescenti-villosi. Limbi 0,13 longi, 0,046 lati. Petioli 0,005 longi. Amenta post anthesim 0,003 crassa. (v. s.)

310. P. NOVO-GRANATENSE, foliis brevissime petiolatis subelliptico-oblongis apice longe protracto-cuspidatis cuspidate mucronulata, basi inæquali rotundatis utrinque glabris rigidulis pellucido-punctulatis, centrali nervo subtus prominulo ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. nervos alternos subadscentes subtus prominulos utrinque 3-4 supremos ad apicem ductos mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum superante, amento maturo quam folium multoties brevior, bractæ pelta semilunari angusta margine dense subaureo-pubescente, bacca glabra obpyramidato-trigona.  $\frac{3}{4}$  In Novæ Granatæ prov. Barbacoas (Triana, n. 30! in h. DC.). Ramuli glabri. Limbi 0,12 longi, 0,055 lati. Petioli 0,003 longi. Amenta matura 0,025 longa, 0,004 crassa. (v. s.)

311. P. PERSICARIÆFOLIUM (Kunth! olim), foliis breviter petiolatis oblongo-lanceolatis lanceolatisque apice angustato-acuminatis, acumine acuto mucronulatoque, basi inæquali obtusis supra subnitidulis scabriusculis glabris subtus ad nervos adpresse pilosulis subcoriaceis opacis demum bullatis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos circ. 12 alternos subadscentes supremos fere ad apicem ductos mittente, petiolo utrinque hirtello basi ima vaginante, pedunculo puberulo petiolum paulo superante, amento cylindrico quam folium multum brevior, bractæ apice peltatæ pelta carnosula triangulari margine hirsuta, bacca glabra obovato-trigona.  $\frac{3}{4}$  In Brasilia meridionali (Sello! in h. Berol.), in hort. Berol. cult. — Kunth! in Linn. v. 13. p. 684. Artanthe persicariæfolia Miq. Syst. p. 499, in Mart. flor. Bras. fasc. 11. p. 55. t. 19 et in Hook.

æquante, amento folium æquante, bractæ apice truncato-peltatæ vertice triangulari nudo, pedicello dorso villosulo, ovario parce villosulo, bacca obovato-trigona. ♀ In Brasilia (Riedell n. 1, et 105 in h. Petrop.). Rami ramulique glabri. Limbi 0,1 longi, 0,036-0,055 lati. Petioli 0,005-0,01 longi. Amenta matura fere 0,004 crassa. (v. s.)

318. *P. VARIEGATUM* (Kunth! in H. et B. nov. gen. v. 1. p. 47), foliis brevissime petiolatis oblongo-ellipticis apice longe acuminatis, acumine acuto basi parum inæquali rotundatis obsolete cordulatis utrinque glabris membranaceis obsolete pellucido-punctulatis subopacis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitudinis nervos alternos subadscendentes utrinque 5-6 supremos fere ad apicem ductos infimos subtiliores mittente, petiolo glabro, pedunculo glabro petiolum paulo superante, amento cylindrico quam folium multum brevius, bractæ apice peltatæ pelta triangulari margine aureo-hirsuta, bacca glabra obovato-trigona subtetragonave. ♀ In sylvis ad flumen Solimões prov. Rio Negro (Mart. ! in monac.), locis umbrosis calidis regni novo-granatensis inter la Egyptiaca et Honda alt. 170 hexap. (Humb. ! l. c. et in h. Willd. n. 643! et pro parte 644!), Cumana (Funck, n. 60! in h. Boiss.), Novæ Granatæ prov. Barbacoas (Triana, n. 36! in h. suo). — Kunth syn. v. 1. p. 104. Schilleria variegata Kunth! in Linn. v. 13. p. 695. Artanthe variegata Miq. ! syst. p. 485, in Linn. v. 20 p. 162. et in Mart. flor. Bras. fasc. 11. p. 52. t. 16. Ramuli glabri. Limbi 0,15-0,175 longi, 0,045-0,07 lati. Petioli 0,005-0,007 lati. Pedunculi 0,01-0,015 longi. Amenta immatura 0,07 longa, 0,002-0,003 crassa. (v. s.)
319. *P. VELUTINUM* (Kunth! in H. et B. nov. gen. v. 1. p. 52), foliis brevissime petiolatis oblongo-ellipticis apice protracto-acuminatis, acumine obtusiusculo mucronulato basi subæquali subattenuato-obtusiusculis utrinque molliter incano-pubescentibus rigidulo-membranaceis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes utrinque 4-5 supremos ad apicem ductos mittente, petiolo villosulo-piloso, pedunculo villosulo-piloso petiolum superante, amento cylindrico brevi quam folium brevius, bractæ apice peltatæ pelta semilunari margine hirtella, stigmatibus filiformibus, bacca obovato-trigona glabra flavo-glandulosa. ♀ In scopulis prope Zaulaca et San Felipe prov. Jaen de Bracamoros et in sylvis Peruvix prope Guancabamba in radicibus montis Paramo de Guamani alt. 700-1100 hexap. (Humb. ! l. c. et in h. Willd. n. 669!), Oaxaca (Buchinger! in h. Lenormand.). — Kunth! synopsis. v. 1. p. 100. Opiz in Reliq. Haenk. fasc. 3. p. 157. Schilleria velutina Kunth! in Linn. v. 13. p. 701. Artanthe velutina Miq. ! Syst. p. 471. Ramuli patentim molliter villosi. Limbi 0,105 longi, 0,05 lati. Petioli 0,004 longi. Pedunculi 0,006 longi. Amenta matura 0,03-longa, 0,003 crassa. (v. s.)
320. *P. FULGINEUM* (Kunth! olim), foliis brevissime petiolatis oblique oblongis apice acuminatis, acumine obtusiusculo, basi inæquali cordulatis cordatisve supra præsertim ad nervos adpresse pilosulis demum striguloso-verrucoso-scabris subtus pilis sordide fuscis crispulis molliter tomentosis coriaceis, nervis subtus prominulis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes latere majore 5 minore 4 supremos utrinque fere ad apicem ductos mittente, petiolo fusce tomentoso, pedunculo fusce pubescente petiolum quater superante, amento cylindrico quam folium brevius, bractæ apice peltatæ pelta triangulari margine densissime aureo-hirsuta, bacca obovata vertice puberula. ♀ In Brasilia meridionali (Sello! in h. Berol.). Steffensia? fulginea Kunth! in Linn. v. 13. p. 655. Artanthe fulginea Miq. ! Syst. p. 455 et in Mart. flor. Bras. fasc. 11. p. 47. Ramuli verruculis minutis scabri. Ramuli pilis crispulis sordide fuscis tomentosi. Limbi 0,06-0,095 longi, 0,025-0,04 lati. Petioli 0,005 longi. Pedunculi 0,02 longi. (v. s.)
321. *P. MOLLI-COMUM* (Kunth! olim), foliis breviter petiolatis oblique oblongo-ovatis vel oblongo-ellipticis apice acuminatis, acumine mucronulato, basi inæquali subrotundatis vel subcordatis supra hirtellis, pilis asperulis demum areolatis et asperis subtus pilis adpressis dense molliter pubescentibus, centrali nervo ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes subtus prominulos utrinque 5-6 supremos ad apicem ductos mittente, petiolo basi vaginante basi dorso et sursum utrinque hirsuto, pedunculo pubescente petiolum bis terve su-

perante, bractæ apice peltatæ pelta triangulari vel elliptica margine hirsuta, bacca obovata apice glanduloso-puberula vel glabra. ♂ In Brasilia (Sello! Chamisso! in h. Berol.), prov. Rio Janeiro (Mart. n. 85! Gaudich. n. 85! et 1115! Casaretto, n. 1670! Guillemín! in h. DC.), sylvis primævis ad Sebastianopolin et prope Corcovado (Mart. in h. monac.), ad Paraíbauna et montem Corcovado (Pohl, n. 20! 5025! in h. Vindob.), Ilheos (Wawra et Maly, n. 243 fide Wawra bot. Ergeb. p. 139), San Gabriel (Spruce, n. 2205 in h. Franco-vil.), Venezuela prope Biscaina (Fendler, n. 1132! in h. DC.), Caracas (Gallmer, n. 325! in h. Berol.), Novæ Granatæ prov. Pasto alt. 2000 m. (Triana, n. 43! in h. suo), Costa-Rica (Hoffmann, n. 349! in h. Berol.), Guatemala (Wars. in h. Boiss.), Panama (Seem. n. 116! in h. Kew.), Cuba (Wright n. 1687! in h. DC.), ins. Java culta (Zollinger, n. 3684! in h. DC.). *Steffensia mollis* Kunth! in Linn. v. 13. p. 648. *Artanthe mollicoma* Miq. 1 Syst. p. 438, in Lion. v. 20. p. 155 et in Mart. flor. Bras. fasc. 11. p. 45. *Art. adunca* Griseb. cat. pl. cub. p. 65 quo ad spec. *Wrightiana*. Ramuli molliter tomentoso-pubescentes. Limbi 0,12 0,14 longi, circ. 0,05 lati. Petioli 0,005 longi. Pedunculi 0,015 longi. Amenta matura usque ad 0,005 crassa. (v. s.)

322. *P. SALICARIÆFOLIUM* (Kunth olim), foliis brevissime petiolatis suboblique oblongo-lanceolatis apice mucronulatis basi æquali obtusiusculis vel acutis supra verruculis minutis obsitis scabriusculis subtus ad nervos piloso-pubescentibus rigido-membranaceis subopacis, nervo centrali subtus prominulo ad  $\frac{1}{4}$  longitud. nervos alternos adscendentes subtus prominulos utrinque 4-5 supremos ad apicem ductos mittente, petiolo hirtello, pedunculo hirtello petiolum superante, bractæ apice peltatæ pelta elliptica vel elliptico-rotunda aut quadrangulari margine hirsuta, bacca plerumque obovato-trigona vertice glandulosa interdum obovato-tetragona. ♂ In Brasilia prope Rio Janeiro (Gaudich. n. 1111! et 1113! in h. Berol. et DC.), ins. Santa Catharinæ (Tweddle! in h. Kew.). *Steffensia salicariæfolia* Kunth! in Linn. v. 13. p. 636. *Artanthe salicariæfolia* Miq. 1 Syst. p. 468, in Mart. flor. Bras. fasc. 11. p. 48 et in Hook. Lond. journ. bot. v. 4. p. 457. Rami ramulique scabriusculi. Limbi 0,1-0,15 longi, 0,025-0,035 lati. Petioli 0,004-0,006 longi. Pedunculi 0,011 longi. Amenta matura 0,05 longa, 0,0035 crassa. (v. s.)

β, *Lehmannianum*, ramuli glabri. — In Peruvia prope Casapi (Mathews, n. 1694! in h. Kew.). *Artanthe Lehmanniana* Miq. msc. in h. Kew.

323. *P. LANCEÆFOLIUM* (Kunth! in H. et B. nov. gen. v. 1. p. 49), foliis brevissime petiolatis suboblique oblongo-lanceolatis longissime acuminatis cuspidatis, cuspidate acuto, basi parum inæquali cordulatis supra glabris bullatis lævibus nitidis subtus molliter pubescentibus coriaceis epunctatis. centrali nervo subtus prominente ad  $\frac{1}{4}$  longitud. nervos alternos adscendentes subtus prominentes utrinque 8 10 supremos ad apicem ductos mittente, petiolo villosa, pedunculo pubescente petiolum multoties superante, bractæ apice peltatæ pelta triangulari margine griseo-lutescente hirsuta, bacca glabra obovato-trigona, stigmatibus 3-4. ♂ In Peruvia (Jameson, n. 1714! in h. Kew.), monte Pichincha (Mathews, n. 380! in h. DC., 714! in h. Kew. et Boiss.), republica æquatoriali (Fraser! in h. DC.), Nova Granata (Linden, in 976! in h. Kew.), sylvis opacis Andium locis udis prope Yanta in prov. Jaen de Bracamoros, ad ripas fluminis Amazonum alt. 200-700 hexap. (Humb. l. c. et in h. Willd. n. 661!) *Schilleria lanceæfolia* Kunth! in Linn. v. 13. p. 702). *Artanthe lanceæfolia* Miq. 1 Syst. p. 433, illust. p. 75. t. 79 et in Hook. Lond. journ. bot. v. 4. p. 455. Ramuli pilis sericeis tecti. Limbi 0,16-0,25 longi, 0,05-0,07 lati. Petioli 0,005-0,01 longi. (v. s.)

β, *acutiusculum*, foliis basi acutiusculis junioribus pubescentibus (Jameson n. 380! in h. Kew.).

γ, *bullosum*, foliis magis coriaceis supra magis bullatis subtus densius molliter pubescentibus. — In Novæ Granatæ prov. Antioquia (Triana n. 41! in h. suo).

324. *P. VILLOSUM* (Cas. DC. in Seem. journ. 1866 p. 212), foliis brevissime petiolatis obovato-lanceolatis apice protracto-acuminatis, acumine acuto, basi æquali subattenuatis cordulatis supra ad bullas subtus ad nervos et ner-

vulus pilis longis villosis ciliatis membranaceis glandulis fuscis subtus spersis epunctulatis supra bullatis haud scabris, centrali nervo ad  $\frac{1}{2}$  longius nervos subalternos subadscendentes utrinque 6 mittente, petiolo villosopetulo glabro petiolum duplo superante, amento quam folium multoties longiore, bractæ apice truncato-peltatæ pelta nuda, bacca obovato-trigona vix glandulosa. ♀ In Novæ Granatæ prov. Barbacoas (Triana n. 61 m DC.). Ramuli juniores villosi. Limbi 0,18 longi, 0,09 lati. Petioli 0,008 longi. Amenta matura 0,05 longa, 0,005 crassa. (v. s.)

325. *P. PSEUDOASPERIFOLIUM*, foliis breviter petiolatis oblique sublanccatis ellipticis apice acuminatis, acumine acuto, basi inæquali obtusis supra pilis pubescentibus firmulo-membranaceis subopacis subtilissimè pellucido-punctulatis, nervis subtus prominulis, centrali fere ad  $\frac{1}{2}$  longius nervos alternos subadscendentes utrinque 6 supremos ad apicem ductos 3-4 minus subtiliores mittente, petiolo basi vaginante basi dorso et sursum utrinque hirsuto, pedunculo hirtello petiolum æquante, amento maturo quam folium paulo brevior, bractæ dorso hirsutæ apice truncato-peltatæ pelta triangulari angusta margine hirsuta, bacca obpyramidato-trigona vix glandulosa. ♀ In Oaxaca (h. Boiss.). Ramuli fuscescentes hirsuti. Limbi 0,18 longi, 0,05 lati. Petioli 0,01 longi. (v. s.)

326. *P. TRIGONOIDÆUM*, foliis subsessilibus oblongo-lanceolatis apice angustè acuminatis, acumine acuto mucronulatoque, basi æquali acutis supra pilis pilosis subtus ad nervos nervulosque densius adpresse hirtello-pubescentibus firmule membranaceis opacis pellucido-punctulatis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitudinis nervos alternos subadscendentes supremos 6 ad apicem ductos mittente, pedunculo hirtello, amento maturo quam folium dimidio brevior, bractæ apice peltatæ pelta triangulari margine fuscè laevi, bacca obpyramidato-trigona nigra glabra. ♀ In Brasiliæ prov. Rio Negro (Mart. n. 377! in h. monac.). Rami adpresse puberuli dein glabrati. Ramuli densius hirtelli. Limbi 0,16 longi, 0,045 lati. Pedunculi 0,01 longi. Amenta matura 0,004 crassa. (v. s.)

327. *P. BLANCHETII*, foliis breviter petiolatis oblique ovato-ellipticis vel obovato-ellipticis apice oblique acuminatis, acumine acuto, basi inæquali rotundius subcordatisve utrinque præsertim ad nervos densius pubescentibus membranaceis subtilissimè pellucido-punctulatis, centrali nervo ad  $\frac{1}{2}$  longitudinis nervos alternos subadscendentes utrinque 5-6 supremos ad apicem ductos mittente, petiolo dense pubescente, pedunculo puberulo quam petiolus brevior, amento cylindrico quam folium brevior, bractæ apice peltatæ pelta triangulari semilunari albo-ciliata, bacca obovato-trigona glabra. ♀ In Brasiliæ meridionali (Blanchet n. 3149 A in h. Deless. fide Miq. syst. p. 173). Baccæ collibus (Salzmänn in h. Deless. fide Miq. l. c.). Artanthe Blanchetii Miq. l. c., illustr. p. 76. t. 81 et in Mart. flor. Bras. fasc. 11. p. 49. Rami glabri vel sparse puberuli. Ramuli densius puberuli. Limbi 0,17-0,18 longi, 0,06-0,085 lati. Petioli 0,015-0,02 longi (ex Miq. l. c.)

328. *P. LAURIFOLIUM* (Kunth! in H. et B. nov. gen. v. 1. p. 47), foliis brevissimè petiolatis ellipticis vel subovato-ellipticis apice acuminatis, acumine acutiusculo, basi æquali obtusis utrinque glabris junioribus supra subtilissimè glanduloso-punctulatis coriaceis nitidis subopacis subtilissimè crebre pellucido-punctulatis, nervis subtus vix prominulis, centrali ad  $\frac{1}{2}$  longitudinis nervos alternos subadscendentes utrinque 4-5 supremos ad apicem ductos sursumque sæpe nervulos validos mittente, petiolo glabro, pedunculo glabro quam petiolus subbrevior, amento quam folium multum brevior cylindrico, bractæ intus et basi dorso hirsutæ apice truncato-peltatæ pelta subquadrangulari glabra angulo exteriori carnosulo-tuberculata, bacca obovato-trigona. ♂ In regione calida Andium peruvianorum locis scopulis inter Chala et fruxillo alt. 150 hexap. (Humb. l. c. et in h. Willd. n. 647). — Kunth synopsis. v. 1. p. 104; Roem. et Schult. syst. v. 1. p. 290. Schüller laurifolia Kunth! in Linn. v. 13. p. 691. Artanthe laurifolia Miq. l. c. p. 520, illustr. p. 80. t. 89. Rami ramulique glabri. Limbi 0,12-0,14 longi, 0,03-0,06 lati. Petioli 0,005 longi. Amenta florentia 0,075 longa, 0,004 crassa. (v. s.)

329. *P. COLURRINUM* (Link Jahrb. v. 3. p. 61, fide Roem. et Schult. mant.), foliis breviter petiolatis oblongo-ellipticis apice breviter attenuatis acutis mucronatisque basi æquali vel subæquali acutis utrinque glabris supra aliquando scabriusculis coriaceis membranaceisve opacis subtus fusce-nigrove-punctulatis, centrali nervo subtus prominulo nervos alternos subadscendentes nervulusque validioribus sæpe intermixtos supremos per anastomoses fere ad apicem ductos mittente, petiolo glabro basi ima vaginante, pedunculo glabro petiolum paullo superante, bractæ apice peltatæ pelta semilunari margine dense hirsuta, bacca obovato-trigona apice glandulosa. ♂ In Peruvia (Poeppig n. 204! in h. Berol. cum *P. heterophyllo confusum*), prope Casapi (Mathews n. 1692! in h. Kew.), Brasilia (Sello! in h. Berol., Riedel n. 292! 24! et 721! Luschinath n. 121! in h. Petrop.), sylvis prope Sebastianopolin et fluv. Itahype (Mart.! in h. monac.), montosis prope Toaca (Schott! in h. Vindob.) et monte Corcovado (Pohl n. 5027, in h. Vindob.), Ilheos (Wawra et Maly n. 311! et 347 fide Wawra bot. Ergeb. p. 139), prope Bahía (Salzman n. 525! in h. DC.; Glocker n. 71 in h. Boiss., Blanchet n. 3151! in h. DC., Mart. n. 1209! in h. DC. et 1211! in h. monac.), vicinibus Para (Spruce! in h. Kew.), Guyanæ sylvis umbrosis (Rich! in h. Francovil.), Caracas (Jose de Vargas n. 471 in h. DC.), Hoffmannsegg in h. Willd. n. 645, — Link enum. alt. v. 1. p. 37. *P. impunctatum* Willd. herb. l. c., Link Jahrb. 1. 3. p. 63 (fide Roem. et Schult. mant.). *P. crassum* Vell. flor. flum. t. 59 (non Bl.). *P. densespicatum* Steud. nom., Miq. syst. p. 535. *P. Willdenowii* Kunth! in h. Berol. *Schilleria colubrina* Kunth! in Linn. v. 13. p. 685. *Artanthe colubrina* Miq.! syst. p. 512, in Linn. v. 20. p. 170 et in Mart. flor. Bras. fasc. 11. p. 56. Art. obovata Miq.! in Hook. Lond. journ. bot. v. 4. p. 468. Ramuli glabri, castaneo-fusci. Limbi 0,16 longi, 0,6 lati. Petioli 0,01 longi. Pedunculi 0,011-0,012 longi. Amenta matura vix 0,005 crassa. (v. s.)

β, *adenophyllum*, foliis minoribus, limbis 0,105 longis, 0,047 latis, amentis 0,045 longis. — In sylvis umbrosis ad Coralfalga (Pohl n. 5026! in h. Vindob.). *Artanthe adenophylla* Miq.! in Linn. v. 20. p. 170 et in Mart. flor. Bras. fasc. 11. p. 57. t. 21. f. 1. (v. s.)

330. *P. LENTICELLOSUM*, foliis breviter petiolatis oblongis apice acuminatis, acumine acuto, basi æquali subattenuatis acutiusculis utrinque glabris coriaceis opacis obsolete pellucido-punctulatis supra sub lente bullulosis nitidulis lævibus, centrali nervo subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes utrinque 4 subtus prominulos supremos ad apicem ductos nervulosque validiores mittente, petiolo glabro, pedunculo glabro petiolum æquante, amento quam folium brevius, bractæ peltatæ pelta subsessili subrotunda centro peltata membranacea nuda, rhachi foveolata, foveolis margine ciliolatis, bacca obovato-trigona glabra vel vertice subtilissime pubesula ♂ In Novæ Granatæ prov. Barbacoas alt. 700 (Triana n. 31! in h. DC.) et prov. Cauca alt. 1000 (Triana n. 29! in h. suo). Ramuli lenticellosi glabri. Limbi 0,115 longi, 0,04 lati. Petioli 0,005 longi. Amenta 0,06 longa, 0,003 crassa, ebullitione aromatica. (v. s.)

## 2° Folia multiplinervia.

331. *P. FUSCUM*, foliis petiolatis inferioribus ovato-oblongo-lanceolatis superioribus oblongo-lanceolatis omnibus apice longiuscule acuminatis, acumine acuto mucronulatoque, basi æquali rotundatis obtusius vel inferioribus cordatis sæpe parce et basi densius fusce pilosis subtus dense fusce piloso-villosulis membranaceis subopacis pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  —  $\frac{2}{3}$  longitudinis nervos alternos subadscendentes utrinque 6-7 nervulus validioribus intermixtos supremos ante apicem per anastomoses evanidos inferos utrinque 2 et basi solutos mittente, petiolo utrinque fusce villosulo basi ima vaginante, pedunculo fusce villosulo petiolum æquante, amento maturo quam folium multoties brevius, bractæ apice truncato-peltatæ pelta triangulari margine subtiliter ciliolata, bacca obovato-trigona nigra glabra. ♂ In Brasilia (Riedel n. 291! in h. Petrop.), in hort. Petrop. culta. Rami verruculis obsiti. Ramuli fusci villosi. Limbi 0,165 longi, 0,045 lati,

Petiolis inferiores 0,015, superiores 0,013 longi. Amenta matura 0,0025 crassa. (v. s.)

- 332: *P. LEPTURUM* (Kunth! olim), foliis breviter petiolatis lanceolatis vel oblongo-lanceolatis apice suboblique acuminatis, acumine acuto mucronulatoque. inferioribus basi æquali cordatis superioribus basi æquali rotundatis vel subattenuatis vel subacutis omnibus utrinque glabris membranaceis pellucidopunctulatis, centrali nervo subtus prominulo nervos alternos adscendentes subtus prominulos nervulis validioribus intermixtos utrinque 4-6 supremos fere ad apicem ductos infimos utrinque 2 vel uno latere 3 e basi solutos magis adscendentes mittente, nervulo obscuro juxta marginem percurrente, petiolo glabro pedunculum glabrum superante, bractæ apice truncato-peltatæ glabræ juniores subtilissime ciliolatæ pelta triangulari-rhombea, bacca obovato-trigona glabra.  $\S$  In Brasilia (Langsdorff n. 33! 34! 109! in h. Petrop.; Mart. n. 1213! in h. DC.), prope Rio Janeiro (Gaudich. n. 1117! in h. Berol.; Meyen! in h. Berol.), humidis umbrosis monte Corcovado (Mart.! in h. monac.), Mexico prope Oaxaca (Buchinger! in h. DC.). *Schilleria leptura* Kunth! in Linn. v. 13. p. 679. *Artanthe meyeniana* Klotzsch! msc. in h. Berol.; Miq.! syst. p. 524 et in Mart. flor. Bras. fasc. 11. p. 60. *Art. leptura* Miq.! syst. p. 523 et in Linn. v. 20. p. 172 et in Mart. flor. Bras. fasc. 11. p. 61. Rami ramulique glabri. Limbi 0,11-0,14 longi, 0,025-0,03 lati. Petioli 0,01, pedunculi 0,003-0,005 longi. Amenta matura 0,05-0,06 longa. 0,002 crassa. (v. s.)
- 333: *P. BARBATUM* (Kunth! in H. et B. nov. gen. 1. p. 55), foliis breviter petiolatis ovato-acuminatis, acumine acuto, basi æquali vel subinæquali rotundatis utrinque glabris subtus ad nervos fimbriulatis firmis opacis, nervis subtus prominulis centrali supra basin nervum utrinque unum alternatim alte ductum et e basi utrinque 2 infimos subtiliores mittente, petiolo margine ciliolato pedunculum superante, amento cylindrico quam folium brevior bractæ apice peltatæ, pelta rotundato-semilunari margine dense luteo-hirsuta stigmatibus 3-4, bacca obovato-globosa glabra.  $\S$  In locis subapricis inter Lloa et Quito alt. 1092 hexap. (Humb. l. c. et h. Willd. n. 687!), in Andibus quintisibus (Jameson n. 409! in h. DC.), Nova Granata (Fraser! in h. DC.). — Kunth synops. v. 1. p. 110. *Schilleria barbata* Kunth! in Linn. v. 13. p. 712. *Artanthe barbata* Miq.! syst. p. 414. Ramuli glabri. Limbi 0,07-0,11 longi, 0,045-0,07 lati. Petioli 0,01-0,02 longi. Amenta matura 0,005 crassa. (v. s.)
- 334: *P. ANMICOLUM* (Kunth in nov. gen. v. 1. p. 55), foliis brevissime petiolatis ovato-acuminatis, acumine acuto, basi æquali cordatis supra ad nervos molliiter pilosulis subtus molliter pubescentibus membranaceo-firmis opacis, centrali nervo subtus prominulo nervos subadscendentes utrinque 3-4 fere e basi solutos subtus prominulos mittente, petiolo puberulo, pedunculo petiolum æquante, bractæ apice peltatæ pelta triangulari margine hirsuta bacca obovata glabra.  $\S$  In prov. Chachapoyas (Mathews n. 3223! in h. Kew.), Quito (Bonpl.! in h. Francovil.), frequentissime in devexis Andium peruvianorum prope Guancabamba et in radicibus montis Guamani alt. 1030 hexap. (Humb. l. c.), Nova Granata (Triana n. 161! in h. Kew.). — Kunth synops. v. 1. p. 110. *Artanthe andicola* Miq.! syst. p. 415 et in Hook. Lond. journ. bot. v. 4. p. 450. Ramuli puberuli. Limbi 0,1 longi, 0,06-0,07 lati. Petioli 0,005-0,006 longi. Amenta matura 0,06-0,07 longa, 0,004 crassa. (v. s.)
- 335: *P. AURITUM* (Kunth! in H. et B. nov. gen. v. 1. p. 54), foliis petiolatis ovato-oblongis apice attenuatis basi valde inæquali cordatis supra dense pilosulis subtus pubescentibus membranaceis, centrali nervo subtus prominulo ad  $\frac{1}{2}$  longitudinis nervos alternos subadscendentes subtus prominulos utrinque 6-7 inferos utrinque circ. 3 e basi divaricantes mittente, petiolo glabro ad limbi basin vaginante, alis linearibus, pedunculo petiolum æquante, bractæ apice peltatæ pelta triangulari margine hirsuta, stam. 3-4 vel apice amenti 2. stigmatibus 3 linearibus bacca trigona vertice glanduloso-puberula.  $\S$  In Nova Granatæ (Triana, n. 11! in h. Triana), istmo Panama (Seemann, n. 1069! in h. Brit. mus. et 737! in h. Kew.), Guatemala (Friedrichsthal, n. 477! in h. Vindob.), Darien meridionali (Miq.! in bot. Herald.), temperatis Novæ Hispaniæ alt. 700 hexap. (Humb. l. c. et in h. Willd. n. 678!), Mexico prope Orizaba (Sallé! in h. DC.), Puebla (Andrieux, n. 93! in h. DC.), sylvis ad



Jalapam (Schiede! in h. Berol.), ad ripas rivulorum prope Mirador et Hacienda de la Laguna alt. 1000-1500 ped. (Galeotti, n. 6016 fide Mart. et Gal. in bull. ac. Brux. v. 10. p. 21.), ins. Santi Thomæ (Friedrichsthal! in h. Lenorm.), Antioquia (Jewse, n. 199! in h. Kew.). — Kunth synops. v. 1. p. 109, Mart. et Gal. l. c. Schilleria aurita Kunth! in Linn. v. 13. p. 713. Artanthe aurita Miq.! syst. p. 400. Art. Seemanniana Miq.! bot. Herald p. 199. t. 39. Ramuli glabri. Limbi 0,16 longi, 0,09 lati. Petioli 0,015, pedunculi 0,025 longi. Amenta mat. 0,004 crassa, ebullitione aromatica. (v. s.)

β, *Schiedeanum*, foliis subtus dense pubescentibus. — In Mexico (Schiede et Deppe! in h. Brit. Mus.).

γ, *amplifolium*, foliis amplioribus 0,35 longis 0,25 latis. — In Costa-Rica (Hoffmann n. 761! in h. Berol.).

336. P. *HILARIANUM*, foliis longe petiolatis rotundato-ovatis ovatisve apice breviter acuminatis, acumine acutiusculo, basi cordulatis supra glabris subtus ad nervos hirtellis membranaceis pellucido-punctulatis subundeninerviis, nervis subtus prominulis centrali paulo supra basin nervum utrinque unum sursum fere tota longitudine nervulos validiores et e basi nervos utrinque 4 subadscendentes mittente, petiolo glabro ad  $\frac{1}{2}$  longitud. vaginante (alis linearibus) pedunculum glabrum multum superante, bractæ apice truncato-peltatæ pelta semilunari ciliolata pedicello cucullato margine ciliolato, bacca obovato-trigona glabra. ♀ In Brasilia ad Yriro (Sello! in h. Berol.). Artanthe Hilariana Miq.! syst. p. 421, illust. p. 74. t. 76, in Mart. flor. Bras. fasc. 11. p. 41. Ramuli glabri. Limbi 0,15 longi. 0,11 lati. Petioli 0,05 longi, pedunculi 0,012 longi. Amenta matura 0,004 crassa. (v. s.)

337. P. *SUBSCUTATUM*, foliis longe petiolatis paulo supra basin peltatis late ovatis apice abrupte falcatis acuminatis, acumine angusto obtusiusculo, basi æquali leviter cordatis utrinque glabris membranaceis pellucido-punctatis 15-17-plinerviis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos alternos utrinque 2 et e basi utrinque 5-6 mittente, petiolo glabro ad  $\frac{1}{2}$  longitud. vaginante, pedunculo quam petiolus brevior, amento maturo folium æquante, bractæ oblongæ vertice truncato triangulari glabro vel parvisime ciliolato pedicello trigono basi pilosulo, rhachi pilosula, bacca obpyramidato-trigona. ♀ In Peruvia (Mathews, n. 1700! in h. Kew.), prope Tarapoto (Spruce, n. 4848! in h. Mart.). Artanthe scutata Miq.! in Hook. Lond. journ. bot. v. 4. p. 444. Ramuli glabri. Limbi 0,26 longi, 0,215 lati. Petioli 0,07-0,09 longi. (v. s.)

338. P. *TENUIFOLIUM*, foliis petiolatis basi ima subpeltatis ovatis apice acuminatis, acumine acuto, inferioribus basi æquali subrotundato-truncatis truncatisve superioribus basi æquali obtusis utrinque glabris membranaceis subpellucidis pellucido-punctulatis septupli-noveninerviis, nervis subtus prominulis centrali supra basin ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitudinis nervos alternos adscendentes utrinque 2-3 nervulosque validiores paucos et e basi nervum utrinque unum mittente, petiolo glabro basi ima vaginante, pedunculo glabro quam petiolus brevior, amento maturo cylindrico folium subæquante, bractæ pelta triangulari margine ciliolata pedicello angusto longiusculo, bacca obpyramidato-trigona vertice glandulosa. ♀ In Nova Hispania (P. tenuifolium Pav. in h. Boiss.). Ramuli glabri. Limbi 0,155 longi, 0,08 lati. Petioli 0,025 longi. Amenta matura vix 0,004 crassa. (v. s.)

### 3° Folia multinervia.

339. P. *POPULIFOLIUM* (Kunth olim), foliis breviter petiolatis subrotundato-ovatis apice acuminatis, acumine obtusiusculo, basi subæqualibus utrinque glabris membranaceis pellucido-punctulatis supra punctulis pluribus minutissimis albidis obsitis 11-13-nerviis, nervis subtus prominulis, petiolo glabro, pedunculo petiolum superante, bractæ lineari-spathulatæ vertice truncato-triangulari margine ciliato, bacca obovato-trigona, stigmat. 3? ♀ In Brasilia (Sello! fide Kunth.). Schilleria populifolia Kunth! in Linn. v. 13. p. 709. Artanthe populifolia Miq. syst. p. 418 et in Mart. flor. Bras. fasc. 11. p. 40. Ramuli læves glabri. Folia 7-8-pollicaria, 5 $\frac{1}{2}$ -7 poll. lata. Petioli 2-3 lin. Pedunculi 9 lin. longi. (v. s.)

b. *Bractea cucullata vertice inflexa vel obovato-cucullata vel obovata.*

1° Folia pennivervia.

\* Centralis nervus tota longitudine nervos nervulosque mittens.

340. *P. EUCALYPTIFOLIUM* (Rudgel! ic. p. 10. t. 6), foliis breviter petiolatis lanceolatis apice acuminatis, acumine acuto, basi æquali cuneatis utrinque glabris rigidulo-membranaceis nitidulis opacis pellucido-punctulatis, centrali nervo tota longitud. nervos subpatulos alternos utrinque 10-11 nervulis validioribus intermixtos et inferiores utrinque 2 subadscendentes mittente, petiolo glabro pedunculo glabro petiolum æquante, amento mucronulato quam folium multoties brevior, bracteæ cucullatæ glabræ vertice inflexo, stam....? stigmatibus brevibus, bacca obteträgona. § In Guyana anglica (Rudge, l. c. et h. Brit. Mus.), Guyana gallica (Poiteau! in h. Deless.), Cayenna (h. DC.). — Mart. syst. mat. med. p. 99. Steffensia eucalyptifolia Kunth in Linn. v. 13. p. 620. Artanthe eucalyptifolia Miq.! syst. p. 505. Ramuli uno latere pilosi. Limbi 0,07-0,11 longi, 0,035-0,045 lati. Petioli 0,005-0,006 longi. (v. s.)

341. *P. LUCÆANUM* (Kunth! olim), foliis breviter petiolatis lanceolatis apice acuminatis, acumine acuto basi æquali acutis utrinque glabris rigidulo-membranaceis creberrime pellucido-punctulatis subopacis, centrali nervo subtus prominulo nervos alternos non procul a basi utrinque 2 subadscendentes sursumque tota longitudine nervos nervulosque validiores patulo-subadscendentes mittente, petiolo glabro, pedunculo glabro petiolum æquante, amento quam folium brevior, bracteæ obovato-cucullatæ vertice inflexo margine subtilissime ciliolato peltam simulante, bacca obovato-trigona. § In Rio Janeiro (Gaudich! in h. Berol.), monte Corcovado et ubique prope Sebastianopolin (Mart.! in h. monac.), Venezuela inter Pentaquire et mare (Fendler, n. 2614! id h. DC.), isthmo Panama (Seem. n. 266! in h. Kew.), Chagres (Fendler, n. 268! in h. Petrop.). — Kunth! in h. Berol. Schilleria lucæana Kunth! in Linn. v. 13. p. 678. Artanthe lucæana Miq.! Syst. p. 523, in Linn. v. 20. p. 172 et in Mart. flor. Bras. fasc. 11. p. 61. Ramuli glabri. Limbi 0,095 longi, 0,025 lati. Petioli 0,008 longi. (v. s.)

β, *magnifolium*, foliis majoribus, limbis 0,15 longis, 0,055 latis, petiolis 0,01 longis. Amenta mat. 0,09-0,1 longa, 0,003 crassa. In Venezuela prope coloniam Tovar (Fendler, n. 1138! in h. DC. et Boiss.)

\*\* Centralis nervus tantum parte longitudinis nervos mittens.

342. *P. TUMIDUM* (Kunth! in nov. gen. v. 1. p. 51), foliis brevissime petiolatis oblongo-ellipticis apice obtusis basi inæquali rotundatis supra scabriusculis subtus præsertim ad nervos hirtellis coriaceis opacis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. nervos suboppositos subadscendentes utrinque 4 supremos ad apicem ductos mittente, petiolo hirtello, pedunculo petiolum superante, amento brevi cylindrico folium æquante, bracteæ cucullatæ extus dorso villosulæ vertice inflexo ciliolato peltam triangularem simulante, bacca ovato-cuneata glabra. § In Andibus Quitensibus alt. 1500 hexap. (Humb. in nov. gen. v. 1. p. 51 et in h. Willd. n. 6681). — Kunth synopsis. v. 1. p. 107. Steffensia? tumida Kunth! in Linn. v. 13. p. 654. Artanthe tumida Miq.! syst. p. 465. Ramuli pilis retrorsis hispiduli. Limbi 0,035-0,04 longi, 0,02-0,03 lati. Amenta matura 0,004 crassa. (v. s.)

343. *P. CORNIFOLIUM* (Kunth! in H. et B. nov. gen. v. 1. p. 52), foliis brevissime petiolatis ovato-ellipticis apice acuminatis, acumine acuto mucronulatoque. basi inæquali cordulatis supra glabris subtus ad nervos adpresse puberulis membranaceis subopacis, centrali nervo subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos adscendentes utrinque 3-4 mittente, petiolo hirtello, pedunculo hirtello petiolum paulo superante, bracteæ cucullato-spathulatæ dorso ciliolato vertice inflexo vel subtruncato peltam triangularem fingente, bacca obovato-trigona vertice glandulosa. § In ripa fluminis Magdalene juxta Pa-

turia, inter La Simitara et Badillas (Humb. l. c. et in h. Willd. n. 6701), Colombia (Cuming, n. 1231! in h. Kew.). — Kunth synops. v. 1. p. 108. Schilleria cornifolia Kunth! in Linn. v. 13. p. 697. Artanthe cornifolia Miq. l. syst. p. 479, illustr. p. 77. t. 83, in Hook. Lond. journ. bot. v. 4. p. 459. Ramuli adpresse hirtelli. Limbi 0,1 longi, 0,05 lati. Petioli 0,003 longi. Pedunculi 0,01 longi. Amenta matura 0,05-0,06 longa, 0,002 crassa. (v. s.)

344. P. KEGELII, foliis brevissime petiolatis elliptico-vel subovato-lanceolatis utrinque glabris membranaceis subpellucidis apice acuminatis, acumine acuto mucronulato, basi inæquali obtusis, nervo centrali ad  $\frac{3}{4}$  longitud. nervos alternos subadscendentes utrinque 5-6 mittente, petiolo glabro, pedunculo glabro petiolum multum superante, amento cylindrico brevi, bractæ cucullatæ vertice inflexo peltam triangularem glabram fingente, bacca obovata glabra nigra. ♀ In Surinam (Hostmann, n. 217! in h. Kew.), sylvis ad Mariepaston (Kegel n. 1329! et 1336! fide Miq. in Linn. v. 22. p. 77). Artanthe hymenophylla Miq. l. syst. p. 532. Ramuli glabri. Limbi 0,09 longi. 0,03-0,045 lati. Petioli 0,003-0,004. Pedunculi 0,02 longi. Amenta matura 0,025 longa, 0,004 crassa. (v. s.)

345. P. CRASSINERVIIUM (Kunth! in H. et B. nov. gen. v. 1. p. 48), foliis breviter petiolatis oblique ovatis acuminatis vel ovato-lanceolatis basi æquali subattenuatis utrinque glabris subtus ad nervos parvisime puberulis subcoriaceis subtiliter glanduloso-punctatis, centrali nervo subtus prominulo ad  $\frac{1}{2}$  longitud. nervos alternos subtus prominulos adscendentes utrinque 4-5 supremos fere ad apicem ductos infimos subtiliores mittente, petiolo glabro vel parvisime puberulo pedunculo quam petiolus multum brevior, bractæ subcalceiformi-peltatæ vertice carnosio semilunari vel subrotundo margine aureo-hirtello pedicello dorso carinato extus hirt. stom. exsertis, bacca obovata tri-tetragona glabra. ♀ In Peruvia (Mathews, n. 1147! in h. Kew.), Brasilia (Sello! in h. Berol.), prope Santo Paulo (Guillemin, n. 376! in h. DC.), Certao Serra d'Estrella (Pohl, n. 1706! et 4774! in h. Vindob.), in regione calidissima ad ripam fluminis Magdalenæ prope Honda alt. 180 hexap. (Humb. l. c. et h. Willd. n. 655!). — Kunth synops. v. 1. 104. P. hydrangeifolium Kunth! in Linn. v. 13. p. 659. Steffensia? crassinervia Kunth! in Linn. v. 13. p. 659. Steff. hydrangeifolia Kunth! l. c. Artanthe Guilleminiana Miq. l. syst. p. 409. Art. enckeoides Miq. l. syst. p. 412. Peltobryon Guilleminianum Miq. l. in Mart. flor. Bras. fasc. 11. p. 30. Artanthe exserens Miq. l. in Linn. v. 20. p. 148. Art. lentaginoides Miq. l. in Hook. Lond. journ. bot. v. 4. p. 468, pro parte. Ramuli glabri. Limbi 0,12-0,14 longi, 0,055-0,06 lati. Petioli 0,01, pedunculi circ. 0,004 longi. Amenta matura 0,05-0,06 longa, 0,005 crassa. (v. s.)

346. P. LAPATHIFOLIUM (Steud. nomencl.), foliis petiolatis oblongo-ovatis apice breviter acuminatis, acumine acuto, basi subinæquali cordatis supra glabris subtus ad nervos hirtellis membranaceis, nervis subtus prominentibus centrali ad  $\frac{3}{4}$  longitud. nervos alternos subadscendentes utrinque 8 supremos fere ad apicem ductos infimos subtiliores magis approximatos mittente, petiolo ad limbi basin usque vaginante dorso hirtello pedunculum superante alis linearibus, amento quam folium multum brevior cylindrico mucronato, bractæ cucullato-obovatæ præsertim postice longe pilosæ vertice inflexo nudo peltam semilunarem simulante, bacca obovata glabra. ♀ In Guyana gallica (Leprieur n. 145! in h. Deless.), sylvis prope Jalapam (Schiedel in h. Berol.), nemoribus humidis et ad rivulos prope Xalapa et Mirador alt. 3000-4500 ped. (Martens et Gal. in bull. ac. Brux. v. 10. p. 21 et Galeotti n. 6012! in h. Deless.), Jamaica (Purdie in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 446). Schilleria laphathifolia Kunth! in Linn. x. 13. p. 714. Artanthe laphathifolia Miq. l. syst. p. 401 et in Hook. Lond. journ. l. c. Art. Lessertiana Miq. l. syst. p. 405. P. Jalapense Mart. et Gal. l. c. Ramuli glabri. Limbi 0,24 longi, 0,11 lati. Petioli 0,04 longi. Amenta ante maturationem 0,06 longa, 0,003-0,004 crassa. (v. s.)

347. P. SUBAURICULATUM, foliis petiolatis ovatis vel superioribus oblongo-ovatis apice acuminatis, acumine obtusiusculo mucronulatoque, basi æquali vel subinæquali rotundatis vel excisis utrinque glabris firmulo-membranaceis subpellucidis creberrime pellucido-punctulatis, nervis subtus vix prominulis centrali ad  $\frac{1}{2}$ - $\frac{2}{3}$  longitud. nervos suboppositos subadscendentes utrinque 4-5

sursumque nervulos validiores mittente nervisque inferis utrinque 2-3 e basi solutis, petiolo glabro basi vaginante, pedunculo glabro petiolum æquante, amento quam folium multum brevior, bractæ obovato-cucullatæ basi cuneatæ extus glabræ intus pilosulæ vertice inflexo triangulari. *stam.* 4 ? filamentis antheras multum superantibus, bacca obovato-trigona. ♀ In Brasilia (Pohl n. 5034! in h. Vindob.). Artanthe subauriculata Miq.! in Linn. v. 20. p. 150. Rami ramulique glabri. Limbi 0,12 longi, 0,045 lati. Petioli 0,01 longi. Amenta matura 0,004 crassa. (v. s.)

348. *P. DENUDANS*, foliis breviter petiolatis oblongo-ovato-lanceolatis apice longe attenuato-acuminatis, acumine acuto, basi subæquali rotundatis supra glabris nitidulis subtus præsertim ad nervos fusce crispulo-pubescentibus membranaceis opacis pellucido-punctulatis, nervis subtus subprominulis centrali ad  $\frac{1}{2}$  longitud. nervos ascendentes utrinque 3-4 nervulosque validiores et nervos utrinque 3 basi solutos ascendentes quorum superiores ad medium ducti infimi subtiliores marginales mittente, petiolo ad limbum usque vaginante dorso hirtello, pedunculo juniore hirtello dein glabrato, amento quam folium paulo brevior, bractæ obovato-cucullatæ dorso et intus pilosulæ vertice subinflexo, bacca subobovato-trigona glabra nigra, rhachi subfoveolata. foveolis subtiliter pilosulis. ♀ In Brasiliæ monte Corcovado (Pohl n. 4118!, 4778!, 5028! in h. Vindob. et Mart.! in h. monac.). Artanthe denudans Miq.! in Linn. v. 20. p. 172 et in Mart. flor. Bras. fasc. 11. p. 60. t. 21. f. 2. Rami glabrati lenticellosi. Ramuli juniores dense pilosi. Limbi 0,12-0,13 longi, fere 0,045 lati. Petioli 0,01 longi. Amenta matura 0,002 crassa. (v. s.)

♂, *angustifolium*, limbis 0,13 longis, 0,025 latis. — In monte Corcovado (Pohl n. 5028!, Mikan n. 29! Endlich. n. 24! in h. Vindob.) *P. lanceolatum* Schott! mss. in h. Vindob.

## 2° Folia multinervia.

349. *P. SCHLECHTENDAHLIANUM*, foliis breviter petiolatis anguste elliptico-lanceolatis lanceolatisve apice longe acuminatis, acumine acuto obtusiusculo mucronulato, basi breviter inæquali obtusis acutisve utrinque glabris firmulo-membranaceis subopacis subtiliter pellucido-punctulatis 5-nerviis, nervis subtus prominulis 3 centralibus ad apicem ductis, petiolo glabro basi ima vaginante, pedunculo glabro petiolum subsuperante juvenili subtilissime puberulo, amento quam folium multoties brevior, bractea cucullato-obovata intus glabra extus basi fusce hirtella, bacca obovata glabra basi rhachi subimmersa, stigmatibus abbreviatis, rhachis foveolis hirtellis. ♂ In Cayenna (Rich.! in h. DC.), Mexico prope Misantla (Schiede! in h. Berol. et Pav.! in h. Boiss.). *P. angustifolium* Pav.! mss. in h. Boiss., certe non *P. angustifolium* Floræ peruvianæ. Enckea Schlechtendahlii Miq.! syst. p. 362. Rami ramulique glabri. Limbi 0,06-0,1 longi, 0,025-0,035 lati. Petioli 0,006 longi. (v. s.)

## § 4. Stigmata 3. Antheræ non articulatæ.

### A. Flores hermaphroditi.

350. *P. SELLOVIANUM* (Kunth! olim), foliis breviter petiolatis oblongo-ovato-ellipticis apice acuminatis, acumine acuto mucronulatoque, basi subæquali rotundatis vel basi ima subinæquali acutiusculis utrinque glabris vel subtus ad nervos ciliolatis membranaceo-rigidulis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervos alternos ascendentes utrinque 5 supremos ad apicem ductos infimos subtiliores mittente, petiolo glabro pedunculum hirtellum superante, amento cylindrico quam folium multum brevior, bractæ apice truncato-peltatæ pelta semilunari margine subaureo-hirsuta pedicello cucullato dorso subaureo-hirsuto, filamentis antheras superantibus, ovario apice attenuato. ♂ In Brasilia (Sello! sive Sellow in h. Berol.). Steffensia Selloviana Kunth! in Linn. v. 13. p. 657. Artanthe Selloviana Miq.! syst. p. 416, in Mart. flor. Bras. fasc. 11. p. 39, in Hook. Lond. journ. bot. v. 4. p. 449.

Ramuli glabri. Limbi 0,18 longi, 0,09 lati. Petioli 0,015 longi. Amenta florentia 0,005 crassa. (v. s.)

351. P. RIEDELII, foliis petiolatis ovatis apice acuminatis, acumine acuto, superioribus basi ima acutis obtusiusculisve inferioribus basi rotundatis breviter cordatis omnibus utrinque glabris membranaceis subopacis pellucido-punctulatis, nervis subtus subprominulis centrali ad  $\frac{1}{4}$  longitud. nervos alternos subadscendentes utrinque 4 infimos e basi solutos sursumque nervulos validiores nervosque subtiliores 2-3 subadscendentes mittente, petiolo glabro ad  $\frac{1}{4}$  longitud. vaginante, amento quam folium multum brevius mucronato densifloro, bractæ apice truncato-peltatæ glabræ pelta triangulari glabra. ♀ In Brasilia (Riedel n. 411! in h. Petrop.). Rami ramulique glabri. Limbi 0,18 longi, 0,15 lati. Petioli 0,025 longi. Amenta florentia 0,08 longa, 0,002 crassa. (v. s.)

#### B. Flores unisexuales.

352. P. NUDUM, foliis inferioribus longiuscule superioribus modice petiolatis inferioribus subrotundo-ovatis apice protracto-acuminatis, acumine obtusiusculo, basi æquali profunde cordatis superioribus ovatis apice acutis subprotracto-acuminatis basi truncato-rotundatis omnibus utrinque glabris membranaceis subpellucidis haud punctulatis inferioribus 9 superioribus 7-nerviis, nervis subtus prominulis, centrali ad apicem ducto sequentibus ante apicem evanidis infimis subtilioribus, petiolo glabro ad  $\frac{1}{4}$  longitud. vaginante alis linearibus, pedunculo glabro petiolum æquante, amento florente folium superante densifloro, bractæ spathulatæ vertice semilunari subinflexo margine exteriori ciliolato, filamentis antheras superantibus. ♀ In Nova Hispania (Pav. in h. Boiss.). Dioicum. Ramuli glabri. Limbi inferiores 0,15 longi, 0,085 lati, superiores 0,095 longi, 0,05 lati. Petioli inferiores 0,025, superiores 0,01 longi. (v. s.)

#### § 5. Stigmata duo lateralía. Flores hermaphroditi.

353. P. POIRETIANUM, foliis inferioribus petiolatis superioribus brevissime petiolatis omnibus elliptico-ovatis apice acuminatis, acumine acuto mucronulatoque, basi æquali obtusis obtusiusculisve utrinque glabris membranaceis subpellucidis subtilissime pellucido-punctulatis, centrali nervo ad  $\frac{1}{4}$  longitud. usque nervos alternos subtus prominulos subadscendentes utrinque 4 supremos per anastomoses ad apicem ductos mittente, petiolo glabro ad limbi basin usque vaginante alis linearibus, pedunculo glabro petiolum æquante, amento cylindrico post anthesin quam folium multoties brevius, bractæ apice truncato-peltatæ pelta elliptica glabra pedicello parce hirtello, ovario ovato glabro sursum in stylum brevem attenuato, stigmatibus recurvulis. ♀ In Brasilia (Riedel n. 1233! in h. Petrop.). Rami ramulique glabri. Limbi 0,07 longi, 0,033 lati. Petioli inferiores 0,011, superiores 0,003 longi. Amenta post anthesin 0,013 longa, 0,003 crassa. (v. s.)
354. P. ARTANTHOPSE (Cas. DC. in Mém. soc. phys. genev. 18. t. 1. fig. 13 organogr.), foliis petiolatis ovato-lanceolatis apice acuminatis, acumine acuto mucronulatoque, basi subæquali rotundatis breviter cordulatis utrinque glabris margine subtus ad nervulum hirtellis membranaceis vel rigidulo-membranaceis opacis subtiliter pellucido-punctulatis, centrali nervo subtus prominulo ad  $\frac{1}{4}$  longitud. usque nervos alternos subadscendentes utrinque 5 supremos remotiores infimos subtiliores mittente, nervulo obscuro subtus hirtello juxta marginem currente, petiolo glabro ad limbi basin usque vaginante alis linearibus, pedunculo glabro petiolum subæquante, amento cylindrico apice mucronato quam folium multoties brevius, bractæ cucullato-subcupuliformis dense carnosæ villosulæ intus pubescentis vertice inflexo triangulari, filamentis epigynis, ovariis rhachi immersis coalitis apice in stylum carnosum attenuatis, stigmatibus oblongis recurvis, baccis coalitis. ♀ In Costa-Rica ad Aguacate (Hoffmann, 687! et 776! in h. Berol.). Ramuli glabri. Limbi 0,17-0,19 longi, 0,06-0,08 lati. Petioli 0,01 longi. Amenta post anthesin 0,02 longa, 0,008 crassa. (v. s.)

355. *P. SINGULARE*, foliis petiolatis ovato-ellipticis apice acuminatis, acumine acuto, basi æquali acutiusculis acutisve utrinque glabris rigidulis opacis pellucido-punctulatis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervos alternos subadscendentes utrinque 5 supremos fere ad apicem ductos mittente, petiolo glabro ad limbi basin usque vaginante alis anguste linearibus, pedunculo glabro petiolum æquante, amento florente quam folium multoties brevior mucronato, bracteæ spathulatæ basi puberulæ vertice nudo inflexo dorso carnosulo peltam triangularem subfingente, antheræ loculis discretis, anthera longiuscule articulata, connectivo supra loculos papilloso-carnoso producto rubello, ovario apice stylifero, stylo ovarium superante, stigmatibus carnosulis applicatis. ♀ In Nova Granata (Triana n. 361! in h. suo). Ramuli glabri. Limbi 0,105 longi, 0,06 lati. Petioli 0,022 longi. (v. s.)

Sect. IV. CARPUNYA. — Pro parte gen. seq.: Schilleria et Enckea Kunth in Linn. v. 13. p. 676 — Piper Kunth olim — Ottonia Spreng. neue Entdeck. v. 1. p. 225. — Artanthe Miq. comm. phyt. p. 40. — Carpunya Presl Epim. p. 229.

Bractea libera. Stamina 3, quorum duo lateralia unum posticum. Amenta oppositifolia. Flores hermaphroditi.

### § 1. Stigmata 4. Antheræ articulatae.

356. *P. RHYNCHOSTACHYUM*, foliis breviter petiolatis lanceolatis apice attenuato-acutis, acumine obtusiusculo, basi leviter inæquali acutis utrinque glabris membranaceis subtilissime pellucido-punctulatis subnitidis supra saturate viridibus subtus fusciscenti-viridibus et ad lentem glandulis minutis punctatis, centrali nervo crassiusculo subtus prominente tota longitudine nervos 12-15 subpatulos mittente, petiolo ad limbum vaginante dorso serie pilorum sensim decidua instructo, amento brevissimo mucronato-rostrato, bractea carnosa inflexo-subconchæformi fere cupulari, stam. 3? stigmatibus teretiusculis obtusis erectis. ♀ In Guyana gallica (Poiteau! in h. Deless. fid. Miq. syst. p. 526). Artanthe rhynchostachya Miq. l. c. Rami cinereo-fusci. Petioli 0,004-0,006 longi. Limbi 0,08-0,03 longi, 0,02-0,03 lati. (ex Miq. l. c.)

357. *P. SPHÆROSTACHYUM* foliis brevissime petiolatis ellipticis apice obtusis acutiusculisve basi æquali acutiusculis acutisve utrinque glabris membranaceis glaucescentibus pellucido-punctulatis, centrali nervo subtus prominulo tota longitudine nervos patulo-subadscendentes subtiles alternos utrinque circ. 12 et e basi utrinque 1-2 (quorum superior sub medio per anastomoses evanidis inferiorque rarior subtilior) mittente, petiolo glabro ad limbi basin usque vaginante, pedunculo glabro petiolum multoties superante, amento folium dimidium æquante, rhachi dense hirtello-pubescente, bractea juniore saccato-galeata postea obovato-cucullata dorso et margine ciliolata paulo infra florem inserta, ovario basi immersulo apice contracto, stigmatibus linearibus. Baccæ subglobosæ glabræ fusciscentis vertice subcomplanato margine extenuato. ♀ In ins. Cuba ad monte Verde (Wright, n. 495! et Linden, n. 1228! in h. DC.). Ottonia sphærostachya Griseb. pl. Wright p. 176. Rami ramulique juniores puberuli lenticellosi. Limbi 0,065 longi, 0,027 lati. Petioli 0,002 longi. Pedunculi 0,012 longi. Planta sicca odorem myricacearum referens. (v. s.)

β, *tenuifolium*, foliis tenuioribus latioribus apice breviter obtuse protracto-acuminatis nervis lateralibus altius ductis. — In ins. Cuba (Wright n. 2272! in h. DC.).

### § 2. Stigmata 3. Antheræ articulatae.

α. Bractea apice peltata vel truncato-peltata vel truncata.

1° Folia penninervia.

358. *P. WEDDELLII*, foliis brevissime petiolatis elliptico-oblongis apice haud longe attenuato-acuminatis, acumine acuto, basi æquali condatis utrinque

glabris rigidis subopacis subtiliter pellucido-punctulatis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes utrinque 7-8 supremos per anastomoses fere ad apicem ductos infimos subtiliores magis patulos mittente, petiolo glabro, pedunculo glabro petiolum paulo superante, amento quam folium brevior, bracteæ glabræ apice truncato-peltatæ vel subcucullatæ vertice triangulari pedicello cucullato. ♀ In Brasilia (Riedel, n. 1961 in h. DC.). Ramuli glabri. Limbi 0,14 longi, 0,065 lati. Petioli 0,004 longi. Amenta florentia 0,0025 crassa. (v. s.)

359. P. BRACHYPODON, foliis brevissime petiolatis amplis ovatis apice acuminatis basi inæquali obtusis rigidis subcoriaceis utrinque glabris tætu scabris, centrali nervo ad  $\frac{1}{2}$  longitud. nervos subadscendentes subtus valde prominentes utrinque 4-6 supremos ad apicem ductos sursumque nervulos validiores mittente, amento brevi breviter pedunculato, bracteæ apice truncato-peltatæ glabriusculæ vertice semilunari glanduloso. ♀ In ins. Gorgoñe (Benth. in bot. of Sulphur, p. 167. t. 54). Artanthe brachypoda Benth. l. c. Folia 0,25, 0,12 lata. Petioli vix 0,005 longi. (ex Benth. l. c.)

360. P. KEWENSE, foliis petiolatis lanceolato-ovatis apice acuminatis, acumine acuto, basin versus subattenuatis basi ima acutis utrinque glabris rigidulo-membranaceis pellucido-punctulatis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes utrinque 5 supremos ad apicem ductos infimos subtiles haud perspicuos mittente, petiolo glabro ad limbi basin usque vaginante, pedunculo glabro petiolum parum superante, amento post anthesin folium dimidium fere æquante, bracteæ apice truncato-peltatæ pelta triangulari nuda, pedicello rhachique hirtellis, bacca obovato-trigona glabra. ♀ In horto Kew. culta. (h. DC.). Rami ramulique glabri. Limbi 0,13 longi, 0,06 lati. Petioli 0,01 longi. (v. s.)

361. P. KARWINSKIANUM (Kunth in h. Lucæano), foliis breviter petiolatis oblique ellipticis acuminatis, acumine obtusiusculo mucronulatoque, basi inæquali rotundatis membranaceis pellucido-punctulatis opacis utrinque molliter puberulis subtus pallidioribus, nervis subtus prominentibus centrali ad  $\frac{1}{2}$  longitud. nervos utrinque 4-5 mittente, petiolo ad  $\frac{1}{2}$  longitud. vaginante basi dorso sursum utrinque pubescente, amento quam folium  $\frac{1}{2}$  brevior brevissime pedunculato, bracteæ apice peltatæ vertice semilunari margine dense hirsuto, bacca subobovato-trigona. ♀ Prope Mexico (Karwinski! in h. Vindeb.). Schilleria Karwinskiana Kunth! in Linn. v. 13. p. 700. Artanthe Karwinskiana Miq. syst. p. 476. Ramuli pubescenti-villosi. Limbi 0,14 longi, 0,055 lati. Petioli 0,006 longi. (v. s.)

362. P. FRIEDRICHSTHALII, foliis brevissime petiolatis lanceolatis apice acuminatis, acumine acute cuspidato, basi inæquali rotundatis supra glabris subtus adpresse puberulis rigidis opacis epunctulatis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitudinis nervos alternos adscendentes supremos ad apicem ductos mittente, petiolo ad limbi basin vaginante dorso pubescente, pedunculo puberulo petiolum æquante, amento maturo folium subæquante, bracteæ apice peltatæ pelta semilunari margine dense albido-villosa, bacca obovato-trigona, stigmatibus linearibus. ♀ In Guatemala Orosi (Friedrichsthal, n. 1424! in h. Vindeb.). Rami pilis minutis retrorsis puberuli fusci. Ramuli adpresse puberuli. Limbi 0,11 longi, 0,02 lati. Petioli 0,002 longi. Amenta matura 0,003 crassa. (v. s.)

363. P. BRONGNIARTII, foliis modice petiolatis ovato-oblongis acuminatis basi æquali rotundatis supra lævibus nitidis glabris subbullatis subtus in nervis puberulis coriaceis subopacis haud pellucido-punctulatis, centrali nervo subtus prominente utrinque nervos 3 alternos infimos fere e basi ortos juxta margines currentes mittente, amentis brevibus, pedunculis brevibus quam petioli paulo brevioribus, bracteæ pelta subrotunda vel transversim oblonga vel semilunari exigua floccis griseo-luteis vel ochraceis ciliata, pedicello sursum incrassato floccoso-puberulo, stam. 3. ♀ In Peruvia (Dombey, n. 907! in h. Mus. par. fide Miq. syst. p. 486). Artanthe Brongniartii Miq. l. c. Ramulus atro-fuscus glaber hic illic scabriusculus. Limbi 0,21-0,23 longi, 0,15 lati. Petioli 0,02 longi (ex Miq. l. c.)

364. *P. ALPINUM*, foliis breviter petiolatis ovatis vel elliptico-ovatis apice attenuato-acuminatis, acumine obtuso, plus minusve inæquilateris basi breviter inæquali rotundatis utrinque glabris membranaceis pellucido-punctulatis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos subtus prominulos subadscendentes utrinque 3-4 infimos e basi solutos mittente, petiolo glabro juniore ciliolato cito glabrato, pedunculo glabro petiolum superante, amento patulo crasso recto folium superante, bracteæ apice peltatæ pelta suborbiculari vel triangulari flavo-hirtociliata, stigmatibus lanceolatis, stam. 37. ♀ In monte Pichincha alt. 10000 ped. (Hall! in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 451). Artanthe alpina Miq. l. c. Rami glabri. Limbi 0,05-0,07 longi, 0,03-0,035 lati. Petioli 0,005-0,01 longi. Amenta matura 0,03-0,06 longa. (Ex Miq. l. c.)

2° Folia multiplinervia.

365. *P. CONCINNUM*, foliis breviter petiolatis elliptico-lanceolatis apice acuminatis, acumine acuto mucronulatoque, basi inæquali acutis utrinque glabris rigidulis subopacis subtilissime pellucido-punctulatis, nervis subtus vix prominulis centrali ad  $\frac{1}{2}$  longitudinis nervos suboppositos subadscendentes utrinque 2 infimos e basi ortos sursum aliquando nervulos validiores mittente, petiolo glabro basi ima vaginante, pedunculo glabro quam petiolus duplo brevior, amento florente quam folium duplo brevior filiformi, bracteæ apice truncato-peltatæ pelta triangulari glabra pedicello basi dorso hirtello, staminibus 3. ♀ In Brasiliæ umbrosis sylvis (Pohl n. 5032! in h. Vindob.). Artanthe concinna Miq. ! in Linn. v. 20. p. 151. et in Mart. flor. Bras. fasc. 11. p. 43. t. 13. f. 2. Rami ramulique glabri. Limbi 0,09 longi, 0,035 lati. Petioli 0,01 longi. (v. s.)

366. *P. COSTARICENSE*, foliis petiolatis lanceolato-oblongis apice longiuscule et obtusiuscule acuminatis basi æquali acutis utrinque glabris nitidulis rigidulo-membranaceis subpellucidis epunctulatis, nervo centrali ad  $\frac{1}{2}$ - $\frac{2}{3}$  longitud. nervos alternos subadscendentes utrinque 4-5 nervulis validioribus sæpe intermixtos supremos ad apicem ductos infimos e basi solutos mittente, nervulo obscuro a basi ad apicem juxta marginem extenuatum percurrente, petiolo glabro, pedunculo glabro quam petiolus paullo brevior amento maturo quam folium fere duplo brevior, bracteæ apice truncato-peltatæ pelta triangulari subtiliter ciliolata, staminibus 3. (uno postico), bacca obpyramidato-trigona glabra. ♀ In Costa-Rica (Hoffmann, 678! in Berol.). Rami ramulique glabri, juniores subtilissime puberuli. Limbi 0,15 longi, 0,04 lati. Petioli 0,012 longi. Amenta matura vix 0,003 crassa. (v. s.)

367. *P. TRISTEMON*, foliis longiuscule petiolatis late ovatis apice brevissime acuminatis, acumine acuto, basi æquali truncato-rotundatis subcordatisve utrinque glabris subtus fusco-punctulatis, nervis subtus haud prominulis centrali nervos suboppositos utrinque 6 supremos per anastomoses ante apicem evanidos superiores subadscendentes inferiores adscendentim subpatulos infimos e basi solutos mittente, petiolo glabro fere ad  $\frac{1}{2}$  longitud. vaginante, alis linearibus, pedunculo glabro petiolum dimidium æquante, amento cylindrico obtuso folium dimidium superante densifloro, bracteæ apice truncato-peltatæ pelta triangulari ciliata pedicello dorso piloso, staminibus 3, rhachi pilosa, ovario glabro, bacca glabra obovato-subtrigona. ♀ In Venezuela prope coloniam Tovar alt. 7500 ped. (Fendler n. 1141! in h. DC.). Ramuli glabri, fusce punctulati. Limbi 0,19 longi, 0,13 lati. Petioli 0,06 longi. (v. s.)

368. *P. OBTUSUM*, foliis breviter petiolatis oblongo-ellipticis apice obtusis basi inæquali obtusis supra glabris subtus basi ad nervos subtiliter puberulis firmulis subopacis pellucido-punctulatis, centrali nervo subtus prominente nervos oppositos subadscendentes utrinque 2 supremos ad medium per anastomoses evanidos infimos e basi solutos sursum tota longitudine nervulos validiores subadscendentim-subpatulos mittente, petiolo ad limbi basin vaginante dorso dense puberulo, pedunculo puberulo petiolum æquante, amento folium subæquante, bracteæ apice truncato-peltatæ pelta semilunari margine hirsuta, ovario glabro. ♀ In ins. Cuba (Wright, n. 2271! in h. DC.). Artanthe geni-



culata Grisebach cat. pl. cub. p. 65. Ramuli glabri. Limbi 0,11 longi, 0,057 lati. Petioli 0,008 longi. Amenta post anthesin fere 0,003 crassa. (v. s.)

369. *P. FOVEOLATUM* (Kunth! olim), foliis breviter petiolatis suboblique oblongo-ovatis apice acuminatis, acumine acuto, basi subæquali rotundatis vel levisime cordulatis utrinque glabris membranaceis opacis pellucido-punctulatis, nervis subtus prominulis centrali supra basin trifido nervum e basi utrinque unum subtiliorem mittente, nervulo obscuro juxta marginem currente, petiolo fere ad limbi basin vaginante dorso subtilissime puberulo glabratove, pedunculo petiolum æquante, amento maturo folium triplo superante, bractæ apice peltatæ pelta subrotunda brevissime pedicellata margine dense hirsuta, rhachis foveolis hirtellis, baccis subremotis obovato-subglobosis basi immersis. ♀ In Guyana gallica (Poiteau! in h. Berol.), Surinam (Kappeler n. 1482! in h. Boiss.), ad ripas flumin. Saramacca (Kegel n. 1364 fide Miq. in Linn. v. 22. p. 76). Enckea dubia Kunth! in Linn. v. 13. p. 606, Miq. in Linn. v. 22. p. 76. Ramuli glabri. Limbi 0,09 longi, 0,045 lati. Petioli 0,012 longi. Amenta matura 0,003 crassa. (v. s.)

370. *P. BLATTARUM* (Spreng. in Neue Entdeck. v. 2. p. 100), foliis breviter petiolatis ellipticis apice acuminatis, acumine obtuso, basi æquali acutis utrinque glabris membranaceis subpellucidis subtiliter pellucido-punctulatis quintuplinerviis, nervis subtus prominulis 3 centralibus supra basin vel interdum fere e basi solutis fere ad apicem usque ductis lateralibus e basi vel paullo supra basin solutis sub medium ductis, petiolo dorso subtilissime hirtello, pedunculo petiolum triplo superante glabro, amento filiformi sublaxifloro folium superante, rhachi hirtella, bractæ apice truncato-peltatæ dorso pubescentis pelta triangulari supra pubescente, pedicello subcucullato, stigmatibus abbreviatis, bacca subglobosa. ♀ In insula Portorico (Wydler n. 407! in h. Berol. et Bertero! in h. DC.), St-Thomas (Wydler n. 407! in h. DC.). *Peperomia blattarum* A. Dietr. sp. v. 1. p. 150. Enckea blattarum Kunth! in Linn. v. 13. p. 600, Miq.! syst. p. 364. Rami glabri. Limbi 0,065-0,09 longi, 0,025-0,035 lati. Petioli 0,003 longi. (v. s.)

♂, *magnifolium*, foliis majoribus firmioribus, limbis 0,11 longis fere 0,05 latis. — In ins. Portorico (Bertero! in h. Berol.). (v. s.)

### β. Bractea subcucullato-obovata. Flores hermaphroditi.

371. *P. CARPUNYA* (Ruiz et Pav. flor. Peruv. 1. p. 37. t. 63. f. b), foliis breviter petiolatis ovato-ellipticis apice modice acuminatis, acumine acuto mucronulatoque, basi subæquali cordatis utrinque glabris supra nitidis coriaceis subtilissime pellucido-punctulatis, centrali nervo subtus prominulo ad  $\frac{3}{4}$  longitud. nervos subalternos adscendentes utrinque 4-5 supremos ad apicem ductos sursumque nervulos validiores mittente, petiolo glabro, pedunculo glabro petiolum æquante, amento densifloro folium subæquante, bractæ subcucullato-obovatæ aureæ intus hirsutæ vertice inflexo, bacca subgloboso-trigona glabra, stigmatibus abbreviatis. ♀ In Peruviz nemoribus Chinchao et Cuchero (R. et P. l. c. et in h. Willd. n. 652! et in h. Berol.), Peruviz prov. Chachapoyas (Mathews n. 1396!, 3222!, 3226! in h. Kew.), montibus Huanuco (Hænke! in h. Berol.), Chili (Poeppig n. 59 vel 1354! in h. Berol. et in h. DC.). — Vahl enum. v. 1. p. 324; Opiz in Reliq. Hænk. fasc. 3. p. 155. *Otonia Carpunya* Miq. l. syst. p. 547, in Linn. v. 20. p. 180, in Mart. flor. Bras. fasc. 11. p. 69, et in Hook. Lond. journ. bot. v. 4. p. 470. *Otonia Lessertiana* Miq. l. syst. p. 546 (ubi sub numero 1696). *Carpunya peruviana* Presl Epim. p. 229. Carp. Lessertiana Presl l. c. Ramuli glabri læves, folia ramulique subaurea. Limbi 0,09-0,15 longi, 0,05-0,075 lati. Petioli fere 0,01 longi. Amenta matura 0,002 crassa. (v. s.)

### § 3. Stigmata 3. Antheræ non articulatz.

372. *P. ABUTILOIDES* (Kunth! olim), foliis longe petiolatis rotundato-ovatis apice breviter acuminatis, acumine acuto mucronulatoque, basi profunde cordatis, lobis conniventibus, supra glabris subtus ad nervos puberulis firmo-membra-

naceis undecinnerviis, nervis subtus prominulis centrali sursum alternatim trifido ad apicem ducto, petiolo glabro ad  $\frac{1}{2}$  longitud. usque vaginante alis linearibus, pedunculo glabro quam petiolum duplo brevior, amento quam folium duplo brevior cylindrico, bractæ spathulatæ intus et extus dorso hirtellæ vertice inflexo pelam fingente, staminibus 3, filamentis antheras ter quaterque superantibus, stigmatibus filiformibus. ♀ In Brasilia meridionali (Sello! in h. Berol.). Schilleria abutiloides Kunth! in Linn. v. 13. p. 720. Artanthe abutiloides Miq. l. syst. p. 384, in Mart. flor. Bras. fasc. 11. p. 33. Ramuli glabri. Limbi 0,14-0,23 longi, 0,1-0,17 lati. Petioli 0,04-0,1 longi. Amenta florentia 0,004 crassa. (v. s.)

373. *P. PARTHENIUM* (Mart. l. syst. mat. med. brasil. p. 100), foliis longe petiolatis e basi cordata vel subrotunda ovatis apice breviter acuminatis, acumine acutiusculo, supra glabris subtus ad nervos adpresse hirtellis pellucido-punctulatis membranaceis novennerviis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. trifido sursumque nervulos validiores mittente proximis fere e basi solutis infimis sæpe obsolete, petiolo ad  $\frac{1}{2}$  longitud. usque vaginante dorso hirtello alis linearibus, pedunculo puberulo quam petiolum brevior, amento cylindrico quam folium brevior, bractæ apice truncato-peltatæ vertice subsemilunari margine hirtello, pedicello margine et basi extus pubescente, filamentis exsertis antheras multum superantibus, bacca obovato-trigona glabra. ♀ In Brasilia meridionali (Sello! in h. Berol. et Boiss.), Rio Grande do Sul (Mart. in h. Mart. et monac.). — Miq. syst. p. 550. Schilleria mikaniæ Kunth! in Linn. v. 13. p. 719. Artanthe mikaniæ Miq. syst. p. 383, in Linn. v. 20. p. 137 et in Mart. flor. Bras. fasc. 11. p. 34. Ramuli adpresse hirtelli. Limbi 0,08-0,1 longi, 0,07-0,09 lati. Petioli 0,03-0,04 longi. Amenta 0,004 crassa. (v. s.)

374. *P. CALADIIFOLIUM*, foliis longe petiolatis ovato-cordatis breviter acuminatis basi vix inæqualibus subcoriaceis tenuissime pellucido-punctulatis, centrali nervo subtus prominente nervos e basi utrinque 3-5 paulo supra basin unam et porro 3-4 nervulos validiores mittente, amento plerumque mucronato, bractæ cucullato-peltatæ pedicello fleccoso-hirsuto apice concavo rotundato membranaceo fimbriolato, stam. 3 an 4? ♀ In Mexico (Kickx fide Miq. syst. p. 387). Artanthe caladiifolia Miq. l. c. Ramuli glabri. Limbi 0,13-0,21 longi, 0,1-0,14 lati (ex Miq. l. c.)

375. *P. PLANTAGINEUM* (Schlecht. non Lam.), foliis petiolatis ovatis apice acuminatis, acumine acuto, basi rotundatis cordulatisve utrinque glabris firmulo-membranaceis septupli-novennerviis, nervis subtus prominulis 3 centralibus fere ad apicem ductis paulo supra basin solutis, petiolo glabro fere ad  $\frac{1}{2}$  longitudinis usque vaginante alis linearibus, pedunculo glabro petiolum æquante, amento cylindrico folium æquante, bractæ cucullato-peltatæ pelta semilunari margine hirtella pedicello ciliolato, filamentis antheras superantibus, stigmatibus carnosulis recurvulis, bacca obovato-tetragona glandulosa. ♀ In Mexico (Schiede n. 1605! in h. Berol.). *P. plantagineum* Schlecht. in Linn. v. 6. p. 353. Steffensia plantaginea Kunth! in Linn. v. 13. p. 672. Artanthe plantaginea Miq. l. syst. p. 389. *P. commutatum* Steud. nom. Ramuli glabri. Limbi 0,09 longi, 0,05 lati. Petioli 0,02 longi. Amenta matura 0,003 longa. Bacca 0,001 longa. (v. s.)

376. *P. SANCTUM* (Schlecht. mss. in h. Schlecht. fide Miq. in Linn. v. 18. p. 714), foliis tenuiter membranaceis pellucido-punctulatis utrinque præsertim subtus ad nervos et secus margines puberulis rotundato-cordatis breviter acuteque acuminatis lobis baseos æqualibus rotundatis sinu dilatato diremptis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos utrinque 5 supremos ad apicem ductos infimos e basi solutos mittente. ♀ Ad Atlacomulco Mexici (Schiede n. 105 in h. Schlecht. fide Miq. l. c.). Artanthe? sancta Miq. l. c. Limbi 0,21 longi, 0,18 lati. An eadem ac Art. plantaginea? (Miq. l. c.). (ex Miq. l. c.)

377. *P. ZACUAPANUM*, foliis longe petiolatis ovatis apice acuminatis, acumine acuto, basi rotundato-cordatis utrinque glabris membranaceis subtiliter pellucido-punctulatis undennerviis, nervis subtus prominulis 3 mediis supra basin solutis centrali ad apicem ducto nervulos validiores fere tota longitudine alternatim mittente, petiolo glabro, pedunculo glabro quam pe-

tiotus multum brevior, amento cylindrico quam folium brevior, bractæa cucullato-peltatæ pelta triangulari-subrotunda margine subtusque subtiliter ciliolata, filamentis antheras ter quaterque superantibus, stigmatibus linearibus. ♀ In Mexico (Schiede n. 1102! in h. Berol.), sylvis ad Zacuapan alt. 2500-3000 ped. (Galeotti n. 6017 hde Mart. et Gal. in bull. ac. Brux. v. 10. p. 2. p. 23). P. tiliaefolium Schlecht. et Cham. in Linn. v. 6. p. 352, non Ham. Ramuli glabri, limbi 0,14 longi, 0,12 lati. Petioli 0,06 longi. Amenta florentia 0,002 crassa. (v. s.)

Sect. V. POTOMORPHE. — Genera : *Heckeria* Kunth in Linn. v. 18. p. 564; *Macropiper* Miq. in comm. phyt. p. 35; *Potomorphe* Miq. in comm. phyt. p. 33.

Stamina 2 vel 3 quorum unum posticum, antheræ articulatæ. Flores hermaphroditi vel unisexuales. Amenta axillaria vel apice ramuli axillaris umbellata.

### A. Flores hermaphroditi. Baccæ obovato-trigonæ.

#### 1° Folia peltata.

378. P. PELTATUM (L! sp. p. 42 et herb.), foliis longe petiolatis ad  $\frac{1}{4}$  longitudinis peltatis rotundato-ovatis apice protracto-acutis basi rotundatis concavo-truncatis vel breviter cordatis utrinque glabris firme membranaceis vel membranaceis subopacis pellucido-punctatis, nervis subtus prominulis, centrali ad apicem ducto nervos alternos vel oppositos adscendentes utrinque supra basin 2 et e basi utrinque 6-5 mittente, nervulis subtransversim parallelis, petiolo ad  $\frac{1}{4}$  longitud. usque vaginante glabro alis linearibus, amentis pedunculo communi axillari quam petiolus multoties brevior glabro 7-10 umbellatis densifloris quam folia fere triplo brevioribus breviter pedunculatis, bractæa apice peltatæ pelta triangulari-semilunari margine hirsuta, bacca glandulosa. ♀ In Peruvia (Poeppig! in h. Petrop.), vicinibus Parâ (Spruce! in h. Kew.; Mart. in h. monac.), in umbris humidis regni Novogranatensis in ripa fluminis Magdalena inter Mompox et Minchiqueo (Humb. et B. nov. gen. v. 1. p. 59 et herb. Willd. n. 704! et 703! et 705!), Nova Granata (Holton n. 233! in h. Kew.), prov. Barbacoas (Triana n. 1! in h. suo), Caracas (Vargas n. 18! in h. DC., Moritz n. 1939! in h. brit. Mus.), Guyana (Schomburgk n. 954! in h. Kew. et Berol., Hostmann n. 42! in h. Kew.), Guyana gallica (Aublet hist. pl. Guy. v. 1. p. 21), in hortis Surinami ubique culta non spontanea (Miq. in Nederl. Kruidk. v. 1. p. 221), Paramaribo (Kegel n. 832 hde Miq. in Linn. v. 22. p. 76), Panama (Sutton et Hayes n. 745!, Fendler n. 267! in h. Kew. 264! in h. Petrop. et 1146! in h. DC.), ins. Trinitatis (h. Kew.), Barbados (Maycock pl. Barb. p. 40), Martinica (Jardin n. 344! in h. Lenormand, Hahn n. 255! in h. brit. Mus.), St Vincent (GUILDING! in h. Kew.), Portorico (Riedley! in h. Deless.), St Thomas (Schreiber! in h. monac.), S<sup>te</sup> Domingo (Mayerhoff n. 339! in h. Berol.), Cuba (Wright n. 496! in h. DC.), Jamaica (Swartz obs. p. 21). Saururus arborescens foliis amplis rotundis Plum. Am. p. 56. t. 74. P. peltatum Hill dict. v. 26. p. 19. t. 19. f. 4, Aublet l. c., Lam. ill. p. 81, Swartz obs. p. 21, Willd. sp. v. 1. p. 166, Vahl enum. v. 1. p. 336, Mart. syst. mat. med. p. 99. P. pruinosum Willd. l. h. n. 705, Kunth! in H. et B. nov. gen. v. 1. p. 59. P. speciosum Willd. h. n. 704, Kunth! l. c. P. scutatum Willd. l. h. n. 703. Heckeria peltata Kunth! in Linn. v. 13. p. 565. Heck. scutata Kunth! l. c. Heck. speciosa Kunth! l. c. Peperomia pruinosa Kunth synops. v. 1. p. 124, A. Dietr. l. c. Pep. speciosa Kunth l. c., Opiz in Reliq. Hænk. fasc. 3. p. 161. Pep. peltata A. Dietr. sp. v. f. p. 142. Pep. scutata A. Dietr. l. c. Pep. speciosa A. Dietr. l. c. Potomorphe peltata Miq. comm. phyt. p. 45. t. 4. f. e, syst. p. 203, in Linn. v. 20. p. 129, in Mart. flor. Bras. fasc. 11. p. 26; Grisebach flor. w. Ind. p. 169 et in Cat. pl. Cub. p. 64. Pot. scutata Miq. comm. phyt. p. 37, syst. p. 206 et in Mart. flor. Bras. l. c. p. 27. Pot. speciosa Miq. comm. phyt. p. 34, syst. p. 205. Poivrier en bouclier (Descourtils flor. Ant. v. 4. p. 14. t. 236). Caulis glabri. Limbi 0,2-0,32 longi latique. Petioli 0,2 longi. Pedunculus communis 0,04 longus. Amenta florentia 0,11-0,12 longa, 0,004 crassa. (v. s.)

β, *Colombianum*, venis foliorum supra hirtis. In Colombia (Benth. bot. of Sulphur p 167 (ex Benth. l. c.)

379. *P. OTTONIS*, foliis longissime petiolatis ad  $\frac{1}{2}$  -  $\frac{1}{3}$  longitud. peltatis rotundato-ovatis apice acuminatis, acumine acuto, basi rotundatis concavo-truncatis cordatisve utrinque glabris membranaceis pellucido-punctatis, centrali nervo ad  $\frac{1}{2}$  longitud. supra basin nervum utrinque unum ascendentem subopposita et a basi utrinque 4 quorum medii sæpe paulo supra basin soluti mittente, nervulis subtransverse parallelis, petiolo glabro ad  $\frac{1}{2}$  longitud. vaginante alis anguste linearibus, amentis pedunculo communi axillari quam petiolus multoties brevior 5-umbellatis breviter pedunculatis densifloris quam folia brevioribus, bractæ apice peltatæ pelta semilunari margine lutescenti-hirsuta. ♀ In Bahía Hongua Veraguas (Seemann n. 899 fide Miq. in bot. Herald p. 198), Panama ad Maone et Gorgone (Wagner! in h. monac.). Venezuela (Otto! in h. Berol.). *Potomorphe Ottonis* Miq.! syst. p. 207, illustr. p. 27. t. 25. Limbi 0,11-0,14 longi, 0,08-0,11 lati. Petioli 0,1-0,15 longi. Pedunculus communis 0,035-0,04 longi. (v. s.)

380. *P. SCUTIPHYLLUM* (Ham. prodr. Ind. occ. p. 3), foliis peltatis cordato-ovatis acuminatis integerrimis basi obliquis multiplinerviis subtredecimnerviis, nervis radiantibus suboctopolicaribus longis subseptempolicaribus latis. In loco udo ad prædium Dezobré prope oppidum capitis Henrici (Hayti) (ex Ham. l. c.). *P. peltatum* L.?

## 2° Folia non peltata.

381. *P. DOMBEYANUM*, foliis longe petiolatis rotundato-reniformibus basi profunde cordatis apice subattenuatis acutiusculis supra subtusque hirtis - puberulis margine ciliolatis membranaceis subpellucidis, nervis subtus prominulis centrali ad apicem ducto paulo supra basin trifido nervos e basi utrinque 8 divaricantes mittente, nervulis subtransversim parallelis, petiolo ad  $\frac{1}{2}$  longitud. usque vaginante dorso hirtis-pubescente, amentis pedunculo communi quam petiolus multoties brevior hirtis puberulo 5-7-umbellatis densifloris quam folia quater brevioribus breviter pedunculatis, bractæ apice peltatæ pelta triangulari margine hirsuta. ♀ In Peruvia (Poeppig n. 57! in h. DC., Mathews n. 170! in h. Kew.), Peruviae provinc. Panatuhuarum nemoribus Cuchero et ad Maichaynio prædium (R. et P. flor. peruv. v. 1. p. 38. t. 59. f. a et Pav. ! in h. Boiss.). *P. peltatum* R. et P. ! l. c. (non L.). *P. umbellatum* Vahl enum. v. 1. p. 337 quoad synonym. flor. peruv. *Potomorphe Dombeyana* Miq.! syst. p. 211, illustr. p. 29. t. 26, in Hook. Lond. journ. bot. v. 4. p. 431, in Linn. v. 20. p. 129. Caules fusco-pubescentes. Limbi 0,21 longi, 0,205 lati. Petioli 0,06-0,1 longi. (v. s.)

382. *P. CORDILLERIANUM*, foliis longissime petiolatis reniformi-orbicularibus apice brevissime acuminatis, acumine acutiusculo, basi profunde cordatis, lobis discretis, supra puberulis subtus pubescentibus et ad nervos ochraceo-tomentosis membranaceis subpellucidis pellucido-punctulatis, nervis subtus prominulis centrali ad apicem ducto ad  $\frac{1}{2}$  longitud. supra basin opposita trifido e basi nervos utrinque 6 divaricantes mittente, petiolo ad  $\frac{1}{2}$  longitud. usque vaginante dorso pilis recurvulis hirtis, amentis pedunculo communi quam petiolus multoties brevior pilis recurvulis puberulo 5-umbellatis, bractæ apice peltatæ pelta triangulari margine hirsuta. ♀ In Cordilleris Mexici locis humidis alt. 3500 ped. prope Vera Cruz (Galeotti n. 6006! in h. Deless.). *Potomorphe Lessertiana* Miq.! syst. p. 210, illustr. p. 28. t. 26. Ramuli pilis recurvulis puberuli. Limbi 0,21 longi, 0,3 lati. Petioli 0,25 longi. Pedunculi communes in eadem axilla binati. (v. s.)

383. *P. UMBELLATUM* (L. sp. p. 43), foliis longe petiolatis rotundato-reniformibus apice breviter attenuatis acutis basi profunde cordatis supra subtusque ad nervos breviter hirtellis membranaceis subpellucidis subtiliter crebre pellucido-glandulosis, nervis subtus prominulis centrali ad apicem ducto usque  $\frac{1}{2}$  -  $\frac{1}{3}$  longitud. supra basin nervos alternos vel suboppositos adscendentes utrinque 2 (quorum inferi fere e basi soluti) et ex ipsa basi nervos utrinque 5-6 divaricantes mittente, nervulis subtransverse parallelis, petiolo circiter ad

$\frac{1}{2}$  longitud. usque vaginante dorso striatim hirtello, amentis pedunculo communi quam petiolus multoties brevior striatim hirtello 4-7-umbellatis breviter pedunculatis quam folia multoties brevioribus, bractæ pelta triangulari-semilunari margine hirsuta, stigmatibus 3 linearibus.  $\frac{3}{4}$  In Brasilia meridionali (Guillemin in h. Deless.), Ilheos (Wawra et Maly n. 199 fide Wawra bot. Ergeb. p. 138), Caracas (Vargas n. 279! in h. DC.), locis calidis inter Cumana et Chara de los Capuchinos (Humb. et B. nov. gen. v. 1. p. 59, et h. Willd. n. 700! fol. 3), locis humidis et ad rupes Colonizæ Mirador alt. 8000 ped. (Galeotti n. 6008 fide Mart. et Gal. in bull. ac. Brux. v. 10. n. 2. p. 23), Xalapa (Galeotti! in h. Kew.), Panama (Sutton et Hayes! in h. Brit. Mus.), in Antillis (Descourtils flor. Ant. v. 1. p. 177. t. 37), Barbadoes (Maycock pl. Barbado. p. 40), Portorico (Bertero! in h. DC.), S<sup>to</sup> Domingo (Poiteau! in h. DC.), Cuba (Wright n. 496! in h. monac.), Jamaica (Swartz! et Murray n. 28! in h. DC.). Aguaxima Piso hist. nat. med. ed. 1658, p. 197 cum ic. P. racemosum malvaceum Sloane! hist. p. 136 et h. n. 86! et 87! Saururus arborescens foliis amplis cordatis non umbilicatis Plum. am. ed. Par. p. 53. t. 73. P. umbellatum Hill dict. v. 26. p. 20. t. 20. f. 2, Aublet l. c., Jacq. ic. rar. t. 216, Lam. ill. p. 81, Vahl enum. v. 1. p. 336 excl. syn. Floræ peruv., Swartz obs. p. 21, Willd. sp. v. 1. p. 167, Kunth! in H. et B. l. c., Opiz in Reliq. Hænk. fasc. 3. p. 161, Vell. flor. flum. t. 54. P. sidæfolium Mart. in syst. mat. med. p. 99. Peperomia umbellata Kunth synops. v. 1. p. 124; A. Dietr. sp. v. 1. p. 142. Heckeria umbellata Kunth! in Linn. v. 13. p. 569. Potomorphe umbellata Miq.! comm. phyt. p. 36, syst. p. 208, in Linn. v. 20. p. 129, in Mart. flor. Bras. fasc. 11. p. 26 (exclusis spec. ad seq. sp. referendis), Grisebach! flor. w. Ind. p. 169 et Cat. pl. Cub. p. 64. Poivrier à ombelles (Descourtils l. c.). Ramuli striatim hirtelli, fusci. Limbi 0,17-0,18 longi, 0,22-0,23 lati. Petioli 0,11 longi. Pedunculi communes (prophyllis fertili), in eadem axilla sæpe binati. (v. s.)

$\beta$ , *majus*, foliis majoribus apice magis attenuatis, petiolis longioribus. — In Venezuela prope Biscaina (Fendler n. 1147! in h. DC. et Moritz n. 283! in h. Brit. Mus.). (v. s.)

$\gamma$ , *glabrum*, petiolis et ramulis glabris pedunculis monostachyis. — In Bahía (Salzmänn n. 1570! in h. Kew.). (v. s.)

384. P. SUBPELTATUM (Willd.! sp. v. 1. p. 166). foliis longe petiolatis rotundato-reniformibus apice brevissime attenuatis acutis basi profunde cordatis supra subtusque ad nervos breviter hirtellis membranaceis suppellucidis subtiliter crebre pellucido-glandulosis, nervis subtus prominulis centrali ad apicem ducto usque ad  $\frac{1}{2}$  longitud. nervos alternos vel suboppositos utrinque 1-2 et e basi utrinque 4-5 divaricantes mittente, nervulis subtransversim parallelis, petiolo fere ad  $\frac{1}{2}$  longitud. usque vaginante basi dorso sursum utrinque veluti-puberulo, amentis pedunculo communi quam petiolus multoties brevior subpresse puberulo 5-7-umbellatis breviter pedunculatis densifloris quam folia multoties brevioribus, bractæ apice peltatæ pelta triangulari-semilunari margine hirsuta, bacca vertice glanduloso-pubescente.  $\frac{3}{4}$  In insulis Java et Celebes (Miq. in Linn. v. 21. p. 481). Sumatra (Miq. in Sumatra p. 186), Philippinis (Cuming n. 441! in h. Kew.), Borneo prope Bangarmassing (Motley n. 605! in h. Kew.), Java (Zollinger n. 840! in h. DC.), commune in locis humidis umbrosis ins. Ceylon usque ad 4000 ped. (Thwaites enum. p. 292. et n. 2714! in h. DC.), prope Penang (Wall. n. 6638a! in h. Wall.), Indiæ montib. Nilghericis (Hook. et Thoms. l. in h. DC.), montib. Travancorensibus (Wall. n. 6638d! in h. Wall.), insul. Maurice (Bouton! in h. DC.), Sieber n. 166! in h. DC., Bory in h. Willd. n. 782!), Bourbon (h. DC.!), Madagascar (Roxburgh! in h. Brit. Mus.), Mayotte (Boivin! in h. Boiss.), Africæ ora orientali prope Montgomery (Martin! in h. DC.), Vahnuna alt. 500 ped. (Kirk! expd. Zambezi in h. Kew.), montib. Cameroon alt. 3000 ped. (Mann n. 1272! in h. Kew.), Fernando Po (Mann n. 70! in h. Kew.), Angola (Welwitsch n. 501! in h. DC.), Gabon (Jardin n. 146! in h. Lenormand). Lamba Rumph. amb. v. 6. p. 133. t. 59. f. 1. P. subpeltatum Vahl enum. v. 1. p. 337 exclus. syn. Lam. P. grandifolium Eckelund apud Afzel. pl. Guin. v. 6. p. 39. (fide Roem. et Schult. mant. v. 1. p. 235). P. Afzelianum Roem. et Schult. l. c. Peperomia grandifolia A. Dietr. sp. v. 1.

p. 166. *Heckeria subpeltata* Kunth! in Linn. v. 13. p. 571. *Potomorphe subpeltata* Miq. ! syst. p. 213, illust. p. 29. t. 26, in Linn. v. 20. p. 129, pl. Jungh. fasc. 1. p. 16, flor. ind. bat. v. 1. pars 2. p. 437, *Sumatra l. c.*, in Linn. v. 21. p. 481, Wight icon. t. 1925. *Ramuli glabri glandulosi*. Limbi 0,16-0,18 longi, 0,22-0,32 lati. Petioli 0,1-0,16 longi. Pedunculus communis circ. 0,02 longus. Pedunculi communes (prophyllis fertili), in eadem axilla sæpe binati. (v. s.)

β, *sidæfolium*, ramulis subtiliter adpresse striatum puberulis. ♂ In Brasilia (Riedel n. 283 ! et 322 ! in h. Petrop. Mart. n. 1207 ! in h. DC., Sello ! in h. Berol.), prope Rio Janeiro (Guillemin n. 3 ! et Gaudich. n. 1103 ! in h. DC.), Bahia (Blanchet n. 221 ! et 524 ! in h. DC.), in udis ad rivulos umbrosos prope Mandiocca prædium et prope Salgado (Mart. in h. monac.), umbrosis humidis ad Besso de Serra et in Via ab Angios dos Rego ad Margaritiva (Pohl in h. Vindob. fide Miq. in Linn. v. 20. p. 129), Colombia (Miq. in Linn. v. 18. p. 713), Caracas (Moritz n. 283 ! in h. Berol., h. Willd. n. 700. f. 1 ! et 2 !) *P. umbellatum* Jacq. ic. rar. v. 2. p. 2. t. 216. *P. latifolium* Lam. ill. p. 81. *Peperomia sidæfolia* A. Dietr. spec. v. 1. p. 141, exclus. syn. Roem. et Schult. *P. sidæfolium* Link et Otto ic. p. 60. t. 6. *Heckeria sidæfolia* Kunth ! in Linn. v. 13. p. 569. *Potomorphe sidæfolia* Miq. ! syst. p. 209, in Linn. v. 20. p. 129, in Mart. flor. Bras. fasc. 11. p. 25 et in Hook. Lond. journ. bot. v. 4. p. 431. Pedunculi communes sæpe bini. (v. s.)

385. *P. PEDUNCULATUM* (Cas. DC. in Seem. journ. 1866. p. 162), foliis longiuscule petiolatis late ellipticis apice subattenuatis basi æquali subattenuatis acutiusculis supra glabris subtus ad nervos puberulis rigidulis opacis epunctulatis, centrali nervo ad apicem ducto subtus prominente ad  $\frac{1}{2}$  longitud. usque nervos alternos subadscendentes subtus prominulos utrinque 6 supremos fere ad apicem ductos mittente, petiolo glabro ad limbi basin usque vaginante alis linearibus sursum in petiolum attenuatis, pedunculo adpresse puberulo petiolum fere duplo superante, amento mucronato quam folium multum brevius axillari solitario, bractæ apice peltatæ pelta triangulari margine hirtella, stam. ...., bacca obovata-trigona glabra, stigmat. 3. ♂ In Nova Granata (Triana! in h. DC.). Ramuli glabri. Limbi 0,135 longi, 0,075 lati. Petioli 0,03 longi. Amenta matura 0,004 crassa. (v. s.)

386. *P. TRIANÆ* (Cas. DC. in Seem. journ. 1866. p. 162), foliis longe petiolatis subrotundato-subovatis apice breviter acuminatis, acumine acuto, basi subinæquali cordatis utrinque glabris membranaceis subtiliter pellucido-punctulatis glandulis fuscis conspersis, centrali nervo ad apicem ducto subtus prominulo ad  $\frac{1}{2}$  longitud. usque nervos alternos subadscendentes subtus prominulos utrinque 7-8 supremos ad apicem ductos mittente, petiolo glabro ad  $\frac{1}{2}$  longitud. usque vaginante alis linearibus apice liberis, amento solitario apice ramuli axillaris petiolum dimidium æquantis filiformis quam folium multoties brevius, pedunculo quam petiolus multoties brevius glabro vel subtilissime puberulo, bractæ apice truncato-estatae carnosæ glabræ pedicello suboculato, ovario apice in stylum longum attenuato, stigmatibus brevibus recurvulis, bacca ovata glandulis fuscis conspersa, rhachi glabra. ♂ In Nova Granatæ prov. Pasto alt. 2500 ped. (Triana n. 2 ! in h. DC.). Ramuli glabri. Limbi 0,16 longi, 0,12 lati. Petioli 0,065 longi. Amenta matura 0,02 longa, 0,006 crassa. (v. s.)

#### B. Flores unisexuales.

387. *P. EXCELSUM* (Forster! prodr. n. 20), foliis petiolatis rotundato-ovatis apice breviter attenuatis acutiusculis basi obtuse truncatis vel cordatis utrinque glabris membranaceis subopacis pellucido-punctulatis septemnerviis, centrali nervo ad apicem ducto lateralibus 2 fere ad apicem 2 supra medium ductis, petiolo glabro ad  $\frac{1}{2}$  longitud. usque vaginante alis linearibus sursum in petiolum attenuatis, amentis pedunculo communi axillari quam petiolus multum breviori plerumque geminatis, maris bractea rotundata centro peltata subsessili glabra, stam. 2 lateralibus vel interdum 3 quorum unum posticum antheris subsessilibus, femine bractea ut in mare, bacca obovato-trigona glabra mu-

crenulata, stigmatibus 3.  $\frac{5}{8}$  In Nova Zelandia (Sinclair ! Lyall ! in h. Kew., Forster ! et Anderson ! in h. Brit. Mus. et DC. Hügel ! in h. Vindob.; Sidney et Park. coll. orig. draw. n. 42. in h. Brit. Mus.; Exped. Astrol. in h. DC.), ins. Norfolk (Cunningham ! in h. Kew.), ins. Lord Blow (Gillivray n. 726 ! et 970 ! in h. Kew.), ins. Pinorum oceani Pacifici (h. Berol. !). — Vahl enum. v. 1. p. 335, Hook. f. flor. Nov. Zeel. v. 1. p. 228. Macropiper excelsum Miq. syst. p. 221, in Hook. Lond. Journ. bot. v. 4. p. 432, in Pip. nov. Nov. Holland. adnot. 9. in Nederl. der konink. Akad. van Vetensch. Afdel. Naturk. 2. Beeks deel. 2. Dioicum vel raro monoicum et tum amenta unisexualia. Ramuli glabri. Limbi 0,09-0,012 longi, 0,07-0,01 lati. Petioli 0,03 longi. Pedunculi 0,004 longi. Amenta florentia 0,04 longa, 0,002 crassa. (v. s.)

$\beta$ , *psittacorum*, foliis majoribus 0,12 longis.  $\frac{5}{8}$  In ins. Norfolk (Cunningham ! Hügel ! Endlich. ! in h. Vindob.). Macropiper psittacorum Endlich. ! prodr. Norfolk. p. 37, Bauer pl. Norf. t. 37, fide Endlich. l. c. (v. s.)

$\gamma$ , *Tahitianum*, foliis apice breviter protracto-acuminatis 9-12-nerviis, petiolo fere ad  $\frac{3}{4}$  longitud. vaginante, amentis maturis pulposis.  $\frac{5}{8}$  In ins. Tahiti (Barclay ! in h. Kew.), ins. Lifuka (Harvey ! in h. Kew.), ins. Corallensis (Beechey ! in h. Kew.), Norfolk (Milne ! in h. Kew.). Limbi 0,07-0,1 longi, 0,05-0,012 lati. Amenta florentia 0,09 longa. (v. s.)

$\delta$ , *Ralphii*, amentis maturis duris, baccis coalitis.  $\frac{5}{8}$  In Nova Zelandia (Ralph n. 51 ! in h. Boiss.). Limbi 0,06 longi, 0,05 lati. Amenta florentia 0,04 longa. (v. s.)

388. P. MACGILLIVRAYI (Cas. DC. in Seem. fl. Vit. p. 262. t. 75), foliis ovato-acuminatis, acumine acuto, basi rotundato-truncatis vel superioribus cordatis supra glabris subtus subpuberulis membranaceis 7-9-nerviis, nervis subtus prominulis 3 centralibus fere ad apicem ductis, petiolo ad  $\frac{1}{2}$  longitud. usque vaginante, amentis femineis solitariis axillaribus maturis pulposis folia æquantibus, ovario oblongo, stigmatibus 3, maris bractea ut in femina, stam. 3, antheris subsessilibus.  $\frac{5}{8}$  In insulis Fijee (Barclay n. 3448 ! in h. Brit. Mus., MacGillivray n. 62 ! in h. Kew., Milne n. 37 ! in h. Kew.), Tongataboo (h. Kew. !). Macropiper puberulum Benth. ! in Hook. Lond. Journ. bot. v. 2. p. 235, Miq. ! l. c. v. 4. p. 431. Dioicum. Limbi 0,1-0,12 longi, 0,06-0,08 lati. Nomen puberulum propter var.  $\gamma$  mutandum. (v. s.)

$\beta$ , *parvifolium*, limbis 0,06 longis, 0,03 latis 5-7-nerviis.  $\frac{5}{8}$  In ins. Fijee (Barclay ! in h. Kew.)

$\gamma$ , *glabrum*, foliis utrinque glabris.  $\frac{5}{8}$  In ins. Fijee (Seemann n. 567 ! in h. DC.). Macrop. puberulum Seem. in Bonpl. 1861. p. 259.

389. P. LATIFOLIUM (L. fil. suppl. p. 91. correct. p. 468), foliis longe petiolatis rotundatis apice brevissime protracto-obtusis basi rotundato-truncatis vel leviter cordatis utrinque glabris membranaceo-rigidulis subopacis subtilissime pellucido-punctulatis 11-12-nerviis, petiolo glabro fere ad limbi basin usque vaginante alis linearibus apicem versus attenuatis, stirpis femineæ amentis axillaribus ternatis cylindricis densifloris, pedunculis filiformibus subtiliter hirtellis, bractea rotundata centro peltata subsessili, rhachi hirtella, bacca obovato-trigona paulo infra apicem margine caroso cincta, stigmatibus 3.  $\frac{5}{8}$  In Ins. Tahiti (Morrenhout ! in h. DC.), ins. Societatis, Amicorum, Novarum Hebridum (Guillemin in Zephir. Tahit. p. 29, Timor (Decaisne in h. Tim. descrip. p. 170). P. methysticum Lam. ill. p. 81. non Forst. P. latifolium Forster ! prodr. n. 22, Parkins. Draw. of Tahit. pl. ined. t. 7, Vahl enum. v. 1. p. 336, Willd. sp. v. 1. p. 161, Hook. et Arn. in Beechey voy. p. 70, Seemann in Flora Vit. pars 7. p. 261. Macropiper latifolium Miq. ! syst. p. 218, illust. p. 29. t. 27. Ava, Ava Irai ab incolis dicta. Dioicum. Limbi 0,09-0,095 longi latique. Petioli 0,04-0,06 longi. Pedunculi 0,02-0,03 longi. Amenta matura 0,07-0,085 longa, 0,005 crassa. (v. s.)

390. P. TRISTACHYON, foliis longe petiolatis rotundatis vel rotundato-ovatis apice breviter protracto-acuminatis basi truncato-rotundatis cordatisve supra glabris subtus hirsuto-pubescentibus membranaceis 11-nerviis, petiolo ad  $\frac{1}{2}$  longitud. usque vaginante extus pubescente, amentis axillaribus ternatis, pedunculis quam petioli triplo brevioribus, bracteis rotundatis centro peltatis subsessi-

libus, stigmatibus 3, baccis oblongo-subtetragonis dense approximatis distinctis. ♀ In ins. Sandwich (Bidwill ! in h. Kew.), Tahiti ubi vulgare (Vieillard ! in h. Lenormand). Dioicum. Ramuli pubescentes. Limbi 0,1 longi, 0,07 lati. Petioli 0,06 longi. Pedunculi 0,02 longi. Amenta matura 0,05 longa, 0,005 crassa. (v. s.)

391. *P. GUAHAMENSE*, foliis longe petiolatis rotundatis vel rotundato-ovatis apice breviter protracto-acuminatis, acumine acutiusculo, basi profunde cordatis, supra glabris subtus ad nervos ochraceo-hirtellis membranaceis rigidulis subopacis subtilissime pellucido-punctulatis 9-11-nerviis, nervis subtus prominulis centrali ad apicem ducto proximis 2 fere ad apicem ductis infimis subtilioribus, petiolo ad  $\frac{1}{2}$  longitud. usque vaginante alis linearibus apicem versus in petiolum attenuatis, stirpis femin. amentis axillaribus solitariis densifloris folia subæquantibus, pedunculis quam petioli dimidio brevioribus sparsim hirtellis, rhachi hirtella, bractea rotundata centro peltata pedicellata, ovario ovato glabro, stigmatibus 3 carnosulis sessilibus, stirpis masc. petiolis glabris, foliis magis membranaceis, amento (in spec. soluto solitario ?) densifloro, rhachi hirtella, stam. 3. ♀ Stirps femin. ex ins. Guaham Archipelagi Mariannum (Chamisso ! in h. Berol.), stirps masc. forsitan specificè distincta ex Mexico (Hænke ! in h. Berol.). *Macropiper methysticum* Miq. ! syst. p. 218. *Macr. latifolium* Miq. ! syst. p. 218, stirps masc., in Hook. Lond. journ. bot. v. 3. p. 431. Dioicum. Ramuli glabri. Limbi 0,11 longi, 0,12 lati. Petioli 0,045 longi. Amenta florentia 0,1 longa, 0,003 crassa. (v. s.)

392. *P. BARCLAYANUM*, foliis petiolatis rotundato-vel elliptico-ovatis apice breviter protracto-attenuatis acutiusculis basi æquali rotundatis truncatis utrinque glabris rigidis opacis trinerviis, nervis subtus prominulis, centrali ad apicem ducto, petiolo glabro, pedunculo glabro petiolum æquante, amentis quam folia multum brevioribus, bractea rotundata centro peltata sessili, stigmatibus 4, baccis obovato-subglobosis glabris subcoalitis. ♀ In ins. Tobia prope Novam Guineam (Barclay n. 4021 ! in h. Brit. Mus.). *Chavica Benthamiana* Miq. ! in Hook. Lond. journ. bot. v. 4. p. 433. Dioicum. Ramuli glabri. Limbi 0,06 longi, 0,05 lati. Petioli 0,012 longi. Baccæ 0,003 crassæ. (v. s.)

Sect. VI. COCCOBYRON. — Genus *Artanthe* Miq. *comm. phyt. p. 40 (pro parte)*. — *Peltobryon* Klotzsch in h. Berol. (*pro parte*). — *Coccobryon* Klotzsch l. c.

Bractea a flore distincta. Stamina 2 lateralialia vel 3 quorum stamen unum positum. Amenta oppositifolia. Flores hermaphroditi. Antheræ articulatae.

### § 1. *Stigmata* 3-4.

#### α. *Bacca stipitata*.

393. *P. CUBENSE*, foliis breviter petiolatis elliptico-lanceolatis, acumine acutiusculo, utrinque acutis supra glabris subtus ad nervos puberulis firmo-membranaceis opacis 5-nerviis, nervis subtus prominulis centralibus 3 per anastomoses ad apicem lateralibus subtilioribus ad medium ductis, nervulis subtransversim parallelis, petiolo glabro basi ima vaginante, pedunculo petiolum multum superante juniore hirtello, amento maturo folium æquante, rhachi hirtella, bractea subrotundato-concava extus subtiliter puberula, staminibus 3, anthera filamentum æquante fusco-glandulosa, ovario oblongo apice in stylum subtetragonum producto, stigmatibus 3 brevissimis ovatis, bacca ellipsoidea apice in stylum brevem basi in pedicellum quam bacca multum breviori contracta. ♀ In ins. Cuba (Wright n. 513 ! in h. Boiss. et DC. et 2213 fide Griseb. Cat. pl. cub. p. 65), prov. Pilos de los Mandones in Cuba (Linden n. 177 ! in h. DC.). *Enckea Amalago* Griseb. Cat. pl. cub. p. 64 quo ad n. 513. *Ottonia punctata* Griseb. l. c. p. 65. Ramuli puberuli. Limbi 0,06 longi, 0,025 lati. Petioli 0,003 longi. Pedunculi 0,03 longi. Amenta matura 0,06 longa. (v. s.)



## β. Bacca sessilis.

## 1° Folia penninervia.

394. *P. SAN-CARLOSIANUM*, foliis brevissime petiolatis lanceolato-ellipticis apice acuminatis, acumine acuto, basi æquali vel subæquali attenuatis acutis supra pilosis subtus molliter adpresse villosis membranaceis opacis, centrali nervo ad apicem usque nervos alternos subadscendenti-patulos alternos utrinque 8-10 mittente, petiolo utrinque (subtus magis) hirsuto basi ima vaginante, pedunculo hirsuto petiolum duplo superante, bractæ apice peltatæ pelta triangulari brevissime pedicellata margine argenteo-villosula, staminibus 2, anthera filamentum valde superante, ovario rhachi immerso et cum eo coalito. ♀ In Brasilia prope San-Carlos ad Rio Janeiro (Spruce n. 3099! in h. Kew. et DC.). Ramuli hirsuti. Limbi 0,08 longi, 0,03 lati. Petioli vix 0,002 longi. Pedunculi 0,003-0,004 longi. Amenta post anthesin 0,015 longa, 0,002 crassa. (v. s.)

395. *P. MALACOPHYLLUM*, foliis breviter petiolatis ovato-oblongis apice angustato-acuminatis basi inæquali cordatis obliquis supra pilis demum plus minus deperditis scabris subtus glandulis transparentibus flavis satis magnis copiosissimis pellucido-punctatis pinnatinerviis, pedunculo quam petiolus duplo longiore, amento elongato cylindrico obtuso quam folium brevior curvato flexuoso, bractea oblonga longe ciliato-hirsuta, staminibus 2, ovario subgloboso subangulato glabro, stigmatibus 3 acutis. ♀ In Brasilia ad Rio Janeiro (Pohl fide Presl l. c.). *Artanthe malacophylla* Presl Epim. bot. p. 227, Miq. in Mart. Flor. Bras. fasc. 11. p. 73. Pubescens, pilis horizontalibus patentibus mollibus brevibus albidis densissimis. Folia 5 poll. longa, 2 poll. lata. Petioli 2 lin. longi. (Ex Presl l. c.)

396. *P. MAROWINIANUM*, foliis breviter petiolatis subelliptico-oblongis apice longiuscule acuminatis, acumine acuto, basi æquali cordulatis supra glabris subtus nigro-punctulatis ad nervos nervulosque villosulis rigidulis subopacis pellucidopunctulatis, nervis subtus prominulis centrali ad  $\frac{1}{2}$ - $\frac{2}{3}$  longitud. nervos alternos subadscendentes utrinque 5 quorum supremi per anastomoses ante apicem evanidi infimi subtiliores sursumque nervulos nonnullos validos mittente, petiolo ad limbi basin usque vaginante dorso villosulo, pedunculo petiolum paulo superante juniore villosulo, amento quam folium multoties brevior densifloro cylindrico apice obtuso, bractea subobovato-subcucullata intus et basi dorso villosula vertice subinflexo nuda, staminibus 2, rhachi villosula, bacca globosa apice in stylum brevem contracta, stigmatibus 3 abbreviatis. ♀ In Brasilia (Riedel n. 1584! in h. Petrop.), et Surinam regionibus interior. ad Rio Marowini (Kappler n. 1885! in h. Boiss.). Ramuli villosi. Limbi 0,13 longi, 0,035 lati. Petioli 0,005 longi. Amenta 0,006 crassa. (v. s.)

397. *P. CYRTOPODON*, foliis brevissime petiolatis oblongo-ellipticis apice breviter acuminatis, acumine acutiusculo, basi subinæquali rotundatis cordulatisque supra glabris subtus pilosulis, nervis subtus prominulis centrali ad  $\frac{1}{2}$ - $\frac{2}{3}$  longitud. usque nervos alternos subadscendentes utrinque 5 nervulis validis intermixtos mittente, petiolo ad limbi basin vaginante dorso piloso-pubescente, pedunculo pilosulo petiolum superante, bractea cucullato-obovata intus et basi extus pilosula apice subinflexa, staminibus 2, ovario apice breviter attenuato, stigmatibus 3, rhachi hirtella. ♀ In vicinibus Obidos provinc. Para (Spruce! in h. Kew. ibid n. 472! in h. Francovil. et n. 4! aut 519! in h. monac.). Peltobryon cyrtopodon Miq. mss. in h. Kew. Ramuli pilosuli. Limbi 0,12 longi, 0,045 lati. Petioli 0,005 longi. Pedunculi 0,012 longi. Amenta florentia 0,035 longa, 0,003 crassa. (v. s.)

398. *P. QUITENSE* (Cas. DC. in Seem. journ. 1866, p. 163), foliis brevissime petiolatis lanceolatis apice acutis mucronulatisque basi acutis utrinque glabris membranaceis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis usque nervos alternos adscendentes ut inque 2 mittente, petiolo glabro, pedunculo glabro petiolum multum superante, bractea glabra lanceolata vertice subinflexo-peltata, staminibus 3 raro 4, filamentum antheram multum superante, ovario oblongo glabro apice in stylum longum attenuato, stigmatibus 3.

♂ In Andibus Quitensibus alt. 8000 ped. (Jameson n. 414! in h. Kew. et DC.). Ramuli glabri. Limbi 0,075 longi, 0,03 lati. Petioli 0,003 longi. Pedunculi 0,01 longi. Amenta brevia, florentia 0,01 longa. (v. s.)

### 2° Folia multiplinervia.

399. *P. EPUNCTULATUM*, foliis longiuscule petiolatis subrotundis apice breviter attenuato-acuminatis, acumine obtuso, basi æquali profunde cordatis supra parce pilosulis ad nervos densius hirtellis dein glabratibus subtus dense molliter tomentosis firmulo-membranaceis subopacis epunctulatis 13-plinerviis, nervis utrinque prominulis centrali ad  $\frac{1}{2}$  longitud. nervum utrinque unum ad apicem usque ductum oppositè mittente cæteris nervis e basi solutis, petiolo ad  $\frac{1}{2}$  longitud. usque vaginante basi dorso et sursum utrinque tomentoso, pedunculo tomentoso petiolum æquante, amento folium subæquante, bractæa oblongo-subspathulata subcucullata intus et basi dorso villosula vertice truncato-carnosula, staminibus 2, filamento exserto antheram multum superante, ovario ovato apice in stylum brevem attenuato basi villosulo, stigmatibus 3 linearibus recurvis. ♂ In provinc. Minas-Geraës (Lindberg n. 486! in h. Mart.). Rami puberuli. Ramuli dense tomentosi. Limbi 0,12 longi, 0,15 lati. Petioli 0,04 longi. Amenta florentia 0,006 crassa. (v. s.)

400. *P. BIRMANICUM* (Cas. DC. in Seem. journ. 1866, p. 163), foliis breviter petiolatis oblongis apice attenuatis acutiusculis vel e basi cuneata obovatis apice breviter attenuatis acutis supra glabris subtus ad nervos puberulis membranaceis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitudinis usque nervos suboppositos subadscendentes utrinque 3 mittente, petiolo hirsuto, pedunculo hirsuto petiolum æquante, bractæa rotundata centro affixa, staminibus 2 interdum 3, filamento antheram multum superante, ovario oblongo apice in stylum brevem attenuato, stigmatibus 3 recurvis, bacca subglobosa apice cuspidata glabra. ♂ In peninsula birmanica prope Malacca (Griffith n. 4414! in h. Kew.). Ramuli apice ochraceo-hirsuti. Limbi 0,1 longi, 0,07 lati. Petioli 0,005 longi. Amenta matura 0,025 longa. (v. s.)

♂, *macrostachyon*. Limbi 0,2 longi, 0,09 lati. Amenta 0,1 longa. — Prope Malacca (Griffith n. 4408! in h. Kew.). (v. s.)

401. *P. MITRE* (Ruiz et Pav. flor. peruv. v. 1. p. 37. t. 58. fig. a), foliis petiolatis ample ovatis apice acuminatis, acumine acuto, basi rotundatis subcordatis supra glabris nitidulis subtus pube stellata dense molliter pubescente firmis opacis subtilissime pellucido-punctulatis quintuplinerviis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervos alternos subadscendentes quorum supremi ad apicem ducti infimi e basi soluti mittente, petiolo fere ad limbi basin usque vaginante basi dorso apice utrinque dense molliter pubescente, amento maturo folium superante crasso, pedunculo dense subfusce pubescente quam petiolus brevior, bractæa truncato-peltata pelta triangulari margine subfusce hirtopubescente pedicello concavo dorso hirtopubescente, staminibus 2, ovario glabro apice in stylum brevem attenuato, stigmatibus 3 linearibus. ♂ In sylvis Cuchero (R. et P. l. c.), Peruvia (Pav. l. in h. Boiss.) — Vahl enum. v. 1. p. 325. *Artanthe mitis* Miq. syst. p. 407. Rami glabri. Limbi 0,22 longi, 0,15 lati. Petioli 0,03 longi. Amenta mat. 0,008 crassa (v. s.)

### 3° Folia multinervia.

402. *P. PAPANTLENSIS*, foliis petiolatis inferioribus ovatis apice subprotracto-acuminatis, acumine acuto mucronulatoque, basi æquali rotundatis, supremis ovato-lanceolatis basi æquali acutis omnibus utrinque glabris opacis epunctulatis 7-vel 5 nerviis, nervis subtus prominulis centrali ad apicem ducto sequentibus ad medium ductis exterioribus subtilioribus, petiolo glabro ad  $\frac{1}{2}$  longitudinis usque vaginante, pedunculo glabro petiolum æquante, amento maturo folium æquante vel paulo superante, bractæa apice truncata vertice triangulari minuto margine subtiliter ciliolato, staminibus 2, filamento antheram multum superante, rhachii foveolata margine foveolarum hirtella, bacca obovato-subquadrangula glabra olivacea, stigmatibus 4 abbreviatis. ♂ In

sylvis Papanilæ (Fischer n. 74! in h. Petrop.). Rami ramulique glabri. Limbi 0,105 longi, 0,05 lati. Petioli 0,012 longi. Amenta matura 0,004 crassa. (v. s.)

## § 2. Stigmata 2.

403. *P. IMPERIALE*, foliis maximis longe petiolatis elliptico-oblongis acutiusculis, basi valde inæquali cordatis, lobo majore petiolum velante, crasse coriaceis supra glabris subtus ad nervos incano-pubescentibus, centrali nervo subtus prominente ad  $\frac{3}{4}$  longitud. usque nervos subadscendentes utrinque 9 quorum supremi ad apicem ducti mittente, amento quam folium breviores, bractee oblongo-obovate dorso ciliolatae vertice inflexo-cucullato, staminibus 2, ovario glabro, stylo teretiusculo, stigmatibus 2 teretiusculis patulis, bacca obovodea. ♂ In ins. Cacagua!, Darien (Seem. n. 1068! in h. Kew.). Artanthe imperialis Miq.! in Seem. bot. Herald p. 198. t. 38. Petiolus tripollicaris. Folium 20 poll. longum, 11  $\frac{1}{2}$  poll. supra medium latum. Pedunculus 2 poll. longus. (Ex Miq. l. c.)

404. *P. PILULIFERUM* (Kunth! in H. et B. nov. gen. v. 1. p. 53), foliis subsessilibus elliptico-lanceolatis apice longe acuminatis, acumine acuto, basi inæquali subattenuatis, lobo majore auriculatum productum, supra glabris subtus ad nervos puberulis, centrali nervo ad  $\frac{3}{4}$  longitud. usque nervos alternos adscendentes utrinque 4-5 mittente, pedunculo hirtello, amento sphaerico densifloro, bractea spatulata apice inflexa glabra, staminibus 2 exsertis, ovario apice in stylum brevem attenuato, stigmatibus 2 lateralibus. ♂ In Andibus Quitensibus alt. 6000 ped. sylvis humidis (Jameson n. 245! in h. DC.). declivitate Andium inter Cuenca et Tambo de Buray schisto micaceo alt. 1400 hexap. (Humb. l. c. et h. Willd. n. 673!). — Kunth synopsis. v. 1. p. 108. Schilleria? pilulifera Kunth! in Linn. v. 13. p. 722. Sphaerostachya Humboldtii Miq.! syst. p. 376, illustr. p. 68. t. 68. Ramuli glabri. Limbi 0,13 longi, 0,04 lati. Pedunculi 0,015 longi. Amenta 0,005 crassa. (v. s.)

405. *P. CAPENSE* (L. fil. suppl. p. 90), foliis longiuscule petiolatis ovatis apice breviter acuminatis, acumine obtusiusculo, basi obtusis vel rotundato-cordulatis supra glabris subtus pilosulis vel glabris membranaceis vel rigidulis 5-7-nerviis, nervis 3 centralibus validioribus ad apicem ductis exterioribus subtilioribus, nervulis transversis, petiolo glabro, pedunculo glabro petiolum superante, amento cylindrico, bractea rotundata glabra centro peltata breviter pedicellata, staminibus 2, filamento antheram superante, ovario ovato, stigmatibus 2 rhachi oppositis, bacca glabra aromatica. ♂ In Promontorio bonæ spei (Drège! in h. Kew.; Meron! Mundt! et Mair! in h. Berol.), montibus ad Puparvalei alt. 2000-4000! (Zeyher! in h. Francov.), ins. St-Thomas ad Fernando-Po et Cameroon montibus alt. 7000 ped. (Mann n. 299! et 2015! in h. Kew.) et 14-15 latitud. (Kirk! in h. Kew.), Natal (Gueinsius! in h. Boiss. et n. 110! in h. Vindob. et h. DC. sine origine). — Thunb. flor. cap. p. 443; Lam. ill. p. 81, Vahl enum. v. 1. p. 329, Willd. sp. p. 159. Coccobryon capense Klotzsch! mss. in h. Berol., Miq.! syst. p. 343, illustr. p. 50. t. 61. Scandens. Ramuli glabri. Limbi 0,08-0,1 longi. (v. s.)

Sect. VII. EUPIPER. — Gen. Piper Bl. in Verh. der bot. Genoots. fasc. 11 (pro parte), Miq. syst. p. 305. — Cubeba. Miq. comm. phyt. p. 35. — Chavica Miq. syst. p. 222 (pro parte).

Bractea a flore distincta. Stam. 2 lateralia vel 3 quorum unum posticum vel rarissime 4; antherae articulatae. Amenta oppositifolia. Flores unisexuales vel basi amentorum hermaphroditi vel in speciebus rarissimis trifidis modo unisexuales modo hermaphroditi.

### α. Bacca stipitata.

#### 1° Folia penninervia.

406. *P. BORBONENSE*, foliis breviter petiolatis oblongo-ovatis apice acuminatis, acumine obtuso, basi subinæqualiter cordulatis subattenuatisve firmis opacis

utrinque glabris, centrali nervo subtus prominente nervos fere e basi solutos ad medium ductos utrinque 1-2 sursumque tota longitudine nervulos alternos patulos utrinque 4-8 mittente, petiolo dorso hirtello, pedunculo glabro petiolum 2<sup>o</sup>-4<sup>o</sup> superante, amento femineo densifloro, bractea obovata subrotundata margine subtusque pubescente, rhachi hirtella, stigmatibus 3, bacca ovata, stipite baccam superante, stirpis masc. foliis et bractea ut in femin. stam. 3 exsertis. ♀ In ins. Bourbon (Du Petit Thouars in h. Willd. n. 636 !), ins. Mauritio (Bouton ! et Bory ! in h. DC.), Madagascar (Boivin ! in h. Vindob.), promont. Bonæ spei (Verreaux ! in h. Deless., Gueinsius ! in h. Vindob.), P. aristolochioides Lam. v. 1. p. 80 ? (Lam. in h. Willd. n. 677. fide Miq. syst. p. 303), Vahl enum. v. 1. p. 326. Cubeba costulata Miq. syst. p. 299. Cubeba borbonensis Miq. syst. p. 301. Dioicum. Supra arbores parasiticum. Rami glabri. Limbi 0,06-0,13 longi, 0,025-0,08 lati. Petioli 0,008-0,01 longi. (v. s.)

407. P. CLUSII, foliis breviter petiolatis vel in femina longius petiolatis ovato-acuminatis, acumine acuto, basi subinæquali cordulatis membranaceo-rigidulis, centrali nervo subtus prominulo fere tota longitudine nervos patulo-adscendentes utrinque 4-5 nervulosque validiores mittente, petiolo glabro pedunculum in femina paullo superante in mare subæquante, floris femin. bractea ovato-rotunda carnosula centro peltata subsessili subtus hirtella, floris masc. bractea ut in femin. stam. 2 exsertis. ♀ In Fernando-Po (Mann n. 162 ! stirpis masc., et Vogel n. 97 ! stips femin. in h. Kew.). Cubeba Clusii Miq. syst. p. 304, Miq. ! in Hook. Lond. journ. bot. v. 4. p. 434. Dioicum, scandens. Ramuli glabri. Limbi 0,08-0,09 longi, 0,04-0,05 lati. Petioli 0,005-0,015 longi. (v. s.)

408. P. VENOSUM, foliis longiuscule petiolatis lanceolato-vel-elliptico-oblongis apice breviter acuminatis basi æquali vel subinæquali emarginatis supra glabris subtus ad nervos numerosos patulos demum subscrobiculatis puberulis subcoriaceis utrinque (præsertim subtus) nigro-punctulatis, amento femineo longiuscule pedunculato, bractea subtus hirtella, bacca brevissime stipitata. ♀ In Java (Miq. pl. Jungh. fasc. 1. p. 295.). Cubeba venosa Miq. l. c. et fl. ind. bat. v. 1. pars 2. p. 450. Dioicum. Scandens.

409. P. PHYLLOSTICTUM, foliis modice petiolatis oblongo-lanceolatis vel lanceolatis apice breviter acuminatis basi inæquali subdimidiata emarginatis submembranaceis subtus glaucescentibus utrinque glabris et nigro-punctatis, nervis numerosis patulis infimis adscendentibus, amento femineo longe pedunculato, bractea decurrenti-adnata, stigmatibus 3 brevibus, bacca globosa stipitem subæquante. In Java (Miq. pl. Jungh. fasc. 1. p. 294), monte Pangerango prope domunculum (Miq. Mus. lugd. bat. fasc. 5. p. 136.), Sumatra in Priaman (Miq. Sumatra p. 186. et 473.). Cubeba phyllosticta Miq. ll. cc. et flor. ind. bat. v. 1. pars 2. p. 450.

410. P. CUBEBA (L. fl. suppl. plant. p. 90), foliis breviter petiolatis stirpis femineæ ovato-ellipticis masculæ oblongo-subovato-ellipticis omnibus apice attenuatis acutis basi subinæquali cordulatis utrinque glabris coriaceis, nervo centrali ad  $\frac{3}{4}$  longitud. usque nervos subadscendentes utrinque 3-4 nervulos que validiores mittente, petiolo glabro, pedunculo glabro petiolum subsuperante, stirpis fem. bractea oblongo-obtusa rhachi adnata margine libera subtus hirtella, ovario oblongo glabro, stigmatibus 4, bacca subglobosa apice subapiculata, stipite baccam paullo superante, stirpis masc. bractea oblongo-rotundata sessili centro affixa, stam. 3. ♀ In Borneo australi (Miq. in Ann. Mus. lugd. bat. v. 1. fasc. 5. p. 136), Java (Junghuhn ! in h. Kew. stirpis fem., Zollinger n. 727 ! stirpis masc. in h. Boiss. et 927 ! stirpis fem. in h. DC., Bl. ! in h. DC. ex h. lugd. bat.), ins. Nusa Kambangan spontanea (Bl. enum. pl. Jav. fasc. 1. p. 70). In Antillis culta (Descourtils flor. Ant. v. 6. p. 201. t. 429). — Lam. ill. p. 81, Vahl enum. v. 1. p. 333, Roem. et Schult. v. 1. p. 312 ? Bl. ! in Verh. der bat. Genoots. v. 11. p. 200. fig. 21 et l. c., Roxb. fl. ind. v. 1. p. 161, Miq. comm. phyt. p. 13. t. 1. et 2. Cubeba officialis Miq. comm. phyt. p. 33, syst. p. 285, illustr. p. 48, in Hook. Lond. journ. bot. v. 4. p. 434, flor. ind. bat. v. 1. pars 2. p. 448, Schnizl.

iconog. t. 81. fig. 18-20. Dioicum. Caulis glaber. Limbi 0,085-0, 13 longi 0,04 lati. Petioli 0,005 longi. Pedunculi 0,008 longi. (v. s.)

β, *parvifolium*, foliis minoribus. — In Java (Zoll. n. 943 ! stirps masc.) (v. s.)

2° Folia multiplinervia.

411. *P. VESTITUM* (Cas. DC. in Seem. journ. 1866, p. 166), foliis petiolatis ovato-rotundatis apice breviter protracto-attenuatis basi cordatis membranaceis utrinque dense pilosis, centrali nervo fere ad  $\frac{1}{2}$  longitud. usque nervos oppositos subadscendentes utrinque 6 quorum infimi subpatuli mittente, nervulis subtransversis, stirpis femin. bacca polygono-globosa, stirpis masc. bractæe triangulari-rotundæ glabræ centro peltatæ pedicello crasso, staminibus 2, antheris subsessilibus. ♀ In littore septentrionali Borneo (Lobb ! in h. Kew.). Dioicum. Limbi 0,25 longi, 0,02 lati. Petioli 0,07 longi. Pedunculi 0,05 longi. (v. s.)

412. *P. CANINUM* (A. Dietr. sp. v. 1. p. 681), foliis breviter petiolatis inferioribus ovato-acuminatis, acumine acuto, basi æquali vel subinæquali obtusis vel interdum cordulatis superioribus et maris omnibus ovato-oblongis apice acute acuminatis, vel lanceolatis basin versus attenuatis basi ima acutis omnibus supra glabris subtus pilosis vel glabris membranaceis tripli-septuplinerviis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos oppositos fere ad apicem ductos utrinque 1-2 mittente, nervulis subtransversis, petiolo pubescente, pedunculo puberulo petiolum subæquante, floris femin. bractea rotundata centro peltata brevissime pedicellata subtus rhachique hirsuta, bacca subglobosa stipitem vix superante, floris masc. bractea rotundata sub centro peltata subsessili subtus ad pedicellum hirtella, staminibus 2, filamentis brevissimis. ♀ In Borneo ad Bangarmassing (Motley 835 et 13 ! stirps fem. in h. kew.), Labuan (Motley n. 210 ! in h. Kew.), Java (Zoll. n. 677. in h. DC. et 710 ! in h. Petrop. et Bl. in h. DC. ex h. lugd. bat.), Poulo Pinang, Sumatra, Moluccis (Miq. flor. ind. bat. v. 1. pars 2. p. 449. et Sumatra p. 186), Ceram (Miq. in Ann. Mus. lugd. bat. v. 1. fasc. 5. p. 141), per totum Archipelagum distributum et valde varians, in singulis insulis paulisper diversum (Miq. in Ann. mus. lugd. bat. l. c. p. 137). Rumph. Amb. v. 5. l. 7. c. 26. t. 28. f. 2 ? Vahl enum. v. 1. p. 332. *P. sylvestre* Lour. flor. Cochinchin. p. 38 ? *P. Cubeba* Roxb. flor. ind. v. 1. p. 159 ? *P. caninum* Bl. ! in Verh. der Bat. Genoots v. 11. p. 214, enum. pl. jav. fasc. 1. p. 72, Miq. comm. phyt. p. 17 et 33. t. 3. *Cubeba canina* Miq. ! syst. p. 293 et in pl. Jungh. fasc. 1. p. 17 et II. cc. Dioicum. Ramuli glabri. Limbi 0,06-0,1 longi, 0,035-0,04 lati. Petioli 0,008 longi. (v. s.)

413. *P. LANATUM* (Roxb. ! flor. ind. v. 1. p. 161), foliis petiolatis oblongo-ellipticis apice acuminatis, acumine acuto, basi subinæquali rotundatis subcoriaceis supra sparse subtus densius hirta-tomentosis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos alternos adscendentes utrinque unum et e basi utrinque 1-2 quorum infimi subtiliores mittente, petiolo hirta-tomentoso, pedunculo petiolum duplo superante, amento femineo quam folium multum brevius, bacca ovata stipitem æquante. ♀ Patria ignota (Roxb. ! in h. Deless.), Sunda Archipel (Miq. flor. ind. bat. v. 1. pars. 2. p. 450). *Peperomia lanata* A. Dietr. sp. v. 1. p. 145. *Cubeba lanata* Miq. ! syst. p. 298 et l. c. Dioicum. Ramuli ochraceo-hirti. Limbi 0,11-0,12 longi, 0,04-0,035 lati. Petioli 0,01-0,015 longi. (v. s.)

414. *P. MURICATUM* (Bl. ! in Verh. der Bat. Genoots v. 11. p. 219. fig. 1), foliis petiolatis inferioribus subovatis subobovatisve apice breviter acuminatis, acumine acuto, basi æquali subcordatis superioribus oblongis apice breviter acuminatis, acumine acuto, basi inæquali subcordatis omnibus supra adpresse pubescentibus subtus densius tomentoso-pubescentibus membranaceis subopacis pellucido-punctulatis inferioribus undeni superioribus quintupli-septuplinerviis, nervis subtus prominulis centrali ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. usque nervos subalternos adscendentes utrinque 1-2 et e basi utrinque 3-4 mittente, petiolo ad  $\frac{1}{2}$  longitud. usque vaginante basi dorso sursum utrinque patentem hirtello, pedunculo hirtello petiolum paullo superante, amento hermaphrodito vel unisexuali quam folium paullo brevius densifloro, bractea rotundata

centro peltata sessili supra glabra subtus pilosula, rhachi hirta, in floribus hermaphroditis staminibus 2, ovario ovato sessili, stigmatibus 3-4 lanceolatis, bacca ovoides glabra, stipite baccam fere æquante. ♀ In ins. Java (h. DC. ex h. Lugd. bat.), locis humilibus ad montem Salak ubi arbores ascendit (Bl. enum. Javæ p. 68), Sumatra occid. (Miq. in Ann. mus. lugd. bat. v. 1. fasc. 5. p. 139). — Miq.! syst. p. 326, ill. p. 55. t. 55, in Hook. Lond. journ. bot. v. 4. p. 438, flor. ind. bat. v. 1. pars 2. p. 454). P. villosum Bl. mss. in pl. Reinw. (fide Miq. in flor. ind. bat.). Trioicum, idest monoicum et tum amenta hermaphrodita, vel dioicum. Rami glabrati, Ramuli dense tomentoso-hirti. Limbi 0,16-0,22 longi, 0,09-0,15 lati. Petioli 0,01 longi. Bacca mat. 0,002 longa. (v. s.)

β, *parvifolium*. Limbi 0,08-0,1 longi, 0,04-0,06 lati. — In ins. Java (h. DC. ex h. lugd. bat.). (v. s.)

γ, *pilosius*, foliis subtus tomentosis, bacca longiuscule stipitata (Miq. in Linn. v. 21. p. 485).

415. *P. arborescens* (Wall. list n. 6637!), foliis petiolatis oblongo-ovatis apice attenuatis acutiusculis basi æquali profunde cordatis utrinque glabris coriaceis, centrali nervo subtus prominulo ad  $\frac{1}{2}$  longitud. nervum utrinque unum et fere e basi utrinque 3 divaricantes mittente, petiolo glabro, pedunculo glabro quam petiolus multum brevior, bacca subpolygono-globosa quam stipites brevior apice mucronulata, stigmatibus 4. ♀ In Chappedong (Wall. n. 6637! in h. DC., 16781 an et 66371 in h. Wall.), Tenasserim et Andaman (Griffith n. 4401! in h. Kew.). — Wall. pl. As. rar. v. 1. p. 79. t. 91. CUBEBA Wallichii Miq.! syst. p. 289, in Hook. Lond. journ. bot. v. 4. p. 434, illustr. p. 47. t. 46 et 47 (non Wright icon. n. 1932 quod ad *P. nigrum* referendum). Dioicum. Ramuli glabri. Limbi 0,25 longi, 0,11 lati. Petioli 0,04 longi. Pedunculi 0,01 longi. (v. s.)

416. *P. LEONENSE* (Cas. DC. in Seem. journ. 1866, p. 163), foliis brevissime petiolatis oblongis apice longiuscule acuminatis, acumine acuto, basi æquali cuneatis utrinque glabris coriaceis triplinerviis, centrali nervo ad  $\frac{1}{2}$  longitud. nervum utrinque unum alterne alteramque utrinque e basi supra medium ductum mittente, petiolo glabro, pedunculo petiolum multum superante, amento quam folium brevior densifloro, stipite baccam subglobosam subsuperante. ♀ In Sierra Leone (Afzelius! in h. Berol. ubi cum *P. CUBEBA* mixtum). Dioicum. Ramuli glabri. Limbi 0,095 longi, 0,025 lati. Petioli 0,005 longi. Pedunculi fere 0,02 longi. Bacca sub 0,003 longa. (v. s.)

417. *P. MOLLISSIMUM* (Bl.! in Verh. der Bat. Genoots. v. 11. p. 223), foliis petiolatis ovatis apice attenuatis acutis mucronulatisque inferioribus basi subæquali superioribus basi inæquali cordatis supra glabris subtus tomentosis coriaceis septuplinerviis, nervis subtus prominulis centrali paullo supra basin nervum utrinque adscendentem et e basi utrinque 2 divaricantes mittente, petiolo ad  $\frac{1}{2}$  longitud. usque vaginante, pedunculo hirtello quam petiolo brevior, amento laxifloro folium superante filiformi, bractea oblonga utrinque obtusa, rhachi puberula, staminibus 3, filamentis crassis persistentibus, antheris caducis. ♀ In ins. Java (Zoll. n. 3001! in h. Boiss. et Bl.! in h. DC. ex h. lugd. bat.), Sumatra occid. (Korthals fide Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 137). Potomorphe? mollissima Miq.! comm. phyt. p. 46. t. 6. fig. A, syst. p. 215. in Linn. v. 20. p. 130, flor. ind. bat. v. 1. pars 2. p. 450. Dioicum. Ramuli hirtelli. Limbi 0,085-0,17 longi, 0,06-0,105 lati. Petioli 0,008 longi. (v. s.)

418. *P. BANKSI* (Miq. in Pip. nov. Novæ Hollandiæ in Meded. der Konink. Akad. van Veten.-ch. Afdee. Naturk. 2<sup>de</sup> Peeks Deel 2), foliis longiuscule petiolatis ovatis acuminatis basi rotundatis leviter emarginatis inferioribus etiam cordatis membranaceis subtus pilosis septuplinerviis, pedunculo quam petiolo brevior, bacca ellipsoidea stipitem æquante. ♀ In Novæ Hollandiæ Nova Cambria australi secus flumen Endeavour (Banks fide Miq. l. c.). P. pubescens Solander mss. fide Miq. l. c. Folia spathamam usque digitum longa in adnot. 7-nervia dicta in speciminib. examinatis septuplinerviis, nervis paullo supra basin series. Amenta baccifera digitum longa.

419. *P. SUMATRANUM*, foliis petiolatis oblongo-ovatis apice acuminatis acutiusculis basi subæquali rotundato-cordatis utrinque glabris coriaceis septuplinerviis, centrali nervo ad  $\frac{1}{2}$  longitud. trifido basi nervos utrinque 2 mittente, petiolo glabro, amento sessili quam folium multum brevior, bacca globosa glabra stipitem æquante. ♀ In ins. Penang (Wall. n. 6646 β! in h. Wall.). Cubeba Sumatrana Miq.! comm. phyt. p. 42. t. 6. fig. A, syst. p. 290, flor. ind. bat. v. 1. pars 2. p. 448, Sumatra p. 186. Piper pseudo-Cubeba Korthals mss. fide Miq. comm. phyt. p. 22. Dioicum. Limbi 0,2 longi, 0,1 circ. lati. Petioli 0,03 longi. Bacca 0,005 crassa. (v. s.)
420. *P. GUINEENSE* (Schum.! pl. guin. p. 19), foliis inferioribus longe petiolatis ovato-acuminatis, acumine acuto mucronulatoque, basi æquali cordatis superioribus breviter petiolatis majoribus oblongo-ellipticis apice acuminatis, acumine acuto mucronulatoque, basi rotundatis vel attenuato-acutiusculis, omnibus utrinque glabris membranaceis firmisve pellucidopunctulatis septuplinerviis, nervis subtus prominulis centralibus 3 supra basin alterne solutis fere ad apicem ductis proximis utrinque 2 fere a basi solutis brevioribus, petiolo glabro, pedunculo glabro petiolum superiorem æquante, amento quam folium brevior densifloro, bractea rotundata glabra centro affixa, stigmatibus 3 carnosulis, bacca globosa apice mucronulata, stipite glabro baccam superante. ♀ In Guinea (Schum.! in h. Petrop.), sylvis primitivis editoribus ad arbores varios scandens Africæ occidentalis regionis Angolensis, distr. Golungo alt. 1000-2400 ped. (Welwitsch n. 508! et 510! in h. DC.). Piper ex Guinea Clus. exot. p. 184. Cubeba Clusii Miq. syst. p. 304. Dioicum. Ramuli glabri. Limbi inferiores 0,05 longi, 0,04 lati. Petioli 0,01-0,008 longi. (v. s.)
421. *P. JAVANICUM*, foliis petiolatis ovato-acuminatis, acumine acuto mucronulatoque, basi æquali vel subinæquali rotundatis vel obtuse subattenuatis vel truncatis vel raro cordulatis supra ad nervos pilosulis subtus pilosis rigidulis quintuplinerviis, centrali nervo paulo supra basin trifido nervis lateralibus ad medium usque arcuatum ductis, petiolo hirtio, pedunculo hirtio petiolum superante, amento femina, quam folium brevior, bacca obovato-globosa glabra stipitem paulo superante, stirpis masc. foliis ut in femina sed minoribus, bractea oblonga utrinque obtusa, staminibus 3, filamentis antheras superantibus. ♀ In ins. Penang (Wall. n. 6647! in h. Wall. et DC.), Singapore (Wall. n. 6645 β! in h. Wall.), Java (Zoll. n. 904! in h. Boiss. stirpis masc.). Dioicum. Ramuli hirti. Limbi 0,12 longi, 0,05 lati. Petioli 0,01-0,02 longi. Bacca 0,003 crassa. (v. s.)
422. *P. GRIFFITHII* (Cas. DC. in Seem. journ. 1866, p. 163), foliis breviter petiolatis ovatis vel ovato-ellipticis apice protracto-acuminatis, acumine acuto, basi obtusis vel subattenuatis acutiusculis utrinque glabris rigidulis quintuplinerviis, nervo centrali paulo supra basin nervos adscendentes utrinque 2 fere ad apicem usque ductos quorum infimi subtiliores mittente, petiolo glabro, pedunculo glabro petiolum paulo superante, amento folium triplo superant. bacca globosa quam stipes brevior. ♀ In Bengalia orientali (Griffith n. 4492! in h. Kew.). Dioicum. Ramuli glabri. Limbi 0,12 longi, 0,07 lati. Petioli 0,01 longi. Amenta matura 0,19 longa. Bacca 0,002 crassa. (v. s.)
423. *P. PEDICELLOSUM* (Wall.! list. n. 6646 A), foliis breviter petiolatis ovato-acuminatis apice obtusiusculis basi cordulatis utrinque glabris coriaceis septuplinerviis, nervis centralibus 3 supra basin solutis ad apicem usque ductis, petiolo glabro, pedunculo glabro quam petiolum duplo brevior, amento quam folium brevior, bacca ellipsoidea apice apiculata stipitem æquante. ♀ In Singapore (Wall. n. 6646 A! in h. Wall. et Lambert! in h. DC.). Cubeba Neesii Miq.! syst. p. 292, flor. ind. bat. v. 1. pars 2. p. 449. Dioicum. Ramuli glabri. Limbi 0,08 longi, 0,05-0,06 lati. Petioli 0,005-0,025 longi. Bacca matura 0,006 longa. (v. s.)
424. *P. NOVÆ HOLLANDIÆ* (Miq. in Pip. nov. Novæ Holland. in Meded. der Konink. Akad. van wetensch. Afdel. Naturk. 2<sup>de</sup> Reeks Deel 2. p. 8), foliis inferioribus cordato-ovatis æquilateris 7-subseptuplinerviis superioribus a basi breviter inæquali-cordata vel oblique rotundato-cordata ovatis vel ovato-ellipticis apice breviter acuminatis, acumine acuto vel obtuso attenuatis, 5-7-plinerviis,

nervis lateralibus supremis haud procul a basi liberis infra apicem evanidis, amentis masculis et femineis longiuscule pedunculatis, stirpis masc. bractea suberibulari centro pedicellata subtus hirtella, staminibus 2 (an et 3?), stirpis femin. bacca ovata quam stipes brevior vel eum æquante. ♀ In Nova Hollandia boreali-orientali prope Keppel Bay (spec. masc. et fem.), prope Moreton Bay (spec. masc.), prope Pine river (spec. sterile) (F. Mueller fide Miq. l. c.). Dioicum, scandens, radicans, glabrum, speciminis sterilis folium 4 poll. longum 3 poll. latum, petiolus  $1\frac{1}{2}$  poll. longus. Speciminum florentium folia plerumque minora, speciminis masc.  $4\frac{1}{2}$ -3 poll. longa,  $2\frac{1}{2}$ - $1\frac{1}{2}$  lata, chartacea, petioli  $\frac{1}{2}$ - $\frac{3}{4}$  poll. longi; speciminis femin. folia ut in mare. Amentum bacciferum  $2\frac{1}{2}$  poll. longum, pedunculus  $1\frac{1}{2}$  poll. longus. Bacca  $1\frac{1}{2}$ -1 $\frac{1}{2}$  lin. longa, stipes  $1\frac{1}{2}$  lin. longus.

425. *P. CRASSIPES* (Korthals mss. fide Miq. comm. phyt. p. 22 et 44), foliis ovatis acutiusculis vel acutis basi obtusis vel subcordatis sextuplinerviis nervis approximatis, 2 superioribus supra basin divergentibus, pedunculo petiololum multum superante, stipite elongato. ♀ In montosis ins. Sumatra ad montem Singalang. Cubeba crassipes Miq. comm. phyt. p. 45, syst. p. 292, flor. ind. bat. v. 1. pars. 2. p. 448, Sumatra p. 186. Dioicum. Petiolus 0,012 longus. Folium 0,1 longum, 0,05 latum.

### 3° Folia multinervia.

426. *P. LOWONG* (Bl. l. in Verh. der Bat. Genoots. v. 11. p. 161), foliis petiolatis inferioribus ovato-oblongis apice suboblique acuminatis, acumine acuto mucronulatoque, basi æquali rotundatis superioribus oblongo-lanceolatis apice ut in inferioribus basi subæquali acutis, omnibus supra ad nervos parce pilosis subtus præsertim ad nervos hirtis membranaceo-rigidulis subopacis pellucido-punctulatis 5-7-nerviis, nervis subtus prominulis, centrali nervo ad apicem ducto, proximis nervis fere ad apicem ductis, infimis subtilioribus brevioribus, petiolo ad limbi basin usque vaginante dorso dense hirtio, pedunculo hirtio petiololum subæquante, amento breviter curvato, bractea apice pubescente oblongiuscula obtusa centro affixa, stigmatibus 4-5, bacca ovali stipitem crassiusculum vix superante. ♀ In media altitudine montis Kadang Javæ occid. (Bl. l. in h. DC. ex h. lugd. bat.). Cubeba Lowong Miq. comm. phyt. p. 33; Miq. l. syst. p. 296, illustr. p. 48. t. 48, flor. ind. bat. v. 1. pars 2. p. 449. Dioicum. Caulis scandens. Ramuli molliter hirtelli. Limbi 0,135 longi, 0,065 lati. Petioli 0,12 longum. (v. s.)

### β. Bacca sessilis.

#### A. Ovarium apice styliforme.

427. *P. RHYNCHOLEPSIS*, foliis brevissime petiolatis maris oblongis apice longe cuspidatis basi cordatis femine ovatis apice longe cuspidatis basi subinæquali cordulatis omnibus utrinque pilosis membranaceis maris 7-nerviis femine 9-nerviis, nervis subtus prominulis centralibus 3 ad apicem ductis proximis supra medium ductis cæteris subtilioribus, petiolo villosio, pedunculo glabro petiololum superante, stirpis masc. amento cylindrico densifloro, bractea ovato-oblonga apice longe subulata sub medio peltata, staminibus 2, filamento longo exserto, anthera minuta, stirpis femin. bractea ovato-subrotunda sub centro peltata pedicellata, ovario glabro, stigmatibus 3. ♀ In ins. Philippinis (Cuming n. 1706! mas et 1697! femina in h. Boiss.). *Rhyncholepis Cumingiana* Miq. l. Syst. p. 282. ill. p. 45. t. 45, in Hook. Lond. journ. bot. v. 4. p. 43, flor. ind. bat. v. 1. pars 2. p. 447. Dioicum. Ramuli rufo-villosi. Limbi 0,15 longi, 0,045 lati, femine 0,15 longi, 0,075 lati. Petioli vix 0,005 longi. (v. s.)

β, *brevicuspe*, foliis ovatis apice subito breviter acuminatis, acumine acuto, basi cordatis 10-nerviis, bractea maris ovato-subrotunda ut in femina. — In ins. malais (Cuming n. 1843! in h. Kew. et Boiss.), in Philippinis ad Manillam (Meyen! in h. Berol.). *Rhyncholepis brevicuspis* Miq. l. syst. p. 283, flor. ind. bat. v. 1. pars 2. p. 447. (v. s.)

428. *P. STYLOSUM* (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 139), foliis



modice petiolatis elliptico-ovatis apice acuminatis basi subæquali vel oblique rotundatis supra glabris subtus adpresse sæpe retrorsum ochraceo-hirsutulis membranaceis septuplinerviis, nervis 3 centralibus ad apicem fere ductis, petiolo adpresse retrorsum ochraceo-hirsutulo, amento brevi cylindrico densifloro hermaphrodito, bractea orbiculari subtus brevissime pilosula centro peltata pedicellata, ovario ovoideo apice in stylum attenuatum, stigmatibus 3-4 lanceolatis, staminibus 2. ♀ In Sumatra (Miq. l. c.). Ramuli adpresse retrorsum ochraceo-hirsutuli. Limbi 5-6 poll. longi. Petioli 2-1 lin. longi.

*B. Stylus nullus.*

*α. Bractea conico-lanceolata.*

429. *P. SUBULATUM* (Cas. DC. in Seem. journ. 1866. p. 165), foliis breviter petiolatis ample ovatis apice acuminatis, acumine longe subulato, basi minime inæquali cordulatis supra parce pilosis subtus densius et præsertim ad nervos fusce pilosis firmulo-membranaceis crebre pellucido-punctulatis, nervis subtus prominentibus centrali ad  $\frac{1}{2}$  longitud. usque nervos alternos subadscendentes utrinque 7-8 quorum medii utrinque 2 ad apicem sequentes alte supra medium ducti infimi subtiles mittente, petiolo dense fusce villosa, pedunculo dense fusce villosa petiolum æquante, amento florente quam folium multoties breviora, bractea conico-lanceolata carnosa apice acutiuscula adpresse hirtella, antheræ loculis discretis connectivo supra loculos apiculato, filamento antheras superante. ♀ In Novæ Granatæ prov. Barbacoas (Triana, n. 18! in h. DC.). Dioicum, femina ignota. Ramuli dense fusce villosi. Limbi 0,3 longi, 0,03 lati. Petioli 0,01 longi. Amenta 0,02 longa. (v. s.)

430. *P. BULLOSUM* (Cas. DC. in Seem. journ. 1866. p. 165), foliis breviter petiolatis ovato-acuminatis, acumine longe cuspidato, basi parum inæquali cordulatis supra glabris subtus fusce villosis subcoriaceis opacis, centrali nervo subtus prominente ad  $\frac{1}{2}$  longitud. usque nervos subalternos subadscendentes utrinque 6 subtus prominulos quorum supremi ad apicem ducti mittente, petiolo fusce villosa, pedunculo fusce villosa petiolum æquante, amento quam folium multoties breviora, bractea conico-lanceolata glabra apice uncinato-acutata, antheræ connectivo supra loculos producta. ♀ In Novæ Granatæ prov. Barbacoas et Tuqueras alt. 1000 m. (Triana, n. 22! in h. DC.). Dioicum, femina ignota. Ramuli juniores villosi. Limbi 0,18 longi, 0,085 lati. Petioli 0,01 longi. Amenta florentia 0,025 longa, 0,004 crassa. (v. s.)

*b. Bractea rotundata centro peltata pedicellata.*

*1° Folia penninervia.*

431. *P. ANDROGYNUM* (Cas. DC. in Seem. journ. bot. 1866. p. 165), foliis longiuscule petiolatis ovatis apice attenuatis acutis? basi æquali subcordatis supra subtiliter hirtellis subtus ad nervos et nervulos adpresse fusce hirtellis rigidis opacis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervos alternos 8-9 subpatulo-adscendentes quorum supremi ad apicem ducti inferi magis approximati mittente, petiolo fere ad limbi basin usque vaginante dorso fusce hirtello, pedunculo fusce hirtello quam petiolus multoties breviora, amento basi hermaphrodito apice masculo, bractea brevissime pedicellata, staminibus 2 lateralibus et in floribus inferioribus hermaphroditis ovario sub-ovato glabro, stigmatibus 3 brevibus. ♀ In Novæ Granatæ prov. Pasto alt. 2600 m. (Triana, n. 51! in h. Triana). (v. s.)

432. *P. TEYSMANII*, foliis breviter petiolatis acutis vel elliptico-ovatis apice breviter acuminatis basi cuneato-rotundatis vel acutiusculis chartaceis supra glabris subtus præsertim ad nervos puberulis, nervis utrinque 5 erecto-patulis arcuatis supremis ad  $\frac{3}{4}$  longitud. circ. solutis per anastomoses ad apicem subcontinuatissimis nervulis transversis, petiolo glabro, pedunculo petiolum circiter æquante, amento cylindrico crasso, bractea peltata, bacca apice acutiuscula. ♀ In Celebes borealis prov. Menado (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 141). Chavica Teysmannii Miq. l. c. Dioicum. Ramuli glabri. Folia

5  $\frac{1}{2}$ –5 poll. longa, 3–2  $\frac{1}{2}$  lata. Petioli 2–3 lin. longi. Amenta feminea 2  $\frac{1}{2}$ –1 poll. longa.

433. *P. CELEBICUM* (Bl. in Verh. der bat. Genoots. v. 11. p. 157), foliis breviter petiolatis oblongo-elliptico-lanceolatis apice acuminatis, acumine acuto, basi leviter inæquali acutis supra glabris subtus pubescentibus, centrali nervo nervos alternos adscendentes utrinque 5 quorum supremi ad apicem ducti mittente nervulis subtransverse parallelis, petiolo dorso pubescente, pedunculo petiolum æquante, amento cylindrico florente quam folium multoties brevior, bractea oblongo-rotundata centro peltata pedicellata, staminibus 2, rhachi albo-pubescente. ♀ In sylvis ins. Celebes monte Lukkou (Miq. in Linn. v. 21. p. 484). — Bl. enum. pl. Jav. fasc. 1. p. 64. *P. ellipticum* Roem. et Schult. (fide Miq. l. c.). *Chavica celebica* Miq. syst. p. 276, illustr. p. 43. t. 41, flor. ind. bat. v. 1. pars 2. p. 446. Dioicum. Ramuli squamulis fuscis conspersi. Limbi 0,13–0,18 longi, 0,06–0,07–0,08 lati. Petioli 0,015–0,01 longi. (Ex Miq. l. c.)

434. *P. AUSTRO-CALEDONICUM*, foliis petiolatis subrotundis apice breviter obtusiuscule apiculatis basi leviter inæquali rotundatis utrinque glabris rigidule membranaceis subpellucidis subtiliter pellucido-punctulatis, nervis subtus vix prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervos alternos subadscendentes utrinque 4 quorum supremi per anastomoses ante apicem evanidi infimi subtiliores e basi soluti mittente, petiolo glabro ad  $\frac{1}{2}$  longitud. usque vaginante glabro, pedunculo glabro petiolum subæquante, amento quam folium  $\frac{1}{2}$  brevior, stirpis masc. bractea rotundata breviter pedicellata pedicello basi pilosulo, staminibus 3, filamentis brevibus, stirpis fem. bractea ut in mare sed longiuscule pedicellata, ovariis lateraliter arcte compressis unde ad aspectu immersis, stigmatibus 3 abbreviatis. ♀ In Nova Caledonia (Vieillard n. 1227! in h. DC. stirp. fem. et h. Lenormand! stirp. masc. et Forster! in h. DC. ex h. Brit. Mus.). *P. Siriboa* Forst. l. prodr. n. 19. Dioicum. Ramuli glabri. Limbi 0,09 longi, 0,07 lati. Petioli 0,017 longi. (v. s.)

435. *P. CORYLISTACHYON*, foliis petiolatis oblongo-lanceolatis apice acuminatis, acumine acuto, basi inæquali subattenuato-obtusiusculis utrinque glabris firmiter membranaceis subtiliter pellucido-punctulatis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervos alternos adscendentes nervulis validioribus intermixtos utrinque 3–4 mittente, petiolo glabro, pedunculo glabro quam petiolum duplo brevior, amento quam folium duplo brevior, bractea rotundata centro peltata pedicellata glabra, staminibus 3, antheris oblongis, connectivo supra loculos peltatim producto. ♀ In ins. Philippinis (Cuming n. 1044! in h. DC.). *Chavica corylistachya* Miq. l. syst. p. 281, flor. ind. bat. v. 1. pars 2. p. 447. Dioicum. Ramuli glabri cinerei. Limbi 0,12 longi, 0,045 lati. Petioli 0,01 longi. (v. s.)

♂, *magnifolium*, foliis 0,2 longis, 0,065 latis, amentis 0,065 longis. — In ins. Philippinis (Cuming n. 1813 fide Miq. in Hook. Lond. journ. bot. v. 4. p. 433). *Chavica corylistachya* Miq. l. c.

436. *P. FUTOKADSURA* (Sieb. flor. jap. fam. nat. p. 107), foliis longiuscule petiolatis in mare ovato-acuminatis, acumine acuto, basi æquali rotundatis leviter cordulatis in femina lanceolatis ovato-lanceolatisve apice acuminatis, acumine acuto, basin versus subattenuatis basi ima acutiusculis acutisve vel obtusis omnibus utrinque glabris rigido-subcoriaceis opacis pellucido-punctulatis septuplinerviis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervum utrinque unum ad apicem ductum adscendentem subopposite et e basi utrinque 2 quorum infimi subtiles mittente, petiolo glabro basi vaginante, maris pedunculo quam petiolum paulo brevior glabro, amento folium æquante densifloro, bractea rotundata centro peltata subsessili glabra margine parce lacero-ciliolata, staminibus 2, filamentis brevissimis, rhachi hirtella, femine pedunculo glabro petiolum æquante, amento quam folium brevior, bractea ut in mare, rhachi hirta, bacca globosa fusco-flavicante. ♀ In Nagasaki (Oldham n. 488! et 669! in h. monac. et h. DC. et Bürger! in h. monac.). *P. foliis septemnerviis* Thunberg flor. Jap. pl. obs. p. 351. *P. arcuatum* forma japonica Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 135. Dioicum. Rami ramulique glabri. Limbi 0,1 longi, 0,04 lati. Petioli 0,025 longi. (v. s.)

437. *P. CHABA* (Bl. ! Verh. der bot. Genoots. p. 168), foliis breviter petiolatis oblongo-elliptico-lanceolatis apice subito acuminatis, acumine acutiusculo mucronulato, basi attenuatis acutis utrinque glabris pellucido-punctulatis membranaceis, nervis subtus prominulis centrali utrinque supra basin usque ad  $\frac{1}{4}$  longitudinis nervos 1-2 adscendentes suboppositos nervumque subtiliorem e basi mittente, petiolo glabro basi ima concavo, pedunculo glabro petiolum multum superante, amento brevi cylindrico crasso, bractea rotundata centro peltata brevissime pedicellata, ovario rhachi immerso, stigmatibus 3-4, baccis obovato-angulosis rubentibus maturitate coalitis glabris. ♀ In Borneo (Korthals! in h. DC. ex h. lugd. bat.), Borneo australi ad Martapoera et ad flumen Doesson et prov. Menado (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 136 et 140), ins. Luzon (Hænk. ! in h. Berol.), Java (h. Berol. ex h. lugd. bat.). — Bl. enum. pl. Jav. v. 1. p. 66. *P. abbreviatum* Opiz ! in Reliq. Hænk. fasc. 3. p. 157. *Chavica Chaba* Miq. ! syst. p. 251, flor. ind. bat. v. 1. pars 2. p. 442. Dioicum. Ramuli glabri. Limbi 0,1 longi, 0,045 lati. Petioli 0,014 longi. Pedunculi 0,02 longi. Amenta matura 0,012 longa, 0,006 crassa. (v. s.)

438. *P. SEEMANNIANUM*, foliis oblongo-ovatis apice acuminatis, acumine acuto, basi inæquali rotundatis subattenuatis utrinque glabris rigidis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos alternos sursumque tota longitudine nervulos validos mittente, petiolo glabro, pedunculo glabro petiolum paullo superante, amento cylindrico sessili, bractea rotundata centro peltata pedicellata subtus pilosa, staminibus 2. ♀ In Nova Irlandia (Barclay n. 3515 ! in h. Brit. Mus.). Dioicum. Ramuli glabri. Limbi 0,18 longi, 0,09 lati. Petioli 0,01 longi. (v. s.)

439. *P. MACULATUM* (Bl. in Verh. der bot. Genoots. v. 11. p. 199. fig. 20), foliis petiolatis oblongis apice breviter acuminatis, acumine acuto, basi suboblique cordatis interdum subpeltatis coriaceis glabris superioribus multo angustioribus, nervo centrali nervos utrinque 7 quorum inferiores oppositi superiores alterni supremi apice proximi et ad hunc ducti mittente, pedunculo quam petiolus brevior, stirpis masc. bractæ peltatæ, pelta ambitu pubescente, staminibus 2. ♀ In Java supra montem calcareum apud prædium Tjampia provinc. Buitenzorg (Bl. l. c.). — Bl. enum. pl. Jav. fasc. 1. p. 70. *Chavica maculata* Miq. syst. p. 256, flor. ind. bat. v. 1. pars 2, p. 443. Dioicum, scandens. Rami juniores purpureo-maculati. Limbi  $4\frac{1}{2}$ -8 poll. longi,  $2\frac{1}{2}$ -3  $\frac{1}{2}$  lati. Petioli inferiores 1-2, superiores  $\frac{1}{2}$  poll. longi. Pedunculi  $\frac{1}{2}$  poll. longi. (Ex Bl. l. c.)

## 2° Folia multiplinervia.

440. *P. AMBOINENSE*, foliis breviter petiolatis e basi subdimidiato-cordata vel rotundata ovato-oblongis summis sublanceolatis omnibus apice breviter acuminatis chartaceis pallide fusciscentibus 11-7-plinerviis, nervis plerisque e basi solutis supremo utrinque ad  $\frac{1}{2}$ - $\frac{1}{2}$  longitud. a basi libero ad apicem producto, amento masculo longe pedunculo elongato, bractæ pelta glabra pedicello rhachique hirtellis, stigmatibus 4-5 brevissimis, baccis glabris coalitis. ♀ In Amboinæ sylvis (Forsten fide Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 134), Ternate (Miq. l. c. p. 140). *Chavica amboinensis* Miq. l. c. Dioicum. Petioli 3-1  $\frac{1}{2}$  lin. longi. Folia 11-6 poll. longa,  $4\frac{1}{2}$ -1  $\frac{1}{2}$  lata. Pedunculi 1-1  $\frac{1}{2}$  poll. longi. Amenta femin. 3  $\frac{1}{2}$ -5 poll. longa. Pedunculi 2-3 poll. longi.

β, *latifolium*, foliis e basi inæquali cordata ovatis acuminatis 11-7  $\frac{1}{2}$  poll. longis, 6  $\frac{1}{2}$ -5 latis, nervis lateralibus 2 ad  $\frac{1}{2}$  longitud. liberis reliquis e basi vel prope basin liberis. In Ternate (Miq. l. p. 140). *Chav. amboinensis* β, *latifolia* Miq. l. c.

441. *P. HONGKONGENSE*, foliis breviter petiolatis oblongo-ellipticis apice obtuse attenuatis basi subæquali cordatis supra glabris subtus molliter pubescentibus rigido-subcoriaceis supra bullatis opacis epunctulatis septuplinerviis, nervis subtus prominulis centrali utrinque paullo supra basin nervum adscendentem ante apicem evanidum opposite mittente, cæteris nervis e basi ortis mediis adscendentibus infimis arcuatis, petiolo dense pubescente, pedunculo pubescente petiolum æquante, amento masculo florente quam folium multo-

ties brevior cylindrico densifloro, rhachi hirtella, bractea rotundata centro peltata breviter pedicellata, staminibus 2 anthera filamentum æquante. ♀ In Hongkong (Seemann! in h. DC.). Rami ramulique hirtello-pubescentes. Limbi 0,1 longi, 0,037 lati. Petioli 0,005 longi (v. s.)

442. P. MALAMIRI (Bl.! in Verh. bat. Genoots. v. 11. p. 203. fig. 22), foliis petiolatis ovato-oblongis aut oblongo-lanceolatis apice acuminatis acutis basi leviter inæquali rotundatis cordulatisve supra parce pilosulis subtus densius molliter stellatim pubescentibus quintupli-septupli-nerviis, nervis subtus prominulis centrali nervos utrinque adscendentes ad apicem ductos mittente cæteris e basi ortis, petiolo ad limbi latus majus vaginante basi dorso et apice utrinque fusce hirtio, stirpis masculæ amento graciliore bractea oblongo-rotundata glabra centro peltata pedicellata, rhachi lana floccosa alba pubescente, staminibus 2, filamentis brevissimis, stirpis femineæ amento paullo brevior crassiore, bractea longius rotundata centro peltata longius pedicellata, stigmata 3-4, bacis minute obovato-angulosis densissime confertis alioquin coalitis aromaticis. ♀ In Javæ sylvis humidis imprimis montanis (Bl.! l. c. et spec. sterile in h. DC. ex h. lugd. bat.), Noessa et Kambanga (Miq. in flor. ind. bat. v. 1. pars 2. p. 442). — Bl. enum. pl. Jav. fasc. 1. p. 70 (non L.). Chavica Blumei Miq. syst. p. 246, flor. ind. bat. l. c. Ramuli pilis crispulis pubescentes. Limbi 0,17 longi, 0,07 lati. Petioli 0,01 longi. (v. s.)

β, *cordatum* (Miq. in Ann. Mus. lugd. bat. v. 1. fasc. 5. p. 136), foliis basi perspicue cordatis. In Sumatra orientali, Borneo australi in Poeloe Lampe.

γ, *hirtellum* (Miq. in Mus. l. c.), foliis densius hirtellis. In monte Pamaton (Korthals fide Miq. l. c.).

443. P. FORSTENII, foliis breviter petiolatis e basi dimidiato-cordata ovato-oblongis apice breviter acuminatis pergamaceis lucidis siccis flavescenti-pubescentibus 5-7-plinerviis, nervo utrinque uno supra basin libero cæteris e basi ortis, amentis elongatis, pedunculo petiolum circiter æquante, bractæ pelta glabra pedicello hirtello. ♀ In sylvis Amboinæ (Forsten fide Miq. in Ann. Mus. lugd. bat. v. 1. fasc. 5. p. 134), ins. Halmakeira (Miq. l. c. p. 140). Chavica imperialis Miq. l. c. Glabrum, ramulis ut videtur herbaceis. Petioli circ. bipollicares. Folia 15-13 poll. longa, 6-4½ poll. lata. Pedunculi tripollicares. Amenta masc. pedalia, crassitie pennæ corvinæ. Folia extant æqualiter cordata probabiliter plantæ sterilis vel novellæ.

444. P. PHILIPPINUM (Miq.! syst. p. 322), foliis breviter petiolatis ovato-ellipticis apice brevissime acuminatis, acumine acutiusculo, utrinque glabris siccis supra nitidulis coriaceis opacis epunciculatis septuplinerviis, nervis subtus prominulis centralibus 3 ad ½ longitud. supra basin solutis infimis subtilioribus, petiolo glabro, stirpis masc. amento densifloro quam folium brevior rhachi hirtella foveolata, bractea rotundata centro peltata breviter pedicellata subtus pilosula, staminibus 2. ♀ In ins Philippinis (Cuming n. 912! in h. Deless.). Dioicum. Ramuli glabri. Limbi 0,13 longi, 0,08 lati. Petioli 0,01 longi. Amenta florentia 0,002 crassa. (v. s.)

445. P. BØHMERIEFOLIUM (Wall.! list), foliis brevissime petiolatis oblongo-lanceolatis apice longe acuminatis, acumine acuto mucronulatoque, basi valde inæquali, latere majore auriculæformi minore acutiusculo, utrinque glabris membranaceis, centrali nervo usque ad ½ longitud. nervos subalternos adscendentes utrinque 4-5 quorum 3 utrinque non procul a basi soluti et infimi subtiliores mittente, petiolo glabro, pedunculo petiolum multum superante glabro, stirpis masc. amento maturo quam folium paullo brevior, bractea rotundata centro peltata breviter pedicellata, stam. 2, filamentis brevibus, stirpis femin. bractea ut in mare, bacca subglobosa glabra. ♀ In Moluccis (Lobb! in h. Kew.), Birma (Griffith n. 4404! et 4416! in h. Kew.), Silhet (Wallich n. 6654 a! in h. Wall. et DC.), mont. Khasia (Hook. et Thoms.! in h. Kew.). Chavica bœhmeriæfolia Miq.! syst. p. 265, illustr. p. 41. t. 37. Dioicum. Rami ramulique glabri. Limbi 0,17 longi, 0,045 lati. Petioli vix 0,005 longi. Amenta femin. matura 0,1 longa. Bacca 0,004 crassa. (v. s.)

- ß, *firmius*, foliis apice acuminatis, acumine acuto, firmioribus. — In Amboina (Dollesh. n. 203! in h. Vindob.)
- γ, *tenuifolium*, foliis tenuius membranaceis infimis basi valde inæquali semicordatis. — In Moluccis (Roxb.! in h. Deless. fide Miq. syst. p. 264). *P. lanceolatum* Roxb. flor. ind. v. 1. p. 150 (non Ruiz et Pav.). Chavica lanceolata Miq. syst. p. 264, illustr. p. 41. t. 36.
446. *P. RAFFLESII*, foliis petiolatis lato-ellipticis vel subovato-ellipticis apice subacutis basi subæquali cordatis supra præter basin nervorum mediorum glabris subtus ad nervos puberulis rigido-membranaceis, centrali nervo ad  $\frac{1}{2}$  longitud. nervos utrinque 9-10 quorum supremi ad apicem ducti et infimi subtiliores mittente, petiolo sparse puberulo. ♀ In ins. Sumatra (Raffles! in h. Deless.), Benekoglen (Miq. in Sumatra p. 186). Chavica Rafflesii Miq.! syst. p. 273, flor. ind. bat. v. 1. pars 2. p. 446, Sumatra l. c. Ramuli fusc. Limbi 0,27-0,29 longi, 0,16-0,17 lati. Petioli 0,025 longi (v. s.)
447. *P. SULCATUM* (Bl.! in Verh. bat. Genoots. v. 11. p. 158. f. 2), foliis breviter petiolatis inferioribus ovatis superioribus in mare suboblique elliptico-oblongis in femina suboblique oblongo-ovatis omnibus apice acuminatis, acumine acuto, basi leviter inæquali inferioribus obtusis rotundatisve superioribus acutiusculis acutisve utrinque glabris membranaceis opacis subtiliter creberrime pellucido-punctulatis septupli-noveninerviis, nervis subtus subprominulis, centrali ad  $\frac{1}{2}$  -  $\frac{3}{4}$  longitud. usque nervum utrinque unum adscendentem ad apicem usque ductum mittente cæteris e basi solutis infimis subtilibus sæpe deficientibus, petiolo glabro ad limbi basin usque vaginante, pedunculo glabro petiolum paullo superante, amento masculo quam folium duplo breviori filiformi, bractea rotundata centro peltata glabra subsessili, staminibus 2, amenti feminei vel hermaphroditi bractea ut in masculo, ovario ovato apice acutato, stigmatibus 3 abbreviato-subglobosis, baccis discretis glabris lateraliter subcompressis. ♀ In ins. Java (Bl.! in h. Mart. spec. hermaph. et h. DC. ex h. lugd. bat. spec. masc. et fem.), sylvis montis Merapi Javæ arbores scandens (Miq. pl. Jungh. fasc. 1. p. 17). — Bl. enum. pl. Jav. p. 64, Miq.! syst. p. 334, flor. ind. bat. v. 1. pars 2. p. 453. *P. nigrescens* Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 137. Monoicum vel dioicum. Rami ramulique glabri. Limbi 0,11 longi, 0,05 lati. Petioli 0,01 longi. Amenta matura 0,004 crassa. (v. s.)
448. *P. ACRE* (Bl.! in Verh. der Bat. Genoots. v. 11. p. 185. f. 13), foliis petiolatis inferioribus oblongo-ovatis apice acuminatis, acumine acuto, basi leviter inæquali rotundatis superioribus oblongo-ellipticis apice acuminatis, acumine acuto, basi subattenuato-acutis supra glabris subtus præsertim ad nervos molliter pubescentibus coriaceis opacis quintuplinerviis, nervis subtus prominulis centrali nervum utrinque unum parum supra basin mittente cæteris e basi solutis ad medium usque ductis, petiolo ad  $\frac{3}{4}$  longitud. usque vaginante basi dorso sursum utrinque fusce tomentoso, pedunculo glabro petiolum æquante, amento masculo quam folium  $\frac{1}{2}$  brevior, bractæ rotundato-s-bquadrangularis centro peltatæ breviter pedicellatæ ciliolatæ pedicello fusce tomentoso, staminibus 2, rhachi fusce tomentella, filamentis antheras superantibus. ♀ In ins. Java (h. DC. ex h. lugd. bat.!), locis humilibus montis Salak ubi arbores adscendit (Bl. l. c.). — Bl. enum. pl. Jav. fasc. 1. p. 68; Miq.! syst. p. 325, flor. ind. bat. v. 1. pars 2. p. 453. Dioicum. Rami ramulique glabri, parce verruculosi. Limbi 0,13 longi, 0,065 lati. Petioli 0,008 longi. (v. s.)
449. *P. KHASIANUM*, foliis petiolatis elliptico-lanceolatis vel oblongo-elliptico-lanceolatis apice acuminatis, acumine acuto mucronulatoque, basi inæquali attenuato-acutis utrinque glabris membranaceo-firmis pellucido-punctulatis septuplinerviis, nervis subtus prominulis 3 centralibus ad apicem ductis infra  $\frac{1}{2}$  longitud. supra basin solutis, proximis e basi solutis sequentibus supra medium ductis, nervulis subtransversim parallelis, petiolo glabro, pedunculo glabro petiolum superante, stirpis masc. amento densifloro filiformi, bractea rotundata centro peltata brevissime pedicellata glabra, rhachi hirtella stam. 2-3 raro 4, filamentis antheras superantibus, stirpis femina. folia ut in mare amento quam folium triplo brevior, pedunculo petiolum paullo

superante, rhachi hirsuta, bractea ut in mare, bacca ovata. ♀ In mont. Khasia et Chittagong (Hook. et Thoms.! n. 5 in h. Kew.), India orient. (Griffith n. 4421! in h. Kew.). Dioicum. Rami glabri. Limbi 0,09 longi, 0,035 lati. Petioli 0,01 longi. Pedunculi 0,015 longi. (v. s.)

450. *P. MAJUSCULUM* (Bl. in Verh. der Bat. Genoots. v. 11. p. 210), foliis brevissime petiolatis oblongo-ellipticis vel oblongo-obovato-ellipticis apice acuminatis, acumine obtusiusculo, basi inæquali cordato-rotundatis, latere majore auriculæformi, utrinque glabris membranaceis epunctulatis 13-16-plinerviis, nervo centrali ad  $\frac{1}{2}$  longitud. usque nervos supra basin utrinque 3 alternos adscendentes mittente, cæteris e basi solutis, nervulis subtransversim parallelis, petiolo glabro, pedunculo glabro petiolum multoties superante, amento folium æquante elongato cylindrico pendulo densifloro, bractea rotundata centro peltata glabra pedicellata subtus ad pedicellum pilosa, baccis coalitis, stigmatibus 3, stirpis masc. bractea ut in femina, staminibus 3-5 vel apice amenti 2. ♀ In Sumatra ad Palembang (Miq. Sumatra p. 186 et 473), Borneo australi (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 136), Java (Miq. in flor. ind. bat. v. 1. pars 2. p. 445, Zoll. n. 2336! in h. DC. et Bl. in h. DC. ex h. lugd. bat.), Amboina (Labillard.! in h. Deless. et h. lugd. bat. in h. DC. et Dolleshott n. 97! in h. Vindob.). Sirium decumanum Rumph. amb. v. 5. p. 451. t. 27? (nervatione diversum). *P. decumanum* L. sp. p. 41 (exclus. syn. Plum.), Lam. ill. p. 80, Hill. dict. v. 26. p. 17. t. 17. f. 37). *P. majusculum* Bl. enum. pl. Jav. fasc. 1. p. 71. *P. methysticum* Roxb. fl. ind. v. 1. p. 159 (exclus. syn. Linn.). *Chavica majuscula* Miq. syst. p. 271 et II. cc. *P. magicum* herb. Reinw. n. 1407 fide Miq. in Linn. v. 21. p. 484. Dioicum. Ramuli glabri. Limbi 0,26 longi, 0,16 lati. Petioli 0,01 longi. Pedunculi 0,1 longi. Amenta mat. 0,2 longa, 0,007 crassa. (v. s.)

451. *P. CERAMICUM*, foliis breviter petiolatis e basi æquali vel vix inæquali oblongo-vel ovato-ellipticis apice breviter acuminatis pergamineis glabris 9-7-nerviis vel 9-7-plinerviis, nervis 3 centralibus paulo supra basin solutis ad apicem ductis omnibus subtus valde prominentibus, amento masc. pedunculato, pedunculo petiolum superante gracili, bractea orbiculari glabra. ♀ In ins. Ceram (Forsten fide Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 125). *Chavica ceramica* Miq. l. c. Dioicum. Folia 9-4  $\frac{1}{2}$  poll. longa. Petioli 3-4 lin. longi. Amenta masc. nondum perfecta pollicaria. *P. miniatum* Bl. cui perquam simile distinguitur foliis basi ut plurimum latioribus et æqualibus vel quam brevissime inæqualibus. (Ex Miq. l. c.)

452. *P. LUZONENSE*, foliis petiolatis rotundato-ovatis apice acuminatis, acumine acuto, basi parum inæquali rotundatis cordatisve vel superioribus obtusis utrinque glabris subcoriaceis opacis 11-plinerviis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  -  $\frac{1}{4}$  longitud. usque nervos alternos subadscendentes per anastomos fere ad apicem ductos utrinque 3 mittente cæteris e basi solutis arcuatis, petiolo glabro ad  $\frac{1}{2}$  longitud. usque vaginante, pedunculo glabro quam petiolus duplo brevior, stirpis masc. amento densifloro florente quam folium duplo brevior, bractæ rotundatæ coriææ glabræ margine extenuatæ centro peltatæ pedicellatæ pedicello glabro, rhachi foveolata glabra, staminibus 2 immersis, anthera filamentum superante, connectivo supra loculos valde producto. ♀ In ins. Luzon (Jayer n. 722! in h. Berol.). Ramuli glabri. Limbi 0,135 longi, 0,1 lati. Petioli 0,02 longi. (v. s.)

453. *P. PEDICELLATUM* (Cas. DC. in Seem. journ. 1866. p. 164), foliis breviter petiolatis ovatis vel elliptico-ovatis apice acuminatis, acumine acutiusculo mucronulatoque, basi rotundatis obtusisve vel subattenuatis obtusiusculis utrinque glabris membranaceis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos subalternos utrinque 4 quorum inferiores utrinque 3 fere a basi soluti mittente, petiolo glabro, pedunculo glabro petiolum duplo superante, stirpis masc. amento gracili folium superante, bractea rotundata supra centrum peltata longiuscule pedicellata, staminibus 2, filamentis brevibus, stirpis femin. amento folium subæquante, bractea ut in mare, bacca obovato-subtetragona glabra. ♀ In Bengalia orientali (Griffith n. 4404! et 4418? in h. Kew.), Sikkim (Hook. et Thoms. in h. Kew. et DC.). Dioicum. Ramuli glabri. Limbi 0,07-0,13 longi.

0,04-0,06 lati. Petioli 0,005 longi. Pedunculi femin. 0,015 longi. Amenta matura  $\delta$ , 1 longa. Bacca 0,004 crassa. (v. s.)

454. *P. POTHOIDES* (Wall.! list n. 6639 A), foliis breviter petiolatis oblongis vel oblongo-lanceolatis apice acuminatis, acumine acuto mucronulatoque, basi subattenuato-acutiusculis cordulatisve subtus sparse hirtellis haud pellucidopunctulatis membranaceis septuplinerviis, nervis subtus prominulis centrali nervos fere e basi utrinque 2 subalterne mittente cæteri e basi salutis, nervulis subtransverse parallelis, petiolo glabro, amento femineo maturo leguminiformi, baccis subglobosis subcoalitis, stigmatibus 3-4.  $\delta$  In Chappedong (Wall. n. 6639 A! in h. Wall. et DC.). Chavica? pothoides Miq.! syst. p. 260, illust. p. 42. t. 38. Dioicum. Ramuli glabri. Limbi 0,19 longi, 0,055 lati. Petioli 0,01-0,015 longi. (v. s.)
455. *P. POTHIFORME* (Wall.! herb. n. 6639  $\beta$  aut 1675), foliis petiolatis oblongis apice acuminatis, acumine acuto mucronulatoque, basi inæquali rotundato-cordulatis utrinque glabris rigidis septuplinerviis, nervo centrali nervos utrinque 3 quorum supremi parum supra basin cæteri fere e basi orti adscendentes supremi fere ad apicem ducti mittente, petiolo glabro, amento maturo pulposo, bacca globosa, baccis plus minusve pulposo-coalitis.  $\delta$  In Chappedong (Wall. l. c.). Dioicum. Ramuli glabri. Limbi 0,23 longi, 0,06 lati. Petioli 0,02 longi. (v. s.)
456. *P. PSEUDO-CHAVICA*, foliis subsessilibus oblongo-ovato-ellipticis apice longe acuminatis, acumine acutiusculo, basi valde inæquali cordatis, latere minore subattenuato majore auriculæformi, utrinque glabris rigidis septupli-vel novenerviis, nervis subtus prominulis centrali paulo supra basin nervos adscendentes utrinque 2 quorum supremi ad apicem ducti mittente, petiolo dorso piloso, pedunculo glabro petiolum multoties superante, bractea rotundata centro peltata pedicellata, staminibus 2, filamentis brevissimis.  $\delta$  In insulis Philippinis (Cuming n. 1342! in h. Brit. Mus. et 1343! in h. Deless.). Chavica Lessertiana Miq.! syst. p. 270, flor. ind. bat. v. 1. pars 2. p. 443. P. Lessertianum Cas. DC. in Seem. journ. bot. 1866, p. 164 (non ejus supra, n. 67). Dioicum. Ramuli pilosi. Limbi 0,2 longi, 0,08 lati. Petioli vix 0,005 longi. Pedunculi 0,055 longi. (v. s.)
457. *P. MICROSTIGMA*, foliis breviuscule petiolatis e basi breviter inæquali obtusa vel subacuta æquilatere-oblongo-ellipticis superioribus sublancoelatis breviter acuminatis coriaceis glabris infra  $\frac{1}{4}$  longitud. a basi septuplinerviis, nervis lateralibus infimis exilibus supremis per anastomoses ad apicem ductis, amento femineo longe pedunculato, bractea pedicellato-peltata rhombo-elliptica, ovario obovoideo rhombeo-tetragono, stigmatibus 3-4 exilissimis ellipsoideis, amentis masculis gracilioribus, staminibus 2.  $\delta$  In Sumatræ occidentalis dist. Doekoe (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 138). Chavica microstigma Miq. l. c. Dioicum. Petioli 4-3 lin. longi. Folia sordide fuscescentia,  $5\frac{1}{2}$ -2  $\frac{1}{2}$  poll. longa. Pedunculi petiolum semper perspicue superantes  $1\frac{1}{4}$ -1  $\frac{1}{4}$  poll. longi. Amentum 3-4 poll. longum, femineum maturum penna corvina crassius.
458. *P. CHAWYA* (Ham.! in Wall. list, n. 6650 c!), foliis breviter petiolatis ovatis apice acuminatis, acumine acuto mucronulatoque, basi inæquali rotundatis utrinque glabris subcoriaceis septuplo-subnovenerviis, centrali nervo prope basin nervos 3 adscendentes quorum 2 fere basi soluti mittente, petiolo glabro, pedunculo glabro petiolum parum superante, amento femineo densifloro folium paulo superante, bractea rotundata centro peltata breviter pedicellata subtus pubescente, stigmatibus 4 filiformibus.  $\delta$  In Gongachora (Wall. n. 6650 c! in h. Wall.). Chavica Chawya Miq.! syst. p. 267, illustr. p. 42. t. 39. Dioicum. Ramuli glabri. Limbi 0,11 longi, 0,075 lati. Petioli 0,01 longi. (v. s.)
459. *P. ZOLLINGERIANUM*, foliis inferioribus longe superioribus breviter petiolatis inferioribus ovatis apice acuminatis, acumine acuto mucronulatoque, basi subcordatis, superioribus elliptico-lanceolatis basi inæquali attenuatis acutis utrinque glabris membranaceo-firmis pellucido-punctulatis septuplinerviis, nervis subtus prominulis, centralibus 3 ad apicem ductis ad  $\frac{1}{4}$  longitud.

subopposite solutis, sequentibus utrinque 2 e basi solutis supra medium exterioribus subtilioribus e basi solutis infra medium ductis, nervulis subtransverse parallelis, petiolo glabro, pedunculo glabro petiolum multum superante, amento quam folium brevius filiformi densifloro, bractea rotundata centro peltata sessili glabra, stigmatibus 3-4 lanceolatis, ovario ovato apice subattenuato, bacca subglobosa apice acutiuscula. ♀ In ins. Java (Zollinger n. 1322! et 1402! in h. Boiss. et DC.). *Chavica sarmentosa* Miq. in Linn. v. 21. p. 482. In Zoll. cat. p. 84 sub nomine pravo *P. diffusum* Vahl. Dioicum. Rami ramulique glabri. Limbi 0,07-0,1 longi, 0,03-0,055 lati. Petioli inferiores 0,035 longi. Pedunculi 0,02 longi. Petioli superiores 0,005 longi. Bacca 0,002 crassa. (v. s.)

460. *P. SARMENTOSUM* (Roxb. l. flor. ind. v. 1. p. 162), foliis inferioribus longe superioribus breviter petiolatis inferioribus lato-ovatis apice obtuse attenuatis basi cordatis superioribus ovatis apice breviter acuminatis, acumine acutiusculo, basi cordatis obtusiusve subattenuatisve utrinque glabris membranaceis vei membranaceo-firmis remotius pellucido-punctulatis quintuplinerviis, nervis subtus subprominulis centrali supra basin nervos 1-2 ascendentes quorum supremi fere ad apicem ducti mittente, lateralibus utrinque 1-2 e basi ortis brevioribus, petiolo glabro, pedunculo petiolum æquante, amento brevi cylindrico densifloro, bractea rotundata membranacea centro peltata pedicellata, baccis glabris dense approximatis, stigmatibus plerumque 4 oblongis. ♀ Femina in collibus calcareis Javæ et in sylvaticis prope Padang (Miq. pl. Jungh. fasc. 1. 16), in India or. (Roxb. l. in h. Mart. Herb. Willd. n. 682! spec. Klein), Sumatra (Miq. in Sumatra, p. 186 et 473, Borneo australi (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 186), Malabar (Wight fide Miq. in Hook. Lond. journ. bot. v. 5. p. 551). — Spreng. Syst. v. 1. p. 117, Wight icon. n. 1929 *P. diffusum*. Bl. fide Miq. in Linn. v. 20, p. 130. *Peperonia sarmentosa* A. Dietr. sp. v. 1. p. 161. *Chavica sarmentosa* Miq. l. Syst. p. 242, in Hook. Lond. journ. bot. v. 4. p. 433, flor. ind. bat. v. 1. pars 2. p. 441, Sumatra, p. 186 et 473. pl. Jungh. fasc. 1. p. 16, in Linn. v. 21. p. 482. Dioicum. Ramuli glabri. Limbi 0,13 longi, 0,11 lati. Petioli 0,06 longi. Amenta matura centim. longa. (v. s.)

461. *P. SUNDAICUM* (Bl. in Verh. der Bat. Genoots. v. 11. p. 187. fig. 14), foliis breviter petiolatis inferioribus ovatis vel oblongo-ovatis summis ellipticis omnibus breviter acuminatis, acumine acuto, basi æquali vel leviter inæquali rotundatis coriaceo-membranaceis utrinque glabris glandulis crebrius conspersis opacis supra saturate viridibus subtus discoloribus majoribus novem-nerviis stirpis masc. minoribus septuplinerviis omnibus præter 3 centrales ad  $\frac{1}{2}$  longitud. solutos ad apicem ductos e basi solutis, pedunculo petiolum circiter æquante, stirpis masc. amento graciliore, bractea orbiculari centro peltata breviter pedicellata supra glabra ad basin puberula, staminibus 2? stirpis femin. bractea rhomboideo-quadrangulari centro peltata pedicellata, ovario glabro, stigmatibus 3-4 lanceolatis brevibus acutis. ♀ In Moluccis, ins. Nussa Kambangan (Bl. l. c.), Amboina (Miq. flor. ind. bat. v. 1. pars 2. p. 446). — Bl. enum. Jav. fasc. 1. p. 68. *Chavica? sundaica* Miq. Syst. p. 274, in Linn. v. 21. p. 483. Dioicum. Ramuli glanduloso-punctati. Limbi 0,32-0,2 longi, 0,165-0,1 lati. Petioli 0,015 longi. (Ex Bl. l. c.)

462. *P. RHOMBOPHYLLUM*, foliis breviter petiolatis oblongo-rhombeis apice acuminatis, acumine obtusiusculo, basi cuneatis acutis utrinque glabris firmo-membranaceis pellucido-punctulatis quintuplinerviis, nervo centrali nervos utrinque 2 quorum infimi e basi soluti supremi supra basin soluti et ad apicem ducti mittente, petiolo glabro, pedunculo glabro petiolum superante, amento femineo quam folium multum brevius densifloro, bractea rotundata centro peltata pedicellata glabra ovario oblongo ovato glabro, stigmatibus plerumque 3 brevibus. ♀ In ins. Philippinis (Cuming n. 834! in h. Boiss. et Berol.). Luzon (Miq. flor. ind. bat. v. 1. pars 2. p. 442). Dioicum. Ramuli glabri. Limbi 0,08 longi, 0,04 lati. Petioli 0,004 longi. Pedunculi 0,01 longi. (v. s.)

463. *P. ARNOTTIANUM*, foliis breviter petiolatis oblongo-ellipticis apice acuminatis, acumine acuto, basi inæquali vel subinæquali obtusis cordulatisve utrinque glabris coriaceis novem-decuplinerviis, nervo centrali non procul a basi



nervos utrinque 2 adscendentes altiusque utrinque 2-3 minus adscendentes nervulosque mittente, petiolo glabro, pedunculo glabro petiolum duplo superante, amento femineo pedunculum paullo superante, bractea rotundata centro peltata, baccis sub-4-angulis subcoalitis, stigmatibus 3. ♀ In horto bot. Calcut. cultum (Wall. n. 6650 E et F! in h. Wall. et h. Finlays. n. 6651 al in h. Wall.), Travancore (n. 6651 β! in h. Wall.). P. Chuvya Hunter! in h. Finlays. *Chavica Arnottiana* Miq.! Syst. p. 268, illustr. p. 43. t. 40. Dioicum. Ramuli glabri. Limbi 0,11 longi, 0,045 lati. Petioli 0,005 longi. Amenta matura 0,03 longa, 0,005 crassa. (v. s.)

464. *P. PENANGENSE*, foliis breviter petiolatis oblongo-lanceolatis apice acuminatis, acumine acuto mucronulatoque, basi æquali vel subinæquali attenuato-acutis utrinque glabris membranaceis septuplinerviis, nervo centrali nervos utrinque 2-3 infimos e basi supremos ad  $\frac{1}{2}$  longitud. solutos fere ad apicem ductos mittente, petiolo glabro, pedunculo glabro petiolum multum superante, amento femineo ovato-globoso densifloro, bractea rotundata centro peltata breviter pedicellata, bacca globosa apice in stylum attenuata, stigmatibus 4 lanceolatis revolutis. ♀ In Penang (Wall. n. 6642 β! in h. Berol.). *Chavica Penangensis* Miq.! Syst. p. 279, illustr. p. 45. t. 44. Dioicum. Ramuli glabri. Limbi 0,1 longi, 0,035 lati. Petioli 0,004 longi. Pedunculi 0,01 longi. Bacca 0,002 crassa. (v. s.)

465. *P. SUBCORDATUM*, foliis modice petiolatis lato-ovatis apice breviter acuminatis sæpe quidquam inæquilateris, basi leviter obliqua rotundatis vel emarginatis subcoriaceis pellucido-punctulatis, 5 vel quintuplinerviis aut adjunctis nervis lateralibus exilissimis sub-7-nerviis nervis supremis per anastomoses ad apicem ductis, pedunculo quam petiolus multum brevior, amento gracili folium æquante vel superante, bractea irregulariter rotundata carnosula glabra subtus hirtella centro breviter pedicellata, staminibus 2. ♀ In Sumatra? (Miq. Ann. bot. mus. lugd. bat. v. 1. fasc. 5. p. 138). *Chavica subcordata* Miq. l. c. ? Dioicum. Ramuli sicci fusciscentes. Petioli pollicares et breviores. Folia supra atro-fusca, subtus pallide fusciscentia vel subglaucina, 4-3 poll. longa. (Ex Miq. l. c.)

466. *P. PHILIPPINENSE*, foliis petiolatis ovatis apice acuminatis, acumine acuto, basi æquali rotundatis utrinque glabris rigidis opacis non pellucido-punctulatis novenerviis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervum utrinque unum adscendentem ad apicem ductum oppositè mittente, cæteris nervis e basi solutis, petiolo glabro ad medium usque vaginante, pedunculo glabro petiolum paullo superante, amento florente quam folium multoties brevior, bractea rotundata centro brevissimè pedicellata glabra, ovariis rhachi immersis coalitis apice liberis tomentosis, stigmatibus 4-6 abbreviatis linearibus. ♀ In Philippinis (Cuming n. 4851 in h. Boiss.). Dioicum. Rami ramulique glabri. Limbi 0,15 longi, 0,05 lati. Petioli 0,02 longi. Amenta florentia 0,002 longa. (v. s.)

467. *P. AUSTRALASICUM*, foliis breviter petiolatis ovatis apice breviter acuminatis, acumine obtuso, basi æquali inferioribus leviter cordatis superioribus rotundatis utrinque glabris rigidis subopacis septuplinerviis, nervis subtus prominulis centrali paullo supra basin nervum utrinque unum fere ad apicem ductum oppositè mittente cæteris e basi solutis, petiolo glabro ad  $\frac{1}{2}$  longitud. usque vaginante pedunculo petiolum æquante, stirpis masc. amento densifloro limbum æquante, bracteæ rotundatæ centro peltatæ pelta margine lacero-ciliata, rhachi pilosula, staminibus 3 anthera filamentum superante. ♀ In Australasia (Hügel! in h. Vindob.). Rami glabri. Limbi 0,055-0,07 longi, 0,03-0,055 lati. Petioli 0,01 longi. (v. s.)

468. *P. PARAMATTENSE*, foliis breviter petiolatis ellipticis apice breviter acuminatis, acumine obtuso, basi æquali attenuato-acutiusculis utrinque glabris membranaceis subopacis epunctulatis, nervis subtus subprominulis septuplinerviis nervo centrali paullo supra basin nervum utrinque unum oppositè mittente cæteris e basi solutis, petiolo glabro, pedunculo glabro petiolum æquante, amento folium dimidium æquante, bracteæ rotundatæ centro peltatæ pelta pedicellata margine lacero-ciliolata pedicello rhachique pilosis, staminibus 3, anthera filamentum superante. ♀ In Australasia inter Anglye et Paramatta

(Hügel! in h. Vindob.). *Chavica Hügeliana* Miq. in Linn. v. 26. p. 218? Dioicum. Caulis glaber herbaceus. Limbi 0,09 longi, 0,05 lati. Petioli 0,005 longi. (v. s.)

369. *P. INSECTIFUGUM*, foliis petiolatis rotundato-ovatis apice acuminatis basi æquali rotundatis subtruncatisve utrinque glabris membranaceis subpellucidis pellucido-punctulatis novenerviis reticulato-nervulosis nervo centrali ad  $\frac{1}{2}$  longitud. usque alterne nervum utrinque unum subadscendentem ante apicem evanidum sursumque nervulos multos validos mittente lateralibus nervis utrinque 3 e basi solutis, petiolo basi subtilissime puberulo basi ima vaginante, maris amento quam folium brevius filiformi, pedunculo glabro quam petiolus brevius, bractea rotundata centro peltata brevissime pedicellata glabra coriacea, staminibus 3, filamento brevi.  $\frac{3}{5}$  In ins. Fijee (Seemann. n. 569! in h. DC.). — Seem. in flor. Viten. p. 262. Wa-Gawa ab incolis dictum. Dioicum. Ramuli glabri. Limbi 0,115 longi, 0,1 lati, petioli 0,016 longi. (v. s.)

### 3° Folia multinervia.

470. *P. METHYSTICUM* (Forster! pl. esc. p. 76), foliis longe petiolatis rotundato-ovatis apice breviter acuminatis, acumine acuto, basi profunde cordatis utrinque glabris ad nervos minutissime puberulis membranaceis 11-13-nerviis, nervis subius prominulis 3 centralibus ad apicem ductis, petiolo glabro, stipulis masc. amento cylindrico quam folium multoties brevius, bractea rotundata margine eroso-subcrenata centro peltata pedicellata, pedicello hirtello, staminibus 2, anthera subglobosa quam filamentum brevius.  $\frac{3}{5}$  In ins. Otahiti (Moerenhout! in h. DC.; Forster in h. Brit. mus.), Nukahiva (Jardin n. 73! in h. Lenormand), Oahu (Chamisso! in h. Berol.), Woahoo (Mac-Rae! in h. Mart.), Sandwich (Seeman n. 1709! in h. DC.). — Forster prodr. n. 21, Parkins draw. of Tahit. pl. ined. in Brit. Mus. t. 6, Willd. sp. v. 1. p. 161, Deless. ic. p. 53. t. 89, Guillem. Zephir. p. 28. *Macropiper latifolium* Miq. in Linn. v. 20. p. 130. *Macr. methysticum* Hook. et Arn. in Beech. bot. p. 96. *P. Ava*, kava ab incolis dictum liquorem inebriantem præbens. Dioicum. Ramuli fusco-maculati. Limbi 0,13 longi, 0,11 lati. Petioli 0,025 longi. (v. s.)

471. *P. RUMPHII*, foliis breviter petiolatis rotundato-ovatis apice subabrupte breviter acuminatis basi haud valde inæquali conniventi-cordatis chartaceo-pergamaceis 11-nerviis nervis 3 centralibus ima basi subconfluentibus nervulis transversis, pedunculo petiolum æquante, amento cylindrico quam folium  $\frac{1}{2}$  brevius, bractea ciliolata peltata subtus pubera, staminibus 2.  $\frac{3}{5}$  In Ternate (Miq. Ann. bot. mus. lugd. bat. v. 1. fasc. 5. p. 141). *Chavica Rumphii* Miq. l. c. Petioli  $1\frac{1}{2}$ -1 poll. longi. Dioicum. Folia 10-11 poll. longa, 7-8 poll. lata. Pedunculi  $1\frac{1}{2}$  poll. longa. Amenta masc.  $4\frac{1}{2}$  poll. longa calamus scriptorium crassa. Hæc species omnino refert iconem Rumph. Amb. 5. t. 27 (ex Miq. l. c.)

472. *P. REINWARDTIANUM*, foliis petiolatis ovato-rotundatis apice brevi acutis basi truncato-cordatis summis rotundatis membranaceis glandulosis glabris vel subtus parce pilosulis 9-7-nerviis, nervis 3 centralibus paullo supra basin solutis, petiolo juniore sparse piloso, amento femineo cylindrico, bacca obovato-pyramidata, stigmatibus 3 raro 4 brevibus.  $\frac{3}{5}$  In ins. Nussa Laut (Miq. in Linn. v. 21. p. 481), in Moluccis, Amboina (Forsten fide Miq. Ann. bot. mus. lugd. bat. v. 1. fasc. 5. p. 136), Ceram (Miq. l. c. p. 140). *Macropiper Reinwardtianum* Miq. in Linn. l. c., flor. ind. bat. v. 1. pars. 2. p. 438. *Chavica Reinwardtiana* Miq. in Ann. bot. l. c. Dioicum. Ramuli sub lente glandulosi. Folium majus 0,16 longum, 0,15 latum, minus 0,1-0,12 longum. Petiolus 0,025-0,05 longus. Amentum maturum 0,11 longum.

473. *P. MINIATUM* (Bl. ! in Verh. der nat. Genoots. v. 11. p. 166), foliis brevissime petiolatis oblongo-ovato-lanceolatis apice longe acuminatis, acumine acuto, basi rotundato-obtusis cordulatisve utrinque glabris subtus basi ad nervos puberulis pellucido-punctulatis membranaceo-rigidulis 5-7-nerviis, nervis subius prominulis 3 centralibus ad apicem usque ductis lateralibus subtilioribus, nervulis crebris subtransversis, petiolo utrinque puberulo hirtellove basi ima vaginante, pedunculo glabro petiolum multum superante,

stirpis masc. amento filiformi foliis multum superante, bractea rotundato-quadrangulati centro peltata breviter pedicellata glabra, staminibus 3 anthera subsessilibus, stirpis femin. amentis et bractea ut in mare pilis floccosis hic illic inter bracteas sparsis, baccis oblongo-cylindrico-pentagonis apice acutiusculis. ♂ In Borneo, Labuan (Molley n. 109! in h. Kew.), Philippinis (Cumming n. 841! in h. DC. et 1718! in h. Boiss. stirpis masc. et h. Berol. stirpis fem.), Java (Zollinger n. 1390! et Hofmannsegg n. 103! in h. DC.), Luzon (Hænke! in h. Berol. et Hænk. in h. monac.), Sumatra (Miq. Sumatrá p. 186 et 473), Singapore (Lobb n. 306! in h. Boiss. et Walker n. 284! in h. Kew.), Amboina (Dolleshatt! in h. Vindob.), Malacca (Griffith n. 4407! in h. Kew.), Ind. or. (Roxb. in h. Deless.). — Bl. enum. pl. Jav. fasc. 1. p. 65. *P. auriculatum* Bl. in Verh. der bot. Genoots. 1. c. p. 171, enum. p. 66. *P. glandulosum* Opiz! in h. Hænk. et Presl! in Reliq. Hænk. p. 158 et A. Dietr. sp. n. 2756. *P. lanceolatum* Roxb. flor. ind. bat. v. 1. p. 159. *Chavica miniata* Miq. syst. p. 234, illustr. p. 32. t. 28, flor. ind. bat. v. 1. pars 2. p. 440. *C. macrostachya* Miq. syst. p. 236. *P. canaliculatum* Klotzsch! in h. Berol. *Chavica lanceolata* Miq. syst. p. 264, flor. ind. bat. v. 1. pars 2. p. 445. *Dioicum*. *Ramuli glabri*. Limbi 0,12-0,15 longi, 0,045-0,06 lati. Petioli 0,003 longi. Pedunculi 0,02 longi. Amenta florentia 0,002 crassa (v. s.)

β, *brevispicatum*, foliis oblongo-ovato-ellipticis apice brevius acuminatis obtusiusculis 7-nerviis 0,19 longis, 0,08 latis, pedunculo petiolum multum superante, amento quam folium multum brevius 0,08 longo. ♂ In ins. Java (Kegel n. 68! in h. Berol.) (v. s.)

γ, *hirtellum*, ramulis novellis petiolisque rufo-hirtis, foliis subtus ad nervos adpresse albedo-pilosulis, bractea masc. medio pulverulenta. ♂ In Sumatræ occ. distr. Pantí (Miq. Sumatra p. 473). *Chavica miniata* β *hirtella* Miq. l. c.

474. *P. LONGUM* (L. l. sp. p. 41 exclus. syn. Rumph.), foliis inferioribus longe superioribus breviter petiolatis subsessilibusque inferioribus ovato-rotundatis apice breviter acuminatis, acumine acuto, basi lato-cordatis superioribus oblongo-ovatis apice attenuatis basi cordatis omnibus membranaceis subpellucidis utrinque glabris subtus ad nervos puberulis 5-7-nerviis, nervis subtus subprominulis, nervis 3 centralibus ad apicem ductis, petiolo puberulo, pedunculo petiolum æquante, amento masc. filiformi bractea oblongo-rotundata centro peltata breviter pedicellata glabra, staminibus 2 antheris subsessilibus, stirpis femin. amento crassiore brevius pedunculum circiter æquante, bractea oblongo-rotundata apice subattenuata centro subsessili glabra, stigmatibus 3-4 lanceolatis, bracteis baccis arcte applicatis, baccis densissime coalitis obpyramidato-ovatis tetragonis vertice liberis. ♂ In Timor (Decaisne herb. Tim. p. 169), Philippinis (h. DC l), locis calidioribus ins. Ceylon ubi rara (Thwaites enum. p. 292 et n. 2953! in h. DC.). Bengalía orientali (Griffith n. 4412! in h. Kew.), Henzeda fl. Irawaddi ubi vix non spontaneum (Wall. n. 6640 G! 6640 H stirpis masc. l. 6640 D! stirpis fem., 6640 J! cultum fem.), Mancha, Caopun (Wall. n. 6640 β! in h. Wall.), Gualpara (n. 6649 c! in h. Wall. utraque stirpis), Parauna (Wall. n. 6640 β! sterilis in h. Wall.), Silhet (n. 6640 F! in h. Wall.), Sikkim alt. 2000 ped. (Hook. et Thoms. l. in h. DC.), Malabar (Belanger n. 336! et 339! in h. Petrop.), montibus Nilgheriosis (Perrottet n. 1069! in h. DC.), Courtallum (Wight l. fide Miq. in Hook. Lond. journ. bot. v. 5. p. 551, icon. t. 1928). *P. longum* Pistolochiæ Pluk. ! almag. p. 297, phyt. t. 104. fig. 4, herb. Pluk. l. v. 97. fol. 105. Cattu Tropolí (Rheede v. 7. p. 27. t. 14). *P. foliis cordatis petiolatis sessilibusque* L. l. flor. Zeyl. n. 30. *P. longum* herb. Burman! v. 4, f. 3, Hill diet. v. 26. p. 17. t. 17. fig. 1, Vahl enum. v. 1. p. 334, Willd. ! spec. v. 1. p. 161, herb. Willd. n. 696! fol. 1, Roxb. flor. ind. v. 1. p. 156, Lour. flor. Cochinch. p. 40. *Chavica Roxburghii* Miq. syst. p. 239, illust. t. 30 icon. n. 256, in Hook. Lond. journ. bot. v. 4. p. 433, in Linn. v. 20. p. 130, flor. ind. bat. v. 1. pars 2. p. 440, Hayne Arnz. Gewach. v. 14. t. 20. *Dioicum*. *Ramuli glabri*. Limbi 0,06 longi, 0,065 lati. Petioli 0,07-0,08 longi. Amenta masc. 0,06 longa. (v. s.)

475. *P. SYLVATICUM* (Roxb. flor. ind. v. 1. p. 158), foliis petiolatis vel inferioribus longiuscule petiolatis inferioribus rotundato-ovatis apice acuminatis, acu-

mine acuto, basi æquali cordatis superioribus ovatis apice acuminatis, acumine acuto, basi rotundatis omnibus glabris 5-7-nerviis, petiolo ad limbum usque vaginante, amento masculo breviter pedunculato filiformi, staminibus 4. ♀ In septentrionalibus provinciis Bengalæ (Wight ic. t. 1930). *Chavica sylvatica* Miq. syst. 248. Folia 3-4 poll. longa, 2-4 lata. Petioli 1-2 poll. longi. (Ex Roxb. l. c.)

476. *P. BOIVINII* (Cas. DC. in Seem. journ. 1866, p. 164), foliis breviter petiolatis ovato-lanceolatis apice acuminatis, acumine acuto mucronulatoque, basi æquali rotundatis truncatisve, utrinque glabris membranaceo-rigidulis, 5-7-nerviis, nervis subtus prominulis 3 centralibus ad apicem ductis, petiolo glabro, pedunculo petiolum æquante, amento quam folium brevius, bractea rotundata glabra centro peltata breviter pedicellata, staminibus 2. ♀ In ins. Mohely (Boivin! in h. Brit. Mus.). Ramuli glabri. Limbi 0,08 longi, 0,05 lati. Petioli 0,006-0,007 longi. (v. s.)

477. *P. BURMANUM* (Miq. ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 141), foliis modice petiolatis ovato-rotundatis apice acuminatis basi breviter cordatis vel subacuminatis chartaceis utrinque glabris 5- sub-7-nerviis, nervis 3 centralibus ad apicem ductis, amento femineo brevissimo pedunculum circiter æquante, bractea peltata irregulariter orbiculata carnosula erosula glabriscula, stigmatibus 4 crasso-teretibus, baccis ovoideis. ♀ In Boerhoe (Miq. l. c.). Dioicum. Petioli  $1\frac{1}{4}$ - $1\frac{1}{2}$  poll. longi. Limbi  $\frac{1}{2}$  ped. longi,  $4\frac{1}{4}$ - $4\frac{1}{2}$  poll. lati. Amenta feminea 2-3 lin. longa, oligocarpa. (ex Miq. l. c.)

### c. Bractea rotundata, centro affixa, sessili.

#### 1° Folia penninervia.

478. *P. OFFICINARUM*, foliis brevissime petiolatis oblongo-ellipticis apice attenuatis subacuminatis, acumine acuto, basi subæquali vel inæquali attenuatis acutis vel cordulatis utrinque glabris coriaceis, nervis subtus prominulis centrali ad  $\frac{3}{4}$  longitud. usque nervos alternos utrinque 3 adscendentes nervulosque validos aliquot mittente, petiolo glabro, pedunculo glabro petiolum superante, stirpis masc. amento folium æquante cylindrico densifloro bractea glabra coriacea staminibus 2-3 antheris sessilibus rhachi glabra, stirpis femin. foliis plerumque majoribus amento quam folium multum brevius bractea ut in mare ovario rhachi immerso et cum ea coalito, baccis subglobosis coalitis stigmatibus 3. ♀ In Celebes ins. distr. Menado (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 140), Timor (Miq. l. c. et h. DC.!), ins. Poeloe Telor (forma amentis valde abbreviatis Miq. l. c.), Sumatra occidentali (Miq. l. c.), ins. Philippinis (Cumming n. 1248! in h. DC. et Berol. stirpis masc.), ins. Java (Zollinger n. 907! in h. DC. et Bl.! in h. DC. ex h. lugd. bat.), China (Miq. flor. ind. bat. v. 1. pars 2. p. 444), hort. Calcutt. cultum (h. monac.!). *P. longum* Rumph. Amb. v. 5. p. 333. t. 116. f. 1 (prava). Pharmacum magnum vulgare Rumph. l. c. p. 42. t. 26. f. 1. ? *P. Amalago* Rheede Mal. v. 7. p. 31. t. 16? (longitudine amentorum diversum). *P. Amalago* L. sp. p. 41 (exclusis syn. Sloane, Plukenet et Rheede?). *P. melamiris* Hill? dict. v. 26. t. 16. fig. 3. *P. longum* Piso mant. arom. p. 182 cum icone. *P. maritimum* Bl.! in h. lugd. bat. *P. longum* Bl.! in Verh. der bat. Genoots. v. 11. p. 197, enum. pl. Jav. v. 1. p. 70. *P. callosum* Opiz! (Haenke n. 73! in h. Mart.). *Chavica officinarum* Miq. l. syst. p. 256, illustr. p. 39. t. 34. *C. maritima* Miq. l. syst. p. 262, illustr. p. 40. t. 35. *C. Labillardieri* Miq. l. syst. p. 263 (h. Deless.). Dioicum. Ramuli glabri. Stirpis masc. limbi 0,085 longi, 0,035 lati. Petioli 0,003-0,006 longi, pedunculi 0,0012 longi, amenta florentia 0,003 crassa. Stirpis fem. limbi 0,14 longi, 0,05 lati, amenta matura 0,025 longa, 0,005-0,008 crassa. (v. s.)

#### 2° Folia multiplinervia.

479. *P. HYPOGLAUCUM* (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 139), foliis petiolatis ovato-ellipticis ellipticisve vel supremis elliptico-lanceolatis omnibus apice acuminatis basi acutis obtusiusculisve subcoriaceis subtus glaucinis supra glabris subtus ad nervum centalem pubescentibus infra  $\frac{1}{2}$

basi 7-pli-5-plinerviis, nervis supremis per anastomoses ad apicem ductis laxè nervulosis, petiolo præsertim dorso pubescente, stirpis femin. pedunculo petiolum superante, amento gracili, rhachi hirtella, bractea irregulariter angulato-rotundata margine ciliolata subtus hirtella centro peltata subsessili, ovario ovoideo, stigmatibus 4 lanceolato-ovatis crassiusculis, baccis ellipsoideo-globosis acutiusculis. ♀ In Sumatra? (Miq. l. c.). Dioicum. Ramuli præter nodos novellos glabriusculi. Limbi 4-3 poll. longi. Petioli 6 lin. longi. Amenta 2-3 poll. longa. Flores densi. Stigmata albedo-pubera.

480. *P. AURANTIACUM* (Wall. ! list p. 6658 A.), foliis petiolatis, stirpis femin. majoribus ellipticis apice longe acuminatis, acumine acuto mucronulatoque, basi acutiusculis, maris ellipticis vel elliptico-lanceolatis apice et basi ut in femina supra glabris subtus sparse pilosis subcoriaceis 5-pli-7-plinerviis, nervo centrali ab 1/2 longitud. trifido, cæteris nervis e basi solutis infimis subtilioribus brevissimis, petiolo glabro, pedunculo glabro petiolum circiter æquante, amento femineo maturo quam folium brevius densifloro, rhachi hirtella, stigmatibus 3, bacca obovato-subtetragona glabra, stirpis masc. amento gracili, bractea rotundata glabra centro peltata subsessili, staminibus 2, filamento antheram superante. ♀ In Javæ monte Ungarum, Merapi (Miq. pl. Jungh. fasc. 1. p. 16), peninsula Birmanica (Griffith ! in h. Kew.), Bengalia orientali (Griffith n. 4417 ! et 4424 ! in h. Kew.), Napalia (Wall. n. 6658 a ! in h. DC. et Kew.). *Chavica Wallichii* Miq. ! syst. p. 254, illust. p. 38. t. 33, flor. ind. bat. v. 1. pars 2. p. 143. Dioicum, scandens. Ramuli glabri. Limbi 0,07-0,1 longi, 0,04-0,05 lati. Petioli 0,015-0,025 longi. Bacca 0,005 crassa. (v. s.)

♂, *glabrum*. *Chavica Wallichii* ♂ glabra Miq. pl. Jungh. fasc. 1. p. 16, flor. ind. bat. v. 1. pars 2. p. 443.

481. *P. THWAITESII*, foliis petiolatis lanceolatis apice acuminatis, acumine acuto mucronulatoque, basi attenuatis acutis vel rotundato-obtusis supra glabris subtus pilosis firmo-membranaceis septuplinerviis, nervis subtus prominulis 3 centralibus ad apicem lateralibus 2 supra medium ductis infimis subtilioribus obsoletis, petiolo hirtello vel glabro, pedunculo glabro quam petiolum paullo brevius, amento quam folium brevius densifloro arcuato, bractea suboblongo-rotundata centro affixa fere omnino libera subtus rhachique tomentoso-hirtellis, ovario ovato, stigmatibus 3 lanceolatis. ♀ In ins. Ceylon (Thwaites n. 2178 ! in h. DC.). *P. arcuatum* Thwaites ! enum. p. 293, non Bl. Dioicum. Ramuli glabri. Limbi 0,06 longi, 0,03 lati. Petioli 0,015 longi. (v. s.)

482. *P. NIGRESCENS* (Bl. ! in Verh. der bat. Genoots. v. 11. p. 161), foliis breviter petiolatis oblongo-ovatis apice breviter acuminatis, acumine acuto, basi subattenuato-obtusis supra glabris subtus basin versus puberulis membranaceis subpellucido-punctatis quintuplinerviis, nervis subtus prominulis centrali supra basin nervos adscendentes utrinque 2 (infimos subtiliores) mittente, petiolo glabro, stirpis femin. pedunculo glabro petiolum duplo superante, rhachi hirtello-puberula, bractea rotundata centro sessili subsessilive, stigmatibus 3 linearibus, stirpis masc. amento quam folium brevius densifloro, bractea et rhachi ut in femina, staminibus 2, filamento antheram æquante. ♀ In ins. Java ad Gœnong Malang et mont. Salak (Miq. in Linn. v. 21. p. 485, Bl. ! in h. DC. ex h. lugd. bat., Zollinger n. 875 ! in h. DC.). — Bl. ! enum. Jav. v. 1. p. 65, Miq. ! syst. p. 325, in Hook. Lond. journ. bot. v. 4. p. 438, flor. ind. bat. v. 1. pars 2. p. 154. Dioicum. Ramuli glabri. Limbi 0,1 longi, 0,05 lati, Petioli 0,005 longi. Pedunculi 0,02 longi. (v. s.)

483. *P. NEPALENSE* (Miq. ! syst. p. 318), foliis petiolatis inferioribus oblongo-ovatis oblongive superioribus elliptico-lanceolatis apice acuminatis, acumine acuto mucronulatoque, inferioribus basi æquali rotundatis superioribus basi breviter inæquali acutis omnibus utrinque glabris subtiliter pellucido-punctulatis 7-pli-5-plinerviis, nervis subtus prominulis centrali supra basin nervum utrinque unum oppositè mittente, lateralibus nervis e basi solutis infimis subtilioribus vel deficientibus, petiolo glabro basi vaginante, pedunculo petiolum superante, stirpis masc. amento quam folium brevius, bractea rotundata vel suboblongo-rotundata centro affixa, rhachi hirtella, staminibus 3, filamento antheram æquante, stirpis femin. amento folium æquante, bractea ut in mare, bacca ovata apice attenuato-acuta, pericarpio glanduloso-rugoso.

♂ In Napalia (Wall. n. 6650 ♂) in h. Berol. stirps femin. et stirps masc. sine num. in h. DC. l.), Courtalloum (Wight, icon. v. 6 et t. 1936 ? haccis globosis diversa). Dioicum. Ramuli glabri. Limbi inferiores 0,12 longi, 0,045 lati. Petioli 0,008 longi. Baccæ 0,004 longæ. (v. s.)

♂, *parvifolium*. Limbi 0,07 longi. In monte Khasia, alt. 4000 ped. (Hook. et Thoms. ! in h. DC.)

γ, *crassifolium* ? foliis paulo crassioribus et validius nervulosis. In Khasia (Griffith in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 438). (Ex Miq. l. c.)

484. P. FRAGILE (Benth. in Hook. Lond. journ. bot. v. 2. p. 234), foliis petiolatis lato-ovatis orbiculatisve apice breviter acuminatis basi cuneatis truncatis vel subpeltatis coriaceis quintuplinerviis, stirpis masc. amento tenui, bractea peltata, staminibus 2, feminae amento quam in mare dimidio brevior, bracteis subconnatis peltatis, stigmatibus 5 recurvis breviter linearibus, haccis confertissimis depresso-globosis subconnatis. ♂ In Nova Guinea (Hinds fide Benth. l. c.). Chavica Benthamiana Miq. syst. p. 253, flor. ind. bat. v. 1. pars 2. p. 440, Benth. sulph. p. 181. Dioicum, scandens. Limbi 2-3-pollicares. Petioli semipollicares. Pedunculi semipollicares. (Ex Benth. l. c.)

485. P. LINXII (Miq. ! syst. p. 317), foliis inferioribus longius petiolatis omnibus elliptico-lanceolatis apice acuminatis acutiusculis basi attenuatis acutis utrinque glabris subcoriaceis tripli-quintuplinerviis, nervis subtus prominulis, centrali supra basin nervos alternos utrinque 1-2 supremos fere ad apicem ductos mittente, petiolo glabro, stirpis masc. pedunculo petiolum æquante, amento folium æquante, bractea obovata basi breviter acuminata centro rhachi adnata margine libera, rhachi glabra, staminibus 3, filamento antheram æquante, stirpis femin. amento et bractea ut in mare, baccæ ovata glabra apice mucronulata. ♂ In ins. Java (Schmidt ! in h. Berol.). — Miq. ! illust. p. 53. t. 53. Dioicum. Rami ramulique glabri. Limbi 0,11 longi, 0,45 lati. Petioli 0,01-0,02 longi. Baccæ 0,004 longæ. (v. s.)

486. P. JAYERI, foliis breviter petiolatis elliptico-oblongis apice acuminatis, acumine acutiusculo, basi æquali obtusis basi ima brevissime cordulatis utrinque glabris nitidulis rigidis opacis, septuplinerviis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervum adscendentem utrinque unum ad apicem per anastomoses fere ductum alterius sæpeque nervulos validiores mittente, lateralibus nervis utrinque 2 e basi solutis adscendentibus quorum medii per anastomoses supra medium ducti, petiolo glabro basi ima vaginante, pedunculo glabro petiolum æquante, stirpis masc. amento florente quam folium circiter duplo brevior densifloro, rhachi pubescente, bractea rotundata centro peltata subsessili glabra, staminibus 2, filamento brevissimo. ♂ In Luzon (Jayer n. 162 ! in h. Berol.). Dioicum. Ramuli glabri. Limbi 0,15 longi, 0,05 lati. Petioli 0,01 longi. (v. s.)

487. P. LONCHITES (Miq. ! syst. p. 331), foliis breviter petiolatis lanceolato-ellipticis apice acuminatis, acumine acuto, basi leviter inæquali inferioribus rotundatis superioribus attenuatis, omnibus utrinque glabris membranaceo-rigidulis subopacis subtiliter pellucido-punctulatis 5-pli-7-plinerviis, nervis subtus prominulis, centrali paulo supra basin nervum utrinque unum adscendentem, fere ad apicem ductum oppositè mittente, lateralibus nervis 1-2 e basi solutis quorum exteriores subtiliores marginales, petiolo villosa-pubescente vel in femina cito glabrato, stirpis masc. pedunculo glabro petiolum triplo superante amento tereti folium subæquante densifloro, bractea rotundata glabra subtus ad rhachin hirtella centro affixa, staminibus 3, filamento brevissimo, stirpis femin. pedunculo bractea et rhachi ut in mare, baccæ ovata glabra basi subimmersa (fid. Miq. l. c.). ♂ In India prope Penang (Wall. n. 6644 ♂ in h. Wall., et in h. DC. spec. masc.). Chavica penangensis Miq. in Hook. Lond. journ. bot. v. 4. p. 433. non syst. Dioicum. Ramuli glabri. Limbi 0,7-0,75 longi, 0,04-0,035 lati. Petioli 0,007 longi. Pedunculi 0,012 longi. (v. s. spec. masc.)

488. P. ARBORESCENS (Roxb. flor. ind. v. 1. p. 161), foliis breviter petiolatis oblongo-ovato-lanceolatis apice acuminatis, acumine acuto, basi subæquali obtusis vel rotundato-cordulatis utrinque glabris coriaceis quintuplinerviis vel

nervis infimis duobus sæpe adjunctis, septuplinerviis, nervis subtilis prominens centrali ad  $\frac{1}{2}$  longitud. usque nervum adscendentem utrinque unum alterne mittente, cæteris nervis e basi solutis, petiolo glabro, pedunculo glabro petiolum multum superante glabro, stirpis masc. amento folium superante, bractea rotundato-ovata centro peltata adnata sessili subtilis rhachique pilosulis, staminibus plerumque 3, filamento brevi, stirpis femin. bractea ut in mare, stigmatibus 4, bacca glabra oblongo-ovata apice acuta. ♂ In Borneo monte Sakœnbang (Miq. Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 137), ins. Java (Zollinger n. 1050 ! in h. DC. et Schmidt ! in h. Berol.), Poulo Pinang (Miq. flor. ind. bat. v. 1. pars 2. p. 452 stirpis femin. excepta), insulæ Ceylon prov. australi (Thwaites n. 2461 ! fide Thwaites enum. p. 293), montibus Nilghericis (Perrottet n. 1070 !, 1071 !, 1072 !, 1073 ! in h. DC.). Sirium arborescens tertium Rumph. Amb. v. 5. cap. 25. t. 28. fig. 1. P. arborescens Bl. in Verh. der bat. Genoots. p. 182. fig. 12, enum. pl. Jav. p. 67, Wight icon. t. 1940 (stirpis femin. excepta), Miq. syst. p. 320, illustr. p. 54. t. 54 (stirpis femin. bractea lineari-oblonga forsan ad P. nigrum referenda), flor. ind. bat. 1. c. Dioicum, alte scandens. Ramuli glabri. Limbi 0,9-0,105 longi, 0,03-0,04 lati. Petioli 0,005 longi. Pedunculi 0,02 longi. (v. s.)

489. P. BETLE (L. ! spec. p. 40), foliis petiolatis et longe petiolatis rotundo-ovatis vel superioribus ovatis omnibus apice acuminatis, acumine acuto, basi sub-inequali vel inequali inferioribus cordatis superioribus rotundatis omnibus utrinque glabris rigidis, 5-pli-9-nonerviis plerumque septuplinerviis, nervis subtilis prominulis, centrali ad  $\frac{1}{2}$  longitud. usque nervos  $\frac{1}{2}$  quorum supremi per anastomoses ad apicem ducti alterne mittente cæteris nervis e basi solutis, petiolo glabro ad  $\frac{1}{2}$ - $\frac{1}{4}$  longitud. usque vaginante, pedunculo glabro in mare petiolum æquante in femina petiolum superante, stirpis masc. amento quam folium triplo brevius, bractea rotundata vel obovata centro peltata sessili, rhachi hirtella, staminibus 2, filamento lato brevi, stirpis femin. amento quam folium multoties brevius, bractea ut in mare, ovario rhachi immerso et cum eo coalito apice tomentoso, stigmatibus 4-5 lanceolatis ♀ In Borneo australi (Korthals fide Miq. ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 136), Philippinis (Blanco fide Hasskarl flora 1864. p. 58), Hongkong (Miq. in Seem. bot. Herald p. 415), Java (Zollinger n. 752 ! et 2285 ! in h. DC. et Hoffmannsegg ! in h. Berol.), Sumatra (Miq. Sumatra p. 186 et 473), Ceylon (Thwaites n. 3692 ! in h. DC.), India orientali (Wall. n. 6652 ♂ ! in h. Wall. stirpis masc.), Silhet (Wall. n. 6652 ♂ !, 6659 ♂ ! ibid. stirpis masc.), Trogia (Wall. n. 6652 c !, 6653 a ! et ♂ ibid. sterilia), Canara (Hohenacker n. 491 ! in h. Boiss. stirpis femin.), Madagascar (Lindley ! in h. Mart. an spontanea ?), late per orientem a Borneo usque ad Bourbon cultum et in occident. insulis Guadaloupa (Balbis ! in h. DC.), Trinitate (Sieber n. 382 ! in h. DC. stirpis femin.) cultum. Herb. Willd. n. 686 !. Sirii folium Rumph. Amb. v. 5. p. 336. t. 116. fig. 2. Beetle Codi Rheede Mal. v. 7. p. 29, t. 15. P. Bette L. ! flor. zeyl. n. 27 h. Burm. v. 3 p. 32 et v. 4. f. 9 ! spec. p. 40, Hill dict. v. 26. p. 16. t. 16. fig. 2, Lam. ! illust. p. 79 (herb. Willd. 1. c.), Vahl enum. v. 1. p. 328, Willd. spec. v. 1. p. 159, Roxb. flor. ind. v. 1. p. 160, Loud. encycl. n. 507. p. 28, Bl. in Verh. der bat. Genoots. v. 11. p. 206, P. Melamiri L. sp. quoad syn. Rumph., Vahl enum. v. 1. p. 327. P. Melamiris Lam. ill. p. 79 ? Chavica Belle Miq. ! syst. p. 228, in Linn. v. 20. p. 130. flor. ind. bat. v. 1. pars 2. p. 439, Sumatra p. 186. Chavica auriculata Miq. syst. p. 269. Artanthe hexagyna Miq. ! syst. p. 412. Dioicum, scandens. Ramuli glabri. Limbi 0,08-0,11 longi, 0,045-0,09 lati. Petioli 0,005-0,015 longi. (v. s.)

♂, *Siriboa*, foliis membranaceo-rigidulis, junioribus supra subtusque pubescentibus, septuplinerviis, stigmatibus 3-6. ♂ Java (Bl. Verh. der bat. Genoots. v. 2. p. 208 et h. DC. ? ex herb. lugd. bat.), Ceylon (Thwaites n. 3690 ! in h. DC.), Bengalia orientali (Griffith n. 4415 ! in h. Vindob.). Betela quam Siri Boa vocant Bont. hist. nat. med. in Piso ed. 1658. p. 91. t. 91. *Siriboa* Rumph. Amb. v. 5. p. 340. t. 117. P. foliis cordatis subseptinerviis venosis L. ! flor. zeyl. n. 29 (herb. Burm. 3. f. 171 et h. Linn. l.). P. *Siriboa* L. ! spec. p. 41, Hill dict. v. 26. p. 17. t. 17. fig. 1, Lam. ill. n. 80, Vahl enum. v. 1. p. 332, Bl. Verh. der Bat. Genoots. v. 11. p. 207, enum. pl. Jav. p. 71. *Chavica Siriboa* Miq. syst. p. 224, in Hook. Lond.

journ. bot. v. 4. p. 433, flor. ind. bat. v. 1. pars 2. p. 438. Dioicum, scandens. Limbi 0,13 longi, 0,075 lati. Petioli 0,025 longi. Pedunculi 0,035 longi. (v. s.)

γ, *densum*, foliis magis ovatis basi magis inæqualibus 5-pli-7-plinerviis membranaceis non pellucido-punctatis subtus ad nervos subtilissime puberulis, stam. 2, stigmatibus 5-7. ♂ In Borneo australi Ternate (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 140), Luzon (Haenke n. 69! in h. Vindob.), Sumatra occ. distr. Siboga prope Batas Sangka, Sungi Pagu (Miq. Sumatra p. 186 et 473), Java (Zolling. n. 724! mar. et fem. et 974! in h. DC. et 927! mar. in h. DC. et Bl. ! in h. DC. ex h. lugd. bat.), Ceylon (Miq. flor. ind. bat. v. 1. pars 2. p. 443). P. densum Bl. ! in Verh. der bat. Genoots. v. 11. p. 193. Chavica densa Miq. ! syst. p. 252, ill. p. 37. t. 32, in Linn. v. 20. p. 131, flor. ind. bat. v. 1. pars 2. p. 443 et in Hook. Lond. journ. bot. v. 4. p. 453 et ll. cc. P. canaliculatum Opiz! in Reliq. Haenk. fasc. 3. p. 156. Dioicum, scandens. Ramuli glabri. Limbi 0,14 longi, 0,09 lati. Petioli 0,03 longi. (v. s.)

δ, *Mariannum*, foliis longiuscule petiolatis utrinque glabris membranaceo-rigidulis subopacis subtilissime pellucido-punctulatis quintuplinerviis, nervo centrali nervum utrinque unum supra basin alterumque e basi opposite mittente. — In ins. Guaham et Mariannis (Haenke! in h. Berol., Gaudich. n. 77! in h. DC.). P. Mariannum Opiz! in Reliq. Haenk. fasc. 3. p. 159. Dioicum, scandens. Limbi 0,09 longi, 0,06 lati. Petioli 0,025 longi. (v. s.)

ε, *monoicum*, foliis breviter petiolatis supra glabris subtus ad nervos subtilissime puberulis firmis non pellucido-punctulatis septuplinerviis, floribus hermaphroditis, bractea coriacea, staminibus 2, rhachi tomentosa, ovario rhachi profunde immerso et cum eo coalito, stigmatibus 3. ♂ In Timor (h. DC. ex h. Mus. par.). Ramuli glabri vel subtilissime puberuli. Limbi 0,14 longi, 0,08 lati. Petioli 0,012 longi. (v. s.)

490. P. ARGUATUM (Bl. ! in Verh. der bat. Genoots. v. 11. p. 180. fig. 1), foliis petiolatis inferioribus ovatis apice acuminatis, acumine acuto, basi æquali rotundatis superioribus lanceolatis apice acuminatis, acumine acuto, basi subattenuato-acutis utrinque glabris coriaceis quintupli-septuplinerviis, nervis subtus prominulis 3 centralibus paullo supra basin oppositis in foliis inferioribus e basi solutis ad apicem ductis sequentibus e basi solutis supra medium ductis infimis brevioribus vel deficientibus, nervulis subtransverse parallelis, petiolo glabro, pedunculo glabro petiolum æquante, stirpis masculo amento densifloro quam folium  $\frac{1}{2}$  brevior recto filiformi, bractea obovato-rotundata sessili centro affixa subtus rhachique tomentosis, staminibus 2, filamentis antheram æquante, stirpis femin. ovario cylindrico arcuato, bractea ut in mare, rhachi foveolata hirtello-tomentosa, ovario foveola basi submerso, stigmatibus 3-4 lineari-acutis, bacca oblongo-globosa rubra. ♂ In Borneo australi (Korthals fide Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 137), Ternate (Miq. l. c.), Hongkong (Miq. in Seem. bot. Herald p. 415), Java (Miq. flor. lugd. bat. v. 1. pars 2. p. 455), Ceylon (Thwaites n. 2177! in h. DC.). Sirium Rumph. ? Amb. v. 5. p. 34. t. 119. fig. 1. P. arcuatum Bl. ! enum. pl. Jav. p. 67, Miq. ! syst. p. 333, ill. p. 57. t. 58, flor. ind. bat. v. 1. pars 2. p. 455. Dioicum. Ramuli glabri. Limbi 0,07-0,05 longi, 0,02-0,03 lati. Petioli 0,006 longi. (v. s.)

β, *angustifolium*, foliis angustioribus, pedunculo femin. petiolum fere duplo superante. — In Java (Bl. ! in h. DC. ex h. lugd. bat.). Dioicum. Ramuli glabri. Limbi 0,065 longi, 0,02 lati. Petioli 0,006 longi. (v. s.)

γ, *quintuplinervium*, foliis lanceolato-oblongis quintuplinerviis, nervis a basi magis remotis. — In sylvis umbrosis Nilgherries in India orientali (Perrottet! in h. Deless.). (v. s.)

δ, *triplinervium*, foliis elliptico-lanceolatis apice breviter oblique acuminatis, acumine acuto, basi subacutis subrotundatis plerumque triplinerviis. — In China (Callery fide Miq. syst. p. 335).

491. P. HAMILTONII, foliis breviter petiolatis ovatis apice attenuatis vel brevissime acuminatis, acumine acuto, inferioribus basi inæquali rotundatis superioribus



basi inæquali acutis vel subattenuatis utrinque glabris coriaceo-membranaceis subopacis subtiliter pellucido-punctatis quintuplinerviis vel septuplinerviis, nervis subtus prominulis centrali parum supra basin nervum utrinque unum adscendentem ad apicem ductum mittente lateralibus nervis utrinque 2 e basi ortis subtilibus marginalibus, petiolo glabro foliorum inferiorum ad  $\frac{3}{4}$  longitud. usque superiorum basi ima vaginante, pedunculo petiolum duplo superante glabro, stirpis masc. amento filiformi, bractea rotundato-subquadrangulari centro affixa glabra subtus rhachique hirtellis, staminibus 3, filamento brevissimo, stirpis femin. bractea ut in mare, stigmatibus 3-4, bacca oblongo-ovata glabra, rhachi hirtella. ♀ In India ad Gualpora (Wall. n. 6659 g! ex h. Ham. in h. Wall. et DC.), Sikkim alt. 3-5000 ped. et Chittagong (Hook. et Thoms. n. 22! in h. DC.). Dioicum. Ramuli glabri. Limbi 0,08-0,09 longi, 0,04-0,05 lati. Petioli 0,01 longi. Pedunculi 0,012 longi. Amenta florentia 0,05-0,06 longa. (v. s.)

492. *P. SIRIUM*, foliis petiolatis ovato-acuminatis, acumine acuto, basi obtuse truncatis vel cordatis utrinque glabris rigidulis septupli-noveninerviis, centrali nervo ad apicem ducto, nervis lateralibus utrinque 2 fere ad apicem 2 ad medium ductis fere omnibus e basi ortis, petiolo glabro, pedunculo glabro petiolum triplo superante, amento maturo folium superante, bractea rotundata centro subsessili, bacca ovata glabra. ♀ In Goalpara (Wall. n. 6642 c! in h. Wall.), Silhet (Wall. n. 6642 D! l. c.). *Sirium* sylvestre Rumph. Amb. v. 5. p. 342. t. 118. fig. 1 et 2? *Chavica Melamiris* Miq. syst. p. 245, flor. ind. bat. v. 1. pars 2. p. 442. *Chav. pseudomelamiris* Miq.! syst. p. 244, flor. ind. bat. v. 1. pars 2. p. 441 exclus. syn. Rheede. *P. Melamiris* Roxb.! flor. ind. v. 1. p. 160 (Roxb. in h. Deless.). Dioicum. Ramuli glabri. Limbi 0,07 longi, 0,05 lati. Petioli 0,01-0,02 longi. (v. s.)

### 3° Folia multinervia.

493. *P. SYLVESTRE* (Lam. ill. p. 79), foliis petiolatis ovatis apice acuminatis, acumine acuto mucronulatoque, basi subinæquali cordatis superioribus obtusis omnibus utrinque glabris membranaceis subtiliter pellucido-punctulatis septemnerviis vel septuplinerviis, nervis subtus subprominulis 3 centralibus ad apicem ductis omnibus fere e basi solutis, nervulis subtransverse parallelis, petiolo glabro, pedunculo glabro petiolum æquante, stirpis masc. amento filiformi densifloro folium æquante, bractea oblongo-obovata glabra centro affixa, staminibus 3, filamento brevissimo, stirpis femin. foliis basi obtusis, bacca ovata, stigmatibus 3-4 lanceolatis (ex Wight icon. t. 1937). ♂ In Ceylon usque ad 2000 ped. alt. (Walker in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 438, Thwaites n. 3688! fide Thwaites enum. p. 293), Courtalloum (Wight fide Miq. l. c. et Wight icon. l. c.), ins. Mauritiæ, Malabar, Philippinis (Vahl enum. v. 1. p. 326), ins. Borbonica (h. DC.), ins. Mauritiæ (Miq. syst. p. 314), h. Willd. n. 685! et 680! *P. sylvestre* Vahl l. c., Miq. syst. II. cc. *P. spurium* Link Jahrb. 3. p. 6 (fide Miq. syst. l. c. et Visiani Hort. bot. Pad. 1842. p. 144). Dioicum, arbores scandens. Limbi 0,075 longi, 0,045 lati. Petioli 0,012 longi. (v. s.)
494. *P. SINENSE*, foliis petiolatis ovatis vel oblongo-ovatis apice attenuatis acutiusculis basi æquali cordatis supra glabris subtus pubescentibus 7-nerviis, centrali nervo ad apicem ducto sequentibus fere ad apicem ductis cæteris brevioribus, petiolo puberulo, pedunculo puberulo quam petiolus brevior, stirpis masc. bractea rotundata glabra membranacea centro subsessili, staminibus 2, anthera subsessili, stirpis femin. bractea ut in mare, ovario glabro, stigmatibus 3, bacca globosa. ♀ Prope St Davis Hongkong (in h. Kew.!). *Chavica sinensis* Champ. in Kew. Journ. Bot. v. 1. p. 106, Benth. flor. Hongk. p. 335. Dioicum, rupes scandens. Ramuli glabri. Limbi 0,09 longi, 0,05 lati. Petioli 0,02 longi. (v. s.)
495. *P. LABILLARDIERI*, foliis petiolatis lato-ovatis apice breviter acuminatis, acumine acuto, basi subinæquali inferioribus late rotundatis superioribus subrotundatis subacutisve omnibus utrinque glabris rigidulis 5-7-nerviis, nervulis subtransverse parallelis, petiolo hirtello, amento filiformi folium

æquante, pedunculo glabro petiolum duplo superante. ♂ In Amboina (Labill. ! in h. Deless.). Chavica consanguinea Miq. ! syst. p. 232, flor. ind. bat. v. 1. pars 2. p. 439. Dioicum. Ramuli glabri. (v. s.)

496. P. LÆVE (Vahl enum. v. 1. p. 332), foliis inferioribus ovatis superioribus oblongis omnibus apice acuminatis basi æqualibus utrinque glabris lævibus sub pallidioribus tenuissimis quinquenerviis, baccis distinctis ovato-oblongis. ♂ In India orientali (Vahl l. c.). — Miq. syst. p. 335. Dioicum. Rami glabri cinerei superne purpurei. Folia quadripollicaria tenera. Petioli læves Amenta tripollicaria. (Ex Vahl l. c.)

d. Bractea rhachi adnata, margine tantum libera.

1° Folia multiplinervia.

497. P. INSULARE, foliis breviter petiolatis lanceolatis apice acuminatis, acumine acuto, basi cuneatis utrinque glabris membranaceis subopacis epunctulatis septuplinerviis, centrali nervo ad  $\frac{1}{2}$  -  $\frac{3}{4}$  longitud. usque nervum utrinque unum adscendentem ad apicem usque ductum alterne mittente, cæteris nervis e basi solutis quorum medii supra medium ducti infimi subtiliores, petiolo glabro, pedunculo glabro petiolum æquante, amento filiformi densifloro folium subæquante, rhachi subtiliter hirtella, bractea oblonga utrinque obtusa sub subtiliter hirtella, staminibus 2, anthera filamentum æquante. ♂ In ins. Ceylon (Thwaites n. 2175 ! in h. DC.). Dioicum. Ramuli glabri. Limbi 0,075 longi, 0,02 lati. Petioli 0,004 longi. (v. s.)
498. P. INDICUM (Miq. ! syst. p. 315), foliis petiolatis oblongo-ovatis apice acuminatis, acumine acuto, basi subæquali obtusis utrinque glabris rigidulis quintuplinerviis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  -  $\frac{3}{4}$  longitud. usque nervum utrinque unum adscendentem mittente, petiolo glabro, pedunculo glabro petiolum duplo superante, stirpis femin. amento folium multum superante, bractea oblonga, stigmatibus 3, bacca subglobosa glabra. ♂ Ad ripas fl. Attran. P. leptostachyum (Wall. l. c. et n. 6649 ! in h. Wall. et DC.) (non Rich.). — Dioicum. Ramuli glabri. Limbi 0,12 longi, 0,04-0,05 lati. Petioli 0,015 longi. Pedunculi 0,02 longi. (v. s.)
499. P. TRINEURON (Miq. in Hook. Lond. journ. bot. v. 5. p. 555), foliis modice petiolatis elliptico-lanceolatis apice breviter acuminatis basi acutis æquilatis coriaceis epunctatis subtrinerviis, nervo centrali crasso lateralibus juxta margines adscendentibus tenuibus ab  $\frac{1}{2}$  longitud. evanidis, stirpis masc. amento breviter pedunculato parumper curvato, bractea oblonga subtus hirtella, staminibus 2. ♂ In ins. Ceylon (Walker n. 1784 in h. Arnott, fide Miq. l. c.). Dioicum. Species certa e specimine nimis incompleto descripta.
500. P. BANTAMENSE (Bl. ! in Verh. der bat. Genoots. v. 11, p. 104), foliis petiolatis oblongo-ellipticis apice acuminatis, acumine acuto, basi æquali vel subæquali subattenuatis obtusis cordulatisve utrinque glabris membranaceo-rigidulis subtilissime pellucido-punctulatis quintuplinerviis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervos utrinque 2 quorum superiores ad apicem ducti alterni infimi e basi orti mittente, nervulis subtransverse parallelis, petiolo glabro, pedunculo glabro petiolum superante, stirpis masc. amento filiformi densifloro quam folium multum brevius, bractea glabra obovata inferne attenuata, staminibus 3-4, stirpis femin. bractea oblongo-obtusula, rhachi pilosula, bacca glabra ovata, stigmatibus 3. ♂ In ins. Java (Bl. ! in h. DC. ex h. lugd. bat., Zollinger n. 754 !, 891 ! in h. DC., 710 ! in h. Kew. et 364 !, 915 ! stirpis femin. in h. DC.), ins. Ceylon (Thwaites n. 35 ! in h. DC.). — Bl. ! enum. pl. Jav. fasc. 1. p. 65. Cubeba ? bantamensis Miq. ! syst. p. 299, in Hook. Lond. journ. bot. v. 4. p. 454, flor. ind. bat. v. 1. pars 2. p. 450. P. arborescens Miq. in Hook. Lond. journ. bot. v. 4. p. 438. Dioicum. alte scandens. Ramuli glabri. Limbi 0,12 longi, 0,055 lati. Petioli 0,01 longi. Amentum femin. maturum 0,15-0,25 longum. Bacca sub 0,003 longa. (v. s.)
501. P. WALKERI (Miq. in Hook. Lond. journ. bot. v. 4. p. 438), foliis petiolatis elliptico-ovato-lanceolatis lanceolatisve æquilatis vel subinæquilatis apice acuminatis, acumine acuto vel obtusiusculo, basi leviter inæquali obtusis cr-

siusculæ membranaceis subtilissime pellucido-punctulatis supra glabris subtus præsertim ad nervos sparsissime pilosulis septuplinerviis, nervis subtus prominentibus 3 centralibus ab  $\frac{1}{4}$ - $\frac{1}{2}$  longitud. solutis ad apicem ductis cæteris e basi solutis quorum inferiores tenuissimis submarginales, petiolo glabro, pedunculo petiolum æquante, stirpis femin. amento filiformi longissimo laxifloro, bractea elongata subtus hirta, stigmatibus 5 crassis brevi-lanceolatis acutis. ♂ In ins. Ceylon (Walker in h. Hook. fide Miq. l. c.). Dioicum. Rami ramulique glabri. Limbi 0,085 longi, 0,025-0,04 lati. Petioli 0,01 longi. Amentum maturum 0,12-0,15 longum. (Ex Miq. l. c.)

502. *P. NIGRUM* (L. l. spec. p. 40), foliis petiolatis inferioribus rotundato-ovatis apice acuminatis, acumine acuto, basi æquali rotundatis vel breviter cordatis truncatisve superioribus ovato ellipticis ellipticisve plerumque inæquilatis apice acuminatis, acumine acuto basi subinæquali obtusis acutiusculisve omnibus utrinque glabris subtus glaucescenti-pallidis coriaceis opacis septuplinerviis, nervis subtus prominulis 3 centralibus supra basin subalterne solutis ad apicem ductis lateralibus utrinque 2-3 e basi solutis quorum exteriores sæpe deficientes, petiolo glabro ad  $\frac{1}{4}$ - $\frac{1}{2}$  longitud. usque vaginante, pedunculo glabro petiolum subæquante, amento sublaxifloro folium æquante hermaphrodito vel unisexuali, rhachi hirtella, bractea oblonga, staminibus 2, stigmatibus 3-4-5, bacca globosa sicca nigra. ♂ In Borneo australi (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 137, an cult. ?), Luzon (Presl epim. p. 224), Sumatra (Miq. Sumatra p. 186), Java (Zollinger n. 3595! in h. DC.), Singapore (Wall. n. 6643 F! in h. Wall.), Ceylon (Thwaites n. 2176! in h. DC. stirpis femin. Walker n. 1397 in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 438), Penang (Wall. n. 6643, E! in h. Wall.), Courtallum (Wall. n. 6643, D! et 6643, C!), Malabar (Wight, fide Miq. in Hook. Lond. journ. bot. v. 5. p. 551), Napalia (Wall. n. 6650 B! in h. Wall.). Late per orbem, in India et Brasilia ad Bahia (h. DC. !), in Antillis (Descourtils flor. ant. v. 6. p. 182. t. 424) cultum. *P. rotundum nigrum* Pluk. alm. p. 297. t. 437. fig. 1, Moris. hist. p. 602. t. 15. fig. 1. Malago Cod Rheede Mal. v. 7. p. 23. t. 12. *P. nigrum* L. l. flor. zeyl. n. 26 h. Burm. v. 3. f. 21 l et 44, et v. 4, f. 21., spec. p. 40, Vahl enum. v. 1. p. 329, Hill dict. v. 26. p. 26. t. 26. fig. 1, Loud. encycl. n. 508. p. 28, Willd. spec. v. 1. p. 159, Mart. syst. mat. med. p. 99, Spach hist. phan. t. 125, Klotzsch in Hayne Arn. gew. v. 14. t. 6, Hassk. pl. Jav. rar. p. 194, Bl. in Verh. der bot. Genoots. v. 11. p. 191, enum. pl. Jav. fasc. 1. p. 69, Miq. l. syst. p. 308, illustr. p. 50. t. 50, in Hook. Lond. journ. bot. v. 4. p. 438. *P. spurium* Link Jahrb. 1. 3. p. 60 (fide R. et Sch. mant. enum. v. 1. p. 37. *P. ? colonum* Presl bot. Bemerk. p. 112. *P. nigrum* Schnizl. Iconog. t. 81. fig. 1-5. Poivrier aromaticum (Descourtils l. c.). Lada aliis Molanga Pis. mant. arom. p. 180 et 181 cum iconib. Dioicum vel monoicum et tum stirpe hermaphrodita sterili trioicum, scandens, radicans. Ramuli glabri. Limbi 0,13 longi, 0,09 lati. Petioli 0,02-0,03 longi. Bacca 0,005 crassa. (v. s.)

β, *trioicum*, foliis angustioribus ad lanceolatam formam tendentibus omnibus minus coriaceis subtus minus pallidis, amentis trioicis. ♂ In Ceylon (Thwaites n. 35? in h. Boiss.), Silhet (Wall. n. 6658? in h. DC.), Bengalia orientali (Griffith n. 4423? in h. Vindob.) Raja Mundri Circar (Roxb. flor. ind. v. 1 p. 151), h. Willd. n. 648 et 680? Late per Asiam et in Guyana (Perrottet in h. Deless.), Brasilia in prov. Para (Mart. ? in h. monac.) et prope Saratopolin (Mart. reise p. 658) cultum. *P. trioicum* Roxb. flor. ind. v. 1. p. 151, Miq. ? syst. p. 310, illustr. p. 51 D. 51 et 52, Wight. icon. t. 1935-6. *P. nigrum* Miq. in Mart. flor. Bras. fasc. 11 p. 71. *P. fallax* Vahl enum. v. 1. p. 335\* (v. s.)

γ, *macrostachyum*, amento folium superante, 0,25 longo. ♂ In monte Khasia (Hook. et Thoms. ? in h. DC.). (v. s.)

503. *P. ATTENUATUM* (Miq. syst. p. 306, ex h. Ham.), foliis longiuscule petiolatis, ovato-acuminatis vel stirpis femin. ovato-ellipticis, acumine acuto, inferioribus basi cordatis vel truncato-rotundatis superioribus basi truncatis, stirpis femin. basi attenuato-acutis omnibus supra glabris subtus ad nervos hirtellis membranaceis pellucidis subtilissime pellucido-punctatis septuplinerviis, nervis subtus prominulis centrali paulo supra basin ærvum utrin-

que unum fere ad apicem ductum opposite mittente lateralibus utrinque 2 e basi ortis quorum exteriores subtiliores, petiolo hirtello pedunculum glabrum multum superante, amento filiformi subdensifloro, rhachi parce hirtella, bractea oblonga utrinque obtusa glabra membranacea, ovario ovato, stigmatibus 4-5, bacca ovato-angulata glabra. ♀ In Java ad Lamadiang (Miq. flor. ind. bat. v. 1 pars 2 p. 451), in India prope Sikkim alt. 4000 ped. et monte Khasia alt. 4000 ped. (Hook. et Thoms. ! in h. DC.), Assam? (Griffith n. 511? 551? 556! fide Miq. in h. Hook. Lond. journ. bot. v. 4. p. 436), monte Himalaya orientali (Griffith n. 4420! in h. DC.). *Sirium frigidum* Rumph. Amb. v. 5. p. 345. t. 119 fig. 2. P. diffusum Vahl enum. v. 1. p. 333, Miq. syst. p. 327. P. attenuatum Miq. syst. p. 306, illust. p. 49 t. 49, flor. ind. bat. v. 1. pars 2. p. 451, Wight. icon. t. 1933? (baccis globosis potius cum P. Zollingeri congruans), Miq.? in Hook. Lond. journ. bot. v. 4. p. 436. *Dioicum*. Ramuli glabri. Limbi 0,06-0,09 longi, 0,055-0,06 lati. Petioli 0,03-0,04 longi. (v. s.)

504. P. ARGYROPHYLLUM (Miq. syst. p. 330), foliis petiolatis elliptico-lanceolatis apice acuminatis, acumine acuto, basi attenuato-acutis utrinque glabris subtus albo-maculatis membranaceis pellucido-punctulatis quintuplinerviis, nervis subtus prominulis, centrali nervo supra basin trifido, lateralibus utrinque 2 fere e basi solutis quorum exterior subtilior brevior, petiolo glabro, pedunculo glabro petiolum æquante, stirpis femin. amento folium multum superante, bractea oblonga utrinque obtusa subtus hirtella, stigmatibus 3 linearilanceolatis, bacca oblongo-ovata, stirpis masc. bractea ut in femina, staminibus 2. ♀ In ins. Ceylon (Thwaites n. 3625! in h. DC. stirps femin.). — Miq. ill. p. 55. t. 56, Wight icon. t. 1941. *Dioicum*. Ramuli glabri. Limbi 0,09 longi, 0,045 lati. Petioli 0,02 longi. Amentum femin. maturum 0,12-0,13 longum. Bacca 0,004 longa. (v. s. et stirps masc. ex Miq. l. c.)

505. P. NILGHIRIANUM, foliis petiolatis ovatis apice breviter acuminatis, acumine acuto mucronulato, basi æquali rotundatis supra glabris subtus piloso-pubescentibus membranaceis opacis septupli-noveninerviis, nervis subtus prominulis centrali paulo supra basin nervos alternos vel oppositos ascendentes alte ductos 2 mittente, stirpis masc. amento filiformi, pedunculo petiolum æquante piloso, bractea oblonga utrinque obtusa, staminibus 4, filamentum antheram superante, stirpis femin. bractea ut in mare, rhachi hirtella, bacca subglobosa, stigmatibus 4. ♀ In India (Wall. n. 6642 H.? in h. DC.), montib. nilghiricis (Hook. et Thoms. ! in h. DC. stirps masc.). P. lanatum Miq. in Hook. Lond. journ. bot. v. 5. p. 553. *Dioicum*. Ramuli pilosi. Limbi 0,09 longi, 0,06 lati. Petioli 0,01 longi. (v. s.)

506. P. HYMENOPHYLLUM (Miq. in Wight icon. t. 1942), foliis petiolatis oblongo-ellipticis apice acuminatis, acumine acuto, basi obtusis acutisve supra glabris subtus ad nervos puberulis membranaceis pellucidis tripli-quintuplinerviis centrali nervo ad apicem ducto lateralibus e basi ortis utrinque 1-2 quorum medii fere ad apicem ducti exteriores subtiliores, stirpis femin. pedunculo petiolum multum superante, amento folium æquante, bractea lineari-oblonga, stigmatibus 3-4, bacca ovato-globosa. ♀ In Courtallum (Wight icon. l. c.). *Dioicum*. Limbi 0,11 longi, 0,04 lati. Petioli 0,01 longi. Pedunculi 0,02 longi. Bacca 0,005 longa. Verisimiliter P. argyrophylli femina.

507. P. ZOLLINGERI, foliis petiolatis inferioribus ovatis vel ovato-lanceolatis apice acuminatis, superioribus elliptico-lanceolatis apice acuminatis, acumine acuto, basi subinæquali obtusis raro cordulatis omnibus supra glabris subtus puberulis pubescentibusve pellucido-punctatis membranaceis vel stirpis femin. coriaceo-membranaceis quintupli-nerviis, nervis subtus prominulis centrali supra basin nervum utrinque unum fere ad apicem ductum mittente, petiolo glabro, pedunculo glabro petiolum æquante stirpis masc. amento filiformi sublaxifloro, bractea oblonga utrinque obtusa, staminibus 3, filamentum antheram æquante, stirpis femin. amento longissimo folium multoties superante, bractea ut in mare, stigmatibus 3-4, bacca globosa glabra nigra. ♀ In ins. Java (Zoll. n. 2653! in h. Boiss. stirps masc. et in h. DC. stirps femin., 304! in h. DC. stirps femin. et 3711! in h. DC.). *Dioicum*. Rami ramulique glabri. Limbi 0,07-0,1 longi, 0,04-0,007 lati. Petioli 0,015 longi. Amentum fem. maturum 0,3 longum. (v. s.)

508. *P. ZUCCARINII*, foliis petiolatis subelliptico-ovatis apice acuminatis, basi æquali acutiusculis utrinque glabris membranaceis subopacis creberrime pellucido-punctulatis septuplinerviis, nervis subtus prominulis centrali nervo ad  $\frac{1}{2}$  longitud. usque nervum utrinque adscendentem ad apicem ductum alterne mittente, lateralibus nervis e basi solutis, petiolo glabro fere ad limbi basin usque vaginante, amento folium subæquante laxifloro, pedunculo glabro petiolum paulum superante, bractea oblonga, rhachi ad basin ovarii hirta, ovario ovato glabro. ♀ In India orientali (Griffith! in h. monac. ex h. Zuccarini). Dioicum. Rami ramulique glabri. Limbi 0,13 longi, 0,07 lati. Petioli 0,01 longi. (v. s.)
509. *P. QUINQUEANGULATUM* (Miq. in Zoll. cat. p. 85), foliis modice petiolatis ellipticis vel ovatis apice breviter acuminatis, acumine acuto, basi breviter inæquali subacutis septupli (vel nervo uno accessorio ad basin adjuncto) octuplinerviis, nervis 3 mediis paulo supra basin solutis fere ad apicem ductis, stirpis masc. amento breviter pedunculato quam folium brevius densifloro, bractea oblonga obtusa coriacea subtus pilosa, stirpis femin. bacca ovata, stigmatibus 4 brevibus.—Miq. flor. ind. bat. v. 1. pars 2. p. 453, Zoll. herb. n. 1233. Dioicum scandens glabrum. Caulis 5-angulatus. Folia 6 poll. longa, 3 poll. lata. (Ex Zoll. l. c.)
510. *P. NAGELII*, foliis petiolatis elliptico-oblongis apice acuminatis, acumine acuto, basi leviter inæquali acutiusculis supra glabris subtus mollior pubescentibus subcoriaceis opacis pellucido-punctulatis quintupli-sextuplinerviis, nervis 5 subtus prominulis, centrali paulo supra basin nervum utrinque unum adscendentem fere ad apicem ductum mittente, lateralibus utrinque 2 e basi solutis quorum exteriores subtiliores sæpe deficientes, petiolo glabro ad  $\frac{1}{2}$  longitud. usque vaginante, stirpis femin. pedunculo glabro petiolum fere duplo superante, amento densifloro folium plus quam duplo superante, rhachi glabra, bractea oblonga glabra, stigmatibus 3-4 ovato-abbreviatis, bacca subglobosa nigra. ♀ In Ins. Java (Nagel n. 65! et 66! in h. Berol.) Dioicum. Rami ramulique glabri. Limbi 0,1 longi, 0,04 lati. Petioli 0,012 longi. Bacca fere 0,004 crassa. (v. s.)
511. *P. KORTHALSH* (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 139), foliis modice petiolatis lato-ellipticis apice breviter acuminatis, acumine acuto, basi leviter inæquali acutis vel subacutis rarius rotundatis subcoriaceis quintupli-nerviis, nervis 3 centralibus prope basin solutis, petiolo glabro, stirpis femin. pedunculo glabro petiolum circiter æquante, amento elongato gracili folium circit. æquante, bractea lineari-oblonga, rhachi hirtella, ovario ovoideo-oblongo angulato, stigmatibus ut plurimum 4 raro 3 lanceolato-teretibus, bacca matura ovoidea acutata subrostrato-attenuata. ♀ In Sumatra occidentali (Miq. l. c.). Dioicum. Ramuli 4-5-angulati glabri. Limbi 7-5 poll. longi,  $3\frac{1}{2}$ -2  $\frac{1}{2}$  lati. Petioli 1- $\frac{3}{4}$  poll. longi. Amentum 5 poll. longum. In specimnib. masc. verisimiliter huc ducendis, folia longius petiolata, elliptico-lanceolata, petioli 2-1 poll. longi, amentum filiforme 8 poll. longum.
512. *P. WIGHTII* (Miq. in Hook. Lond. journ. bot. v. 5. p. 552), foliis petiolatis ovatis vel elliptico-ovatis apice breviter acuminatis basi leviter inæquali vel æquali rotundatis supra glabris lævibus subtus junioribus ad nervulos hirtellis cito glabratibus coriaceo-membranaceis pellucido-punctatis septemnerviis vel (nervis 3 centralibus paulo supra basin solutis) subseptuplinerviis, stirpis femin. pedunculo petiolum superante patente, amento dein elongato folium fere æquante, bractea oblongo-lineari submembranacea subtus hirtella, stigmatibus 3-4, bacca elliptica subglobosa. ♀ In India (Wight fide Miq. l. c.), montibus nilghiricis (Hook. et Thoms. l in h. Berol.). Dioicum. Ramuli glabri. Limbi 0,08-0,11 longi, 0,05-0,07 lati. Petioli 0,015-0,01 longi. Pedunculi 0,015-0,02 longi. (Rx Miq. l. c. et spec. cit.)
513. *P. TRIANDRUM* (Ferd. Mull. l fragm. 5), foliis petiolatis ovatis apice acuminatis, acumine acuto, basi leviter inæquali obtusis utrinque glabris membranaceis subopacis pellucido-punctulatis septuplinerviis, nervis subtus subprominulis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervum utrinque unum fere ad apicem usque ductum adscendentem alterne mittente, cæteris e basi solutis adscendentibus, petiolo glabro basi ima vaginante, stirpis masc. amentis

filiformibus, bractea oblonga utrinque obtusa, staminibus 3, filamento antheram æquante. — In Australia ad Mackay river (Ferd. Muill. f in h. DC.) (v. s.)

2° Folia multinervia.

514. *P. HOOKERI* (Miq. in Hook. Lond. journ. bot. v. 4. p. 437), foliis petiolatis ovatis æquilateris apice breviter acuminatis, acumine obtuso, basi æquali cordatis vel rotundatis coriaceo-membranaceis pellucido-punctatis supra glabris subtus ad nervos hirtellis 7-nerviis, vel nervis 3 mediis paullo supra basin solutis, septupli-noveninerviis, nervis subtus prominentibus centralibus 3 ad apicem ductis cæteris tenuibus, nervulis subtransverse parallelis, petiolo hirtello, pedunculo petiolum superante, stirpis femin. amento folium superante, bractea oblonga subtus hirtella cito glabrata, ovario ovato, stigmatibus 4. ♀ In Bombay (Lambert in h. Hook. fide Miq. l. c.). Dioicum. Ramuli pubescenti-hirtelli. Limbi 0,08-0,1 longi, 0,04-0,065 lati. Petioli 0,005-0,01 longi. Amentum maturum 0,08 longum.

515. *P. CUMINGIANUM* (Miq. l. syst. p. 329), foliis petiolatis elliptico-lanceolatis apice acuminatis, acumine mucronulato, basi æquali rotundatis vel superioribus attenuato-acutiusculis utrinque glabris membranaceis subtilissime pellucido-punctulatis 5-nerviis, nervis subtus prominulis 3 centralibus ad apicem ductis lateralibus brevioribus sæpe deficientibus, nervulis subtransverse parallelis, petiolo glabro, pedunculo glabro, petiolum æquante, stirpis masc. amento filiformi subaxifloro, bractea oblonga utrinque obtusa glabra, staminibus 3, filamento antheram æquante, stirpis femin. amento subaxifloro folium superante, rhachi glabra, bractea ut in mare, stigmatibus 3 obtusis, bacca ovata glabra. ♀ In ins. Philippinis (Cuming n. 1190! in h. DC. stirps fem. ? et h. Berol. stirps femin., 1624! stirps femin. in h. Deless.). — Miq. flor. ind. bat. v. 1. pars 2. p. 454 *P. Philippinum* Miq. syst. p. 322, flor. ind. bat. v. 1. pars 2. p. 453. Dioicum. Ramuli glabri. Limbi 0,07 longi, 0,035 lati. Petioli 0,008 longi. Amentum femin. florens 0,12-0,14 longum. (v. s.)

516. *P. ZEYLANICUM* (Miq. in Hook. Lond. journ. bot. v. 4. p. 436), foliis petiolatis ovatis æquilateris apice acuminatis, basi inæquali cordatis rotundatisve rigide coriaceis epunctatis utrinque glabris 5-6-nerviis, nervis subtus prominentibus, centrali ad apicem ducto, sequentibus nervis fere ad apicem ductis exterioribus brevioribus, petiolo glabro, stirpis masc. pedunculo glabro petiolum superante, amento brevi densifloro, bractea oblonga subtus rhachique hirtis, staminibus 2, filamento crasso, stirpis femin. bractea ut in mare, stigmatibus 4 raro 5 reflexis, bacca globosa. ♀ In Ceylan (Walker n. 31. in h. Hook. fide Miq. l. c.). Dioicum, ramosissimum, an scandens? Limbi 0,03-0,06 longi, 0,02-0,03 lati. Petioli 0,005-0,015 longi. Forma major extat cum limbis oblongis 0,08-0,1 longis, 0,04-0,045 latis, amentis masc. longioribus 0,01 longis. In Mayburan (Adam in h. Hook. fide Miq. l. c.).

Sect. VIII. APOPIPER. — Bractea a flore distincta, stamina 2. Amenta oppositifolia. Antheræ non articulatae persistentes.

517. *P. HEDERACEUM*, foliis petiolatis ovato-ellipticis apice attenuatis acutis, basi subattenuatis acutiusculis utrinque glabris subcoriaceis opacis quintupli-septupli-nerviis nervis subtus subprominulis centrali paullo supra basin nervos utrinque 2-3 adscendentes mittente, petiolo glabro, pedunculo glabro petiolum æquante, amento filiformi densifloro, bractea rotundata glabra ciliolata centro subpeltata subsessili, rhachi foveolata ciliolata, staminibus 2, filamentis brevibus, antheris non caducis. ♀ In Nova Hollandia (Cunningham! in h. DC. et Kew. Hügel! in h. Vindob.). *Cubeba hederacea* Miq. l. mss. in h. Kew. in Hook. Journ. bot. v. 4. p. 436. Dioicum, scandens. Ramuli glabri. Limbi 0,06-0,07 longi, 0,03-0,045 lati. Petioli 0,015 longi. (v. s.)

518. *P. MEMBRANACEUM*, foliis petiolatis inferioribus ovatis apice acuminatis. acumine acuto, basi cordatis superioribus suboblongo-ovatis apice acuminatis. acumine acuto, basi cordulatis vel rotundato-truncatis vel et subattenuatis omnibus utrinque glabris firmulis opacis epunctulatis inferioribus 9-nerviis

superioribus 4-nerviis, nervis subtus prominulis, petiolo glabro ad  $\frac{1}{2}$  longitud. usque vaginante, amento maturo folium dimidium superante densifloro, pedunculo glabro quam petiolus  $\frac{1}{2}$  brevior, bractæ truncato-peltatæ pelta semilunari ciliolata, staminibus 2, antheris non articulatis persistentibus, ovario glabro, stigmatibus 3, bacca obovata glabra. ♀ In Nova Hispania (Pav. in h. Boiss. ubi sub nom. inedit. *P. membranaceum*). Monoicum, flores hermaphroditi. Ramuli glabri. Limbi inferiores 0,1 longi, 0,06 lati. Petioli 0,025 longi. Amenta 0,004 crassa. (v. s.)

Sect. IX. NEMATANTHERA. — Genus Nematanthera *Miq. in Linn.*  
v. 18. p. 606. t. 2.

Bractea a flore distincta. Stamen unicum. Flores hermaphroditi vel unisexuales. Amenta oppositifolia. Plantæ mopoicæ. Antheræ articulatæ.

519. *P. NEMATANTHERA*, foliis breviter petiolatis oblongo-subobovato-ellipticis apice acuminatis, acumine acuto, basi inæquali subattenuato-obtusis, lobo majore auriculæformi, utrinque glabris membranaceis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos patulo-subadscendentes utrinque 7-8 quorum supremi fere ad apicem ducti nervulosque validos mittente, petiolo glabro, pedunculo petiolum æquante, floribus hermaphroditis, bractæ obovato-cucullatæ basi contractæ dorso sub apice flavide hirtellæ vertice inflexo nudo peltam triangularem margine puberulam simulante, stamine unico laterali oblongo. anthera valde elongata exserta, connectivo supra loculos producto, ovario glabro ovato, stigmatibus 3 raro 2 recurvis brevibus, amento cylindrico. ♀. In Surinam (Hostmann n. 19. in h. Kew. et Kappler n. 1438! in h. Boiss.). Ramuli glabri. Limbi 0,13 longi, 0,05 lati. Petioli 0,01 longi. Amenta florentia 0,01 longa, 0,003 crassa. (v. s.)

520. *P. SAGORI*, foliis subsessilibus oblongo-ellipticis apice acuminatis, acumine acuto, basi inæquali obtusis utrinque glabris membranaceis, centrali nervo subtus prominulo tota longitudine nervos patulo-adscendentes alternos subtus prominulos utrinque 13-15 nervulosque validos mittente, pedunculo glabro, amentis unisexualibus, amenti masculi globosi bractea cucullata brevi extus puberula, stamine unico oblongo, anthera filamentum multum superante, connectivo supra loculos producto incrassato, amenti feminei cylindrici brevis bractea obovato-cucullata extus puberula, ovario globoso, stigmatibus 2 rhachi oppositis. ♀ In Guyana gallica ad Maroni (Sagot n. 1255! in h. Kew. et Berol. ubi *Peltobotrya* dicta. Ramuli glabri. Limbi 0,05 longi, 0,02 lati. Pedunculi 0,008 longi. (v. s.)

### *Species non satis notæ.*

† Amenta oppositifolia. Flores hermaphroditi.

#### *A. Americanæ.*

521. *P. CHILENSE*, foliis alternis petiolatis cordatis planis glabris novemnerviis, petiolis vaginantibus alatis glabris, stipulis ovato-lanceolatis sessilibus trinerviis glabris, amentis pedunculatis. ♀ In Chili. *Peperomia plantaginifolia* Opiz in Reliq. Hænk. fasc. 3. p. 164, A. Dietr. v. 1. p. 161. *Artanthe dubia* *Miq.* syst. p. 535. Caulis herbaceus glaber articulatus, articulis nodosus. Folia *Plantaginis* majoris.

522. *P. CASAPIENSE*, foliis petiolatis oblongis vel obovato-oblongis apice breviter acuminatis, acumine acuto, basi inæquali rotundatis crassiuscule membranaceis pellucido-punctatis utrinque glandulis fuscis conspersis glabris, centrali nervo tota longitudine nervos 13-14 subadscendentes quorum supremi fere ad apicem ducti mittente, petiolo glabro, amento valde juvenili. ♀ In Casapi Peruvizæ (Mathews n. 1712 fide *Miq.* l. c.). *Artanthe casapiensis* *Miq.* in HOOK. Lond. journ. bot. v. 4. p. 463. Rami glabri minute verruculosi. Limbi 0,28 longi, 0,115 lati. Petioli 0,02 longi.

523. *P. CURVATUM* (Ruiz et Pav. *flor. peruv.* v. 1. p. 36. t. 54. fig. a), foliis petio-

- latis oblongis apice subito acuminatis, acumine acuto, basi leviter inæquali acutis subcoriaceis utrinque glabris opacis, centrali nervo tota longitudine nervos patulo-subadscendentes utrinque circiter 12 nervulis validis intermixtos mittente, pedunculo quam petiolus paullo brevior, amento maturo quam folium brevior, bractea subreniformi (?). ♂ In nemoribus Cuchero et Chinchac (R. et P. l. c. et Endlich. in h. Vindob.), in Peruvia orientali prope Tarapoto (Spruce n. 4137! in h. DC.). — Vahl enum. v. 1. p. 315. *Artanthe curvata* Miq. syst. p. 522. Limbi 0,13 longi, 0,045 lati. Petioli 0,015 longi. Amenta 0,005 longa (ex R. et P. l. c.).
524. *P. EXCAVATUM* (R. et P. flor. peruv. v. 1. p. 34. t. 54. fig. b), foliis petiolatis oblongo-subovatis apice protracto-acuminatis, acumine acuto, basi breviter subattenuatis acutis, centrali nervo nervos subadscendentes alternos lat. majore 6 minore 5 quorum supremi ad apicem ducti mittente, pedunculo petiolum æquante, amento cylindrico quam folium fere duplo brevior, bractea obovata glabra margine membranacea, staminibus quam ovarium longioribus. ♂ In Chinchao nemoribus (R. et P. l. c.). *Artanthe excavata* Miq. syst. p. 534. Glabrum. Limbi 0,155 longi, 0,068 lati. Petioli 0,01 longi.
525. *P. SCABRUM* (R. et P. flor. Peruv. v. 1. p. 34. t. 55. fig. b), foliis breviter petiolatis suboblongo-subovatis apice acuminatis, acumine acuto, basi inæquali acutis supra asperis subtus pubescentibus, centrali nervo ad  $\frac{1}{2}$  longitud. nervos alternos subadscendentes quorum supremi ad apicem ducti mittente, petiolo pubescente, pedunculo pubescente petiolum æquante, amento cylindrico crasso folium dimidio æquante, staminibus 2, stigmatibus 3, stylo nullo. ♂ In Huassahuassi nemoribus (R. et P. l. c.). *P. obesum* Vahl enum. v. 1. p. 321. *Artanthe obesa* Miq. syst. p. 470. Rami scabri. Limbi 0,08 longi, 0,04 lati. Petioli 0,007 longi. (v. s.)
526. *P. PANGOENSE*, foliis petiolatis rotundato-ovatis summis lato-ovatis omnibus æquilateris vel inæquilateris apice subabrupte brevi-acuminatis, acumine obtuso, basi lato-truncatis utrinque glabris rigide membranaceis 9-sub-7-nerviis, nervis 5 centralibus fere ad apicem ductis, petiolo glabro, pedunculo petiolum subæquante, bractea conchæformi glabriuscula? vertice subpeltata?, stigmatibus 4-5, staminibus.... ♂ In Peruviz montibus ad Pangoa (Mathews n. 1143! in h. Kew., Spruce n. 4157! in h. Kew.). *Artanthe Ruiziana* Miq. in Hook. Lond. journ. bot. v. 4. p. 444. Ramuli nascentes glabriusculi. Limbi inferiores 0,25-0,28 longi, 0,23-0,19 lati, superiores 0,16-0,19 longi, 0,1-0,15 lati. Petioli 0,02-0,01 longi. (v. s.)
527. *P. LEHMANNIANUM*, foliis brevissime petiolatis lanceolato-oblongis æquilateris vel inæquilateris apice longe attenuato-acuminatis, acumine acuto, basi leviter inæquali rotundatis vel acutis crebre pellucido-punctatis supra glabris subtus ad nervos adpresse hirtellis, centrali nervo ad  $\frac{1}{2}$  -  $\frac{1}{4}$  longitud. nervos 5-6 adscendentes tenues mittente, petiolo glabro, pedunculo petiolum paullo superante, bractea inflexo-peltata glabra tenerrime ciliolata. ♂ In Peruvia ad Casapi (Mathews n. 1694! in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 446). *Artanthe Lehmanniana* Miq. l. c. Ramuli verruculoso-punctati, juniores pilosuli. Limbi 0,1-0,125 longi, 0,03-0,035 lati. Petioli 0,002-0,004 longi (ex Miq. l. c.)
528. *P. ALVEOLATUM* (Opiz in Reliq. Hænk. fasc. 3. p. 155), foliis petiolatis maximis acuminatis cordatis membranaceis supra rugoso-bullatis glabris subtus lacunosis ad nervos nervulosque pilosiusculis margine pubescentibus alterne nervoso-venosis, petiolo membranaceo-alato verrucoso-scabro, amento cylindrico mucronato quam folium dimidio brevior, pedunculo quam petiolus multum brevior glabro. ♂ In Peruviz montibus Huanocensibus (Opiz l. c.). Rami verrucosi.
529. *P. SCABRIUSCULUM* (A. Dietr. sp. v. 1. p. 647), foliis brevissime petiolatis ovatis apice acuminatis basi inæquali angustatis nervuloso-reticulatis subtus ad nervulos adpresse pilosis, amentis oppositifoliis brevissime pedunculatis cylindricis obtusiusculis quam folia multum brevioribus, pedunculis teretibus. ♂ In montanis Huanocensibus Peruviz (Opiz in Reliq. Hænk. fasc. 3. p. 151).



- P. scabrum* Opiz l. c. *P. scabridum* Flora 1829. p. 229 (solum nomen). *Artanthe scabruscula* Miq. syst. p. 447. Rami scabriusculi.
530. *P. CASSINOIDES* (Opiz in Reliq. Hænk. fasc. 3. p. 153), foliis rotundato-ovatis apice acuminatis, acumine acuto, coriaceis venoso-nervosis supra margineque glabris subtus ad nervos et nervulos villosis, petiolis stipulaceis, stipulis brevissimis obtusis, pedunculis glabris. § In montibus Huanocensibus Peruviae (Opiz l. c.). *Artanthe cassinoides* Miq. syst. p. 490. Quid syn. *P. cassinoides* herb. Willd. ? sec. Presl. Rami glabri. (Ex Opiz l. c.).
531. *P. OPIZIANUM* (Flora, 1829, p. 228), foliis brevissime petiolatis ovato-oblongis utrinque attenuatis basi inæqualibus venosis lineatis utrinque pilis sparsis obtectis margine pilis longis sparsis ciliatis, petiolis glabris incrassatis brevissimis, amentis arcuatis mucronatis folia æquantibus oppositifoliis mucrone piloso. § In peruvianis montibus (Opiz l. c.). *P. arcuatum* Opiz in Reliq. Hænk. fasc. 3. p. 151, non Bl. *P. falcatum* A. Dietr. sp. 1. p. 639. *Artanthe arcuata* Miq. syst. p. 533. Caulis fruticosus ramosus. Rami punctis minutis elevatis scaberrimis. (Ex Opiz l. c.).
532. *P. OBLONGATUM* (Opiz in Reliq. Hænk. fasc. 3. p. 153), foliis brevissime petiolatis patentibus ovato-lanceolatis longe acuminatis basi inæqualibus glabris coriaceis venoso-nervosis, amentis pedunculatis cylindricis acuminatis oppositifoliis folio multo brevioribus, pedunculis brevissimis glabris. § In montibus Huanocensibus Peruviae (Opiz l. c.). *Artanthe oblongata* Miq. syst. p. 535. *Carpunya oblongata* Presl. Epim. p. 229. Rami glabri. Petioli brevissimi glabri. (Ex Opiz l. c.).
533. *P. PATENS* (Opiz in Reliq. Hænk. fasc. 3. p. 154), foliis ovatis acuminatis basi inæquali subrotundatis coriaceis supra rugosis scabris pilis brevissimis adpressis obtectis margine scabris subtus reticulato-venosis pilosis, nervis nervulisque patentissime pilosis, petiolis villosis, amentis patentibus oppositifoliis cylindricis pedunculatis mucronatis quam folia duplo brevioribus, pedunculis quam petioli brevioribus pilosis. § In montibus Huanocensibus Peruviae (Opiz l. c.). *Artanthe patula* Miq. syst. p. 554. Rami pilosi exasperati. (Ex Opiz l. c.).
534. *P. PHYTOLACÆFOLIUM* (Opiz in Reliq. Hænk. fasc. 3. p. 151), foliis ovato-oblongis utrinque attenuatis glabris basi inæqualibus, petiolis canaliculatis glabris, amentis folio triplo brevioribus obtusis oppositifoliis, pedunculis glabris. § In Guayaquil (Opiz l. c.). *Artanthe phytolacæfolia* Miq. syst. p. 534, in Linn. v. 20. p. 174. *Peltobryon phytolacæfolium* Presl. epim. p. 224. Rami glabri. (Ex Opiz l. c.).
535. *P. APPENDICULATUM*, foliis subsessilibus obovati-oblongis apice longe acuminatis, acumine obtusiusculo, basi angustatis hinc resectis illinc auriculato-rotundatis pellucido-punctatis valde bullatis scabro-hirtellis subtus longe ferrugineo-pilosis, petiolo longe ferrugineo-piloso, pedunculo longe ferrugineo-piloso, amento quam pedunculus brevior, bractea conchæformi-peltata ciliato-hirta, filamentis exsertis, staminibus 3? ovario ovoideo glabro apice in stylum brevem attenuato. § In sylvis Nanegal declivitatibus occidentalis Andium Quitensium (Benth. pl. Hartw. p. 254). *Artanthe appendiculata* Benth. l. c. Frutex pilis longis ferrugineis insignis. Folia 7-9 poll. longa, supra medium 2½-3 poll. lata.
536. *P. HARTWEGIANUM*, foliis amplis petiolatis sub medio peltatis ovatis apice acuminatis basi rotundatis et leviter excavatis multinerviis costatisque bullato-rugosissimis scabriusculis subtus ferrugineo-villosis subcoriaceis, nervis circiter 9 e petiolo divaricantibus, centrali nervos 4-5 mittente, petiolo alato, amentis elongatis floribus densissimis, bractea clavata fere undique hirta, staminibus 3 (vel 4?), stigmatibus 3-4 vix recurvis, bacca oblonga perspicue angulata. § In sylvis juxta Popayan (Benth. l. c.). *Artanthe Hartwegiana* Benth. pl. Hartw. p. 254. Petioli 1½ poll. longi. Folia pedalia, 7 poll. lata. Amenta florentia 8 poll. longa, vix crassitie pennæ anserinæ.
537. *P. ALBIDUM* (Kunth in H. B. nov. gen. v. 1. p. 55), foliis petiolatis ovatis apice acuminatis basi inæquali rotundatis membranaceis reticulato-septuplinerviis supra viridibus tenuissime pubescentibus subtus canescentibus tomen-

- tosopubescentibus, petiolo pubescente, amento crassitie pennæ et pedunculo pubescente quam petiolus brevior. 5 In regione calidis Carthagera de India et Turbaco (Kunth l. c.). — Kunth synops. v. Artanthe albidia Miq. syst. p. 415. Rami tomentosopubescentes. pollicaria. Petiolisemipollicares. Pedunculi 4 lin. longi. (Ex Kunth l.
538. *P. HISPIDUM* (Kunth in H. B. nov. gen. v. 1. p. 50), foliis petioloblongis apice longe acuminatis basi parum angustatis inæqualiter lineatis rigidis supra scaberrimis subtus pilosiusculis ad nervulos hinculato-nervulos, nervis primariis parallelis, amento quam folium cylindraceo crassitie pennæ columbinæ, pedunculo petiolum parum pilis raris obsito. 5 In locis temperatis frondosis inter Turbaco et C de India, alt. 180 hexap. (Kunth, l. c.). — Kunth synops. v. 1. p. 106. controversa Miq. syst. p. 448. Rami hirti pilis reflexo-patentibus. 5 quepollicaria-semipedalia, tres pollices fere lata. Petioli quatuor lin Spadices 3-4-pollicares. (Ex Kunth l. c.)
539. *P. CARIPENSE* (Kunth in H. B. nov. gen. v. 1. p. 48), foliis brevitiolatis oblongis apice longe acuminatis, acumine obtuso, basi æquali glabris, amentis folio quadruplo brevioribus. 5 In crepidinibus umbr Cœnobium Caripense prov. Cumanensis. — Kunth synops. v. 1. 5 tanthe caripensis Miq. syst. p. 522. Rami læves flexuosi glaberr 4-5-pollicaria, 2 poll. lata. Petioli 3 lin. longi. (Ex Kunth l. c.)
540. *P. UMBROSUM* (Kunth in H. B. nov. gen. v. 1. p. 51), foliis oblique ovatis apice acuminatis basi inæquali rotundatis utrinque p membranaceis opacis pellucido-punctulatis, centrali nervo ad ½ long nervos subadscendentes alternos utrinque 5-6 quorum supremi usque ducti mittente, petiolo piloso-hirto. 5 Ad flumen Magda Buenavista et isla de Bruxas in ripa paludosa (Humb. et Bon Humb. in h. Willd. n. 667 ! unicum folium). — Kunth synops. v Artanthe umbrosa Miq. syst. p. 470. Limbus 0,125 longus, 0,075 l latus 0,015 longus. (Ex Kunth l. c.)
541. *P. PSILOSTACHYUM* (Kunth in H. B. nov. gen. v. 1. p. 50), f petiolatis oblongis apice acuminatis basi subæquali rotundatis utri reticulato-venosis, centrali nervo subtus prominente, petiolo pubesc folium æquante crassitie caudæ murinæ, pedunculo petiolum r perante. 5 In regione calidissima in ripa fluminis Magdalenæ prop de Carare alt. 140 hexap. (Kunth l. c.). — Kunth syn. v. 1. p. 1 psilostachya Miq. syst. p. 448. Rami pubescentes. Folia 3-4-pollie 2 lin. longi. Pedunculi tripollicares. Amenta quadripollicaria. (Ex
542. *P. GRANDE* (Vahl, Eclog. am. fasc. 2. p. 3. t. 11), foliis peti oblongis apice breviter acuminatis, acumine acuto, inferioribus subcordatis superioribus basi subæquali subacutis utrinque brevi nervo tota longitudine nervos alternos patulo-subadscendentes u 12 nervulosque validos mittente. 5 Ad St. Martham (Von Rol l. c.). — Vahl enum. v. 1. p. 321. Chavica Lessertiana Miq. syst. p glabris, crassitie pennæ anserinæ. Folia semipedalia, 6-7 poll. lata. careas. (Ex Vahl l. c.)
543. *P. DECUMANUM* (L. sp. p. 41, excl. syn. Rumph.), foliis petio ovatis apice acuminatis basi profunde cordatis utrinque glabris r lucidis parce pellucido-punctulatis 9-nerviis, nervis subtus promin libus 5 per anastomoses fere ad apicem ductis, lateralibus fere r alte ductis, infimis subtilioribus sub medio evanidis, nervuliis parallelis, petiolo glabro. 5 In locis scopulosis ad radices montis L Guayra prope Caracas (Humb. et Bonpl. nov. gen. v. 1. p. 57), in dent. (Bredemeyer in h. Willd. n. 693 !), Cajenna ? (Aubl. hist. p. p. 21, exclus. syn. Rumph.). — Herb. Willd. l. c., Aublet ? l. c., Opiz in Reliq. Hænk. fasc. 3. p. 160. Limbi 0,29 longi, 0,08 lati. longi. Confer ad num. 450. (v. s.)
544. *P. SCOPULORUM* (Kunth, olim); foliis longe petiolatis oblique ov apice acuminatis basi oblique sinuato-cordatis lobis rotundatis inæ

membranaceis haud pellucido-punctulatis glaberrimis supra punctulis alissimis nigris dense conspersis et opacis subtus pallidioribus, nervis prominulis, centrali nervos 5-7 quorum inferiores approximati mittunt, petiolo glabro quam folium dimidio breviori glabro, bractea cucullato-exa subtriangulari margine ciliato-villosa, bacca subtetragona lateraliter apressiuscula. ♀ In Brasilia meridionali ad Serra dos Orgaos (Luschath fide Anth. l. c.). *Steffensia scopulorum* Kunth in Linn. v. 13. p. 669. *Artanthe pulorum* Miq. syst. p. 396, in Mart. flor. Bras. fasc. 11. p. 36. Ramuli glabri. Abi 0,27-0,29 longi, fere 0,12 lati. Petioli 0,05-0,065 longi. Amenta 0,15-0,16 longa. Pedunculi 0,025 longi. (Ex Kunth l. c.)

*P. DIMIDIATUM*, foliis petiolatis oblongo-lanceolatis acutissimis basi valde aequali acutis subdividiatisque utrinque glaberrimis coriaceis opacis imbricatis pinnatinerviis reticulato-venosis, petiolo marginate, pedunculo triolum equante, amento cylindrico obtuso erecto-patente quam folium plus am triplo breviori, bractea longiore peltato-cochleariformi densissime imbricato-hirsuta, stigmatibus 3, ovario glabro angulato. ♀ In Brasilia prope o-Janeiro (Lhotzky fide Presl. Epimel. bot. p. 225). *Artanthe dimidiata* Presl. l. c., Miq. in Mart. flor. Bras. fasc. 11. p. 72. Ramuli glaberrimi. Folia que 9 poll. longa usque semipollicem lata. (Ex Presl. l. c.)

*P. SUBÆQUALE*, foliis petiolatis oblongo-lanceolatis apice acutis basi inaequali acutis utrinque glaberrimis coriaceis minutissime et creberrime ellucido-punctatis, pinnatinerviis pedunculis compressis petiolo glaberrimo abduple longioribus, amentis cylindricis obtusis folium subæquantibus, bractea apice peltata hirsuto-ciliata, ovario subgloboso glabro angulato, stigmatibus 3 puberulis minutis. ♀ In Brasilia ad Rio de Janeiro (Mikan fide Presl. in Epimel. bot. p. 226). *Artanthe subæqualis* Presl. l. c. Miq. in Mart. flor. Bras. fasc. 11. p. 72. Ramuli glaberrimi. Folia 3-5 pollices longa, 1-1½ poll. lata. Petioli 3 lin. longi. (Ex Presl. l. c.)

*P. DIVERGENS*, foliis breviter petiolatis oblongo-lanceolatis apice angustato-acuminatis obtusis basi acutis utrinque glaberrimis tenuiter coriaceis pellucido-punctatissimis crebre pinnatinerviis reticulato-venosis, petiolo glaberrimo, pedunculo compresso quam petiolus subduplo longiore, amentis cylindricis obtusis folium æquantibus, bractea cochleariformi a basi longe dilata, bacca glabra rhomboidali, stigmatibus 3. ♀ In Brasilia ad Rio-Janeiro Presl. Epim. bot. p. 227). *Artanthe divergens* Presl. l. c. Miq. in Mart. flor. Bras. fasc. 11. p. 74. Ramuli glaberrimi. Folia usque 5-poll. longa, 2 poll. lata. Petioli 3-4 lin. longi. (Ex Presl. l. c.)

*P. MACROSTACHYUM*, foliis petiolatis late ovato-ellipticis apice abrupte acuminatis basi aequali acutis utrinque glaberrimis minute pellucido-punctatissimis innatinerviis, nervis lateralibus utrinque 5 nervulis validis intermixtis, pedunculo petiolum æquante, amento cylindrico mutico quam folium breviori, bractea obovata longe denseque ciliolata, fructibus glabris obconicis triquetris. ♀ In Brasilia ad Rio de Janeiro (Mikan in Presl. Epim. bot. p. 227). *Artanthe macrostachya* Presl. l. c. Miq. in Mart. flor. bras. fasc. 11. p. 74. Folia usque 6 poll. longa, usque semiquartum pollicem lata. Petioli 6-8 lin. longi. (Ex Presl. l. c.)

*P. HIRTULUM*, foliis petiolatis ovato-oblongis apice acuminatis basi inaequali obtusatis subtus ad nervos in axillis nervorum hirsutis minutissime pellucido-punctulatis pinnatinerviis, nervis lateralibus utrinque 5, petiolis pilis albidis patentissimis hirsutis, pedunculis petiolo subduplo longioribus, amentis cylindricis apiculatis folio plus duplo longioribus, bractea peltata albida ciliata. ♀ In Brasilia ad Bahiam (Lhotzky fide Presl. in Epim. bot. p. 226). *Artanthe hirtula* Presl. l. c., Miq. in Mart. flor. Bras. fasc. 11. p. 73. Rami hirtuli. Folia usque 7 pollices longa, 3 poll. lata. Petioli 3-6 lin. longi. (Ex Presl. l. c.)

*P. BAKIENSE*, foliis breviter petiolatis oblongo-lanceolatis apice acuminatis basi oblique et inaequaliter cordatis rugosis muriculato-scabris subtus ad nervos nervosque hirsutis pellucido-punctatissimis pinnatinerviis reticulato-venosis, nervis lateralibus latere majore 6-7 minore 5-6, pedunculis petiolo vix longioribus, amentis elongatis cylindricis apiculatis rectis curvatisve. ♀ In

- Brasilia* ad Bahiam (Lhotzky fide Presl. l. c.). *Artanthe Bahiensis* Presl. Epimel. bot. p. 227, Miq. in Mart. flor. Bras. fasc. 11. p. 73. Rami glabri, multi pilis densissimis rigidulis brevissimis scabri. Folia usque ultra  $\frac{1}{2}$  poll. longa,  $\frac{1}{4}$  poll. lata. Petioli 1-3 lin. longi. (Ex Presl. l. c.)
551. *P. AURICULATUM*, foliis brevissime petiolatis lato-vel obovato-ellipticis brevissime apiculatis basin versus subattenuatis basi ima inæquali obliquo lobo majore auriculatim producto petiolum velante supra glabris nitidis subtus præsertim ad nervos et nervulos molliter hirtello-pubescentibus mule membranaceis opacis subtiliter pellucido-punctulatis, nervis ad prominulis, centrali tota longitudine nervos alternos subalternosve petiolo subadscendentes utrinque circ. 13 mittente, petiolo ad limbi basin usque vaginante dorso pubescente.  $\hookrightarrow$  In sylvis ad ostia fluminis Rio dos Engenheiros dicti in flumen Japura provinc. Rio Negro (Mart. in h. monac.). *Artanthe auriculata* Miq. l. in Linn. v. 20. p. 167, in Mart. flor. Bras. fasc. 11. p. 73. Rami juniores pilis recurvulis densissime hirti. Limbi 0,15 longi, 0,08 lati. Petioli 0,003 longi. (v. s.)
552. *P. KEGELIANUM*, foliis breviter petiolatis elliptico-vel ovato-oblongis breviter acuminatis basi subæquali obtusis membranaceis subpunctulatis supra atro-viridibus nitidis glabris subtus ad nervos hirtellis, costulis utrinque 6-7 distinctioribus aliisque venulis interpositis patulo-adscendentibus, pedunculo petiolum haud superante, amentis folio brevioribus tenuibus, bracteis pedicellato-peltatis hirtis-ciliatis.  $\hookrightarrow$  In sylvis Paraensibus prope Klein Onoribo (Leg. n. 659 fid. Miq. in Linn. v. 22. p. 77). *Artanthe Kegeliana* Miq. l. c. Rami glabrescentes. Petioli 1-2 lin. longi. Folia  $5\frac{1}{2}$ -6 poll. longa, 2-2 $\frac{1}{2}$  poll. lata. Amentum junius 3 poll. longum rectum.
553. *P. ABUTILIFOLIUM*, foliis longe petiolatis cordato-subrotundis apice acuminatis, acumine acuto, basi lobis divergentibus, utrinque glabris membranaceis pellucido-punctulatis ciliolatis 9-nerviis, nervis imprimis subtus prominentibus 3 centralibus ad apicem usque ductis sequentibus sensim brevioribus et tenuioribus, petiolo glabro ad limbi basin usque vaginante, amento recto quam folium fere dimidio breviori, pedunculo quam petiolum multoties breviori glabro, bractea e stipite tenui spatulata glabra vertice antico concavo in petalum trigonum pubescentem incrassato, bacca obovato-trigona vel tetragona, staminum vestigiis.  $\hookrightarrow$  In *Brasilia* (in h. Mus. Par. fide Miq. syst. p. 385). *Artanthe abutilifolia* Miq. l. c. et in Mart. flor. Bras. fasc. 11. p. 33. t. 6. fig. 1. Ramuli brevissimi glabri. Limbi 0,25 longi, 0,19 lati. Petioli 0,06 longi. Amenta 0,1 longa, 0,003 crassa.
554. *P. TOMENTOSUM*, foliis brevissime petiolatis suboblongo-ellipticis apice acuminatis, acumine acuto, basi inæquali rotundatis utrinque dense ad nervos densius molliter tomentosus rigido-coriaceis opacis subtiliter pellucido-punctulatis, nervis subtus prominulis, centrali ad  $\frac{1}{2}$  longitud. usque nervos alternos subadscendentes utrinque 6 quorum supremi ad apicem ducti infimi subliores mittente, petiolo utrinque dense molliter tomentoso ad  $\frac{1}{2}$  longitud. usque vaginante (amentum in specimine nondum efformato).  $\hookrightarrow$  In *Brasilia* Pohl! in h. Vindob. fide Miq. in Linn. v. 20. p. 159). *Artanthe Schottii* Miq. l. c. Rami puberuli dein glabrati. Ramuli pilis retrorsis tomentosi. Limbi 0,13 longi, 0,05 lati. Petioli 0,006 longi. (v. s.)
555. *P. MULTINERVIUM* (Mart. et Gal. in bull. Acad. Brux. v. 10. n. 2, p. 20), foliis subsessilibus ovato-lanceolatis acuminatis basi inæqualibus supra scaberrimis subtus villosis-pubescentibus brevibus 13-15 nerviis, spicis tenuibus elongatis incurvis pedunculatis.  $\hookrightarrow$  Prope Ialapa et Mirador alt. 2500-4000 ped. caulis glabriusculus. Folia 5-6 poll. longa 2 poll. lata. (et Mart. et Gal. l. c.)
556. *P. HOFFMANSEGGIANUM* (Rœm. et Sch. Mant. 1. p. 242), foliis petiolatis lanceolatis apice acuminatis venosis glaberrimis concoloribus, amentis brevipedunculatis.  $\hookrightarrow$  In *Brasilia* (Hoffmansegg in h. Willd. fide Rœm. et Sch.). *P. citrifolium* Link Jahrb. 1. 3. p. 63 (fide R. et Sch. l. c.). Caulis glaber. Folia fere 4 poll. longa,  $\frac{1}{2}$  poll. lata. Pedunculi 4 lin. longi. Amenta 2 poll. longa.
557. *P. JAVITENSE* (Kunth in H. et B. nov. gen. v. 1. p. 54), foliis petiolatis ova-

*acutis basi inæquali cordatis coriaceis reticulato-venosis rugosis supra glabris subtus incanis tomentoso-hirtis, petiolo margine membranaceo tenuissime cano-tomentoso, amento quam folium brevius cylindrico crassitie pennæ lubrinæ, pedunculo tomentoso-hirto quam petiolus multum brevius.* ♀ In cis opacatis frondosis calidis in ripa fluminis Tuamini prope Javita, Missio-nem del alto Orinoco, alt. 160 hexap. Scandens. — Kunth synopsis. v. 1. p. 110. *Artanthe Javitiensis* Miq. syst. p. 401. Rami pubescentes. Folia 9-10 poll. longa, 6-7 poll. lata. Petioli 2 poll. longi. Pedunculi  $\frac{1}{2}$  poll. longi.

*P. ALARE* (Ham. prodr. pl. Ind. occ. p. 3), foliis alternis sub-9-nerviis proinde cordatis lato-ovatis apice acuminatis, acumine acuto, leviter asperis, petiolis membranaceo-alatis, amento longissime funiculari pedicellato. ♀ In Cayenna. *Artanthe alaris* Miq. syst. p. 406. (Ex Ham. l. c.)

*P. RUGOSUM* (Lam. ill. 1. p. 81), foliis breviter petiolatis ovato-lanceolatis rugosis supra nitidis subtus ramisque villosis, petiolis villosis. ♀ In ins. Sio-Domingo, in Cayenna (Vahl enum. v. 1. p. 313), Trinitate (Griseb. flor. W. Ind. p. 171). *P. bullatum* Vahl l. c. *Artanthe rugosa* Miq. syst. p. 489, Griseb. l. c. Rami scabri. Folia 2-4 poll. longa. (Ex Lam. et Vahl. l. c.)

*P. MICROSTACHYON* (Vahl eclog. amer. fasc. 2. p. 3), foliis oblongis acuminatis utrinque acutis basi æquali lævissimis supra obscure subtus tenuissime venosis, amento brevissimo mucronato. ♀ In Cayenna (Von Rohr fide Vahl l. c.). — Vahl enum. v. 1. p. 315. *Artanthe microstachya* Miq. syst. p. 527. Rami apicem versus leviter pubescentes. Folia 3 poll. longa. Pedunculi quam petioli parum longiores.

*P. RUDGEMANUM*, foliis inferioribus longe superioribus breviter petiolatis ovato-vel-oblongo-lanceolatis attenuato-acutis ætate obtusatis rectis vel subobliquis basi subæquali attenuatis vel acutis utrinque elevato-et pellucido-punctatis subtus pallidis venulis subpatulis utrinque pluribus, amento pedunculum paullo superante brevissimo obtuso deflexo, bractea conchiformi-peltata recte glabra margine antico rotundato subciliolatis, ovario ovato aliquod attenuato, stigmatibus 3. In sylvis maximis prope Loango. *Artanthe Rudgemana* Miq. in Linn. v. 22. p. 77. Folia 5-3 poll. longa, 12-6 lin. lata. Petioli 0-1 poll. longi. Pedunculi 1-1½ poll. longi. Amenta 2-3 lin. longa.

*P. SALICIFOLIUM* (Vahl enum. v. 1. p. 312), foliis petiolatis lanceolatis acutis basi obtusis supra scabris subtus ad nervos villosis subtus nervosis venosis, nervis transversis simplicibus, petiolis villosis, pedunculo solitario quam petiolus parum longiore. ♀ In Surinamo (Vahl l. c.). *Artanthe salicifolia* Miq. syst. p. 533. Rami superne villosi. Folia 2-3 poll. longa. Petioli 2 lin. longi.

*P. DIVARICATUM* (Meyer prim. flor. Essequibo p. 15), foliis erecto-patentibus oblongis acutis basi vix inæqualibus nervoso-venosis utrinque tuberculis minutis punctatis glabris, amentis divaricatis et leviter curvatis, pedunculo petiolum duplo superante. ♀ In sylvaticis Essequibo (Meyer l. c.). *Artanthe? Meyeri* Miq. syst. p. 454. Rami superne granulis minutis dense adpersi glabri. Folia 5 poll. longa, 2 poll. lata.

*P. PARKERIANUM*, foliis brevissime petiolatis lanceolato-oblongis æquilatis apice acuminatis, acumine ipso obtusiusculo vel acuto, basi æquali vel subinæquali acutis crassiuscule membranaceis pellucido-punctatis supra glabris subtus ad nervos hirtellis, centrali nervo subtus prominente nervos subpatulos utrinque 12-16 subtus prominulos infimos 2 magis adscendentes mittente, petiolo dorso hirtello postea glabrescente, pedunculo petiolum bis-quaterve superante, amento brevi cylindrico obtuso, bractea conchiformi recte truncata subpuberula glabrescente, stigmatibus 3 brevibus. ♀ In Dennerara (Parker in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 467). *Artanthe Parkeriana* Miq. l. c. Rami glabri. Ramuli duplici serie pilorum patulorum instructi. Limbi 0,13-0,16 longi, 0,04-0,06 lati. Petioli 0,002-0,005 longi.

*P. CRASSIPICATUM* (Opiz in Reliq. Hænk. fasc. 3. p. 152), foliis ovato-oblongis apice acuminatis basi æquali rotundatis coriaceis glabris reticulato-venosis venis prominentibus, petiolis membranaceo-alatis vaginantibus glabris, amentis pedunculatis cylindricis quam folia duplo fere brevioribus obtusis

- crassitie pennæ anserinæ, pedunculo quam petiolus duplo brevior glabro.  $\frac{5}{2}$  In Panama (Opiz l. c.). *Artanthe crassispicata* Miq. syst. p. 521. Rami glabri.
566. *P. PILOSIUSCULUM* (Opiz in Reliq. Hænk. fasc. 3. p. 154. t. 29. fig. 1), foliis brevissime petiolatis ovatis apice acuminatis, acumine acuto, basi inæquali rotundatis utrinque margineque pilosiusculis, centrali nervo ad  $\frac{3}{4}$  longitud. usque nervos alternos subadscendentes utrinque 6 quorum supremi fere ad apicem ducti mittente, pedunculis brevissimis pilosiusculis, bacca obpyramido-trigona.  $\frac{5}{2}$  In Panama (Opiz l. c.). *Artanthe pilosiuscula* Miq. syst. p. 474. Rami villosi. Limbi 0,19 longi, 0,08 lati. Petioli 0,003 longi.
567. *P. DARIENENSE*, foliis petiolatis ovatis vel ovato-oblongis apice longiuscule acuminatis, acumine acuto, basi æquali acutis vel obtusis membranaceis pellucido-punctatis glabris supra læte viridibus subtus glaucis, nervulis patulis utrinque 6-7 tenuissimis, amentis brevibus pedunculatis (in specim. nondum efformatis).  $\frac{5}{2}$  In nemoribus Darien (Miq. in Seem. bot. Herald p. 200). *Otonia glaucescens* Miq. l. c. Adultum glabrum sed ramuli juniores vix omnino glabri. Folia 4-5  $\frac{1}{2}$  poll. longa, 1  $\frac{1}{2}$ -2  $\frac{1}{2}$  poll. lata. Petioli 2-2  $\frac{1}{2}$  lin. longi.
568. *P. PATULUM* (Bertol. flor. Guatemala p. 7. t. 1), foliis petiolatis inferioribus ovatis apice acuminatis, acumine acuto, basi subinæquali cordatis superioribus oblongo-ovatis apice acuminatis, acumine acuto, basi subæquali, supra glabris subtus ad nervos tantum puberulis 9-nerviis, amento folium superante densifloro.  $\frac{5}{2}$  In Esquintla Guatimalæ (Bertol. l. c.). Caulis glaber. Limbi inferiores 0,15 longi, 0,09 lati. Petioli 0,02 longi.
569. *P. RETICULOSUM* (Opiz in Reliq. Hænk. fasc. 3. p. 155), foliis ovatis acuminatis basi inæquali rotundatis utrinque pilis exasperatis subrugosis margine fimbriato-ciliatis nervoso-venosis subtus lacunosis, petiolis dense villosis, amentis pedunculatis quam folia brevioribus.  $\frac{5}{2}$  In Mexico, Guayaquil (Opiz l. c.). *Artanthe reticulosa* Miq. syst. p. 554. Rami villosi.
570. *P. SCABRIFOLIUM* (Hook. et Arn. in Beech. Voy. p. 310), foliis brevi-petiolatis oblongo-ovatis apice brevi-acuminatis basi inæqualibus subcoriaceis venosis utrinque scabris, amentis brevi-pedunculatis elongatis cylindraceis folio brevioribus.  $\frac{5}{2}$  In Mexico.
571. *P. NERVOSUM*, foliis ovato-lanceolatis apice acuminatis basi vix inæquali acutis utrinque molliter pubescentibus membranaceis nervosis, amentis brevi-pedunculatis cylindraceis quam folium dimidio brevioribus.  $\frac{5}{2}$  In Mexico. *P. patens* Hook. et Arn. in Beech. Voy. p. 310, non Opiz.
572. *P. SALICINUM* (Opiz in Reliq. Hænk. fasc. 3. p. 150), foliis breviter petiolatis lanceolatis glabris nervoso-venosis, petiolis ciliatis.  $\frac{5}{2}$  In Mexico et Panama.
573. *P. CARDIOPHYLLUM*, foliis longe petiolatis cordatis, sinu obtusissimo, acuminatis multinerviis glaberrimis tenerimis pellucido-punctatis reticulato-venosis margine pilis minutis confertissimis ciliatis, amento longe pedunculo cylindrico apice sensim attenuato folium æquante arcuato, pedunculo glabro.  $\frac{5}{2}$  In Mexico (Opiz l. c.). *P. populifolium* Opiz in Reliq. Hænk. fasc. 3. p. 160, non Kunth. *Artanthe cardiophylla* Presl. epim. bot. p. 225 (solum nomen), Miq. syst. p. 391. Rami glabri. Petioli glabri, supra medium membranaceo-alati, vaginantes.
574. *P. ELLIPTICUM* (Hook. et Arn. in Beech. Voy. p. 443), foliis oblongo-ellipticis basi valde inæquali utrinque rotundatis obtusissimis nervosis subtus reticulatis, petiolo brevi amplexante, amento longitudine foliorum.  $\frac{5}{2}$  Ad Realejo regni Mexicani (Hook. et Arn. l. c.). Caulis fruticosus, tuberculatus, maculatus. Folia 4-5 poll. longa.
575. *P. BEGONIÆFOLIUM* (Hook. et Arn. in Beech. Voy. p. 310), foliis oblongo-ovatis basi valde inæqualibus utrinque rotundatis, amentis elongatis cylindraceis brevi-pedunculatis folium æquantibus. In Mexico. Glaberrimum.
576. *P. NITIDULUM* (Opiz in Reliq. Hænk. fasc. 3. p. 154. t. 27. fig. 1), foliis breviter petiolatis ovato-oblongis basi æquali rotundatis apice acuminatis supra glabris subtus ad nervos nervulosque pubescentibus, centrali nervo tota longitudine nervos utrinque 8 nervulosque mittente, amentis cylindricis

mucronulatis quam folium multum brevioribus, pedunculis quam petiolus brevioribus puberulis. ♀ In regno Mexicano (Opiz l. c.). *Artanthe nitidula* Miq. syst. p. 534. Rami juniores pubescentes. Limbi 0,12 longi, 0,042 lati. Petioli 0,005 longi.

577. *P. STIPULACEUM* (Opiz! in Reliq. Hænk. fasc. 3. p. 160), foliis longiuscule petiolatis ovato-subrotundis apice acuminatis, acumine obtuso, basi cordatis utrinque canescenti-tomentosis membranaceis subopacis epunctulatis 11-nerviis, nervis 3 centralibus per anastomoses ad apicem ductis, petiolo dorso canescenti-tomentoso. ♀ In Mexico (Hance! in h. Berol.). *Enckea? stipulacea* Kunth! in Linn. v. 13. p. 600. *Enckea stipulacea* Miq.! syst. p. 366. *Otonia stipulacea* Presl. epimel. p. 231. Ramuli canescenti-tomentosi. Limbi 0,12 longi, 0,09 lati. Petioli 0,03 longi.
578. *P. TRIQUETRUM* (Opiz in Reliq. Hænk. fasc. 3. p. 160), foliis petiolatis ovatis subrotundo-cordatis apice acuminatis glabris ad nervos pilosis 5-nerviis, amento pedunculo cylindraceo quam folium breviori, pedunculo pubescente. ♀ In Mexico (Opiz l. c.). *Enckea californica* Miq. syst. p. 364. ? Rami glabri, juniores leviter pubescentes.
579. *P. PLATYPHYLLUM*, foliis petiolatis amplis ovato-orbiculatis apice acuminatis basi oblique vel inæquali-truncatis subcordatisve subtilissime pellucido-punctatis 9-nerviis, nervis subtus prominentibus, 3 centralibus ad apicem usque ductis, amento baccifero quam folium 3-4-plo-breviore, bractea parva recurva ciliata, staminibus 4? bacca subglobosa pressione subtetragona. ♀ In Manzanilla Bay (Benth. bot. Sulph. p. 167). *Enckea platyphylla* Benth. l. c. Folia usque 10 poll. longa, 8 poll. lata, membranacea, supra demum nitidula, juniora tactu scabriuscula. Petioli 6-9 lin. longi. Amentum 3 poll. longum.
580. *P. MICONIÆFOLIUM*, foliis ovato-oblongis acuminatis basi inæquali cuneatis vel hinc rotundatis subcoriaceis membranaceis subtilissime pellucido-punctatis late viridibus nitidis 5-nerviis, bractea minuta concava, bacca ovoideo-oblonga. ♀ In Salango, nempe littore ditionis Ecuador (Benth. bot. Sulph. p. 167). *Enckea miconiæfolia* Benth. l. c. Rami glabri. Folia 3-4 poll. longa, 1-2 poll. lata. Amenta matura 2-3 poll. longa, baccis distinctis subdistantibus sessilibus.
581. *P. CELTIDIFOLIUM* (Ham. prodr. pl. ind. occ. p. 3), foliis 3-5-nerviis oblique cordatis lanceolatis acutis margine tenuissime cartilagineis glaberrimis. ♀ In Antillis. *P. dubium* A. Dietr. sp. v. 1. p. 694. (Ex Dietr. et Ham. l. c.).
582. *P. SAURUROIDES* (Ham. prodr. pl. ind. occ. p. 3), foliis septemnerviis subobliquis lato-ovatis acuminatis subtus glaberrimis, fructibus dense dispositis submucronatis. In Porto Rico. *Artanthe saururoides* Miq. syst. p. 414.
583. *P. LINGUÆFOLIUM* (Ham. prodr. pl. ind. occ. p. 3), foliis oblique lanceolatis subreticulatis rugosis, petiolis brevissimis marginatis, amentis subincurvis. In montibus Hispaniolæ (Ham. l. c.). *Artanthe? linguæfolia* Miq. syst. p. 533. Caulis incano-rugosus.
584. *P. CUSPIDATUM* (Ham. prodr. pl. ind. occ. p. 3), foliis lato-ovatis acuminatis infra attenuatis obliquis subtus nervoso-reticulatis nervis hirtis. In montibus Hispaniolæ. *Artanthe Wydleriana* Miq. syst. p. 487?
585. *P. LANCIPOLIUM* (Ham. prodr. pl. ind. occ. p. 3), foliis lanceolatis acutis basi oblique rotundatis subtus supraque rugosis nervis magis pubescentibus, amentis rectis. In Jamaica. *P. lanceatum* A. Dietr. sp. v. 7. p. 638. *Artanthe lanceata* Miq. syst. p. 674 (Ex Ham. l. c.).

#### B. Asiaticæ.

586. *P. CHINENSE* (Miq. in Hook. Lond. journ. bot. v. 4. p. 430), foliis petiolatis ovato-vel lato-ellipticis subinæquilateralis apice acute acuminatis basi ima acutiusculis utrinque glabris tenuiter membranaceis epunctatis 5-nerviis, nervis tenuissimis, centrali nervo ad apicem usque lateralibus 2 per anastomoses ad apicem ductis, petiolo glabro, pedunculo petiolum circiter æquante, amentis hermaphroditis cylindricis densifloris, bractea clavato-peltata? ovario ovato, stigmatibus 3 brevibus erectis, staminibus 2 vel pluribus ovario circumposi-

tis, bacca ovato-acuta stigmatum rudimentis coronata. ♀ In China (Millet in h. Hook. fide Miq. in Hook. Lond. Journ. bot. v. 4. p. 430). Ramuli glabri. Limbi 0,11-0,13 longi, 0,05-0,07 lati. Petioli 0,01 longi.

†† Amenta oppositifolia. Plantæ dioicæ.

A. *Species Australasicæ.*

587. P. HUGELIANUM, foliis superioribus modice petiolatis ovatis apice breviter oblique acuminatis basi parum inæqualibus membranaceis obsolete pellucido-punctatis præter nervulum utrinque e basi unum tenuiter quintuplinerviis vix venosis, amento masculino crassiusculo rectiusculo quam folium brevius, pedunculo petiolum circiter æquante, bractea orbiculari margine suberosula extus glabra subtus hirtula centro peltata pedicellata. ♀ In Nova Hollandia ad Port Jackson (Hügel in h. Zuccarini fide Miq. in Linn. v. 26. p. 218). Chavica Hugeliana Miq. l. c. An P. Paramattense C. DC. supra sub n. 468. ? Folia 2½-2 poll. longa, 10-11 lin. lata. Petioli 2-2½ lin. longi. Amenta 1½ poll. longa. (Ex Miq. l. c.)

B. *Ex Asia vel archipelago indico.*

588. P. POTAMOGETONIFOLIUM (Opiz reliq. Hænk. fasc. 3. p. 156), foliis petiolatis cordato-ovatis apice acuminatis basi æqualibus, lobis approximatis, coriaceis glabris nitidis venoso-reticulatis, petiolo vaginante quam folium multo brevius, amento cylindrico quam folium multo brevius mucronato, pedunculo glabro. ♀ In ins. Mariannis (ex Opiz l. c.). Macropiper potamogetonifolium Miq. syst. p. 221.
589. P. LAXUM (Vahl enum. v. 1. p. 326), foliis ovato-oblongis apice acuminatis basi æqualibus 5-nerviis venosis utrinque glaberrimis, petiolis glabris, pedunculis pendentibus, floribus remotiusculis globosis magnitudine P. nigri. ♀ In Philippinis et ins. Franciæ (Vahl l. c.). — Miq. syst. p. 335. Rami glabri. Folia 3-4 pollicaria basi sæpe pollicem lata interdum latiora.
590. P. PARVIFOLIUM (Blanco flor. philip. ed. 1. p. 23) foliis basi cordatis ovato-lanceolatis 3-5-nerviis integris imberbibus, floribus in receptaculo conico magno squamis rotundis margine attenuatis invicem imbricantibus, fructu maturescente parum patentibus tecto. Squamæ illæ sunt stigmata (!! certo bractæ). Stamina 2. Filamenta subnulla. Stylus. In ins. Philippinis (Blanco l. c.) Chavica? celtidiformis Miq. syst. p. 276 pro parte. Chavica? parvifolia Hasskarl in Flora 1864 p. 59.
591. P. MUCRONULATUM (Bl. in Verh. der bat. Genoots. v. 11. p. 218 fig. 27), foliis cordatis obtusiusculis cum mucronulo lobis subæqualibus approximatis infra reticulato-quinquenerviis coriaceis glabris. In Nussa Kambangan (ex Bl. l. c.). Stolones steriles an alicujus Cubeæ speciei? Species delenda (Miq. in Ann. Bot. Mus. lugd. bat. v. 1. fasc. 5. p. 138).
592. P. CRASSUM (Bl. in Verh. der bat. Genoots. v. 11. p. 189. fig. 15), caule suffruticoso striato repente, foliis longiter petiolatis ovatis acuminatis septemnerviis venosis coriaceis glabris. In monte Salak (ex Bl. l. c.). Stolo luxurians sterilis probabiliter ipsius P. nigri. Species delenda (Miq. in Ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 137).
593. P. ALBUM (Vahl enum. v. 1. p. 334), foliis cordatis attenuatis septemnerviis duobus superioribus glabris, petiolis æqualibus. In Java (Vahl l. c.). Rami glabri. Folia 6-7 poll. longa. Petioli 1½ poll. longi. Amenta 3 poll. longa.
594. P. PRIANAMENSE, foliis breviuscule petiolatis e basi rotundata leviter emarginata æquali ovatis vel oblongo-ovatis acuminatis subtus in nervis versus basin densius villosulis 11-nerviis, nervis 5 mediis ad acumen productis infimis parvis transverse venosis, amentis longiuscule vel breviter pedunculatis, pedunculo petiolum superante, baccis bracteisque glabris. In Sumatra occ. prov. Prianam (Miq. in Sumatra p. 473). Chavica multinervia Miq. l. c. Ramuli petiolique densius villosuli. Folia 6-5 poll. longa. Amenta 4-3 poll. longa.



595. *P. ALBIDO-PUNCTATUM* (Bl. in Verh. der bat. Genoots. v. 11. p. 190), foliis petiolatis inferioribus subrotundo-ovatis superioribus oblongo-ovatis oblique acuminatis 5-subquintupli-nerviis coriaceis supra glabris subtus pilis floccosis microscopicis hic illic adpersis albido-punctatis, nervis subtus prominentibus centralibus 3 supra basin ortis duobus e basi ortis infimis tenuissimis brevioribus, pedunculo quam petiolus brevior, amento filiformi. ♂ Supra arborum truncos scandens, in insula Nussa Kambangan ad oram meridionalem Javæ non procul a littore (Bl. in h. lugd. bat. fide Miq. syst. p. 316). *P. albido-punctatum* Bl. enum. pl. Jav. fasc. 1. p. 68, Miq. syst. p. 316, flor. ind. bat. v. 1. pars 2. p. 452. Limbi inferiores 0,11-0,12 longi, 0,08-0,09 lati, superiores 0,14 longi, 0,08-0,07 lati. Petioli 0,025-0,015 longi. (Ex Miq. syst. l. c.). Species in system. delenda ad ramulum sterilem foliosum constituta probabiliter Mulderæ cordatæ vel recurvæ (Piper supra num. 2 et 3) adnumeranda (Miq. ann. bot. Mus. lugd. bat. v. 1. fasc. 5. p. 137).
596. *P. ROSTRATUM* (Roxb. flor. ind. v. 1. p. 162), foliis breviter petiolatis ovato-cuneatis obtuse acuminatis villosis 5-7-nerviis, amentis breviter pedunculatis, baccis rostratis. In Moluccis. Rhyncholepis? Roxburghii Miq. syst. p. 284, flor. ind. bat. v. 1. pars 2. p. 447. Rami villosi. (Ex Roxb.)
597. *P. RUFINERVE* (Opiz Reliq. Hænk. fasc. 3. p. 159), foliis rotundato-ovatis basi subcordatis æqualibus acuminatis glabris 5-nerviis rufis, petiolis glabris? ♂ In ins. Luzon. Rami glabri. *P. rufinerve* Miq. syst. p. 336.
598. *P. DENUDATUM* (Opiz Reliq. Hænk. fasc. 3. p. 158), foliis ovato-oblongis elongatis junioribus dense villosis adultioribus supra glaberrimis subtus ad nervos pilosis sextuplinerviis, petiolis dense longe villosis, amento cylindrico quam folium triplo brevior, pedunculo quam petiolus longiore glabro, baccis globosis distinctis sessilibus. ♂ In ins. Luzon. Rami villosi. Petioli abbreviati. Rhyncholepis Hænkeana Miq. syst. p. 284.
599. *P. HÆNKEANUM* (Opiz Reliq. Hænk. fasc. 3. p. 159), foliis inferioribus exacte cordatis superioribus cordato-ovatis oblongis lanceolatisque 5-nerviis utrinque hirsutis, amento filiformi, pedunculo hirsuto. Ad portum Sorzogon ins. Luzon. ♂ Rami hirsuti.
600. *P. VIMINALE* (Opiz Reliq. Hænk. fasc. 3. p. 150. t. 26. fig. 1), foliis breviter petiolatis lineari-lanceolatis, acumine obtusiusculo, utrinque acutis junioribus basin versus ad nervum puberulis cito glabratissimis membranaceis subtiliter pellucido-punctulatis triplinerviis, centrali nervo paulo supra basin nervum utrinque adscendentem alterne sursumque nervulos validos patulos numerosos mittente, petiolo antice ciliolato, amento masc. brevi, bractea peltata. In ins. Luzon (Hænke n. 78 in h. Vindob.). — Miq. l. syst. p. 336. Chavica? viminalis Miq. l. in Linn. v. 20. p. 131, flor. ind. bat. v. 1. pars 2. p. 444. Rami subverrucosi. Limbi 0,06-0,1 longi, 0,008-0,015 lati. Petioli 0,003-0,004 longi. (v. s.)
601. *P. CELTIDIFORME* (Opiz Reliq. Hænk. fasc. 3. p. 152 t. 26. fig. a), foliis breviter petiolatis ovato-oblongis basi inæqualiter rotundatis acuminatis glabris reticulato-venosis, nervis prominentibus, centrali nervo fere tota longitudine nervos utrinque 6 subalternos quorum inferiores magis approximati mittente, amentis abbreviatis obtusis. ♂ In ins. Luzon (Opiz l. c.). Chavica? celtidiformis Miq. syst. p. 276. Rami glabri. Limbi 0,13 longi, 0,047 lati. Petioli 0,006 longi. (Ex Opiz l. c.)
602. *P. INTERRUPTUM* (Opiz Reliq. Hænk. fasc. 3. p. 157), foliis petiolatis late ovatis subæqualibus 5-nerviis reticulato-venosis glabris subtus pallidioribus, petiolo glabro, amento cylindrico laxo interrupto quam folium multo longiore, pedunculo glabro, fructu lævi patentissimo. ♂ In ins. Luzon ad portum Sorzogon (Opiz l. c.). *P. interruptum* Miq. syst. p. 336. Rami glabri.
603. *P. SUBSEPTEMNERVIUM*, foliis longiuscule petiolatis inferioribus majoribus late subrotundato-ovatis subacuminatis æquilateris vel vix inæquilateris glabris minute pellucido-punctulatis membranaceis 7-nerviis, centrali nervo saltem ad apicem producto passim superne unicostulato infimis basilariis exilibus parum venosis, foliis superioribus e basi obtusa vel acuta inæquali ovatis vel

- ellipticis acutis vel acute apiculatis 5-nerviis vel (accedente nervulo basilar) sub-7-nerviis, bractearum peltis exilibus centro verruculosus glabris ambitu subpuberis. In Pulo Pisang prope Padang. Chavica subseptemnervia Miq. Sumatra p. 186 et 473. Fruticosum, scandens. Folia  $4\frac{1}{4}$ -4 poll. longa. (Ex Miq. l. c.)
604. *P. VENUSTUM* (Wall. list n. 6666!), foliis alternis subsessilibus ovatis basi rotundato-cordatis utrinque glabris pellucidis membranaceis 9-nerviis nervulis transversis. In Singapore. Herba longe repens. Caulis filiformis glaber. Limbi 0,012 longi, 0,01 lati. Amenta ignota. (v. s.)
605. *P. CLYPEATUM* (Wall. list n. 6665  $\alpha$  et  $\beta$ 1), foliis brevissime petiolatis rotundatis aut ovato-rotundatis basi cordatis supra subtusque ad nervos nervulosque hirtellis coriaceis 9-nerviis. In Penang et Singapore (Wall. l. c.), Pulo Pinang! (Gaudich. 1837 in h. Deless.), Manilla (Gaudich. n. 1836! ibid.). *P. obtusissimum* Miq. syst. p. 337. *P. Manillanum* Miq. syst. p. 339. Ramus 0,005 crassus, glaber. Limbi 0,12-0,15 longi latique. (v. s.)
606. *P. FINLAYSONIANUM* (Wall. list n. 6657!), foliis rameis longe petiolatis ramulinis breviter petiolatis e basi cordata lanceolatis apice obtusiusculis supra glabris subtus dense pilosulis quintuplinerviis, centrali nervo supra basin trifido, lateralibus subtilibus, petiolo hirtello, pedunculo masculo hirtello petiolum æquante. Ramuli hirtelli. Limbi 0,055-0,07 longi, 0,02-0,03 lati. Petioli 0,007-0,03 longi. (v. s.)
607. *P. THERMALE* (Vahl enum. v. 1. p. 329), foliis ovato-oblongis attenuatis acutis altero latere baseos parum brevioribus sæpe æqualibus glabris obscure venosis 5-nerviis, bacca globosa quam *P. nigrum* parum minore.  $\S$  In vicinibus thermarum Tranquebarensium copiose solo humido (Konig fide Vahl l. c.). Chavica? thermalis Miq. syst. p. 276. Scandens. Rami glabri. Folia 4 poll. longa. Petioli pollicares. Amenta 4-5 poll. longa. Pedunculi pollicares. (Ex Vahl l. c.)
608. *P. EXASPERATUM* (Vahl enum. v. 1. p. 332), foliis ovato-oblongis parum acuminatis basi inæqualibus supra vix venosis tenuissime alternatim nervosis subtus rugosis, pedunculo quam folium dimidio longiore et crassiore, floribus confertissimis, stigmatibus 3 sessilibus.  $\S$  In India orientali. Chavica? exasperata Miq. syst. p. 271. Rami inferne et punctis minutis elevatis scaberrimis. Folia 3-4 poll. longa. Amentum pollicare. Petioli unguiculares. (Ex Vahl l. c.)
609. *P. RETROFRACTUM* (Vahl enum. v. 1. p. 314), foliis elliptico-lanceolatis valde attenuatis basi obtusis altero latere parum brevioribus obliquis utrinque glabris supra vix venosis alterne tenuissime venosis, pedunculo quam petiolus duplo crassiore et longiore floribus confertissimis.  $\S$  In India orientali. Chavica? retrofracta Miq. syst. p. 275. Folia 5-6 poll. longa,  $1\frac{1}{4}$  poll. lata. Petiolus vix unguicularis. Spica  $1\frac{1}{4}$  poll. longa, obtusissima, penna anserina parum crassior.
610. *P. SYRINGÆFOLIUM* (Vahl enum. v. 1. p. 328), foliis rhombeo-ovatis attenuatis acutis basi æqualibus obtusis glaberrimis lævibus supra nitidiusculis subtus pallidioribus 5-nerviis, pedunculo amentum æquante.  $\S$  In India orientali? Chavica? syringæfolia Miq. syst. p. 275. Rami glabri. Folia 4 poll. longa et ultra, fere 3 poll. lata. Amenta poll. lata, crassitie pennæ columbinæ. (Ex Vahl l. c.)
611. *P. AVANUM* (Wall. h. n. 6661!), foliis petiolatis lanceolatis apice longe acuminatis acutis mucronulatisque basi attenuatis acutiusculis utrinque glabris membranaceis quintupli-nerviis, nervis subtus prominulis, centrali nervo parum supra basin trifido. In Toong Dong Ramuli glabri. Limbi 0,15 longi, 0,06 lati. Petioli 0,03 longi. (v. s.)
612. *P. SUPIGUA* (Don prodr. flor. Nep. p. 20 ex Ham. mss.), caule volubili, foliis ovatis basi rotundatis 5-nerviis glabris, amentis filiformibus folium dimidium æquantibus. In Napalia (ex Don l. c.). Chavica? Suipigua Miq. syst. p. 275.
613. *P. MULLESUA* (Don prodr. flor. Nep. p. 20 ex Ham. mss.), foliis ellipticis acuminatis basi attenuatis acutis trinerviis glabris, amentis globosis peduncu-

latis, petiolo longioribus. In Napalia (ex Don l. c.). Chavica Mullesua Miq. syst. p. 280.

614. P. GUIGAL (Don prodr. flor. Nep. p. 20 ex Ham. mss.), foliis ovatis longe acuminatis basi rotundatis 5-nerviis glabris, amentis globosis pedunculatis petiolo æquantibus. In Napalia (ex Don l. c.). Sirium terrestre Rumph. Amb. v. 5. p. 344. t. 119. fig. 1. (Ex Miq. l. c.)

C. *Species Mascarenenses.*

615. P. MADAGASCARIENSE, foliis alternis petiolatis oblongis æquilateris basi acutis apice acuminatis coriaceis subtilissime pellucido-punctatis glabris multcostulatis costa una utrinque e basi orta, amentis ignotis. In ins. Madagascar. copiose circa Tanimbales in sylvis maximis ubi altissime ad arborum truncos scandit. Piperis albi nomine incolis nota. Folia constricta odorem gratum aromaticum spirant. (Goudot n. 34 in h. Deless. fide Miq. syst. p. 277). Chavica? Madagascariensis Miq. l. c. Ramulus sterilis cylindraceus glaber. Petioli 0,03-0,035 longi. Limbi 0,17-0,18 longi, 0,06-0,075 lati. (Ex Miq. l. c.)
616. P. PERROTTEI, foliis petiolatis e basi rotundato-cordata ovato-acuminatis, acumine obtusiusculo, supra glabris subtus ad nervos hirtellis rigidis opacis pellucido-punctatis 7-9-nerviis, centrali nervo ad  $\frac{1}{2}$  longitud. nervum utrinque patulo-adscendentem subalterne mittente, lateralibus e basi ortis subtilioribus, petiolo hirtello ad limbum usque vaginante. In ins. Bourbon (Perrottet in h. Deless.). P. dubium Miq. syst. p. 339. non Dietr. Scandens, glaber. Limbi 0,05 longi, 0,03 lati. Petioli 0,017 longi. (v. s.)
617. P. PYRIFOLIUM (Vahl enum. v. 1. p. 327), foliis breviter petiolatis ovato-oblongis glabris attenuatis basi subinæqualibus 5-nerviis vix venosis, baccis distinctis oblongis sessilibus semine sinapeos duplo longioribus.  $\frac{3}{4}$  In ins. Bourbon (Commerson, Thouin, fide Vahl l. c.). Chavica? pyrifolia Miq. syst. p. 275. Rami glabri. Folia 3-poll. Amenta  $1\frac{1}{2}$  poll. longa, crassitie pennæ anserinæ, obtusissima. (Ex Vahl l. c.)
618. P. RADICANS (Vahl enum. v. 1. p. 333), foliis petiolatis inferioribus cordato-ovatis acutis obscure 5-nerviis venosis superioribus oblongis acuminatis altero latere baseos parum brevioribus, nervis subtus petiolis cauleque villosis.  $\frac{3}{4}$  In ins. Franciæ (Commerson, fide Vahl l. c.), ins. Luzon (Opiz Reliq. Hænk. fasc. 3. p. 159). P. radicans Opiz l. c. Coccobryon capense Miq. syst. p. 343? Caulis geniculis radicans, crassitie pennæ anserinæ. (Ex Vahl l. c.)

D. *Patriæ ignotæ.*

619. P. BREVISPICATUM (Opiz Reliq. Hænk. fasc. 3. p. 151. t. 28. fig. 1), foliis brevissime petiolatis ovato-oblongis utrinque attenuatis acutis basi subæqualibus supra subtusque pilis minutis oboeitis, centrali nervo fere tota longitudine nervos alternos subadscendentes utrinque 5 nervulosque validos mittente, amentis brevissimis.  $\frac{3}{4}$  Artanthe brevispicata Miq. syst. p. 508. Rami petiolique glabri. (Ex Opiz l. c.)
620. P. RENIFORME (Poir. dict. encycl. v. 5. p. 465), foliis ovato-cordatis glabris acutis basi reniformibus appendiculatis quinquenerviis, amentis lateralibus. In Indiis.

*Species admodum dubiæ aut excludendæ.*

621. P. TILLEFOLIUM (Ham. prodr. fl. Ind. occ. p. 4), foliis oppositis lato-cordato-ovatis acuminatis trinerviis supra nitidis subtus pubescentibus, spadici-bus subgeminis axillaribus difformibus latere irregulariter sulcatis. In Cayenna. — Miq. syst. p. 551. Verisimiliter excludendum.
622. P. BOTRYITES (Vahl enum. v. 1. p. 321), foliis petiolatis ovatis ovato-oblongis elliptico-lanceolatisque attenuatis basi subæqualibus interdum subcordatis venosis lævibus, nervis subtus villosis, amento petiolum parum superante.  $\frac{3}{4}$  In America (h. Jacquin fide Vahl l. c.) Saururus racemosus seu

*Botryites minor* Plum. nov. gen. p. 51. *P. glabrum* Mill. dict. n. 12. *Artanthe botrytis* Miq. syst. p. 533. Rami crassitie pennæ cycnæ teretes, geniculis nodosis. Amentum crassitie pennæ columbinæ. Petiolus semipollicaris vel parum longior. Pedunculus petiolo longior. (Ex Vahl l. c.)

623. *P. MILLII*, foliis cordato-ovatis nervosis acuminatis, spicis reflexis. ♀ *P. decumanum* Mill. dict. n. 9. — *Planta ignota*.
624. *P. RACEMOSUM* (Mill. dict. n. 13). — *Planta ignota*.
625. *P. OBIUSIFOLIUM* (Hill dict. v. 26. p. 19. t. 19. fig. 1), foliis alternis petiolatis obovatis apice breviter mucronulatis eveniis, amentis terminalibus folia æquantibus. ♂ In America meridionali. — *Planta ignota*.
626. *P. ROTUNDIFOLIUM* (Hill l. c. fig. 2), foliis alternis petiolatis rotundatis septemnerviis, amentis sparsis folia æquantibus. *Planta minuta*. In Jamaica. — *Planta ignota*.
627. *P. MACULOSUM* (Hill l. c. fig. 3), foliis alternis longiuscule petiolatis fere medio peltatis fere 8-nerviis. ♂ In America meridionali. Forsan *Peperomia variegata* Ruiz et Pav.?
628. *P. DISTACHYON* (Hill l. c. t. 20. fig. 1), foliis alternis ovatis apice acutis, amentis apice caulis cymose confertis alternis folia æquantibus. Scandens. In America meridionali. — *Planta ignota*.
629. *P. REVOLUTUM* (Rich. fide Kunth nov. gen. v. 1. t. 3. Analysis excellens sine descriptione. — *Planta ignota*.
630. *P. DISCOLORUM* (Desf. Cat. hort. Par. ed. 3. p. 414), foliis alternis cordatis acutis rugulosis superne nitidis subtus pallidis integerrimis quinquenerviis omnibus basi inæqualibus, amentis gracilibus pendulis uncias 3-5-longis. — *Planta ignota*.
631. *P. RHOMBOIDALE* (Desf. l. c.), foliis oppositis rhomboideis 8-9 lin. longis crassiusculis, caule herbaceo semipedali et ultra erecto simplici purpurecente lenti vitreo-pubescentibus. — *Planta ignota*.
632. *P. PURPURASCENS* (Desf. l. c.), foliis quaternis-quinis parvis ovatis ellipticisque integerrimis lævibus carnosissimis crassiusculis subtus convexis purpureis breviter petiolatis, caule herbaceo erecto fere filiformi ramoso villosio purpurascente 5-6-unciali. Forsan *Peperomia rubeola* Hook. et Arn.?
633. *P. CHINANTLENSE* (Mart. et Gal. bull. Acad. Brux. v. 10. n. 2. p. 21), foliis alternis ovato-lanceolatis acuminatis basi inæqualiter subcordatis nervoso-venosis coriaceis nitidis, petiolo alato amplexicauli, amentis axillaribus solitariis breviter pedunculatis petiolo sesquolongioribus. ♀ In sylvis Chinantlæ regione ad declivitatem orientalem Cordilleræ Oaxaca alt. 3000 ped. (Gal. n. 6020? fide Mart. et Gal. l. c.)
634. *P. MIRADORENSE*, foliis alternis patulis oblongo-lanceolatis acuminatis in petiolum longe attenuatis glabris, pedunculo terminali polystachyo gracili, amentis geminis alternis filiformibus distantibus breviter pedicellatis. Ad rupes humidus Mirador alt. 3000 ped. (Gal. n. 6001 fide Martens et Gal. l. c.) *Piper patulum* Martens et Gal. l. c. non Opiz. *Peperomia* species. Caulis carnosus-sulcatus. Folia 3-7 poll. longa. Pedunculus terminalis semipedalis et longior. (Ex Mart. et Gal. l. c.)
- 635 (1). *P. LEPTOSTACHYUM* (Rich. fide Griseb. cat. pl. Cub. p. 65). *Enckea articulata* Griseb. l. c.

*Species e Pipere exclusæ.*

- P. ACUMINATUM* Forst. = *Peperomia rhomboidea* Hook. et Arn.  
*P. ACUMINATUM* Hill = *Peperomia acuminata* Ruiz et Pav.  
*P. ACUMINATUM* Sw. = *Peperomia obliqua* Ruiz et Pav.  
*P. ACUMINATUM* L. = *Peperomia nemorosa* C. DC. et *Pep. Guadaloupensis* C. DC.  
*P. ADSCENDENS* Endl. = *Peperomia Baueriana* Miq.

---

(1) Numerus specierum est revera 634, propter sp. sub num. 101 deletam.

- P. *ÆMULUM* Endl. = *Peperomia reflexa* β.  
 P. *AFFINE* Martens et Gal. = *Peperomia quadrifolia* Kunth.  
 P. *AGGREGATUM* Berg = *Lacistema myricoides* (Prodr. 17. s. 2. p. 591).  
 P. *ALATUM* Vahl = *Peperomia alata* Ruiz et Pav.  
 P. *ALPINUM* Sw. = *Peperomia alpina* A. Dietr.  
 P. *ALSINOIDES* Lam. = *Peperomia alsinoides* A. Dietr.  
 P. *AMPLEXICAULE* Sw. = *Peperomia amplexicaulis* A. Dietr.  
 P. *AMPLEXIFOLIUM* Link = *Peperomia amplexifolia* A. Dietr.  
 P. *ANGULATUM* Rœm. et Sch. = *Peperomia angulata* Kunth.  
 P. *ANGUSTIFOLIUM* Roxb. = *Peperomia Roxburghiana* A. Dietr.  
 P. *AROMATICUM* Willd. = *Peperomia inæqualifolia* Ruiz et Pav.  
 P. *BASELLÆFOLIUM* Rœm. & Sch. = *Peperomia basellæfolia* Kunth.  
 P. *BASELLÆFOLIUM* Wall. = *Peperomia Wallichii* Miq.  
 P. *BILINEATUM* Bl. = *Peperomia bilineata* Miq.  
 P. *BIUNCIALE* Rœm. et Sch. = *Peperomia biuncialis* Kunth.  
 P. *BLANDUM* Jacq. = *Peperomia blanda* Kunth.  
 P. *BØEHMERIÆFOLIUM* Wall. = *Chavica sphærostachya* Miq.  
 P. *BRACHYPHYLLUM* Spr. = *Peperomia trinervis* Ruiz et Pav.  
 P. *BRACHYSTACHYUM* Vahl = *Peperomia filiformis* Ruiz et Pav.  
 P. *BRACHYSTACHYON* Wall. = *Chavica brachystachya* Miq.  
 P. *BRACTEATUM* Thoms. = *Peperomia scandens* Ruiz et Pav.  
 P. *CANDIDUM* Miq. Fl. = *Peperomia candida* Miq. Syst.  
 P. *CILIATUM* Rœm. et Sch. = *Peperomia ciliata* Kunth.  
 P. *CIRCINATUM* Rœm. et Sch. = *Peperomia circinata* Link.  
 P. *CIRCULARE* h. Willd. = *Peperomia circinata* Link.  
 P. *CLUSIÆFOLIUM* Jacq. = *Peperomia obtusifolia*, β.  
 P. *COLORATUM* Rœm. et Sch. = *Peperomia colorata* Kunth.  
 P. *CONCINNUM* Haw. = *Peperomia concinna* A. Dietr.  
 P. *CONJUGATUM* Rœm. et Sch. = *Peperomia conjugata* Kunth.  
 P. *CONVEXUM* Bl. = *Peperomia convexa* Miq.  
 P. *CUNEIFOLIUM* Jacq. = *Peperomia obtusifolia* C. DC.  
 P. *DEPENDENS* h. Willd. = *Peperomia dependens* R. et Pav.  
 P. *DISSIMILE* Rœm. et Sch. = *Peperomia blanda*, β.  
 P. *DISTACHYON* L. = *Peperomia distachya* A. Dietr.  
 P. *DIVERSIFOLIUM* Rœm. et Sch. = *Peperomia diversifolia* Kunth.  
 P. *DOLABRIFORME* Rœm. et Sch. = *Peperomia dolabriformis* Kunth.  
 P. *ELLIPTICUM* Lam. = *Peperomia elliptica* A. Dietr.  
 P. *EMARGINATUM* Vahl = *Peperomia emarginata* Ruiz et Pav.  
 P. *EMARGINELLUM* Sw. = *Peperomia emarginella* C. DC.  
 P. *EXIGUUM* Bl. = *Peperomia hymenophylla* Miq.  
 P. *EXTENSUM* Rœm. et Sch. = *Peperomia elongata* Kunth.  
 P. *FASCICULARE* Rudge ic. t. 4 = *Lacistema myricoides* (Prodr. 17. s. 2. p. 592).  
 P. *FILIFORME* Pers. = *Peperomia filiformis* Ruiz et Pav.  
 P. *FILIFORME* Sw. = *Peperomia Swartziana* Miq.  
 P. *FOLIIFLORUM* Vahl = *Peperomia foliiflora* Ruiz et Pav.  
 P. *FOLIOSUM* Rœm. et Sch. = *Peperomia foliosa* Kunth.

- P. FONTINALE Rœm. et Sch. = *Peperomia fontinalis* Kunth.  
 P. FREIREIFOLIUM Hochst. = *Peperomia hymenophylla*, β.  
 P. GALEOTTIANUM Martens et Gal. = *Peperomia dendrophila* Sehl.  
 P. GALIODES Rœm. et Sch. = *Peperomia galioides* Kunth.  
 P. GLABELLUM Sw. = *Peperomia caulibarbis* Miq. et *Peper. glabella*.  
 P. GLOBOSUM Poir. = *Peperomia umbilicata* Rœm. et Sch.  
 P. HERBACEUM Bot. Mag. in Miq. syst. p. 96, tanquam synonym. citatum, non adest  
 in Bot. Mag., nec nisi error typ. esse videtur.  
 P. HERNANDIÆFOLIUM Vahl = *Peperomia hernandiæfolia* A. Dietr.  
 P. HISPIDULUM Sw. = *Peperomia hispidula* A. Dietr.  
 P. HISPIDUM Sw. = *Peperomia hispidula* Griseb.  
 P. HUMILE Vahl = *Peperomia humilis* A. Dietr.  
 P. HYALINUM Wall. = *Peperomia hymenophylla* Miq.  
 P. HYDROCOTYLIFOLIUM Martens et Gal. = *Peperomia umbilicata* Ruiz. et Pav.  
 P. INÆQUALIFOLIUM Vahl = *Peperomia inæqualifolia* Ruiz et Pav.  
 P. INCANUM Spr. = *Peperomia incana* A. Dietr.  
 P. INCISUM Rich. ined. = *Peperomia quadrifolia* Kunth.  
 P. LÆVIFOLIUM Bl. = *Peperomia lævifolia* Miq.  
 P. LAXIFLORUM Spr. = *Peperomia laxiflora* Kunth.  
 P. MACRORHIZON Rœm. et Sch. = *Peperomia macrorhiza* Kunth.  
 P. MACULOSUM L. = *Peperomia maculosa* A. Dietr.  
 P. MAGNOLIÆFOLIUM Jacq. = *Peperomia magnoliæfolia* C. DC.  
 P. MARGINATUM Haw. = *Peperomia obtusifolia*, β.  
 P. MAURITIANUM Rœm. et Sch. = *Peperomia elliptica* A. Dietr.  
 P. MAYPURENSE Rœm. et Sch. = *Peperomia Maypurensis* Kunth.  
 P. MICROPHYLLUM Rœm. et Sch. = *Peperomia microphylla* Kunth.  
 P. MOLLE Rœm. et Sch. = *Peperomia mollis* Kunth.  
 P. MONOSTACHYON Vell. = *Peperomia major* C. DC.  
 P. MONOSTACHYON Vahl = *Peperomia monostachya* Ruiz et Pav.  
 P. MUCRONATUM h. Francov. = *Peperomia melanostigma* Miq.  
 P. MURINUM Rœm. et Sch. = *Peperomia pyrifolia* Humb. et B.  
 P. MUSCOSUM Rœm. et Sch. = *Peperomia muscosa* Link.  
 P. MYOSUROIDES Rudge = *Peperomia myosuroides* A. Dietr.  
 P. MYRTIFOLIUM Vahl = *Peperomia myrtifolia* A. Dietr.  
 P. NEMATOSTACHYON Rœm. et Sch. = *Peperomia nematostachyon* Link.  
 P. NEMOROSUM Vahl = *Peperomia nemorosa* C. DC.  
 P. NERVO-VENOSUM Bl. = *Peperomia nervo-venosa* Miq.  
 P. NUMMULARIFOLIUM Lam. et Willd. herb. 726! = *Filix sterilis*.  
 P. OBTUSIFOLIUM Jacq. = *Peperomia polystachya* Miq.  
 P. OBTUSIFOLIUM Lam. = *Peperomia obtusifolia* β.  
 P. OBTUSIFOLIUM L. = *Peperomia obtusifolia* C. DC.  
 P. OBVERSUM Vahl = *Peperomia obversa* Griseb.  
 P. ORBICULATUM Poir. = *Peperomia orbiculata* Dietr.  
 P. OVALIFOLIUM Heyne = *Peperomia Dendygulensis* Miq.  
 P. PALLENS Rœm. et Sch. = *Peperomia pallens* Kunth.  
 P. PARVULUM Martens et Gal. = *Peperomia Mexicana* Miq.

- P. PEEPULOIDES Rœm. et Sch. = *Peperomia peepuloides* Kunth.  
 P. PEEPULOIDES Roxb. = *Chavica peepuloides* Miq.  
 P. PELLUCIDUM L. = *Peperomia pellucida* Kunth.  
 P. PELTOIDEUM Rœm. et Sch. = *Peperomia peltoides* Kunth.  
 P. PERESKIAEFOLIUM Jacq. = *Peperomia pereskiaefolia* Kunth.  
 P. PHYLLANTHUS h. ber. = *Peperomia foliiflora* Ruiz et Pav.  
 P. PINNATUM Lour. fl. Coch. p. 38; Vahl en. v. 1. p. 351; Miq. syst. p. 350.  
 recte a cl. Miq. exclusum = ign.  
 P. POLYBOTRYON Spr. = *Peperomia polybotrya* Kunth.  
 P. POLYSTACHYUM Vahl = *Peperomia polystachya* Miq.  
 P. PORTULACÆFOLIUM Rœm. et Sch. = *Peperomia portulacæfolia* Kunth.  
 P. PORTULACOIDES Lam. = *Peperomia portulacoides* A. Dietr.  
 P. PUBESCENS Pers. = *Peperomia pubescens* Ruiz et Pav.  
 P. PUBESCENS Vahl = *Peperomia Vahlia* A. Dietr.  
 P. PULCHELLUM Vahl = *Peperomia pulchella* A. Dietr.  
 P. PURPUREUM Vahl = *Peperomia purpurea* Ruiz et Pav.  
 P. PUSILLUM Bl. = *Peperomia reflexa* A. Dietr.  
 P. QUADRIFOLIUM Vell. = *Peperomia reflexa* var. *t.*  
 P. RACEMOSUM Vahl = *Peperomia secunda* Ruiz et Pav.  
 P. RECURVATUM Bl. = *Peperomia recurvata* Miq.  
 P. REFLEXUM L. f. = *Peperomia reflexa* A. Dietr.  
 P. RENIFORME Willd. = *Verhuellia elegans* Miq.  
 P. RETUSUM L. f. = *Peperomia retusa* A. Dietr.  
 P. RHOMBEUM Vahl = *Peperomia rhombea* Ruiz et Pav.  
 P. ROLLINGO Rœm. et Sch. = *Peperomia Rollingo* Link.  
 P. RUBIODES Rœm. et Sch. = *Peperomia rubioides* Kunth.  
 P. RUBRICANTE Link = *Peperomia pereskiaefolia* Kunth.  
 P. RUSCIFOLIUM Rœm. et Sch. = *Peperomia ruscifolia* Kunth.  
 P. SALIGNUM Rœm. et Sch. = *Peperomia saligna* Kunth.  
 P. SAXATILE Wall. = *Peperomia reflexa* A. Dietr.  
 P. SAXICOLUM Vahl = *Peperomia pubescens* Ruiz et Pav.  
 P. SCANDENS Sw. = *Peperomia glabella* Miq.  
 P. SCUTELLÆFOLIUM Vahl = *Peperomia scutellæfolia* Ruiz et Pav.  
 P. SCUTELLATUM Pers. = *Peperomia scutellæfolia* Ruiz et Pav.  
 P. SECUNDUM Poir. = *Peperomia secunda* Ruiz et Pav.  
 P. SEPTEMNERVE Vahl = *Peperomia septemnervis* Ruiz et Pav.  
 P. SERPENS Sw. = *Verhuellia serpens* Miq.  
 P. SERPYLLIFOLIUM Vahl = *Filix sterilis*, et (pro parte) *Peperomia serpyllifolia*  
 A. Dietr.  
 P. SIMPLEX Endl. = *Peperomia Endlicheri* C. DC.  
 P. STELLATUM Jacq. = *Peperomia stellata* A. Dietr.  
 P. STELLATUM Vell. = *Peperomia pereskiaefolia* Kunth.  
 P. STOLONIFERUM Rœm. et Sch. = *Peperomia stolonifera* Kunth.  
 P. SUAVEOLENS Desf. = *Peperomia suaveolens* Ham.  
 P. SUBROTUNDUM Bl. = *Peperomia Reinwardtiana* Miq.  
 P. TALINIFOLIUM Rœm. et Sch. = *Peperomia talinifolia* Kunth.

- P. TENELLUM Sw. = *Peperomia tenella* A. Dietr.  
 P. TETRAGONUM Vahl = *Peperomia tetragona* Ruiz et Pav.  
 P. TETRAPHYLLUM Forst. = *Peperomia reflexa* A. Dietr.  
 P. TITHYMALOIDES Vahl = *Peperomia tithymaloides* A. Dietr.  
 P. TOMENTOSUM Vahl = *Peperomia tomentosa* A. Dietr.  
 P. TRICARINATUM Haw. = *Peperomia tricarinata* A. Dietr.  
 P. TRIFOLIUM L. = *Peperomia trifolia* A. Dietr.  
 P. TRIFOLIUM Willd. herb. = *Peperomia portulacoides* A. Dietr.  
 P. TRINERVE Vahl = *Peperomia trinervis* Ruiz et Pav.  
 P. UMBILICATUM Vahl = *Peperomia umbilicata* Ruiz et Pav.  
 P. UMBROSUM Bl. = *Peperomia nervo-venosa* Miq.  
 P. VARIEGATUM Pers. = *Peperomia maculosa* A. Dietr.  
 P. VERTICILLATUM L. = *Peperomia pulchella* A. Dietr.  
 P. VILLOSUM Rich. ined. = *Peperomia rotundata* Kunth.

### *Index specierum et synonymorum.*

(Numeri ad numerum speciei respondent.)

P. *abbreviatum* Opiz 437, *abutilifolium* C. DC. 553, *abutiloides* Kunth 372, *Acarouanyanum* C. DC. 303, *acre* Bl. 448, *aculeatum* C. DC. 254, *acuminatum* Bot. mag. 635, *acutifolium* R. et P. 114, *adenandrum* C. DC. 130, *adenophorum* C. DC. 134, *aduncum* L. 184, *æquale* Vahl 299, *affine* C. DC. 188, *Afetianum* Röm. et Sch. 384, *alare* Ham. 558, *albido-punctatum* Bl. 595, *albidum* Kunth 537, *album* Vahl 593, *alnoides* Kunth 179, *alpinum* C. DC. 364, *alveolatum* Opz 528, *Amalago* L. 26, *Amalago* Rheed. 478, *Amazonicum* C. DC. 126, *Amboinense* C. DC. 440, *amplectens* C. DC. 215, *amplifolium* C. DC. 88, *andicolum* Kunth 334, *androgynum* C. DC. 431, *angustifolium* Lam. 230, *angustifolium* Pav. 349, *angustifolium* R. et P. 185 et? 119, *anisatum* Kunth 20, *anonæfolium* Kunth 244, *apiculatum* C. DC. 133, *appendiculatum* C. DC. 535, *arborescens* Miq. 500, *arborescens* Roxb. 488, *arcuatum* Bl. 490, *arcuatum* var. Miq. 436, *arcuatum* Opiz 531, *arcuatum* Thw. 481, *areolatum* C. DC. 267, *argyrophyllum* Miq. 504, *aristolochioides* Lam. 406, *Arnottianum* C. DC. 463, *Artanthe* C. DC. 1:1, *artanthopse* C. DC. 354, *asperifolium* R. et P. 314, *asperiusculum* Kunth 145, *attenuatum* Miq. 503, *attenuatum* h. Willd. 13, *augustum* Rudge 292, *aurantiacum* Wall. 480, *auriculatum* C. DC. 551, *auriculatum* Bl. 473, *auritum* Kunth 335, *Australasicum* C. DC. 467, *Austro-caledonicum* C. DC. 434, *Avanum* Wall. 611, *avellanum* C. DC. 258.

P. *baccans* C. DC. 157, *baccatum* Bl. 3, *Bahiense* C. DC. 550, *Banksii* Miq. 418, *Bantamense* Bl. 500, *Baptisianum* C. DC. 48, *barbatum* Kunth 333, *Barclayanum* C. DC. 392, *Bartlingianum* C. DC. 65, *Baryanum* C. DC. 125, *begoniæfolium* Hook. et Arn. 575, *Bennetianum* C. DC. 189, *Benthamianum* C. DC. 95, *Berlandieri* C. DC. 229, *Berteroanum* C. DC. 30, *Bette* L. 489, *Birmanicum* C. DC. 400, *Blanchetii* C. DC. 327, *blattarum* Spreng. 370, *boehmeriæfolium* Wall. 445, *Bogotense* C. DC. 221, *Boissierianum* C. DC. 276, *Boivinii* C. DC. 476, *Bolivianum* C. DC. 159, *Bontianum* C. DC. 313, *Borbonense* C. DC. 406, *Botryies* Vahl 622, *Boucheanum* C. DC. 50, *brachypodon* C. DC. 359, *brachystachyum* C. DC. 231, *Brasiliense* C. DC. 73, *Bredemeyeri* Jacq. 122, *brevispicatum* Opiz 619, *Brongniartii* C. DC. 363, *bullatum* Vahl 559, *bullosum* C. DC. 423, *Burmanum* Miq. 477.

P. *Cabellense* C. DC. 277, *Cabralanum* C. DC. 171, *caladiifolium* C. DC. 374, *calceolatum* C. DC. 284, *Californicum* Opiz 33, *callosum* Opiz 478, *callosum* Ruiz et Pav. 77, *calocoma* C. DC. 96, *Cambessedesii* C. DC. 123, *canaliculatum* Kl. 473, *canaliculatum* Opiz 489, *caninum* A. Dietr. 412, *Capense* L. f. 405, *Caraboboense* C. DC. 74, *Caracasenum* Bredem. 279, *cardiophyllum* C. DC. 573, *Caripense* Kunth 539, *carpinifolium* C. LC. 57, *Carpunya* R. et P. 371, *Casapiense* C. DC. 522, *cassinoides* Opiz 530, *catalpæfolium* Kunth 20, *caudatum* Vahl 20,



ceanothifolium Kunth 24, Celebicum Bl. 433, *celtidifolium* Desf. 34, *celtidifolium* Ham. 581, *celtidifolium* Kunth 184, celtidiforme Opiz 601, Ceramicum C. DC. 451, cernuum Vell. 280, *cernuum* Opiz 545, Ceylanicum C. DC. 8, Chaba Bl. 437, Chamissonis Steud. 174, chavicoide C. DC. 282, Chawya Ham. 458, chimonanthifolium Kunth 156, Chilense C. DC. 521, Chinantlense Mart. et Gal. 633, Chinense Miq. 586, Churumayu R. et P. 265, *Churya* Hunter, 463, cinereum C. DC. 228, citrifolium Lam. 118, *citrifolium* Link 556, Clausenianum C. DC. 143, Clusii C. DC. 407, clypeatum Wall. 605, coccoloboides Kunth 195, *colonom* Presl 500, colubrinum Link 329, *colubrinum* var. Miq. 255, *commutatum* Steud. 375, concinnum C. DC. 365, confusum C. DC. 117, consanguineum Kunth 230, Corafalgense C. DC. 104, Corcovadense C. DC. 55, cordatum C. DC. 98, Cordillerianum C. DC. 382, cordulatum C. DC. 241, cornifolium Kunth 343, coruscans Kunth 211, *corylifolium* Kunth 312, corylistachyon C. DC. 435, Cosmianum C. DC. 63, Costaricense C. DC. 366, crassifolium C. DC. 255, crassinervium Kunth 345, crassipes Korthals 425, crassispicatum Opiz 565, crassum Bl. 592, *crassum* Vell. 329, crocatum R. et P. 116, Cubeba L. fil. 410, Cubense C. DC. 393, Cumanense Kunth 316, Cumaralense C. DC. 87, Cumbotianum C. DC. 109, Cumingianum Miq. 515, curvatum R. et P. 523, cuspidatum Ham. 584, cyrtopodon C. DC. 397.

P. Darienense C. DC. 567, dasyoura C. DC. 113, dasypodium C. DC. 270, decipiens C. DC. 132, decrescens C. DC. 39, *decumanum* Aubl. et Willd. 20, *decumanum* L. 543, *decumanum* L. 450, *decumanum* Mill 623, decurrens C. DC. 261, Demeraranum C. DC. 243, *densespicatum* Steud. 329, *densum* Bl. 489, denudans C. DC. 348, denudatum Opiz 598, Descourtilsianum C. DC. 167, dichotomum R. et P. 79, *diffusum* Bl. 460, *diffusum* Vahl 503, dilatatum Rich. 312, 312, dimidiatum C. DC. 545, diospyrifolium Kunth 252, *discolor* Sw. 23, discolorum Desf. 630, distachyon Hill 628, divaricatum Meyer 563, divergens C. DC. 547, Dombeyanum C. DC. 381, *dubium* A. Dietr. 581, *dubium* Miq. 616, Duchassaingii C. DC. 40, dumosum Rudge 111.

P. Egense C. DC. 94, ellipticum Hook. et Arn. 574, *ellipticum* Rœm. et Sch. 433, elongatum C. DC. 11, *elongatum* Vahl 185, Enckea C. DC. 15, epunctulatum C. DC. 399, eriopodon C. DC. 183, Estrellense C. DC. 127, eucalyptifolium Rudge 340, eucalyptiphyllum C. DC. 43, *evonymifolium* Kunth 92, exasperatum Vahl 608, excavatum R. et P. 524, excelsum Forster 387, *eximium* Kunth 281 β, exserens C. DC. 82.

P. Fadyeni C. DC. 181, *falcatum* A. Dietr. 531, *fallax* Vahl, 502, Fendlerianum C. DC. 120, Finlaysonianum Wall. 606, firmum C. DC. 6, Fischerianum C. DC. 148, fistulosum C. DC. 164, *flagellare* Kunth 281, flavicans C. DC. 216, flexicaule C. DC. 289, flexuosum Jacq. 149, *flexuosum* Rudge 289, Flumnense C. CC. 285, formosum C. DC. 187, Forstenii C. DC. 443, foveolatum Kuntz 369, fragile Benth. 484, Francovilleianum C. DC. 68, Fraseri C. DC. 112, Friedrichsthali C. DC. 362, *fruticosum* Rich. 314, fuligineum Kunth 320, Funckii C. DC. 297, fuscescens C. DC. 208, fuscum C. DC. 331, Futokadsura Sieb. 436.

P. Gabrielianum C. DC. 108, galeatum C. DC. 5, Gardnerii C. DC. 51, Gaudichaudianum Kunth 147, geniculatum Sw. 105, gigantifolium C. DC. 278, glabellum C. DC. 305, glabrescens C. DC. 124, *glabrum* Mill. 621, glanduligerum C. DC. 240, *glandulosum* Opiz 473, *glaucescens* Jacq. 34, gracile R. et P. 182, Granatense C. DC. 266, grande Vahl 542, grandifolium Kunth 296, *grandifolium* Ekelund 384, *granulosum* Ruiz et Pav. 185, Griffithii C. DC. 422, griseum C. DC. 169, Guahamense C. DC. 391, Guianense C. DC. 253, Guigal Don 614, Guineense Schum. 420.

P. Haenkeanum Opiz 599, halesiaefolium Kunth 290, Hamiltonii C. DC. 491, Hartwegianum C. DC. 536, Hayneanum C. DC. 45, hederaceum C. DC. 517, heptandrum C. DC. 16, heterophyllum R. et P. 307, Hilarianum C. DC. 336, Hillianum C. DC. 202, hirsutum Sw. 145, hirtellum C. DC. 52, hirtulum C. DC. 549, hispidum Kunth 538, *hispidum* Martens et Gal. 144, Hoffmanseggianum Rœm. et Sch. 556, Hohenackeri C. DC. 12, Holtonii C. DC. 249, Hongkongense C. DC. 441, Hookeri Miq. 514, Hookerianum C. DC. 213, Hostmannianum C. DC. 191, Hugelianum C. DC. 587, *hydrangeifolium* Kunth 345, hymenophyllum Miq. 506, hypoglauca Miq. 479.

*P. imperiale* C. DC. 403, *impunctatum* Willd. 329, *inæquale* C. DC. 256, *incurvum* Sieb. 223, *indecorum* Kunth 91, *indicum* C. DC. 498, *insectifugum* C. DC. 469, *insigne* Kunth 198, *insulare* C. DC. 497, *interruptum* Opiz 602.

*P. Jaborandi* Vell. 41, *Jacquemontianum* Kunth 118, *Jacquinii* C. DC. 161, *Jacquianum* Roem. et Schult. 149, *Jalapense* C. DC. 146, *Jalapense* Mart. et Gal. 346, *Japurense* C. DC. 90, *Javanicum* C. DC. 421, *Javitense* Kunth 557, *Jayeri* C. DC. 486, *juliflorum* Nees et Mart. 237, *Jussæanum* Kunth 292.

*P. Kappleri* C. DC. 302, *Karwinskianum* Kunth 361, *Kegelianum* C. DC. 552, *Kegelii* C. DC. 344, *Kewense* C. DC. 360, *Khasianum* C. DC. 449, *Klotzschianum* C. DC. 62, *Korthalsii* Miq. 511, *Kunthii* C. DC. 37.

*P. Labillardieri* C. DC. 495, *lacunosum* Kunth 263, *lætum* C. DC. 60, *læve* Vahl 496, *lævigatum* Kunth 13, *lanatum* Roxb. 413, *lanatum* Miq. 505, *lance-folium* Schott 348, *lanceæfolium* Kunth 323, *lanceatum* A. Dietr. 585, *lanceolatum* Roxb. 446, 473, *lanceolatum* R. et P. 92, *lanceifolium* Ham. 585, *Langdorffianum* C. DC. 201, *lapathifolium* Steud. 346, *latifolium* Lam. 384, *latifolium* L. f. 389, *lætum* Kunth 18, *laurifolium* Kunth 328, *laxum* Vahl 559, *Lechlerianum* C. DC. 115, *Ledebourii* C. DC. 166, *Lehmannianum* C. DC. 527, *leio-phyllum* C. DC. 238, *Lenormandianum* C. DC. 246, *lentiginoides* Kunth 306, *lenticellosum* C. DC. 330, *Leonense* C. DC. 416, *leptophyllum* C. DC. 155, *leptostachyum* Griseb. 14, *leptostachyum* Wall. 498, *leptostachyum* Rich. 635, *lepturum* Kunth 332, *Lessertianum* C. DC. 67, *Lessertianum* C. DC. olim 456, *leucophyllum* C. DC. 151, *Lhotzkyanum* C. DC. 193, *Lindbergii* C. DC. 234, *Lindenii* C. DC. 28, *lineatum* R. et P. 301, *linguæfolium* Ham. 583, *Linkii* Miq. 485, *lonchites* Miq. 487, *longifolium* R. et P. 72, *longum* L. 474, *longum* Piso 478, *longum* Bl. 478, *longum* Rumph. 478, *Lowong* Bl. 426, *Lucæanum* Kunth 341, *Luzonense* C. DC. 452.

*P. Machadoense* C. DC. 49, *Macgillivrayi* C. DC. 388, *macrophyllum* Kunth 128, *macrophyllum* Loud. 232, *macrophyllum* Sw. 105, *macrorhynchon* C. DC. 89, *macrostachyum* C. DC. 548, *macrourum* Kunth 103, *maculatum* Bl. 439, *maculosum* Hill 627, *Madagascariense* C. DC. 615, *magicum* Reinw. 450, *majusculum* Bl. 450, *Malabarenses* C. DC. 7, *malacophyllum* C. DC. 395, *Malami-ri* Bl. 442, *Mandonii* C. DC. 142, *Manillanum* Miq. 605, *Marequitense* C. DC. 204, *marginatum* Jacq. 20, *Mariannum* Opiz 489, *maritimum* Bl. 478, *Maro-winianum* C. DC. 396, *Marowynense* C. DC. 107, *Martensianum* C. DC. 38, *Martianum* Kunth 207, *Mastersianum* C. DC. 170, *Mathewsii* C. DC. 83, *Mayna-sianum* C. DC. 86, *medium* Jacq. 23, *Melamiri* L. 489, *Melantris* Hill 478, *Melamiris* Lam. 489, *Melamiris* Roxb. 492, *melanostictum* C. DC. 137, *melas-tomoides* Cham. et Schl. 19, *membranaceum* C. DC. 518, *methysticum* Forster 470, *methysticum* Lam. 389, *methysticum* Roxb. 450, *Mexicanum* C. DC. 144, *miconiæfolium* C. DC. 580, *microstachyon* Vahl 560, *microstigma* C. DC. 457, *Miersium* C. DC. 150, *Millii* C. DC. 623, *Minarum* C. DC. 269, *miniatum* C. DC. 473, *Miquelianum* C. DC. 53, *Miradorense* C. DC. 634, *Misanthense* C. DC. 186, *mite* R. et P. 401, *modestum* C. DC. 140, *mollicomum* Kunth 321, *mollipilosum* C. DC. 85, *mollissimum* Bl. 417, *montanum* C. DC. 205, *Morisonianum* C. DC. 300, *Moritzianum* C. DC. 196, *mucronulatum* Bl. 591, *Muelleri* C. DC. 10, *Mullesua* Don 613, *multinervium* Mart. et Gal. 555, *multiplernvium* C. DC. 218, *Munchanum* C. DC. 70, *muricatum* Bl. 414, *Murrayanum* C. DC. 163.

*P. Nagelii* C. DC. 510, *Neesianum* C. DC. 58, *Negroense* C. DC. 235, *nema-tanthera* C. DC. 519, *Nepalense* Miq. 483, *nervosum* C. DC. 571, *nervulosum* C. DC. 110, *nigrescens* Bl. 482, *nigrescens* Miq. 447, *nigrum* L. 502, *nigrum* Miq., *Schnizl.* 502, *Nilghirianum* C. DC. 505, *nitidulum* Opiz 576, *nitidum* Sw. 105, *nitidum* Vahl 138, *nobile* C. DC. 197, *nodulosum* Linn. 105, *Novæ-Hispaniæ* C. DC. 257, *Novæ-Hollandiæ* Miq. 422, *Novo-Granatense* C. DC. 310, *nubigenum* Kunth 116, *nudum* C. DC. 352, *nummularifolium* Lam. 636, *nutans* Opiz 103.

*P. Oaxacanum* C. DC. 136, *obesum* Vahl 525, *obliquum* R. et P. 281, *oblon-gatum* Opiz 532, *oblongifolium* C. DC. 131, *oblongum* Cham. et Schl. 174, *oblon-gum* Kunth 304, *obovatum* R. et P. 106, *obtusifolium* Hill 625, *obtusissimum* Miq. 605, *obtusum* C. DC. 368, *obumbratum* C. DC. 102, *officinatum* C. DC.

478, *Olfersianum* Kunth 145 ♂, *Opizianum* Flora 531, *orthostachyum* Kunth 25, *Ottoneifolium* C. DC. 206, *Ottonis* C. DC. 379, *ovale* C. DC. 80, *ovalifolium* C. DC. 274, *ovatum* Vahl 47.

*P. pachystachyon* C. DC. 268, *padifolium* C. DC. 46, *paludosum* C. DC. 177, *palustre* C. DC. 217, *Panamense* C. DC. 129, *panduratum* C. DC. 36, *Pangoense* C. DC. 526, *Papantlense* C. DC. 402, *Paraense* C. DC. 203, *Paramaribense* C. DC. 236, *Paramattense* C. DC. 468, 587, *Parkerianum* C. DC. 564, *Parthenium* Mart. 373, *parvifolium* Planco 590, *patens* Hook. et Arn. 571, *patens* Opiz 533, *patulum* Bertol. 568, *patulum* Martens et Gal. 634, *Paulianum* C. DC. 222, *Pavoni* C. DC. 220, *pedicellare* C. DC. 294, *pedicellatum* C. DC. 453, *pedicellosum* Wall. 423, *pedunculare* C. DC. 209, *pedunculatum* C. DC. 385, *peltatum* L. 378, *peltatum* Ruiz et P. 381, *P. Peltobryon* C. DC. 71, *Penangense* C. DC. 464, *Perrottetii* C. DC. 616, *persicariæfolium* Kunth 311, *Peruvianum* C. DC. 175, *petiolare* C. DC. 160, *Philippinense* C. DC. 466, *Philippinum* Miq. 444, *Philippinum* Miq. syst. 515, *phthiotricon* C. DC. 242, *phyllostictum* C. DC. 409, *phytolaccæfolium* Opiz 534, *pilosiusculum* Opiz 566, *piluliferum* Kunth 404, *Pisoense* C. DC. 152, *plantagineum* Lam. 26, *plantagineum* Schl. 375, *platyphyllum* C. DC. 579, *Plukenetianum* C. DC. 317, *Pœppigii* C. DC. 78, *Pohlmanum* C. DC. 61, *Poiretianum* C. DC. 353, *Poiteanum* Kunth 369, *populifolium* Kunth 339, *populifolium* Opiz 573, *potamogetonitifolium* Opiz 588, *pothifolium* Kunth 200, *pothiforme* Wall. 455, *pothoides* Wall. 454, *Potomorphe* C. DC. 286, *præmorsum* Roth 138, *pregnans* Rich. herb. 191, *Preslii* C. DC. 66, *Priaramense* C. DC. 594, *propinquum* C. DC. 81, *pruinosum* Willd. 378, *prunifolium* Jacq. 158, *pseudo-asperifolium* C. DC. 325, *pseudo-Chavica* C. DC. 456, *pseudo-Churumayu* C. DC. 192, *pseudo-Cubeba* Korth. 419, *pseudo-mollicomplum* C. DC. 153, *pseudo-peruvianum* C. DC. 173, *pseudo-pothifolium* C. DC. 199, *pseudo-velutinum* C. DC. 172, *psilostachyum* Kunth 541, *pteropodon* C. DC. 56, *pubescens* Sol. 418, *pulchrum* C. DC. 275, *punctatissimum* C. DC. 54, *punctatum* Ruiz et P. 295, 299?, *Purdiei* C. DC. 247, *purpurascens* A. Dietr. 185, *purpurascens* Desf. 632, *pyrifolium* Opiz 34, *pyrifolium* Vahl 617.

*P. quinqueangulatum* Miq. 509, *Quitense* C. DC. 398.

*P. racemosum* Mill 624, *radicans* Vahl 618, *radula* Kunth 122, *Rafflesii* C. DC. 446, *rarum* C. DC. 180, *recurvum* Bl. 2, *Regelii* C. DC. 291, *Regnellii* C. DC. 283, *Reinwardtianum* C. DC. 472, *reniforme* Poir. 620, *reticulatum* L. 227, *reticulatum* Vell. 24, *reticulosum* Opiz 569, *retrofractum* Vahl 609, *revolutum* Rich. 629, *rhododendrifolium* Kunth 244, *rhomboidale* Desf. 631, *rhombophyllum* C. DC. 462, *rhyncholepis* C. DC. 427, *rhynchostachyum* C. DC. 356, *ribesoides* Wall. 415, *richardiæfolium* Kunth 272, *Richardianum* C. DC. 32, *Riedelii* C. DC. 351, *rigidum* C. DC. 99, *riparium* Kunth 296, *rivinoide* Kunth 306, *Rohrii* C. DC. 232, *rostratum* Roxb. 596, *rotundifolium* Hill 626, *rubescens* C. DC. 251, *rude* Kunth 178, *Rudgeanum* C. DC. 561, *rufinerve* Opiz 597, *rugosum* Lam. 559, *rugosum* Opiz 116, *P. rugosum* Vahl 118, *rugulosum* C. DC. 168, *Ruizianum* C. DC. 287, *rumicifolium* C. DC. 239, *Rumphii* C. DC. 471.

*P. Sagoti* C. DC. 520, *Sagranum* C. DC. 212, *salicariæfolium* Kunth 322, *salicifolium* Vahl 562, *salicinum* Opiz 572, *salviæfolium* C. DC. 119, *Salzmanni* C. DC. 327, *San-Carlosianum* C. DC. 394, *sanctum* Schlecht. 376, *Santaremsense* C. DC. 139, *sarmentosum* Roxb. 460, *saururoides* Ham. 582, *scabridum* Flora 529, *scabrifolium* Hook. et Arn. 560, *scabriusculum* A. Dietr. 529, *scabrum* R. et P. 525, *scabrum* Lam. 184, *scabrum* Opiz 529, *scabrum* Sw. 145, *Schachii* C. DC. 194, *Schizonephos* C. DC. 1, *Schlechtendahlia* Kunth C. DC. 349, *Schlimii* C. DC. 135, *Schottii* C. DC. 315, *scopulorum* Kunth 544, *scutatum* Willd. 378, *scutiphyllum* Ham. 380 et 378?, *Sebastianopolitense* C. DC. 44, *secundum* R. et P. 250, *Seemannianum* C. DC. 438, *Sellovianum* Kunth 350, *Selloi* C. DC. 42, *septuplennervium* C. DC. 308, *sidæfolium* Link et O. 384 var. β, *sidæfolium* Mart. 383, *Sieberi* C. DC. 27, *Sicensense* C. DC. 494, *singulare* C. DC. 355, *Siriboa* Forst. 434, *Siriboa* L. 489, *Sirium* C. DC. 492, *smilacifolium* Kunth 17, *Solmsianum* C. DC. 210, *Sororium* C. DC. 225, *speciosum* Willd. 378, *sphærostachyum* C. DC. 357, *Sprengelianum* C. DC. 176, *Spruceanum* C. DC. 245, *spurium* Link 493, 502, *stellipilum* C. DC. 84, *stipulaceum* Opiz 577, *stuppeosum* A. Dietr. 116,

stylosum Miq. 428, subæquale C. DC. 546, subandinum C. DC. 141, subauriculatum C. DC. 347, subcordatum C. DC. 465, subflavum C. DC. 154, subfuscum C. DC. 273, subpeltatum Willd. 384, subscutatum C. DC. 337, subseptemnervium C. DC. 603, subulatum C. DC. 429, Suipigua Don 612, sulcatum Bl. 447, Sumatranum C. DC. 419, Sundaicum Bl. 461, Surinamense C. DC. 223, Swartzianum C. DC. 100, sylvaticum Roxb. 475, sylvestre Lam. 493, *sylocstre* Vell. 232, syringæfolium Vahl 610, Swartzianum C. DC. 100.

P. Tarapotianum C. DC. 21, Teapense C. DC. 75, tectoniæfolium Kunth 260, Tejucense C. DC. 190, tenue Kunth 14, tenuisuspe C. DC. 271, tenuifolium C. DC. 338, tenuinerve C. DC. 309, tenuistylum C. DC. 76, Tequendanense C. DC. 248, *terminale* Kunth 34, Teysmanii C. DC. 432, thermale Vahl 607, Twaitesi C. DC. 481, Tigerianum C. DC. 35, *tiliæfolium* Ham. 621 (1), *tiliæfolium* Cham. et Schl. 377, Tocacheanum C. DC. 97, tomentosum C. DC. 554, Trianæ C. DC. 386, riandrum Ferd. Mull. 513, trichoneuron C. DC. 165, trichostachyon C. DC. 4, tricuspe C. DC. 93, trigonioideum C. DC. 326, trigonum C. DC. 298, trinerve C. DC. 214, trineuron Miq. 499, *trioicum* Roxb. 502, triquetrum Opiz 578, tristachyon C. DC. 390, tristemon C. DC. 367, *truncatum* Vell. 200, tuberculatum Jacq. 103, tumidum Kunth 342.

P. Uhdei C. DC. 29, umbellatum L. 383, *umbellatum* Jacq. 384 var.  $\beta$ , *umbellatum* Vahl 381, umbrosum Kunth 540, undeninervium C. DC. 226, Ungaramense C. DC. 9, unguiculatum R. et P. 34.

P. Variegatum Kunth 318, variifolium C. DC. 288, velutinum Kunth 319, Venezuelense C. DC. 262, venosum C. DC. 408, venustum Wall. 604, Veraguense C. DC. 224, verbascifolium C. DC. 69, *verrucosum* Sw. 105, *verrucosum* Willd. 105, 312, verruculosum C. DC. 254, vestitum C. DC. 411, Victorianum C. DC. 22, villosum C. DC. 324, *villosum* Bl. mss. 414, viminale Opiz 600.

P. Wagneri C. DC. 259, Walkeri Miq. 501, Waracaboura C. DC. 64, Weddellii C. DC. 358, Wightii Miq. 512, *Willdenowii* Kunth 329, Wulfschlaegelii C. DC. 31, Wyderianum C. DC. 162.

P. xylostoeides Steud. 219.

P. Zacuapanum C. DC. 377, Zeylanicum Miq. 516, Zippelia C. DC. 59, Zollingeri C. DC. 507, Zollingerianum C. DC. 459, Zuccarinii C. DC. 508.

VII. CHAVICA Miq. syst. p. 222, *pro parte*. — Piper Wall. list, *pro parte*.

Bractea libera. Ovarium unicum. Stamina 2-4. Anthera matura bivalvata, extrorsum dehiscens. — Frutices asiatici vel sundaici.

1. C. FRUSTRATA (Miq. l. in Zoll. cat. p. 84), foliis breviter petiolatis oblongis apice acuminatis, acumine acuto et mucronulato, basi subinæquali cordatis supra glabris subtus stellatim molliter pubescentibus coriaceis opacis 8-septulinerviis, nervis subtus prominentibus, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervum utrinque unum fere ad apicem ductum alterne mittente, lateralibus nervis utrinque 2 e basi alterne ortis fere ad medium ductis, petiolo stellatim pubescente, stirpis masculæ pedunculo pubescente petiolum superante, amento densifloro filiformi quam folium brevior, bractea obovata basi subacuta centro petolata subsessili, staminibus 2, anthera oblonga, filamentis brevissimo rhachi immerso.  $\gamma$  In ins. Java (Zoll. n. 2624! in h. DC.) — Miq. pl. Jungh. fasc. 1. p. 16. Ch. Blumei  $\beta$  frustrata Miq. flor. ind. bat. v. 1. pars 2. p. 442. Dioica. Ramuli stellatim pubescentes. Limbi 0,12 longi, 0,04 lati. Petioli 0,005 longi. Amenta florentia 0,05 longa, 0,002 crassa. (v. s.)
2. C. SPHÆROSTACHYA (Miq. l. syst. p. 278), foliis breviter petiolatis superioribus elliptico-lanceolatis apice acuminatis, acumine obtusiusculo, basi cuneatis inferioribus subrotundatis apice acuminatis, acumine acuto, basi cordatis omnibus utrinque glabris coriaceo-membranaceis coriaceisve opacis non

(1) Errore sub num 621 admissum, quoniam sit Laureia tiliæfolia Gaudich. supra pag. 235<sup>14</sup>.

pellucido-punctatis, quintupli-septuplinerviis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervos alternos adscendentes utrinque 2-3 quorum inferiores fere e basi orti marginales mittente, petiolo glabro pedunculo superante, stirpis masc. amento filiformi densifloro, bractea rotundata centro affixa subsessili glabra, rhachi hirtula, staminibus 2, anthera filamentum superante semilunari, stirpis femin. amento globoso, bractea ut in mare, stigmatibus 3 brevibus, bacca obovato-subtetragona glabra basi immersa. ♀ In Moluccis (Miq. in flor. ind. bat. v. 1. pars 2. p. 446 (exclus. synon. Rumph.), Java (Miq. l. c. Zollinger n. 1322! in h. DC. et Schmidt! in h. Berol.), Pulo Pinang (Miq. l. c.), Courtallum (Wight fide Miq. in Hooker Lond. journ. bot. v. 5. p. 557), Bengalia orientali (Griffith n. 4409! in h. Kew. et DC.), Napalia (Wall. n. 6656! in h. DC. et h. Wall. et? 6654 β et C. in h. Wall. et n. 6660! in h. Wall. sub nomine *P. vasculosum* Ham.), Sikkim (Hook. et Thoms. in h. Kew.), monte Khasia alt. 4-6000 ped. (Hook. et Thoms. in h. DC.), montibus nilghericis (Perrottet in h. DC. Hohenacker n. 1325! in h. Boiss.). — Miq. l. illustr. t. 42 et 43 et Wight's icon t. 1931. Piper brachystachyon Wall. list n. 6656, majore parte. Piper bœhmiaefolium Wall. list n. 6654 β. Chavica Neesiana Miq. l. syst. p. 249. Dioica. Ramuli glabri. Limbi 0,08-0,09 longi, 0,025-0,03 lati. Petioli circ. 0,01 longi. Amenta masc. florentia 0,02 longa, vix 0,001 crassa, feminea matura 0,003-0,008 crassa. (v. s.)

3. C. PETIOLATA, foliis longe petiolatis ovato-acuminatis vel ovato-oblongis apice acuminatis, acumine acuto, basi aequali rotundatis utrinque glabris membranaceis 5-vel quintuplinerviis, centrali nervo ad  $\frac{1}{2}$  longitud. trifido, petiolo glabro, pedunculo glabro, stirpis masc. amento florente quam pedunculus multum brevior, bractea rotundata centro peltata subsessili glabra, staminibus 2, anthera subglobosa, filamentum antheram æquante, stirpis femin. bractea ut in mare, bacca globosa amento brevissimo. ♀ In Bengalia orientali (Griffith n. 4405! et 4410! in h. Kew. stirpis femin.), monte Khasia (Hook. et Thoms. in h. Kew. stirpis masc.). Dioica. Ramuli glabri. Limbi 0,12 longi, 0,07 lati. Petioli 0,02-0,06 longi. (v. s.)
4. C. THOMSONII, foliis petiolatis elliptico-oblongis apice acuminatis, acumine acuto, basi leviter inæquali obtusis utrinque glabris membranaceis subopacis pellucido-punctulatis septuplinerviis, nervis subtus prominulis centrali ad  $\frac{1}{2}$  longitud. usque nervum utrinque unum adscendentem opposite mittente, nervis lateralibus e basi ortis subtilioribus, petiolo glabro basi ima vaginante, pedunculo quam petiolus brevior, stirpis masc. amento filiformi densifloro florente quam folium triplo brevior, rhachi hirtella, bractea rotundata glabra centro peltata subsessili, staminibus 4, stirpis femin. amento brevissimo, baccis subglobosis discretis densis nigris. ♀ In Monte Khasia (Hook. et Thoms. n. 181 in h. Berol.). Dioica. Ramuli glabri. Limbi 0,13 longi, 0,043 lati. Petioli 0,01 longi. Amenta feminea 0,006 longa. (v. s.)
5. C. PEEPULOIDES (Miq. syst. p. 237), foliis breviter petiolatis elliptico-lanceolatis vel ovato-acuminatis, acumine acuto mucronulatoque, basi rotundato-obtusis summis oblongo-lanceolatis basi acutis, omnibus utrinque glabris firme membranaceis quintuplinerviis, nervis 3 centralibus ad apicem, usque ductis supra basin solutis, petiolo glabro, pedunculo petiolum subæquante, stirpis masc. amento quam folium multoties brevior filiformi, bractea rotundata centro peltata pedicellata, staminibus 4-3-2, stirpis femin. amento quam folium duplo brevior, bractea ut in mare, baccis subglobosis nigris discretis dense approximatis. ♀ In Bengalia orientali (Griffith n. 4411! in h. Kew.), Silhet (Wall. n. 6650 a! in h. Wall.), monte Khasia alt. 3-6000 ped. et Chittagong alt. 1000 ped. (Hook. et Thoms. in h. Kew.). Peperomia peepuloides A. Dietr. sp. v. 1. p. 145. Piper peepuloides Roxb. flor. ind. v. 1. p. 159. Dioica. Ramuli glabri. Limbi 0,08 longi, 0,04-0,06 lati. Petioli 0,05-0,01 longi. (v. s.)

### *Species exclusæ.*

C. AMBOINENSIS Miq. = Piper Amboinense C. DC.

C. ARNOTTIANA Miq. = Piper Arnottianum C. DC.,

- C. AURICULATA Miq. = Piper Betle L.  
 C. BENTHAMIANA Miq. = Piper fragile Benth.  
 C. BETLE Miq. = Piper Betle L.  
 C. BLUMEI Miq. = Piper Melamiri Bl.  
 C. BÖHMERIÆFOLIA Miq. = Piper bœhmeriæfolium Wall.  
 C. CELEBICA Miq. = Piper Celebicum Bl.  
 C. CELTIDIFORMIS Miq. = Piper celtidiforme Opiz.  
 C. CERAMICA Miq. = Piper Ceramicum Bl.  
 C. CHABA Miq. = Piper Chaba Bl.  
 C. CHAWYA Miq. = Piper Chawya Ham.  
 C. CONSANGUINEA Miq. = Piper Labillardieri C. DC.  
 C. CORYLISTACHYA Miq. = Piper corylistachyum C. DC.  
 C. DENSA Miq. = Piper Betle L. var. β.  
 C. EXASPERATA Miq. = Piper exasperatum Vahl.  
 C. HUGELIANA Miq. = Piper Hugelii C. DC. et Piper Hugelianum C. DC.  
 C. LABILLARDIERI Miq. = Piper officinarum C. DC.  
 C. LANCEOLATA Miq. = Piper bœhmeriæfolium β et Piper miniatum ~~et~~ (pro p.)  
 Piper Roxburghii.  
 C. LESSERTIANA Miq. = Piper grande Vahl.  
 C. MACROSTACHYA Miq. = Piper miniatum Bl.  
 C. MACULATA Miq. = Piper maculatum Bl.  
 C. MADAGASCARIENSIS Miq. = Piper Madagascariense.  
 C. MAJUSCULA Miq. = Piper majusculum Bl.  
 C. MELAMIRIS Miq. = Piper Sirium C. DC.  
 C. MICROSTIGMA Miq. = Piper microstigma C. DC.  
 C. MINIATA Miq. = Piper miniatum Bl.  
 C. MULLESUA Miq. = Piper Mullesua Don.  
 C. MULTINERVIA Miq. = Piper Prianamense.  
 C. OFFICINARUM Miq. = Piper officinarum C. DC.  
 C. PENANGENSIS Miq. = Piper Penangense C. DC. et Piper lonchites Miq.  
 C. POTHOIDES Miq. = Piper pothoides Wall.  
 C. PSEUDOMELAMIRIS Miq. = Piper Sirium C. DC.  
 C. PYRIFOLIA Miq. = Piper pyrifolium Vahl.  
 C. REINWARDTIANA Miq. = Piper Reinwardtianum C. DC.  
 C. RETROFRACTA Miq. = Piper retrofractum Vahl.  
 C. ROXBURGHII Miq. = Piper longum L.  
 C. RUMPHII Miq. = Piper Rumphii C. DC.  
 C. SARMENTOSA Miq. = Piper sarmentosum Roxb.  
 C. SINENSIS Champ. = Piper Sinense C. DC.  
 C. SIRIBOA Miq. = Piper Betle L.  
 C. SUBCORDATA Miq. = Piper subcordatum.  
 C. SUBSEPTEMNERVIA Miq. = Piper subseptemnervium C. DC.  
 C. SUPIGUA Miq. = Piper Suipigua Don, C. DC.  
 C. SUNDAICA Miq. = Piper Sundaicum Bl.

- C. SYLVATICA Miq. = *Piper sylvaticum* Roxb.  
 C. SYRINGÆFOLIA Miq. = *P. syringæfolium* Vahl.  
 C. TEYSMANNII Miq. = *Piper Teysmannii* C. DC.  
 C. THERMALIS Miq. = *Piper thermale* Vahl.  
 C. VIMINALIS Miq. = *Piper viminale* Poir.  
 C. WALLICHII Miq. = *Piper aurantiacum* Wall.

### Tribus III. PEPEROMIÆ Miq. syst. Pip. (mutatis char.)

Systema fibro-vasculare simplex, sparsum Ovarium unicum uniloculare. — Herbæ vel frutices.

VIII. VERHUELLIA Miq. syst. p. 47, pro parte, ill. p. 5. t. 1. fig. a, Cas. DC. in Mem. soc. phys. et hist. nat. genev. 18. pars 2. t. 1. f. 58. — *Piperis* sp. Sw. flor. ind. occ. v. 1, Vahl. enum. v. 1, Spreng. syst. v. 1, quoad speciem unicum. — *Peperomia* sp. A. Dietr. sp. v. 1, quoad speciem unicum, Schott in Bot. Zeit. 1851. — *Mildea* Griseb. Cat. pl. Cub. — *Piperis* subgen. *Piperoides* Cas. DC. in Seem. journ. 1866, p. 161.

Bractea libera. Stamina 2; anthera matura quadrivalvata. Stigmata 4. — Herbula ex America et Africa.

1. V. ELEGANS (Miq. syst. p. 48), foliis 3-5-6-verticillatis, in eodem verticillo inæqualibus, longe petiolatis reniformi-cordulatis subtus glabris junioribus supra remote pilosulis tenuissime membranaceis subtiliter pellucido-punctulatis 3-5-nerviis, nervis subtilibus, centrali ad apicem ducto supra basin trifido, lateralibus ad medium ductis fere e basi bifidis, foliis ideo fere quintuplinerviis nervulosis, petiolo glabro membranaceo, amentis axillaribus longe pedunculatis, pedunculo petiolum evolutum æquante, amentis tenuiter filiformibus laxifloris folia triplo superantibus modo unisexualibus modo bisexualibus basi femineis apice masculinis, floribus raro hermaphroditis, bractea suboblongo-elliptica infra medium peltata breviter pedicellata, staminibus 2, filamentis brevissimis antheras æquantibus, ovario basi in pedicellum brevem in floribus hermaphroditis stamina gerentem contracto ovato-subgloboso hirsuto apice breviter styliifero, stigmatibus ovarium æquantibus filiformibus. ① In ins. Sto Domingo (Rich. in h. Francovil.), Cuba (Wright n. 2264! in h. DC.). — Miq. illustr. p. 5. t. 1. fig. a. *Piper* reniforme Willd. herb. n. 742! Spreng. syst. v. 1. p. 117. *Peperomia* reniformis A. Dietr. sp. v. 1. p. 162. V. serpens Griseb. flor. west Ind. p. 164. *Mildea elegans* Griseb. Cat. pl. Cub. p. 63. Herbula repens. Caulis glaber. Limbi 0,008 longi, 0,013 lati. Petioli 0,25 circ. longi. (v. s.)
2. V. CORDIFOLIA, foliis 3-5-verticillatis petiolatis valde inæqualibus orbiculato-cordatis supra remote pilosulis subtus glabris fusco-punctulatis tenuissime membranaceis pellucidis 3-5-nerviis, petiolo glabro, amentis axillaribus filiformibus unisexualibus vel basi femineis apice masculinis, femineis laxifloris, pedunculis petiolos æquantibus, staminibus 2, ovario basi in pedicellum brevem contracto ovato-globoso hirsuto apice breviter styliifero, stigmatibus 4. ② In ins. Cuba (Wright n. 2266! in h. DC.). *Mildea hydrocotylifolia* Griseb.! Cat. pl. Cub. p. 63. Herbula pellucida, inter muscos repens, humilis. Limbi circ. 0,003 longi latique. Petioli 0,005 longi (v. s.)
3. V. KNOBLECHERIANA, foliis alternis petiolatis rotundato-ovatis apice acutis basi breviter cordatis utrinque glabris supra punctulatis, centrali nervo ad  $\frac{3}{4}$  longitud. usque nervos oppositos subadscendentes utrinque 3 quorum inferiores longiores supra medium ducti mittente, amentis axillaribus solitariis quam folium brevioribus laxifloris, floribus hermaphroditis, bractea ovata sessili, staminibus 2 (ex figura 4-valvis), ovario ovoideo sessili apice attenuato apice imo stigma penicillatum gerente. ③ In Africa ad ripas Nili albi (Knoblecher

fide Schott in Bot. Zeit. 1851. p. 225. t. 5). *Peperomia Knoblechteriana* Schott l. c. *Herbula simplex*, tenerrima, glabra, 0,05 alta. Limbi 0,01 longi, 0,007 lati. Petioli 0,001 longi. (Ex Schott l. c.)

*Species dubia.*

- 4 *V. SERPENS* (Miq. syst. p. 49), foliis alternis petiolatis subrotundis sed magis latis basi quasi retusis, acumine brevissimo, obtusis planis subsucculentis eveniis utrinque glabris subtus pallidioribus, amentis axillaribus solitariis pedunculatis, pedunculis folia superantibus glabris, floribus hermaphroditis minutissimis, bracteis ovatis, filamentis vix ullis, staminibus 2 lateralibus, ovario ovato acuto, stylo nullo, stigmatibus 3, fructu minutissimo acuto sessili. ☉ In petrosis sylvaticis muscosis Jamaicae frigidioris. *Piper serpens* Swartz prodr. p. 16, flor. Ind. occ. v. 1. p. 69, Willd. sp. v. 1. p. 165, Vahl enum. v. 1. p. 347, Spreng. syst. v. 1. p. 118. *Acrocarpidium serpens* Miq. in diar. inst. Nederl. 1843. *Herba*. Caulis repens filiformis glaber (Ex Swartz ll. cc.)

*Species exclusa.*

*V. BRASILIENSIS* Miq. = *Peperomia Verhuellia* C. DC.

IX. *PEPEROMIA* Ruiz et Pav. prodr. flor. Per. p. 8, flor. Per. v. 1. p. 29, Kunth in H. et B. nov. gen. v. 1. p. 60, Opiz reliq. Hænk. fasc. 3, A. Dietr. sp. v. 1, Meissn. gen. p. 335, Miq. syst. Pip., Lindl. t. 352. — *Piper* L. gen. n. 832, sp. p. 40, pro parte, Jacq. icon. rar. et coll., Jacq. f. eclog. pro parte Lam. ill. pro parte, Willd. sp. pro parte, Vahl enum. pro parte, Ræm. et Sch. syst. pro parte, Sprengel syst. pro parte, Bl. monogr. der Oost. Ind. peps. in Verh. der Bat. Genoots. v. 11 et enum. pl. Jav., pro parte, Loud. encycl., pro parte, Endlich. gen. n. 1820, pro parte. — *Micropiper* Miq. Comm. phyt. p. 37. — *Acrocarpidium*, *Tildenia* et *Erasmia* Miq. in Diar. Inst. Nederl. 1842. — *Phyllobryon* Miq. syst. pip. — *Dugagelia* ? Gaudich. voy. Uran. p. 5137.

Bractea libera. Flos hermaphroditus. Stamina 2, lateralia. Antheræ non articulatae, bivalvatae, extrorsum dehiscentes. Stigma unicum. — Herbæ vel frutices totius orbis calidioris.

CONSPECTUS GENERIS.

§ 1. *Ovarium stylosum.*

a. Bractea peltata .....	Confer. ad spec.	1
1° Folia peltata .....		1
2° Folia non peltata .....		8
b. Bractea non peltata .....		16

§ 2. *Stylus nullus. Bractea peltata.*

A. Folia alterna.

† Ovarium apice summo stigmatiferum .....	17
α. Bacca stipitata .....	17
β. Bacca sessilis .....	22
* Stigma conicum .....	22
** Stigma discoideo-bilobulatum .....	23
*** Stigma simplex, penicillatum vel punctiforme .....	25
1° Folia penninervia .....	25
2° — multiplinervia .....	26
3° — multinervia .....	30



†† Ovarium apice oblique stigmatiferum..... Confer. ad spec.	64
1° Folia penninervia .....	64
2° — multiplinervia .....	79
3° — multinervia .....	89
†† Ovarium antice stigmatiferum, vertice oblique complanatum vel scutello auctum, sursum rostratum.....	162
1° Folia penninervia .....	162
2° — multiplinervia.....	189
3° — multinervia .....	197
Species affines, non satis notæ.....	221
B. Folia opposita.	
1° Folia penninervia .....	243
2° — multiplinervia.....	246
3° — multinervia .....	249
Species affines, non satis notæ .....	281
C. Folia verticillata.	
† Ovarium apice summo stigmatiferum .....	287
†† Ovarium apice oblique stigmatiferum .....	331
Species affines, non satis notæ.....	371
Peperomiæ non satis notæ et incertæ sedis .....	384

### § 1. *Ovarium styliiferum.*

#### a. Bractea peltata.

##### 1° Folia peltata.

1. *P. MACRORHIZA* (Kunth in H. et B. nov. gen. v. 1. p. 72), foliis longissime petiolatis orbicularibus centro peltatis utrinque glabris rigidis eveniis, petiolo glabro limbum multoties superante, amento radicali longe pedunculo, pedunculo glabro petiolum subæquante amentum valde superante, amento densifloro, bractea subsessili. ④ In devexis Andium peruvianorum locis scopulosus inter urbem Caxamarcae et Cerro de Cunturcagua alt. 900 hexap. (Humb. l. c., Rich. ! in h. Francovil.). — Kunth synopsis v. 1. p. 123. *Piper macrorrhizon* Rœm. et Sch. syst. v. 1. p. 336. Herba acaulis, pedunculi et petioli densissime e radice cæspitose orti. Limbi 0,006 longi latique. Petioli 0,05 longi. (v. s.)
2. *P. PARVIFOLIA* (Cas. DC. in Seem. journ. 1866. p. 133), foliis superioribus parvis ovato-orbicularibus longissime petiolatis sub medio peltatis subcucullatis glabris coriaceis enerviis, inferioribus scariosis incompletis dense approximatis, amentis longe pedunculatis densifloris, bractea rotundato-peltata, ovario emerso apice substyloso carnosulo, stigmate capitellato. ④ In Peruvia (Pav. ! in h. Boiss.), Bolivia (Pentland n. 12850 ! in h. Kew.), Bolivia prov. Larecaja viciniis Sorata Cabezeras de Chilcani Lacatia in graminosis regionis alpinæ alt. 3600-4200 m. (Mandon n. 1123 ! in h. DC.). Planta cæspitosa, caule subnullo. Radix fibrosa. Limbi 0,002 longi. Petioli 0,015 longi. (v. s.)
3. *P. UMBILICATA* (Ruiz et Pav. ! flor. per. v. 1. p. 30. t. 45. fig. b), foliis longissime petiolatis orbicularibus vel suborbicularibus paullo sub medio peltatis glabris epunctatis pellucido-membranaceis a petiolo inconspicue 7-nerviis, nervulo obscuro juxta marginem currente, petiolo glabro, amentis longe pedunculatis subulxifloris, bractea ovato-acuminata, ovario emerso apice stylium carnosulum gerente, stigmate capitellato. ④ In Peruvia (Pav. ! in h. Boiss., Gaudich. n. 150 ! in h. DC.), circa lacum Tititacam (Miq. in Nov. act. nat. cur. n. 19 suppl. p. 483), Bolivia prov. Larecaja viciniis Sorata colli Ullontigi supra rupes humo refertos regionis temperatæ alt. 2800 m. (Mandon n. 1122 ! in h. DC.), locis excelsis scopulosis frigidis regni mexicani prope Santa

Rosa de la Sierra alt. 1300 hexap. (Humb. et Bonpl. nov. gen. p. 72. t. 15. fig. 1, Humb. in h. Berol. et h. Willd. n. 764!), inter saxa in montibus calcareis aridis et cactiferis ad Tehuacan alt. 7000 ped. (Galeotti n. 6023 fide Mart. et Gal. bull. ac. Brux. v. 10. p. 19), locis petrosis Sierræ Yavezia ad Oaxaca alt. 7000 ped. (Galeotti n. 6021 l. c.). Piper umbilicatum Vahl enum. v. 1. p. 356, Martens et Gal. l. c. P. g'obosum Poir. P. hydrocotylifolium Martens et Gal. l. c. Tildenia peruviana Miq. in Nov. act. nat. cur. l. c. Pep. umbilicata Kunth in H. et B. nov. gen. l. c., synops. v. 1. p. 123, Miq. syst. p. 69, in Linn. v. 20. p. 118. Herba acaulis. Radix tuberosa, tuber 0,015 circ. crassum. Limbi 0,01-0,015 longi. Petioli 0,05 circ. longi. Pedunculi 0,05 longi. (v. s.)

α, *macrophylla*, limbis majoribus apice brevissime subacutatis 0,03-0,035-0,08 in diam. Petioli et pedunculi 0,16 circ. longi. ① In Nova Hispania (Pav. in h. Boiss.). Herba acaulis. Radix tuberosa. Tuber 0,01 crassum. (v. s.)

β, *subacutifolia*, foliis subovato-orbicularibus apice breviter obtuse subattenuatis. ① In Mexico (Uhhe n. 254! in h. Berol.). Limbi sæpe 0,03 longi, 0,025 lati. (v. s.)

4. P. SCUTELLÆFOLIA (Ruiz et Pav. flor. per. v. 1. p. 29. t. 44. fig. b.), foliis longissime petiolatis ovatis apice obtusis basi rotundatis concavis ad  $\frac{1}{4}$  longitud. peltatis utrinque glabris supra punctatis 5-7-nerviis, petiolo glabro, amentis radicalibus, pedunculis glabris quam petioli paullo brevioribus folia duplo superantibus subdensifloris, bractea ovata centro peltata concava. ① In Peruvixæ collibus ad Atiquipa. Piper scutellæfolium Vahl enum. v. 1. p. 356. P. scutellatum Pers. syn. p. 34. Pep. scutellæfolia Miq. syst. p. 71. Herba acaulis. Radix tuberosa. Limbi 0,03 longi, 0,02 lati. Petioli 0,06 longi. (Ex R et P. l. c.)

5. P. OVATO-PELTATA (Cas. DC. in Seem. journ. 1866, p. 132), foliis alternis longe petiolatis ad  $\frac{1}{2}$  longitud. peltatis ovato-acutis apice acuminatis basi rotundatis utrinque glabris pellucido-membranaceis pellucido-punctatis 7-9-nerviis, nervulo juxta marginem currente, nervis subtus subprominulis, petiolo glabro, amentis longe pedunculatis subdensifloris, pedunculo glabro anguste alato, bractea ovato-longe-acuminata centro peltata, ovario emerso, stylo carnosulo apice stigma capitellatum gerente. ① In Nova Hispania (Pav. in h. Boiss.); Costa-Rica (Hoffman n. 521! in h. Berol.) Herba subacaulis. Radix tuberosa. Limbi 0,055 longi. Petioli pedunculique circ. 0,1 longi. (v. s.)

6. P. SPRUCHII (Cas. DC. in Seem. journ. 1866, p. 133), foliis petiolatis ovatis vel ovato-rotundatis interdum basi subpeltatis subcordatisque apice obtusis membranaceo-pellucidis pellucido-punctatis 5-nerviis, centrali nervo ad apicem usque ducto, lateralibus ad marginem fere currentibus remote anastomosantibus, petiolo glabro, amentis longiuscule pedunculatis filiformibus subulxifloris, bractea rotundata centro peltata, ovario emerso, stylo apice stigma penicillatum gerente, bacca ovata apice breviter retellata. ① In Peruvia orientali prope Tarapoto (Spruce n. 4981! in h. Kew.). Herba acaulis. Limbi 0,025-0,03 longi, 0,025 lati. Petioli 0,01 longi. Pedunculi 0,035 longi. (v. s.)

7. P. MEXICANA (Miq. syst. p. 75), foliis longiuscule petiolatis basi ima peltatis ovatis vel ovato-oblongis apice acuminatis basi rotundatis membranaceis glabris trinerviis, centrali nervo ad apicem usque ducto, lateralibus per anastomoses fere ad apicem ductis, amentis folia superantibus longe pedunculatis subulxifloris, bractea ovato-acuminata infra medium peltata, ovario emerso, stylo ovarium æquante apice imo stigma capitellatum gerente. ① Prope Xalapa (Galeotti in h. Kew.), Cordillera Vera-Cruz (Galeotti n. 7111! in h. Deless.), ad saxa secundum rivulos in humidissimis nemoribus Mirador alt. 3000 ped. (Martens in bull. ac. Brux. v. 10. n. 2. p. 20), Miradores (Linden n. 127! in h. Boiss.), America æquinox. (Kickx! in h. Boiss.). — Miq. illust. p. 12. t. 6. Piper parvulum Mart. et Gal. l. c. Herba acaulis. (v. s.)

## 2° Folia non peltata.

8. *P. MIRADORESIANA*, foliis petiolatis lanceolatis utrinque acutis et glabris membranaceis subpellucidis subtiliter pellucido-punctulatis, centrali nervo subtus prominulo ad  $\frac{1}{2}$  longitud. usque nervos alternos subadscedentes utrinque 9 quorum supremi per anastomoses ad apicem usque ducti mittente, petiolo glabro, amentis apice caulis aphylli terminalibus vel axillaribus foliis abortivis fultis folia æquantibus, pedunculo quam petiolus brevior, amentis filiformibus subaxillifloris, bractea subrotundata glabra centro peltata brevissime pedicellata, ovario basi rhachi immersulo oblongo, stylo brevi apice imo stigma puberulum gerente, bacca cylindrica apice mucronata basi immersula. (4) In *Miradores* (Linden n. 126! in h. Boiss.). Herba caulescens. Caulis glaber. Limbi 0,15 longi, 0,035 lati. Petioli 0,02 longi. Bacca matura 0,0015 longa. (v. s.)
9. *P. STOLONIFERA* (Kunth! in H. et B. nov. gen. v. 1. p. 71), foliis longe petiolatis ovato-subrotundatis apice obtusis basi leviter cordatis utrinque hirsuto-pubescentibus subtus nigro-punctulatis membranaceis subobscuris 7-nerviis, petiolo hirsuto-pubescente. In valle arcta umbrosa subcalida juxta pagum *Pandi* et *Puente Natural* de *Icononzo* 480 hexap. in *Andibus novo-granatensibus* (Humb.! l. c. et h. Willd. n. 763! et in h. Berol.). — Kunth syn. v. 1. p. 123, Miq.! syst. p. 74. *Piper stoloniferum*. Roem. et Sch. syst. v. 1. p. 336. Radix fibrosa. Caulis nodoso-tuberosus, contractus, apice folia et amenta exserens. Limbi 0,044-0,06 longi, 0,03-0,04 lati. Petioli 0,045 longi. (v. s.)
10. *P. MUSCOPHILA* (Cas. DC. in Seem. journ. of bot. 1866. p. 133), foliis petiolatis superioribus brevioribus ovato-rotundatis apice acutis inferioribus rotundato-reniformibus omnibus supra puberulis subtus hirsuto-puberulis pellucido-membranaceis 5-7-nerviis, petiolo hirsuto-puberulo, amentis axillaribus subaxillifloris, bractea rotundata centro peltata, ovario emerso, stylo brevi apice imo stigma capitellatum gerente. (4) In *Nova Hispania* inter muscos (Pav.! in h. Boiss. et DC. ex h. Boiss.). Caulis brevis, superne hirsuto-puberulus, pellucidus, 0,02 circ. altus. Limbi inferiores petiolique 0,006 longi. (v. s.)
11. *P. GALEOTTIANA* (Hook. icon. plant. t. 327), foliis longe petiolatis ovato-acuminatis glaberrimis trinerviis, petiolo gracili, scapis petiolum æquantibus, amentis elongatis gracilibus radicalibus folia superantibus. In *Xalapá Mexici* (Galeotti in h. Hook.). *Pep. mexicana* Miq.? (ex Hook. l. c.)
12. *P. MANDONII*, foliis ternis petiolatis ovatis vel subrotundo-ovatis apice obtusis basi rotundatis vel subtruncato-rotundatis utrinque glabris margine remote ciliolatis tenuissime membranaceis pellucidis 5-nerviis nervuloque apice juxta marginem currente, petiolo glabro, amentis terminalibus solitariis vel ramulos axillares ternatim terminalibus lateralibus foliis minoribus fultis, bractea ovato-subrotunda sub medio peltata brevissime pedicellata, ovario apice in stylum subalato, stigmatibus apice styli inserto, bacca oblonga apice subalata basi in stipitem brevissimum contracta. (4) In *Bolivæ* prov. *Larecaja* vicinibus *Sorata* inter *Ylabaya* et *Paracallo* ad rivulos in umbrosis alt. 3200 m. (Mandon n. 1124! in h. DC.). *Herbula* pellucida glabra. Limbi 0,012 longi, 0,01 lati. Petioli 0,003 longi. (v. s.)
13. *P. SCHOMBURGKII*, foliis quaternis longiuscule petiolatis suborbicularibus apice emarginulatis utrinque glabris ciliolatis coriaceis opacis fusco-subflaviscantibus obscure trinerviis, petiolo glabro subtiliter hirtello, amentis terminalibus longe pedunculatis maturis folia triplo superantibus densifloris, pedunculo glabro, bractea rotundata centro peltata breviter pedicellata, rhachi glabra foveolata, ovario ovato in stylum brevem apice stigmatiferum acutato. (4) In *Guyana anglica* (R. Schomburgk n. 406! in h. Berol.). Repens, ramosa. Ramuli coriacei, glabri. Limbi 0,012 longi, 0,01 lati. Petioli 0,006 longi. Pedunculi 0,015 longi. (v. s.)
14. *P. DIFFUSA*, foliis plerumque quinis breviter petiolatis oblongis apice obtusis basin versus attenuatis acutiusculis utrinque glabris rigidulis pellucidis pellucido-punctulatis fere eveniis subtiliter trinerviis, centrali nervo ad apicem usque ducto, lateralibus supra medium ductis, nervulo sæpe deficiente vel

interrupto juxta marginem currente, petiolo glabro, amentis terminalibus filiformibus subaxillaribus folia multoties superantibus, pedunculo glabro petiolum fere duplo superante, bractea rotundata centro peltata subsessili glandulosa, ovario immenso ovato stylo longiuscule protracto acuminato apice imo stigma carnosulum gerente, bacca oblonga basi immersa apice mucronata. ♀ In Venezuela prope coloniam Tovar alt. 3000-4000 (Fendler n. 11781 in h. DC.). Repens ramulosa glabra. Ramuli sicci subtetragoni, anguste subulati, fere 0,003 crassi. Limbi 0,015 longi, 0,005 lati. Petioli 0,002 longi. Amenta matura 0,002 crassa. (v. s.)

15. P. *HOFFMANNII* (C. DC. in Seem. journ. 1866. p. 133), foliis quaternis brevissime petiolatis obovatis apice emarginatis basi cuneatis utrinque glabris rigulidis subpellucidis pellucido-punctulatis uninerviis, nervo sæpe inconspicuo utrinque nervulos 1-2 patulo-subadscendentes mittente, nervulo obscuro apice juxta marginem currente, amentis terminalibus solitariis longiuscule pedunculatis quam pedunculi brevioribus densifloris, bractea rotundata, centro peltata subsessili, bacca ovato-oblonga, stylo apice imo stigmatifero. ☉ In Costa-Rica (Hoffmann n. 415! in h. Berol.). Herba repens, filiformis, glabra, ramulos erectos 0,04 altos mittens. Limbi 0,005 longi, 0,004 lati. (v. s.)

#### b Bractea non peltata.

16. P. *FOLIIFLORA* (Ruiz et Pav.! flor. peruv. v. 1. p. 30. t. 45. fig. c.), foliis caulinis sessilibus auriculæformibus tenuissime pellucidis ramulinis sessilibus e basi cordata ovato-acutis utrinque glabris tenuiter membranaceis pellucidis 9-nerviis, centrali nervo ad apicem ducto, lateralibus 2 fere ad apicem ductis cæteris divaricantibus exterioribus subtilioribus nervulo tenui apice juxta marginem currente, amentis 4-6 in axilla folii umbellatis foliis abortivis squamosis ovato-acutis basi cordatis? fultis, floribus bifariam dispositis, bractea sessili oblonga apice obtusa tenuissima pellucida, ovario oblongo-ovato glabro per anthesin pedicellato, stylo apice imo stigma discoideopapillosum gerente. ④ In Andium peruvianorum nemoribus ad arborum truncos (Ruiz in h. Willd. n. 761!). Piper foliiflorum Vahl enum. v. 1. p. 356. Phyllanthus herb. Willd.! l. c. Peperomia phyllantha Opiz in Reliq. Haenk. fasc. 3. p. 162. Phyllobryon Pavoni Miq.! syst. p. 50, illust. p. 7. t. 2. in Hook. Lond. journ. bot. v. 4. p. 410. Herba circ. 0,1 alta, glabra, basi radicans. Folia caulina apice caulis dense imbricata. Ramulus quisque apice folium sessile, 0,025 longum, 0,021 latum, in axilla umbellam foveentem gerente. Ramuli 0,04 longi. (v. s.)

### § 2. *Stylus nullus, bractea peltata breviter vel brevissime pedicellata.*

#### A. — *Folia alterna.*

#### † Ovarium apice summo stigmatiferum.

##### α. Bacca stipitata.

17. P. *TENELLA* (A. Dietr. sp. v. 1. p. 153), foliis brevissime petiolatis ovatis apice attenuato obtusis, apice imo sæpe emarginulato, utrinque sparsim pilosiusculis ciliolatis demum glabratis rigidulo-membranaceis membranaceisve subpellucidis eventis uninerviis nervo ad apicem usque ducto, nervuloque juxta marginem apice currente, petiolo pilosiusculo, amentis terminalibus folia superantibus longiuscule pedunculatis, pedunculo glabro, bractea rotundata centro peltata, ovario oblongo emerso, stigmate capitellato. ☉ In ins. Guadaloupa ubi interdum parasiticum in arbusculis prati montis sulphurigeri (Rich.! in h. Francovii), Cuba (Wright n. 523! in h. DC.), Jamaica (Griseb. flor. w. Ind. p. 165). — Griseb. Cat. pl. Cub. p. 63. Piper tenellum Sw. prodr. p. 16, flor. ind. occ. v. 1. p. 65, Vahl enum. v. 1. p. 345, Willd. sp. v. 1. p. 163. Acrocarpidium tenellum Miq. syst. p. 33. Herba. Caulis 3-4-uncialis

simplex, pilosiusculus, basi decumbens. Limbi 0,015 longi, 0,006 lati. Petioli vix 0,002 longi. (v. s.)

β. *glabra*, foliis utrinque glabris apice imo ciliolatis coriaceis obscuris, ovario rhachi impresso, bacca apice mucronulata. — In Venezuela inter Maracai et Choroni (Fendler n. 1818! in h. DC.). Limbi 0,008 longi. Petioli 0,001-0,002 longi. Amenta 0,03-0,04 longa. Pedunculi vix 0,005 longi. (v. s.)

18. *P. HISPIDULA* (A. Dietr. sp. v. 1. p. 165), foliis petiolatis triangulari-vel reniformi-rotundatis vel semi-orbicularibus utrinque setis albis articulatis instructis pellucido-membranaceis 5-nerviis parce et pellucide reticulato-nervulosis, petiolo setuloso, amentis axillaribus terminalibusque filiformibus laxifloris longe pedunculatis, bractea rotundata centro peltata, ovario emerso apice contracto, stigmatibus carnosulo, bacca ovata apice mucronulata. ☉ In Brasilia (Sellow! in h. Berol. et h. Willd. n. 731! specim. duo infer. et Riedel! in h. Petropol.), Petropoli ad arbores (Wawra et Maly n. 4690! fide Wawra Ergeb. bot. p. 131), Venezuela prope Tovar (Fendler n. 1248! in h. DC.), Caracas (Gallmer n. 2331 in h. Berol.), Colombia (Moritz n. 866! et 1630! in h. Kew.), Costa-Rica prope Vulcan de Barba (Hoffmann n. 54! in h. Berol.), Jamaica summitatibus montis albi in sylvis humidis (Griseb. flor. w. Ind. p. 165). — Griseb. flor. w. Ind. p. 65. *Piper hispidulum* Sw. prodr. p. 15, icon. t. 4, flor. Ind. occ. v. 1. p. 63, Vahl enum. v. 1. p. 347, Willd. v. 1. p. 163, Spreng. syst. v. 1. p. 117. *Acrocarpidium Sellowianum* Miq.! syst. p. 55, ill. p. 9. t. 3. fig. a, in Hook. Lond. Journ. bot. v. 4. p. 411. *Acrocarpidium* Miq. syst. p. 54, Schnizl. ic. t. 81. fig. 6. *Pep. tenera* Miq.! in Mart. flor. Bras. fasc. 11. p. 19. t. 1. fig. 1. Radix fasciculato-fibrosa. Caulis plures erecti, ramosi, glabri, 0,18-0,2 longi. Limbi 0,006-0,015 longi, 0,007-0,012 lati. Petioli 0,01 longi. (v. s.)

19. *P. FERNANDOPOANA*, foliis petiolatis lanceolato-acuminatis versus basin rotundatis basi ima acutis membranaceis opacis glabris 5-nerviis, centrali nervo ad apicem usque ducto, lateralibus nervis tenuioribus alte ductis exterioribus subtilioribus, petiolo glabro amentis axillaribus terminalibusque solitariis filiformibus pedunculatis, pedunculo glabro, bractea rotundata centro peltata, bacca ovata, stipite immerso. ☉ In Fernando Po (Manu n. 394! in h. Kew.). Herba procumbens. Caulis radicans, glaber. Limbi 0,05 longi, 0,025 lati. Petioli 0,01 longi. (v. s.)

β. *subopacifolia*, foliis subopacis obsolete pellucido-punctulatis apice acutiusculis. — In sylvis editis densioribus insulæ S.-Thomas alt. 1800 ped. crescit gregaria (Welwitsch n. 507! in h. Kew.), rupestribus umbrosissimis ad Matu de Pungo præsidii Pungo Andongo alt. 2400-3500 ped. (Welwitsch n. 505! in h. DC.)

20. *P. BREVIPES*, foliis brevissime petiolatis orbiculatis carnosius punctatis junioribus hirsutis adultis glabris ciliatis uninerviis, amentis terminalibus vel oppositifoliis (ramis abortientibus) gracilibus confertifloris, pedunculo hirsuto, bacca ovoidea, stipite brevissimo, bractea rotundata centro peltata. — In Conchagua Americae occid. tropic. (Benth. bot. of Sulph. p. 166). *Acrocarpidium brevipes* Benth. l. c. Petiolus semilineam raro excedens.

21. *P. EXILIS* (Griseb. flor. w. Ind. p. 164), foliis petiolatis orbicularibus minutis membranaceis pellucidis utrinque glabris margine remote fimbriulatis trinerviis vel enerviis, amentis terminalibus filiformibus folia multoties superantibus, bractea rotundata centro peltata subsessili glandulosa, stigmatibus penicillato, bacca ovoidea breviter stipitata. ☉ In India occident. (Poiteau? in h. Deless.), Trinitate, Portorico, Jamaica (Griseb. l. c.). *Acrocarpidium exile* Miq.! syst. p. 56. Herba repens. Caulis filiformis, glaber, tenuissimus. Limbi 0,003 in diam. Petioli 0,002 longi. (v. s.)

β. Bacca sessilis.

\* Stigma conicum.

22. *P. ERASMA*, foliis petiolatis lanceolatis vix puberulis membranaceis penninerviis, centrali nervo ad apicem ducto nervos adscendentes utrinque 10

mittente, nervulis reticulatis, amentis geminatis pedunculo communi longo sustentis (h. kew!) vel pedunculo communi paniculatum 9-confertis (fide Miq. syst. p. 200), bractea rotundata peltata apice subattenuata, ovario longo exserto cylindrico apice acuto, stigmatē conico quam ovarium ter quaterque breviorē. In Mexico prope Xalapa (Galeotti! in h. Kew.). *Erasmia floribunda* Miq. in diar. instit. reg. Nederl. a. 1842, syst. l. c., ill. p. 26. t. 24, in Hook. Lond. journ. bot. v. 4. p. 430. Herba? Limbi 0,13 (spec. kew.), 0,22 (fide Miq.) longi. Petioli 0,02 longi. (v. s.)

\*\* Stigma discoideo-bilobulatum.

23. *P. VERHUELLIA*, foliis longe petiolatis paullo supra basin peltatis ovatis vel rotundato-ovatis basi rotundatis cordulatisve supra pilosulis subtus glabris tenuiter membraceis pellucidis reticulato-nervulosis 5-7-nerviis, nervis centralibus 3 fere ad apicem usque ductis, nervulo juxta marginem ad apicem currente, petiolo glabro pellucido, amentis axillaribus terminalibusve solitariis filiformibus folia superantibus laxifloris longe pedunculatis, pedunculis petiolo circ. æquantibus glabris pellucidis, bractea rotundata centro peltata breviter pedicellata, ovario glabro rhachi impresso ovato, stigmate bilobulato discoideo, bacca ovato-globosa apice mucronulata. ☉ In Brasilia (Sello! in h. Berol.) *Verhuellia Brasiliensis* Miq. l. syst. p. 48, ill. p. 6. t. 1 fig. b, in Mart. flor. Bras. fasc. 11. p. 10. t. 1. fig. 6. *Herbula repens*. Caulis filiformis, glaber. Limbi 0,01-0,02 longi, 0,01-0,015 lati. Petioli 0,04-0,05 longi. (v. s.)

24. *P. FRASERI* (C. DC. in Seem. Journ. bot. 1866. p. 134), foliis sparsis longiuscule petiolatis cordato-lanceolatis glabris pellucido-punctatis 7-9-nerviis subreticulato-nervulosis, petiolo glabro membranaceo, amentis apice caulis elongati aphylli paniculato-confertis densifloris post anthesim pedunculatis, bractea rotundata centro peltata, ovario granuloso emerso, stigmate orbiculari bicrenato granuloso-puberulo. ☉ In Republica Æquatoriali (Fraser! in h. DC. ex Spruce n. 5552! in h. DC.), Nova Hispania (Pav. in h. Boiss.). Herba erecta simplex. Limbi 0,05 longi. Petioli 0,045 longi. Folia basi caulis 4-5-approximata, subverticillata, apice caulis opposita vel subopposita vel alterna. (v. s.)

\*\*\* Stigma simplex penicillatum vel punctiforme.

1° Folia penninervia.

25. *P. PANICULATA* (Regel! in Bull. Acad. Mosc. 1858. p. 542. t. 3), foliis petiolatis subobovato-ellipticis apice acuminatis, acumine acuto, basi in petiolum attenuatis vel lanceolato-oblongis utrinque glabris membranaceis subopacis, centrali nervo ad apicem ducto validiore subtus prominulo fere tota longitudine nervos subascendentes subtiliores utrinque 4-5 suboppositos mittente, petiolo glabro, amentis apice caulis spicatum confertis quam folia multoties brevioribus densifloris, bractea orbiculari centro peltata glandulosa, ovario oblongo, bacca oblongo-cylindrica basi immersula. ☉ In saxosis prope Castel Novo in Brasilia (Riedel! in h. Petrop.). Herba erecta, cum panicula 0,17 alta. Limbi 0,135-0,17 longi, 0,05-0,045 lati. Petioli 0,02 longi. Amenta matura 0,01 longa. Pedunculi 0,003 longi. (v. s.)

2° Folia multiplinervia.

26. *P. ASARIFOLIA* (A. Dietr. sp. v. 1. p. 106), foliis longe petiolatis subovato-rotundis apice obtusis vel acutiusculis basi æquali rotundo-cordatis utrinque glabris membranaceis subpellucidis subtilissime pellucido-punctulatis duodenonerviis reticulato-nervulosis nervuloque juxta marginem currente, nervis paullo supra basin solutis, petiolo glabro, amentis axillaribus terminalibusque folia multoties superantibus, pedunculo glabro quam petiolo breviorē, bractea subrotunda centro peltata breviter pedicellata fusco-glandulosa ovario subimmerso glanduloso, stigmate penicillato. ☉ In Mexico ad rupes humidās (Schiedel! in h. Berol., Galeotti n. 6022! in h. Deless.), locis humidis ad dunas Oceani Pacifici inter Tchuantepec et Acapulco (Mart. et Gal. in bull. Acad. Brux. v. 10. p. 18). In horto Berol. culta. — Miq. l. syst. p. 73, in Linn. v. 5.

- p. 75. Herba glabra. Caulis brevis. Limbi 0,04 longi. Petioli 0,045 longi. Pedunculi 0,005 longi. Amenta florentia 0,09 longa. (v. s.)
27. P. TRIANÆ (Cas. DC. in Seem. Journ. 1866. p. 135), foliis petiolatis subelliptico-lanceolatis apice acutis mucronulatisque basi in petiolum cuneatim subdecurrentibus utrinque glabris junioribus supra parce subtusque ad nervos parce appresso-pubescentibus membranaceis pellucidis enervulosis septuplinerviis, centrali nervo ad apicem ducto paullo supra basin nervos adscendentes subalternos utrinque 3 quorum medii alte supra medium ducti mittente, petiolo juniore villosulo, amentis oppositifoliis folia æquantibus pediculatis foveolatis filiformibus densifloris, pedunculo petiolum æquante juniore appresse pubescente, bractea rotundata centro peltata breviter pedicellata, ovario immerso subovato vertice complanato, stigmate puberulo, bacca subglobosa. ♀ In prov. Antioquia Novæ Granatæ alt. 1500 ped. (Triana n. 65! in h. DC.). Ramuli elongati glabri. Limbi 0,006 longi, 0,027 lati. Petioli 0,006 longi. Amenta matura 0,0025 crassa. (v. s.)
28. P. MINIMA, foliis petiolatis rotundis globosis ciliolatis pellucido-membranaceis enervulosis vel tenuissime triplinerviis, petiolo glabro, amentis terminalibus sublaxifloris, bractea rotundata centro peltata, ovario emerso stigmate carnosulo-puberulo. (1) In ins. Cuba (Wright n. 519! in h. DC.), Portorico (Blau-mer n. 205! in h. DC.). Pep. exigua Griseb. Cat. pl. cub. p. 63, pl. Wright. pars 1. p. 175. Herbula repens glabra. Caulis filiformis. Limbi 0,002 in diam. Petioli 0,002-0,003 longi. Verisimiliter eadem ac Pep. exilis Griseb. sed bacca ignota. (v. s.)
29. P. ALISMÆFOLIA (Presl Epim. bot. p. 222), foliis radicalibus longe petiolatis ovatis obtusis basi rotundatis pellucido-punctatis, pilis tenuibus ciliatis opacis supra pilis brevibus pubescentibus subtus glabris 5-nerviis, centrali nervo supra basin nervum lateralem mittente (id est verisimiliter septuplinerviis) reticulato-nervulosis, amentis radicalibus oppositifoliis cylindricis obtusis quam petiolum duplo brevioribus, bractea peltata ciliato-fimbriata, ovario subgloboso, stigmate orbiculato-umbilicato. ♀ In Cordillera Peruvix. Rhizoma repens, horizontalis, dentatus, pennam columbinam crassus. Folia 21-30 lin. longa, 11-20 lin. lata. Petioli sesqui-bipollicares, glaberrimi.

## 3° Folia multinervia.

30. P. PSEUDOFURCATA, foliis alternis longiuscule petiolatis cordatis lanceolatis apice obtusiusculis glabris membranaceo-pellucidis 7-9-nerviis et reticulato-nervulosis, petiolo glabro, amentis oppositifoliis solitariis, bractea rotundata centro peltata, ovario impresso, stigmate discoideo. ☉ In Peruvia orientali prope Tarapoto (Spruce n. 4009! in h. DC.). Herba basi radicans, an repens? glabra. Limbi vix 0,045 longi. Petioli 0,03 longi. Pedunculi 0,02 longi. Foliis non ciliatis a Pep. furcata Opiz diversa. (v. s.)
31. P. HYDROCOTYLOIDES (Miq. in Linn. v. 20. p. 118. t. 1. fig. 1), foliis omnibus radicalibus longe petiolatis ad  $\frac{1}{3}$  longitud. peltatis rotundato-ovatis vel subrotundis basi lato-rotundatis subtruncatisve membranaceo-pellucidis utrinque glabris 7-nerviis reticulato-nervulosis nervuloque obscuro juxta marginem percursis, petiolo glabro membranaceo pellucido, amentis folia superantibus densifloris, pedunculo petiolum superante membranaceo, bractea rotundata centro peltata subsessili, ovario rhachi impresso obovato-subgloboso vertice complanato. ☉ In locis umbrosis prov. S. Pauli Brasiliæ (Martius! in h. monac.). — Miq. in Mart. flor. Bras. fasc. 11. p. 7. t. 2. fig. 8. Herba acaulis. Limbi 0,023 longi latique. Petioli 0,02-0,03 longi. Radix parva, tuberosa, subglobosa? (v. s.)
32. P. TRANSPARENS (Miq. in Linn. v. 20. p. 121), foliis longe petiolatis paullo supra basin peltatis ovatis apice attenuato-acutiusculis basi rotundatis utrinque glabris tenuiter pellucido-membranaceis 7-nerviis reticulato-nervulosis nervuloque juxta marginem percursis, centrali nervo ad apicem ducto, lateralibus fere ad apicem, sequentibus ad medium ductis, exterioribus subtilioribus, petiolo glabro membranaceo, amentis longissime pedunculatis folia æquantibus maturis laxifloris, pedunculo glabro petiolum superante membra-

naceo, bractea rotundata centro peltata breviter pedicellata, ovario ovato.  $\frac{1}{2}$  In Brasiliæ prov. Minas-Geraes (Mart. in h. monac.). — Miq. in Mart. flor. Bras. fasc. 11. p. 10. Herba radicans, ramosa. Limbi 0,045 longi, 0,03 lati. Petioli 0,08-0,12 longi. (v. s.)

33. P. SANDERSII, foliis longe petiolatis ad  $\frac{1}{2}$  longitud. peltatis ovato-vel cordato-rotundatis apice acutis 7-9-nerviis utrinque glabris supra læte viridibus subtus pallidis, petiolo rubro glabro, amento longe pedunculato terminali vel laterali limbum æquante densifloro, pedunculo petiolum æquante glabro, bractea rotundata centro peltata breviter pedicellata. In Caldariis prope Chelsea culta. Pep. arifolia var. argyreia Miq. in Bot. mag. t. 5634 et Reliq. Hortie. 1867 ex c. ic. Acaulis, glaberrima. Limbi 0,12 longi, 0,093 lati. Petioli 0,16 longi. (Ex Bot. mag. l. c.).
34. P. ARIFOLIA (Miq. syst. p. 72), foliis longe petiolatis paullo supra basin peltatis rotundato-ovatis ovatisve apice acutiusculis basi leviter cordatis vel retuso-truncatis membranaceis subpellucidis pellucido-punctulatis 7-9-nerviis reticulato-nervulosis nervulose juxta marginem percursis, amento terminali longe pedunculato subdensifloro folium multum superante, bractea obovato-rotundata supra medium peltata breviter pedicellata, ovario emerso ovato stigmate orbiculari pilosulo. ☉ In Brasilia (Riedel in h. Kew. et Berol., Fischer in h. DC. ex horto Petrop.), in horto Berolin. culta (h. Berol.). — Miq. ill. p. 12. t. 4. Planta succulenta glabra. Caulis abbreviatus. Limbi 0,06-0,085 longi, 0,05-0,06 lati. Petioli 0,05 longi. Pedunculi 0,095 longi. Amenta florentia 0,115 longa. (v. s.)
35. P. WEDDELLII (Cas. DC. in Seem. journ. 1866, p. 135), foliis longe petiolatis paullo supra basin peltatis ovatis apice breviter acuminatis, acumine acuto, basi rotundatis utrinque glabris membranaceis pellucidis reticulato-nervulosis 7-9-nerviis, centrali nervo ad apicem ducto lateralibus 2 fere ad apicem 2 alte supra medium ductis cæteris brevioribus, nervulo juxta marginem currente, petiolo glabro membranaceo, amentis apice caulis oppositifoliis (foliis sæpe in squamas mutatis) longe pedunculatis, pedunculo glabro petiolum æquante membranaceo, amentis quam folia brevioribus? bractea rotundata centro peltata breviter pedicellata, ovario emerso claviformi, stigmate penicillato. ☉ In Brasilia prope Rio Janeiro (Weddell n. 762! in h. DC.). Herba inter muscos repens. Caulis filiformis, glaber, vix 0,05 altus. Limbi 0,04 longi, 0,035 lati. Petioli 0,06 longi. Amenta florentia 0,001 crassa. (v. s.)
36. P. CLAYTONIODES (Kunth in Ind. semin. horti Berol. 1847), foliis longe petiolatis parum sub centro peltatis ovato-orbicularibus apice acutiusculis obsolete 5-nerviis carnosiss glaberrimis subtus glaucescenti-viridibus, petiolo glabro rubro lineato-punctulato, amentis apice scapi radicalis geminis (uno terminali altero laterali squama fulto) longe pedunculatis folia parum superantibus subaxilloribus, bractea ovata apice acuminata centro peltata, stigmate apice imo ovarii sessili capitellato. ☉ In Guatemala (h. Berol. a F. G. H. comm.). Acaulis. Radix tuberosa, fusiformis. Limbi 10 lin. longi, 9 lin. lati. (v. s.)
37. P. GUAYAQUILENSIS, foliis petiolatis subpeltatis orbicularibus glabris rigidulis ciliatis enervulosis, amentis axillaribus pedunculatis, bractea rotundata centro peltata, ovario emerso, stigmate minuto capitellato. ☉ In prov. Guayaquil ad arbores (Jameson! in h. Kew.). Acrocarpidium Guayaquilense Miq. in Hook. Lond. journ. bot. v. 4, p. 410. Herba repens, scandens. Caulis filiformis, glaber. Limbi 0,0012 in diam. Petioli 0,008 longi. Pedunculi 0,01 longi. (v. s.)
38. P. SECUNDA (Ruiz et Pav. fl. Per. 1. p. 30. t. 45. fig. a), foliis apice caulis confertis longe petiolatis cordato-rotundatis apice acutiusculis obtusisve basi subpeltatis ciliolatis subtus præsertim ad nervos tomentosis pellucido-membranaceis pellucido-punctatis 3-7-nerviis reticulato-nervulosis, nervulo juxta marginem currente, petiolo patentim incano-tomentoso subpellucido, amentis apice caulis elongati aphylli unilateraliter paniculato-confertis squamis lanceolatis fultis filiformibus densifloris, bractea rotundata centro peltata tenuissime membranacea, ovario emerso, stigmate puberulo, bacca rostellata.



- ⊙ In montibus Peruviae (Pav. in h. Boiss. et h. Willd. n. 706!). — A. Dietr. sp. v. 1. p. 163, Miq. l. syst. p. 177, illustr. p. 23. Piper racemosum Vahl enum. v. 1. p. 337, Spreng. syst. v. 1. p. 117. P. paniculatum Willd. l. h. l. c. P. secundum Poir. Herba erecta. Caulis superne aphyllus foliis (in squamis mutatis) glabrior, inferne tomentosus. Limbi 0,06 longi latique. Petioli usque ad 0,09 longi. Amenta pedunculis communibus nascentia, pedunculi communes cauli unilateraliter inserti. (v. s.)
39. P. COTYLEDON (Benth. l. plant. Hartw. p. 145), foliis longe petiolatis orbicularibus glabris infra centrum peltatis pellucidis pellucido-punctatis 5-7-nerviis reticulato-nervulosis, nervulo juxta marginem currente, petiolo tenuissime puberulo, amentis apice scapi communis radicalis paniculatis multifariam dispositis squamis lanceolatis fultis, bractea oblongo-lanceolata centro peltata, ovario subimmerso, stigmatibus papillosulo-carnosulo. ⊙ In Peruviae prov. Quito (Hartweg n. 833! in h. Boiss.). Herba acaulis. Radix fibrosa. Limbi 0,035 longi latique. Petioli circ. 0,09 longi. Amenta pedunculis communibus 0,09-0,1 longis nascentia. (v. s.)
40. P. UMBELLATA (Miq. l. in Hook. Lond. journ. bot. v. 4. p. 427), foliis longiuscule petiolatis rotundatis basi cordatis membranaceis 5-6-nerviis glabris, amentis apice ramulorum circ. 10-umbellatis basi squamis ovato-lanceolatis minutis fultis subaxillariis filiformibus, bractea rotundata centro peltata, ovario emerso. ⊙ In Peruvia (Mathews n. 3230! in h. Kew.). Herba erecta? glabra. Limbi 0,02 longi. Petioli 0,02-0,05 longi. (v. s.)
41. P. CONJUGATA (Kunth l. in H. et B. nov. gen. t. 9. p. 63), foliis longiuscule petiolatis oblongo-subovato-ellipticis apice obtusis basi obtusiusculis acutiusculis utrinque glabris rigidulis subpellucidis trinerviis, centrali nervo ad apicem usque ducto, lateralibus fere ad apicem ductis, nervulo juxta marginem apice currente, petiolo glabro, amentis filiformibus quam folium longioribus subaxillariis apice caulis spicato-confertis. ⊙ In montosis frigidis Andium inter Collessey et Ingatambo alt. 1250 hexap. (Kunth l. c. et Bonpl. l. in h. Francoville ubi P. conjugatum B. dicta). — A. Dietr. sp. v. 1. p. 146, Miq. syst. p. 101. Piper conjugatum R. et Sch. syst. v. 1. p. 326. Caulis glaber. Limbi 0,04 longi, 0,015 lati. Petioli 0,015 longi. (v. s.)
42. P. VILLOSA (Cas. DC. in Seem. journ. 1866, p. 135), foliis longiuscule petiolatis cordato-ovatis apice obtusis supra villosis subtus ad nervos villosis membranaceo-pellucidis 7-nerviis, nervulo juxta marginem currente, petiolo villosulo, amentis axillaribus terminalibusque, pedunculo villosulo, bractea rotundata centro peltata, ovario subimmerso, stigmatibus carnosulo-puberulo. ⊙ In sylvis Andium Quitensium alt. 9000 ped. (Jameson n. 24! in h. DC.). Herba villosa, basi procumbens radicans. Limbi 0,035 longi. Petioli 0,03 longi. Pedunculi 0,035 longi. (v. s.)
43. P. TUBEROSA (Opiz in Reliq. Hænk. fasc. 3. p. 164), foliis radicalibus cordatis 7-nerviis glabris obtusiusculis margine tenuissime pubescentibus, petiolo glabro quam limbus longiore, amentis scapo longioribus, bractea peltata. ¼ In Mexico. — Miq. syst. p. 76. Acaulis. Radix tuberosa cylindrica obtuse grosseque dentata.
44. P. GEMELLA (Miq. l. flor. Ind. bat. v. 1. pars 2. p. 432), foliis petiolatis e basi subcordata rotundatave lato-ovatis apice breviter acuminatis vel obtusiusculis utrinque glabris apicem versus ciliatis fusco-punctatis pellucidis 5-nerviis, centrali nervo ad apicem usque lateralibus supra medium ductis, amentis axillaribus terminalibusque pedunculatis foveolatis, bractea oblonga vel rotundata centro peltata brevissime pedicellata glabra, ovario subimmerso glabro, bacca ovato-attenuata rugulosa. ⊙ In ins. Java rupibus cataractæ prope Tji Kapundung (Zoll. n. 2013! in h. DC.). — Miq. in Zoll. Verz. p. 83, 85. Herba glabra. Limbi 0,02-0,03 longi. Petioli 0,005-0,009 longi. Pedunculi 0,012-0,015 longi. Amenta vix 0,05 longa (v. s.)
45. P. GARDNERIANA (Miq. l. syst. p. 73), foliis longe petiolatis rotundato-ovatis apice obtusis basi cordatis utrinque glabris membranaceis pellucidis 7-9-13-nerviis, nervis 7 centralibus conspicuis cæteris subtilioribus, petiolo glabro

subpellucido, amentis terminalibus axillaribusque pedunculatis frequenter geminatis cylindricis densifloris, bractea rotundata centro peltata ovario ovato apice imo stigma puberulum gerente. ☉ In prov. Geyaz (Gardner n. 2480! in h. Kew.), Tejuco (Pohl n. 4795! in h. Vindob.). — Miq. illustr. p. 12. t. 15, Miq. in Hook. Lond. journ. bot. v. 4. p. 413 et in Linn. v. 20. p. 119. Herba erecta. Caulis simplex. Limbi 0,06 longi. Pedunculi 0,008 longi. (v. s.)

46. *P. SERPENTARIOIDES* (Miq. ! in Linn. v. 20. p. 120), foliis longe petiolatis ovatis apice acuminatis, acumine acuto, basi cordatis utrinque glabris membranaceis subpellucidis 7-nerviis, reticulato-nervuleis centrali nervo ad apicem ducto lateralibus 2 fere ad apicem 3 ad medium ductis exterioribus brevioribus nervuloque obscuro juxta marginem currente, petiolo glabro, amentis oppositifoliis folia æquantibus densifloris longe pedunculatis, pedunculo glabro petiolum paullo superante, bractea rotundata centro peltata subsessili, ovario rhachi impresso obovato vertice complanato, stigmate punctiformi. ☿ In Brasiliæ orientalis sylvis aboriginibus ad terram (Nees ab Esenb. ! in h. Mart.). — Miq. ! in Mart. flor. Bras. fasc. 11. p. 9. Radix rhizomatosa nec tamen crassa. Caulis 0,1 prope modum altus, simplex, glaber. Limbi 0,09 longi, 0,065 lati. Petioli 0,065 longi. (v. s.)

47. *P. SPRUCEANA* (Benth. ! in h. Brit. Mus.), foliis longiuscule petiolatis ovato-rotundatis apice obtusis supra inter nervos pilis longis lineas formentibus vestitis subius glabris membranaceo-pellucidis 5-nerviis reticulato-nervuleis. petiolo glabro vel puberulo, amentis axillaribus? bractea oblonga flayq-pauculata centro peltata, stigmate capitellato. ☿ In Brasilia boreali prope San Gabriel da Cocheira ad Rio Negro (Spruce n. 2246! in h. DC.). Herba repens. Caulis filiformis glaber. Limbi 0,02 longi, 0,015-0,02 longi. (v. s.)

48. *P. PELLUCIDA* (Kunth ! in H. et B. nov. gen. v. 1. p. 64), foliis petiolatis deltoideo-cordatis apice acuminatis obtusiusculis glabris pellucidis pellucido-punctatis membranaceis 7-nerviis, centrali nervo ad apicem ducto lateralibus 2 alte supra medium 2 ad medium ductis exterioribus brevioribus, petiolo basi amplexicauli, amentis oppositifoliis laxifloris glabris, pedunculo glabro, bractea rotundata centro peltata brevissime pedicellata, ovario emerso, stigmate carnosulo puberulo, bacca ovata rostellata. ☉ In Peruvia (Poeppig n. 3112! in h. Berol. et DC.), prope Guayaquil (Jameson n. 352! aut 582! in h. Brit. Mus.), Andibus Æquatorialibus (Spruce n. 5461! in h. DC.), Caracas (Humb. ! in h. Willd. n. 725!), Venezuela prope La Vitoria (Fendler n. 1813! in h. DC.), Brasiliæ prov. Para (Poeppig n. 1990! in h. Vindob. ad Belmont (Menke ! in h. Mart.), Orinoco (Ayres ! in h. Berol.), Guyana gallica (Sagot ! in h. Francovil.), Surinamo (Hostman n. 65! in h. Boiss. Kegel n. 476, fide Miq. in Linn. v. 22. p. 76), Guyana anglica (Schomburgk n. 939! in h. Kew.), Barbadoes (Maycock pl. Barb. p. 40), Trinitate (Sieber n. 207! in h. DC.), Jamaica (Griseb. flor. W. Ind. p. 165), Africa prope Senna (Livingston ! Zamb. exped. in h. Kew.), sylvis umbrosissimis primitivis juxta rivulos alt. 1800-2000 ped. (Welwitsch n. 504! in h. DC.), ins. S. Thomas (Mann ! in h. Kew.). Saururus minor procumbens Plum. Am. ed. Par. p. 54. t. 72. Piper foliis cordatis caule procumbente L. Hort. Cliff. t. 4. *P. pellucidum* L. ! sp. p. 42 (et h. L. !), Hill dict. v. 26. p. 18. t. 18. fig. 3, Aublet hist. pl. Guy. v. 1. p. 21, Lam. ill. p. 82, Vahl enum. v. 1. p. 346, Willd. sp. v. 1. p. 163, Spreng. syst. p. 117. *Peperomia pellucida* Kunth synopsis. v. 117, plant. æq. v. 2. p. 53, A. Dietr. sp. v. 1. p. 164, Miq. ! syst. p. 79. in Hook. Lond. journ. bot. v. 4. p. 121, in Linn. ? v. 20. p. 120. *Micropiper pellucidum* Miq. comm. phyt. p. 54. Herba pedalis vel semipedalis vel minor, pellucida, glabra, ramulosa. Ramuli post 3 internodia plerumque amentis oppositifoliis terminati. Limbi 0,025 longi. Petioli 0,015 longi. (v. s.)

49. *P. PARNASSIEFOLIA* (Miq. ! in Linn. v. 20. p. 119), foliis longiuscule petiolatis caule abbreviato condensatis rotundato-ovatis apice obtusis basi rotundatis leviter cordatis utrinque glabris subcoriaceis subopacis 7-nerviis, reticulato-nervuleis centrali nervo ad apicem usque ducto lateralibus 2 fere ad apicem usque ductis exterioribus brevissimis subtilibus, nervulo interrupto juxta marginem currente, amento longiuscule pedunculato folium æquante densifloro, pedunculo glabro petiolum paullo superante, bractea rotundata

- centro peltata subsessili, ovario rhachi impresso clavato, stigmate punctiformi. ☉ In Brasilia ad arborum truncos (Mart.) in h. Mart.). — Miq.† in Mart. flor. Bras. fasc. 11. p. 9. Herba glabra. Caulis brevissimus. Radix tuberosa, subcylindrica. Limbi 0,03 longi latique. Petioli 0,03 longi. (v. s.)
50. *P. tenuiflora* (Opiz in Reliq. Haenk. fasc. 2. p. 164), foliis petiolatis subrotundo-cordatis basi latioribus apice obtusiusculis glaberrimis pellucidis 6-7-nerviis, petiolo glabro, amentis oppositifoliis folia superantibus capillaribus laxifloris, pedunculo glabro, bractea peltata, bacca ovata sessili mucronata. ☉ In Mexico. Radix fibrosa, annua, Caulis herbaceus, erectus, glaber.
51. *P. Vogelii* (Miq.† in Hook. Lond. Journ. bot. v. 4. 413), foliis petiolatis rotundato-vel lato-ovato-cordatis apice lato-rotundatis summis apice obtusiusculis pellucidis tenuiter pellucido-punctulatis utrinque glabris 6-7-nerviis parce reticulato-nervulosis, amentis axillaribus filiformibus dein laxifloris, bractea rotundata centro peltata pedicellata, ovario cylindrico recto, bacca ovato-globosa rostellata. ☉ In Quorra (Vogel exped. Niger n. 1921 in h. Kew.). Herba erecta, glabra, pauciramosa, succulenta, 0,07 alta. Radix fibrillosa. Limbi 0,074 longi, 0,025 lati. Petioli 0,005-0,015 longi. (v. s.)
52. *P. exiua* (Miq.† syst. p. 77), foliis petiolatis deltoideo-reniformibus basi truncatis rarius reniformibus basi rotundato-cordulatis cordatisve utrinque glabris tenuiter pellucidis 5-7-nerviis reticulato-nervulosis, centrali nervo ad apicem ducto lateralibus 2 supra medium 2 ad medium usque ductis, nervulo interrupto juxta marginem currente, amentis oppositifoliis terminalibusque folia paulo superantibus æquantibusve, bractea rotundata centro peltata, ovario rhachi impresso, stigmate papillosulo-piloso, bacca basi immersa subglobosa apice rostellata. ☉ In Archip. Moluccano (Miq. ann. bot. mus. Lugd. Bat. v. 1. fasc. 5. p. 137), Manilla (Miq. in nov. act. nat. cur. v. 19. p. 494 et Meyen† in h. Berol.), ins. Java (Zollinger n. 3745† in h. DC. et n. 2711† in h. Francovil.), India orient. monte Prome (Wall. n. 6662† in h. DC. et in h. Wall.), Africa ad Factory† (Lenorm. in h. DC.) P. by-nenophylla Miq.† syst. p. 78, flor. ind. bat. v. 1. pars 2. p. 432, plant. Jungh. fasc. 1. p. 14. Piper hyalinum Wall. l. c. Piper exiguum Bl. in Verh. der Bat. Genoots. v. 11. p. 233. f. 36. t. 7, enum. pl. Jav. fasc. 1. p. 75. Micropiper exiguum Miq. comm. phyt. p. 55. t. 9. fig. d. Herbula repens, pellucida. Limbi 0,01 longi, 0,015 lati. Petioli 0,008 longi. (v. s.)
53. *P. freireifolia*, foliis tenuissimis, nervis nervulisque tenuissimis. ☉ In Abyssiniæ umbrosis subrupibus in valle fluvii Toqaze (Schimper n. 1942† in h. DC.) et gregaria ad rupes humidæ juxta rivulum ad occidentem præsidii Pungo Andango alt. 2400-3800 ped. (Welwitsch n. 5031 in h. DC.). P. freireifolium Hochst. (sive Schimper l. c.). Herbula pellucida, humilis. (v. s.)
54. *P. laxiflora* (Kunth† in H. et B. nov. gen. v. 1. p. 63. t. 10), foliis petiolatis ovato-acuminatis, acumine obtusiusculo, basi rotundatis cordulatisve utrinque glabris subtus parce glandulosis 5-nerviis parce nervulosis, centrali nervo ad apicem ducto lateralibus supra medium ductis exterioribus brevioribus, nervulo juxta marginem currente, petiolo glabro, amentis oppositifoliis longiuscule pedunculatis folia multoties superantibus filiformibus laxifloris, pedunculo petiolum duplo superante glabro, bractea rotundata centro peltata subsessili, ovario rhachi impresso obovato basi in pedicellum brevissimum contracto, stigmate puberulo. ☉ In locis temperatis umbrosis regni Novogranatensis prope Mariquita ad truncos Caryocaris amygdaliferi alt. 270 hexap. (Kunth† l. c. et Bonpl.† in h. Berol.). — Kunth synopsis. v. 1. p. 116, A. Dietr. sp. v. 1. p. 163. Piper laxiflorum Spreng. syst. p. 116. Caulis 0,36 altus. Limbi 0,04 longi, 0,02 lati. Petioli 0,007 longi. (v. s.)
55. *P. hlabriana* (Miq.† syst. p. 89), foliis petiolatis ovato-lanceolatis lanceolatisve apice acutis basi acutis acutiusculisve apicem versus supra pilosulis glabrativè subtus glabris ciliolatis utrinque parce reticulato-nervulosis 3-nerviis, petiolo glabro sparse nigro-punctulato, amentis filiformibus subaxillaribus folia æquantibus, pedunculo glabro petiolum æquante, bractea rotundata centro peltata margine pallidior, ovario emerse ovato, stigmate puberulo, bacca globosa apice breviter rostellata. ☉ In Brasilia (Sello† in h. Berol.).

- Miq. in Mart. Flor. Bras. fasc. 11. p. 11. Planta succulenta, spithamea, basi repens radicans stolonifera. Rami erecti, simplices vel pauciramosi. Caulis ramique glaberrimi, nascentes pilis adpressis instructi. Limbi 0,03 longi, 0,016 lati. Petioli 0,01 longi. (v. s.)
55. P. ANDINACEA (Cas. DC. in Seem. journ. 1866. p. 134), foliis petiolatis orbicularibus minutissimis subtus glabris supra pilosulis pellucidis subtiliter 3-nerviis, petiolo glabro, amentis terminalibus densifloris, bractea rotundata membranacea centro peltata subsessili, ovario rhachi semi-immerso, stigmate carnosulo. ④ In Andibus Quitensibus (Jameson! in h. Kew.). Herbula caespitosa, tenerrima. Caulis tenuiter filiformis, glaber. Limbi 0,001 in diam. Petioli 0,002 longi. Amenta immatura 0,01 longa. (v. s.)
56. P. TOVARIANA, foliis petiolatis ovatis basi subcordulatis truncatisve vel subreniformibus apice obtusis supra pilosulis ciliolatisve membranaceo-pellucidis indistincte 3-nerviis vel enerviis, amentis terminalibus? laxifloris, bractea rotundata centro peltata, ovario emerso ovato-attenuato, stigmate discoideo-papilloso. ④ In Venezuela prope coloniam Tovar (Fendler n. 1168! in h. DC. Moritz n. 1941! in h. Brit. Mus.). Herbula repens. Caulis filiformis pilosulus. Limbi 0,003-0,008 longi, 0,004 lati. Petioli 0,0015 longi. (v. s.)
57. P. HAYCOCKII, foliis breviter petiolatis ovato-ellipticis apice obtusis basi attenuatis utrinque pilosulis membranaceis pellucidis 3-nerviis remote reticulato-nervulosis, petiolo pilosulo, amento terminali breviter pedunculato glabro, bractea rotundata centro peltata, ovario rhachi impresso, stigmate discoideo-fimbriato. ④ In ins. Ceylan (Haycock! in h. Kew.). Herbula repens. Caulis filiformis pilosulus. Limbi 0,02 longi. Petioli 0,003-0,004 longi. (v. s.)
58. P. CALEDONICA (Cas. DC. in Seem. journ. 1866, p. 135), foliis petiolatis ellipticis utrinque acutiusculis et pilosulo-pubescentibus membranaceis subopacis pellucido-punctulatis 3-nerviis fere enervulosis nervis subtilibus centrali ad apicem ducto lateralibus alte ductis nervuloque juxta marginem apice currente, petiolo pilosulo, amentis terminalibus, solitariis filiformibus folia paullo superantibus subdensifloris pedunculatis pedunculo petiolum paullo superante, rhachi glabra, bractea rotundata centro peltata subsessili, bacca subglobosa basi rhachi impressa apice subacutata brevissime mucronulata. ④ In Nova Caledonia (Delaplanche! in h. DC.). Herbula repens. Caulis 0,03 alti, pilosulo-pubescentes, filiformes, tenuissimi. Limbi 0,12 longi, 0,07 lati. Petioli 0,002 longi. Amenta vix 0,0002 crassa. (v. s.)
59. P. BANGROOANA (Cas. DC. in Seem. journ. 1866, p. 134), foliis petiolatis orbicularibus utrinque pilosulis margine ciliolatis rigidulis subobscuris enervulosis? amentis terminalibus pedunculatis, bractea rotundata centro peltata breviter pedicellata, pedicello rhachi immerso, ovario rhachi profunde immerso, stigmate carnosulo, bacca subglobosa sessili. ④ In Africa tropicali occidentali ad flumen Bangroo (Mann n. 905! in h. Kew.). Herba repens. Caulis filiformis, subtiliter hirtellus. Limbi 0,008 in diam. Petioli 0,003 longi. (v. s.)
60. P. MASCARENA (Cas. DC. in Seem. journ. 1866, p. 135), foliis brevissime petiolatis rotundis utrinque et margine ciliolatis pellucidis tenuissime membranaceis enervulosis nervulo tamen juxta marginem currente, amentis axillaribus terminalibusve brevissimis densifloris, pedunculo amentum superante, bractea rotundata centro peltata subsessili, ovario rhachi subimpresso obtuso, stigmate minuto, bacca globosa basi rhachi immersa. ④ In ins. Madagascar (Rexburgh! in h. Brit. Mus.), in valle Olifantschoek inter ostia fluvii or. Zondagriri et Boshman (Zeyher! in h. Francovill.). Herbula repens. Caulis filiformis. Limbi 0,005 longi. Amenta 0,004 longa. Pedunculi 0,007 longi. (v. s.)
61. P. NANA (Cas. DC. in Seem. journ. 1866. p. 135), foliis petiolatis orbicularibus vel subreniformibus apice obtusis basi rotundatis truncatisve utrinque glabris tenuissimis pellucidis enervulosis 3-nerviis, centrali nervo ad apicem ducto lateralibus subtilioribus, petiolo pellucido glabro, amentis oppositifoliis subaxillifloris, bacca subimmersula ovato-acuta. ④ In ins. Mohey

- (Boivin! in h. Brit. Mus.). Herbula pellucida, glabra, 0,02 alta. Limbi petiolique 0,003 longi. (v. s.)
62. *P. VINCENTIANA* (Miq.! in Hook. Lond. journ. bot. v. 4. p. 415), foliis breviter petiolatis ellipticis basi subattenuatis apice obtusis glabris membranaceo-rigidulis uninerviis, petiolo glabro, amentis terminalibus solitariis brevibus breviter pedunculatis, bractea rotundata centro peltata apice subattenuata. ☉ In ins. Saint-Vincent (h. Kew!). Herbula repens. Caulis filiformis, glaber. Limbi 0,007 longi. Petioli 0,001 longi. (v. s.)
63. *P. ABYSSINICA* (Miq.! in Hook. Lond. journ. bot. v. 4. p. 419), foliis brevissime petiolatis ellipticis utrinque attenuatis apice imo emarginulatis utrinque glabris apice ciliolatis carnosulis enervulosis 3-5-nerviis nervuloque plus minusve interrupto juxta marginem currente, petiolo glabro, amentis terminalibus sublaevifloris glabris, bractea rotundata carnosulo-papillosula. ☉ In Abyssiniæ rupibus vallis Mashiha (Schimper n. 1319! in h. DC.): ins. Madagascar (Bory! in h. Petrop.). Herba repens, glabra. Internodia vix 0,015 longa. Limbi 0,035 longi. Petioli 0,001-0,002 longi (v. s.)

†† Ovarium apice oblique stigmatiferum.

1° Folia penninervia.

64. *P. STRIATA* (Ruizet Pav. flor. Peruv. v. 1. p. 32. t. 52. fig. b), foliis petiolatis subovato-lanceolatis apice longiuscule acuminatis, acumine acuto, basin versus attenuato-acutis utrinque pubescentibus, centrali nervo nervos alternatos patulo-subadscendentes utrinque circiter 20 mittente, petiolo subtus striato pubescente, amentis apice caulis paniculatum confertis folia æquantibus densifloris, pedunculo quam petiolus dimidio brevior punctis aureis asperso, ovario apice mucronulato. ¼ In nemoribus Cuchero et Chinchao locis umbrosis (R. et P. l. c.), Peruvia (Poeppig! in h. Petrop.). — Miq. syst. p. 87. Planta pedalis. Radix fibrosa. Caulis simplicissimus, erectus, pubescens, striatus, geniculatus, e nodis radicans. Limbi 0,12 longi, 0,058 lati. Petioli 0,022 longi. (v. s.)
65. *P. PILOSA* (Ruiz et Pav.! flor. Per. 1. p. 32. f. 50 fig. a), foliis oblango-lanceolatis basi inæqualibus subcordatisque pellucido-membranaceis supra remote pilosis subtus fusco-punctatis centrali nervo nervos utrinque circiter 10 mittente, petiolo glabro fusco-punctato, amentis axillaribus solitariis primum densifloris, pedunculo puberulo, bractea rotundata peltata fusco-punctata, ovario emerso obovato fusco-punctato stigmate minuto inconspicuo. ☉ In Peruvia (Pav.! in h. Boiss.), in sylvis umbrosis ad Pozuzo (R. et P. l. c.). — Miq. syst. p. 91. (v. s.)
66. *PERNAMBUCENSIS* (Miq.! in Hook. Lond. journ. bot. v. 4. p. 420), foliis petiolatis lato-ellipticis apice breviter attenuatis acutis basi attenuatis in petiolum decurrentibus utrinque glabris, centrali nervo subtus subprominulo nervulos adscendentes subtiles utrinque 7-8 mittente; reticulato-nervulosis, amentis in axillis foliorum rhachi communi cymose alternatim confertis squamis minutis caducis fultis brevibus pedunculatis, bractea rotundata centro peltata, ovario emerso, stigmate minuto, bacca subrotunda minima. ☉ In sylva super magnæ arboris truncum prope Pernambuco (Gardner n. 1157! in h. Kew.), ins. Trinitatis (Griseb. flor. W. Ind. p. 167). — Miq.! in Mart. flor. Bras. fasc. 11. p. 13. Caulis glaber, radicans. Limbi 0,17 longi, 0,075 lati. Petioli 0,035 longi. Pedunculi 0,004 longi. Amenta matura 0,006-0,007 longa. (v. s.)
67. *P. LONGIFOLIA*, foliis longiuscule petiolatis oblango-lanceolatis apice acuminatis acumine acuto mucronulatoque, basi in petiolum decurrentibus utrinque glabris tenuiter membranaceis subpellucidis epunctulatis, centrali nervo fere tota longitudine nervos subtiles alternos subadscendentes utrinque 5-7 mittente, subtiliter nervulosis, petiolo glabro, amentis rhachi communi terminali glabra folia æquante ternatim nascentibus basi squamis lineari-lanceolatis cito deciduis fultis breviter pedunculatis densifloris, bractea rotundata centro peltata subsessili glandulosa, ovario rhachii impresso ovato, stigmate puberulo, bacca cylindrico-ovata. ☉ Ad arborum truncos in sylvis Guyanæ gallicæ (Rich.! in

- h. Francov.), près Cayennæ (Léprieur in h. Francov. ubi *Pep. multipes* dicta). Herba glabra Caulis brevis radicans. Limbi 0,2 longi, 0,046 lati. Petioli 0,04 longi. Pedunculi 0,005 longi. (v. s.)
68. *P. AGUMINATA* (Ruiz et Pav. flor. Peruv. l. p. 32. t. 51. fig. 2), foliis petiolatis obovato-oblongis apice acuminatis, acumine acuto, centrali nervo fere tota longitudine nervos circiter 5 subadscendentes mittente, amento terminali crasso folia superante. ☉ In Muna, Pillab et Macora. *Piper acuminatum* Hill dict. v. 26. p. 18. t. 18. fig. 4? Herba erecta. Folia carnosa. Limbi 0,08 longi. Petioli fere 0,005 longi. (Ex R. et P. l. c.)
69. *P. LARECAJANA*, foliis breviter petiolatis obovato-lanceolatis apice acuminatis, acumine acuto, basi cuneatis in petiolum subdecurrentibus utrinque glabris coriaceis pellucido-punctulatis opacis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos inconspicuos alternos subadscendentes utrinque circiter 14 mittente, petiolo glabro, amentis terminalibus solitariis folia duplo superantibus densifloris crassis, bractea rotundata centro peltata breviter pedicellata, ovario rhachi immerso apice oblique subscutatum complanato, bacca basi flavicante oblonga obtusa. ☿ In Bolivia: prov. Larecacha vicinis Anomea Cerro de Silusani in nemoribus regionis temperatæ (Mandon n. 1115! in h. DC.) Suffrutesculus glaber. Caulis coriaceus. Limbi 0,145 longi, 0,055 lati. Petioli 0,012 longi. (v. s.)
70. *P. ASCENDENS* (Cas. DC. in Seem. journ. 1866, p. 140), foliis petiolatis oblongo-elliptico-lanceolatis apice acutis basi cuneatis in petiolum decurrentibus utrinque glabris coriaceis subopacis pellucido-punctulatis centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos ascendentes subtiles utrinque 8-10 mittente nervuloque obscuro juxta marginem extenuatum currente, petiolo glabro, amento terminali folia æquante crasse filiformi densifloro, pedunculo glabro petiolum æquante, bractea rotundata centro peltata breviter pedicellata, ovario semi-immerso subovato apice oblique subcomplanato, stigmatibus mucronulo puberulo, bacca oblonga basi flavicante. ☿ In Venezuela prope coloniam Tovar alt. 6500 ped. (Fendler n. 1158! in h. DG. et 1135! in h. Boiss.; Poiteau n. 393! in h. Boiss.). Ad arborum truncos scandens. Ramuli glabri, ruguloso-plicatuli, 0,01 crassi. Limbi 0,26 longi, 0,09 lati. Petioli 0,025 longi. Amenta ante maturitatem 0,006 crassa. Bacca 0,001 longa. (v. s.)
71. *P. TALINFOLIA* (Kunth in H. B. nov. gen. l. p. 62. t. 8), foliis brevissime petiolatis oblongo-sublanceolatis apice attenuatis acutiusculisve basi subcuneatis utrinque glabris carnosio-subcoriaceis, centrali nervo fere tota longitudine nervos patulo-subadscendentes utrinque 6 mittente, petiolo glabro, amentis axillaribus terminalibusque solitariis vel geminatis folia æquantibus filiformibus. ☉ In frigidis montibus Quindiu inter los Volcancitos et la Garita del Paríamo alt. 1700 hexap. (Humb. et Bonpl.), ins. Trinitatis (Griseb. flor. W. Ind. p. 167). — Kunth synopsis. v. l. p. 116, Miq. syst. p. 193, Griseb. l. c. *Piper talinifolium* Røem. et Sch. syst. v. 1. p. 325. Caulis 0,26 altus erectus. simplex vel ramosus, carnosus. Limbi 0,085 longi, 0,033 lati. Petioli via 0,005 longi. (v. s.)
72. *P. BASELÆFOLIA* (Kunth! in nov. gen. v. l. p. 66.), foliis petiolatis lanceolatis apice protracto-acuminatis, acumine arcuato, basi cuneatis in petiolum decurrentibus subcoriaceis pellucidis centrali nervo nervulos 7-8 adscendentes mittente, petiolo glabro membranaceo, amentis terminalibus solitariis folia valde superantibus, bractea rotundata centro peltata breviter pedicellata, filamentis elongatis, ovario semi-immerso apice subscutatum complanato, stigmatibus minuto, bacca ovata obtusa vel vix oblique mucronulata. ☉ In Peruvia (Martens n. 1838! in h. Kew.), in umbrasis et scopulosis Peruviae prope Lox alt. 1400 hexap. (Humb. l. c. et in h. Berol.), prov. Popayan alt. 1700 hexap. (Triana n. 61!), Venezuela prope coloniam Tovar (Fendler n. 1154-55! in h. DC.), Caracas (Gallmer n. 337! in h. Berol.) — Kunth synopsis. v. l. p. 119. A. Dietr. sp. v. l. p. 167. *Piper baselæfolium* Røem. et Sch. syst. v. l. p. 333. Suffrutesculus? ramosus glaber. Limbi 0,9 longi. Petioli 0,01 longi. Amenta 0,12 longa, 0,005 crassa. (v. s.)
73. *P. LYALLI* (Cas. DC. in Seem. journ. 1866. p. 150), foliis petiolatis elliptico-oblongis basi cuneatis in petiolum decurrentibus utrinque glabris membra-

necesse subobscuris centrali nervo nervulos 4-5 adscendentes mittente, petiolo glabro, amentis axillaribus solitariis pedunculatis subdensifloris, pedunculo glabro, bractea rotundata centro peltata subsessili, ovario basi immerso obovato, stigmatibus minuto. ♀ In ins. Madagascar (L'Yall n. 308? in h. Kew.). Suffrutescens glaber. Limbi 0,07 longi. Petioli 0,007-0,008 longi. Pedunculi 0,015 longi. (v. s.)

74. *P. MYOSUROIDES* (A. Dietr. v. 1. p. 157), foliis petiolatis oblongo-ovato-ellipticis apice acuminatis, acumine acuto, basi æquali rotundatis utrinque glabris membranaceis centrali nervo nervos subtiles raras utrinque mittente reticulato-venosis, petiolo glabro, amentis folia multoties superantibus filiformibus densifloris, bractea rotundato-subquadrangula centro peltata margine frangulata subsessili, bacca ovata apice mucronulata. ☉ In Guyana (Mart. in h. Rudge in h. Brit. Mus.). — Miq. syst. p. 184. *Piper myosuroides* Rudge! pl. rar. Guy. p. 4. t. 5. Spreng. syst. v. 1. p. 116. Ramuli glabri. Limbi 0,09 longi, 0,045 lati. Petioli 0,008 longi. (v. s.)

75. *P. LIGNESCENS* (Cas. DC. in Seem. Journ. 1866. p. 127), foliis longiuscule petiolatis oblongis apice acuminatis, acumine acuto, basi cordulatis supra glabris subtus ad nervos subtiliter hirtellis utrinque fusco-punctulatis membranaceis subpellucidis novemnerviis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos subalternos adscendentes 4 mittente, petiolo subtiliter hirtello glabrato-ve, amentis axillaribus terminalibusque solitariis folia paulo superantibus densifloris filiformibus, pedunculo quam petiolum brevioris glabro, bractea rotundata aut fusco-punctulata centro peltata subsessili, ovario rhachi immerso ovato apice oblique complanato. ♀ In Costa Rica (Hoffman in h. Berol.) Planta scandens? e nodis infer. radicans. Caulis fere tetragonus, 0,006 crassus. Epidermis canescenti-rugulosa. Limbi 0,055 longi, 0,02 lati. Petioli 0,02 longi. (v. s.)

76. *P. JAPURENSIS*, foliis petiolatis subelliptico-lanceolatis apice acuminatis, acumine acutiusculo, basi acutis utrinque glabris apicem versus ciliolatis membranaceis subopacis pellucido-punctulatis, centrali nervo parum supra basin nervos adscendentes suboppositos utrinque 2 sursumque ad  $\frac{1}{2}$ - $\frac{3}{4}$  longitud. nervum utrinque unum alterne mittente, reticulato-nervulosus nervulogque obscuro juxta marginem currente, petiolo glabro superne ciliato, amentis apice caulibus binatis, uno terminali altero laterali folio rudimentario scarioso fulto, filiformibus densifloris folia duplo superantibus, bractea rotundata centro peltata subsessili, ovario rhachi impresso acutato, bacca oblonga patente vertice oblique complanata mucronulataque basi immersula. ☉ Ad arbores scandens ad parum Miranharum ditionis japurensis (Mart. in h. Monac.). Pep. elongata Miq. in Linn. v. 20. p. 127 et in Mart. flor. Bras. fasc. 11. p. 22, non Kunth. Herba scandens glabra. Caulis e nodis radicans. Limbi 0,075 longi, 0,025 lati. Petioli 0,01 longi. (v. s.)

77. *P. FRAGRANS* (Cas. DC. in Seem. Journ. bot. 1866. p. 140), foliis subsessilibus suboblongo-elliptico-lanceolatis apice longe acuminatis, acumine acuto, basi in petiolum brevissimum cuneatim decurrentibus margine hyalino-extenuatis utrinque glabris membranaceis pellucidis pellucido-punctulatis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos utrinque 4 subalternos adscendentes mittente, reticulato-nervulosus nervulogque sæpe interrupto juxta marginem currente, amentis axillaribus terminalibusque apice ramulorum approximatis filiformibus densifloris folia fere triplo superantibus, bractea rotundata centro peltata subsessili, ovario rhachi impresso, stigmatibus minuto. ☉ In Venezuela inter ripes ad ripas fluminis Tuy et Maya alt. 3000 ped. (Fendler n. 1156! in h. DC.). Glabra. Caulis subpellucidus, quum secatus aromaticè fragrans. Limbi 0,085 longi, 0,023 lati. Amenta vix 0,002 crassa. (v. s.)

78. *P. ELLIPTICO-RHOMBEA*, foliis petiolatis elliptico-rhombeis apice obtusiusculis basi subattenuatis supra ad nervum centralem pilosulis subtus glabris nigropunctulatis coriaceo-membranaceis subopacis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos subadscendentes alternos utrinque 2-3 mittente, petiolo superne pilosulo, amentis axillaribus terminalibusque solitariis filiformibus, pedunculo petiolum subæquante vel paulo superante glabro, bractea ovato-rotundata centro peltata, ovario immerso, bacca ovato-mucronulata basi immersa. ☉ In

Peruvia (Mathews n. 1835 ! in h. Kew.). Caulis sub-4-angularis, pubescens, 0,002 crassus. Limbi vix 0,05 longi, 0,035 lati. Petioli pedunculique vix 0,01 longi. Amenta matura 0,14 longa, 0,0015 crassa. (v. s.)

2° Folia multiplinervia.

79. P. LEVIS (Cas. DC. in Seem. journ. 1866. p. 139), foliis petiolatis inferioribus subrhombeo-lanceolatis superioribus elliptico-lanceolatis apice obtusiusculis brevissime emarginulatis basi subacutis utrinque glabris subtilissime ciliolatis rigidulo-membranaceis subpellucidis reticulato-nervulosis novemnerviis, centrali nervo ad apicem usque ducto, lateralibus utrinque 2 ad  $\frac{1}{2}$  longitud. circiter alterne solutis, cæteris vel et omnibus e basi solutis, nervulo juxta marginem currente, petiolo glabro, amentis terminalibus solitariis densifloris filiformibus folia paullo superantibus, pedunculo glabro petiolum æquante, bractea rotundata centro peltata subsessili, ovario subimmerso apice breviter rostellato, stigmate minuto. ☉ In Venezuela prope coloniam Tovar alt. 6500 ped. (Fendler n. 1165 ! et 1164 ! in h. DC.). Fruticulus lævis, glaber, basi radicans, procumbens, ramulos suberectos vix 0,1 altos mittens. Folia apice imo ramulorum opposita. Limbi 0,047 longi, 0,024 lati. Petioli 0,008 longi. Amenta florentia 0,002 crassa. (v. s.)
80. P. SURINAMENSIS, foliis petiolatis lanceolatis apice acuminatis, acumine acuto, basi cuneatis utrinque glabris apicem versus ciliolatis membranaceis subpellucidis sparse nervulosis septuplinerviis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos alternos utrinque 3 mittente nervuloque juxta marginem currente, petiolo superne ciliolato, amentis terminalibus solitariis pedunculatis folia subsuperantibus densifloris filiformibus, bractea rotundata centro peltata breviter pedicellata, ovario semi-immerso, bacca cylindrico-ovata patente nigra apice oblique subacutata. ♀ In agris prope urbem Paramaribo (Kappler n. 1577 ! in h. DC.), Acarouany (Sagot n. 538 ! in h. Lenorm.). Suffrutescens scandens, glaber, e nodis radicans. Limbi 0,07-0,08 longi, 0,02 lati. Petioli 0,015 longi. Baccæ 0,001 longæ. Caulis apicem versus ciliolatus. (v. s.)
81. P. CARLOSIANA (Cas. DC. in Seem. journ. bot. 1866. p. 140), foliis approximatis longe petiolatis ovato-lanceolatis apice acuminatis basi subcordatis cordatisve membranaceis pellucidis fusco-punctulatis glabris septuplinerviis centrali nervo validiore ad apicem usque ducto lateralibus paullo supra basin solutis omnibus subprominulis nervulo juxta marginem currente, petiolo glabro, amentis axillaribus terminalibusque, bractea rotundata centro peltata fusco-punctulata, ovario emerso obtuso, stigmate minuto. ☉ In Venezuela ad flumen San Carlos (Fendler n. 1148 ! in h. DC.). Herba stolonifera ? Limbi 0,08 longi, 0,035 lati. Petioli 0,05-0,06 longi (v. s.)
82. P. VELLOZIANA (Miq. ! syst. p. 88), foliis breviter petiolatis oblongo-vel rhombeo-lanceolatis apice attenuato-obtusis obtusiusculisve basin versus cuneatim in petiolum contractis utrinque glabris membranaceis subpellucidis septuplinerviis sparse reticulato-nervulosis, centrali nervo nervos adscendentes suboppositos utrinque 3 mittente nervuloque interrupto juxta marginem currente, petiolo glabro, pedunculo glabro petiolum æquante, amentis axillaribus terminalibus filiformibus densifloris sæpe paniculatim confertis folia duplo triplove superantibus, bractea obovato-rotundata margine pallidior supra centrum peltata, stigmate minuto, bacca ovata breviter rostellata. ☉ In Brasilia (Sello ! in h. Berol.). Planta succulenta 0,35-0,4 alta. Caulis glaber, inferne aphyllus, ramos paucos majores patentes vel erectos mittens. Limbi 0,055 longi, 0,022 lati. Petioli 0,004 longi. (v. s.)
- β, *polysticta* (Miq. in Linn. v. 18. p. 226). Tota glandulis fuscis crebris confersa. Statura paullo minor. ☉ In arboribus vetustis prope Surinam (Miq. l. c.) et locis putridis prope Tourtione (Kegel n. 437 a, fide Miq. in Linn. v. 22. p. 76).
83. P. MELANOSTIGMA (Miq. ! syst. p. 90), foliis petiolatis lanceolatis vel subrhombeo-lanceolatis utrinque glabris et nigro-punctulatis membranaceo-rigidis haud nervulosis obscure quintuplinerviis, lateralibus nervis fere e basi solutis, amentis axillaribus terminalibusque folia multoties superantibus subaxillaribus



filiformibus, pedunculo glabro petiolum æquante vel superante, bractea ob-ovata nigro-punctulata centro peltata subsessili, ovario subimpresso apice acutiusculo, bacca ovato-acuta rostellata. ☉ In Surinam (Hostmann n. 437! in h. Berol.), Cuba (Sagra! in h. Francov., ubi *Piper mucronatum* Rich. dicta, Santos! in h. DC.), Martinica (Belanger! in h. Francov.), Costa-Rica (Belanger! in h. Francov.). — Miq. l in Hook. Lond. journ. bot. v. 4. p. 415. *Micropiper melanostigma* Miq. comm. phyt. p. 5. t. 9. fig. A. Caulis glaber. Limbi 0,055 longi, vix 0,02 lati. Petioli 0,01 longi. Amenta immatura 0,15 longa, vix 0,001 crassa. (v. s.)

β, *nervulosa*, foliis sparse reticulato-nervulosis, nervulo juxta marginem currente. ☉ In Surinam (Hostmann n. 437! spec. sinistr. in h. Berol.).

γ, *glabrior*, foliis glabrioribus rigidioribus apicem versus ciliolatis. ☉ In Guadalupe (Krauss! in h. DC.), Jamaica (Swartz! in h. monac.), in arborum truncis prope Paramaribo (Kegel n. 438 fide Miq. in Linn. v. 22. p. 76), in Brasilia (Riedel et Langsd. ! in h. Petrop.) (v. s.)

84. P. PATULA (Cas. DC. in Seem. journ. 1866. p. 139), foliis petiolatis elliptico-lanceolatis apice obtusiusculis obtusivse basi cuneatis acutis utrinque pubescentibus membranaceis quintuplinerviis reticulato-nervulosis, centrali nervo paullo supra basin nervos 2 quorum medii ad medium exteriores infra medium ducti mittente, petiolo dense pubescente, amentis plerumque apice ramulorum binatis breviter pedunculatis subaxillaris folia fere triplo superantibus, bractea rotundata centro peltata subsessili, ovario immerso, bacca ovato-globosa apice oblique substrostellata. ☉ In Venezuela prope coloniam Tovar alt. 6500 ped. (Fendler n. 1166! in h. DC.). Herba ad arborum truncos parasitica. Caulis patule ramosus, pubescens. Folia tenuissima. Limbi 0,07 longi, 0,036 lati. Petioli 0,01 longi. (v. s.)

85. P. DENDROPHILA (Schlecht. ! in Linn. v. 5. p. 74), foliis petiolatis elliptico-oblongo-ovatis vel obovato-subrhombis basi cuneato-attenuatis apice paullo protractis obtusiusculis utrinque glabris membranaceo-subpellucidis quintupli-septuplinerviis, parce reticulato-nervulosis, nervis omnibus fere e basi solutis, nervulo interrupto juxta marginem currente, amentis axillaribus terminalibusve longe pedunculatis filiformibus densifloris folia multoties superantibus, bractea rotundata apice acuta centro peltata, ovario subimmerso, stigmatibus minutis, bacca ovato-acuta semi-immersa apice breviter mucronulata. ☉ In Mexico prope Xalapa (Schiede! in h. Berol., Galeotti n. 6002! in h. Deless.), Miradores (Linden n. 129! in h. Boiss.), Costarica (Hoffman n. 56! in h. Berol.). — Miq. ! syst. p. 89, illustr. p. 13. t. 7, in Hook. Lond. journ. bot. v. 4. p. 415, Griseb. flor. W. Ind. p. 165. *Piper Galeottianum* Mart. et Gal. in bull. Acad. Brux. v. 10. p. 18. Herba basi radicans, glabra. Limbi 0,04 longi. Petioli 0,01 longi. Pedunculi 0,015-0,03 longi. (v. s.)

86. P. NIGRO-PUNCTATA (Miq. ! syst. p. 188), foliis petiolatis elliptico-lanceolatis apice suboblique acuminatis, acumine acuto, basi subinæquali acutiusculis utrinque glabris crebre nigro-punctulatis membranaceis pellucidis haud nervulosis quintuplinerviis, centrali nervo paullo supra basin nervos suboppositos utrinque 2 mittente nervuloque juxta marginem currente, petiolo glabro nigro-punctulato, amentis axillaribus solitariis filiformibus subdensifloris maturis folia triplo-quadruplo superantibus, pedunculo petiolum fere æquante nigro-punctulato, rachis nigro-punctulata, bractea rotundata nigro-punctulata centro peltata, ovario semiimmerso ovato apice acuto nigro-punctulato, stigmatibus carnosulo puberulo. ☉ In Martinica (Sieber n. 6! in h. Berol.), rupibus saxosis sylvarum prædii La Perle ins. S. Thomas (Rich. ! in h. Francov.), Cuba (Wright n. 503! pro parte in h. DC.), Antigua (Nicholson in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 429). *Pep. glabella* Griseb. flor. v. Ind. p. 165. Herba basi decumbens 0,2-0,3 alta, tota glaberrima et nigro-punctulata, erecta. Limbi 0,065-0,08 longi, 0,026-0,045 lati. Petioli 0,01-0,015 longi. Amenta 0,001 crassa. (v. s.)

87. P. CAULIBARBIS (Miq. ! syst. p. 98), foliis petiolatis ovato-ellipticis vel elliptico-rhombis apice breviter protracto-obtusiusculis basi subcuneatis utrinque glabris crebre nigro-punctatis membranaceo-subcoriaceis tripli-quintupliner-

vis, centrali nervo validiore subtus prominulo, ceteris tenuioribus fere e basi vel paulo supra basin subopposite ortis, nervulo juxta marginem currente, petiolo ciliato, amentis axillaribus vel subpaniculatis confertis (et tam pedunculis squamis fultis) folia multoties superantibus filiformibus densifloris elongatis, bractea rotundata centro peltata, ovario subsemimmerso, stigmate minuto. ☉ In ins. S<sup>a</sup> Catharina (Gaudich. n. 287! in h. DC. et Deless.), Brastilia (Cham. et Sello! in h. Berol.), Ilheos super arbores parasitica (Wawra et Maly n. 251! fide Wawra bot. Ergeb. p. 132. t. 52. fig. 1), ad Tomaca (Schott! in h. Vindob.), Portorico (Blauher n. 203! in h. DC.), S<sup>a</sup> Domingo (Mayerhoff p. 344 in h. Berol. et Rich. I in h. Francov.), Cuba (Sagra! in h. Francov. ubi *Pep. linatifolia* Rich. dicta). — Miq. in Linn. v. 20. p. 123 et in Mart. flor. Bras. fasc. 11. p. 12, Wawra l. c. Herba basi procumbens radicans. Caulis uno latere serie una pilorum e petiolis decurrentibus instructus. Limbi 0,045 longi. Petioli 0,005-0,007 longi. Amenta 0,09-0,12 longa. (v. s.)

88. *P. COLORATA* (Kunth! in H. et B. nov. gen. v. 1. p. 64. t. 11), foliis petiolatis oblongo-ellipticis utrinque acutiusculis utrinque glabris epuncatulis rigidulis subopacis triplinerviis lateralibus nervis fere e basi solutis nervulo juxta marginem currente, petiolo glabro, pedunculo glabro petiolum paulo superante, amentis apice caulis fasciculato-congestis folia multoties superantibus filiformibus densifloris, bractea rotundata centro peltata subsessili, ovario subimpresso obovato, stigmate discoideo minuto. ☉ In saxosis temperatis prov. Popayan juxta vicum La Erre inter montem Paraguay et altam planitiem Sacandoy alt. 1050 hexap. (Kunth l. c. et Bonpl. I in h. Francov.), Caracas (Gallmer n. 336! in h. Berol.). — A. Dietr. sp. v. 1. p. 150, Miq. syst. p. 95. *Piper coloratum* Roem. et Sch. syst. v. 1. p. 319. Caulis glaber, subquadrangulus. Limbi fere 0,03 longi, 0,02 lati. Petioli 0,005 longi. Amenta florentia 0,075 longa. (v. s.)

### 3<sup>a</sup> Folia multinervia.

89. *P. POLYBOTRYA* (Kunth! in H. et B. nov. gen. v. 1. p. 60. t. 5), foliis petiolatis ad  $\frac{1}{2}$  longitud. peltatis subrotundo-ovatis apice acuminatis, acumine acuto, basi rotundatis utrinque glabris rigidulis pellucido-punctulatis 11-nerviis parce nervulosis, petiolo glabro, amentis terminalibus axillaribusve brevissime pedunculatis filiformibus densifloris quam folia quadruplo brevioribus paniculatim confertis squamis caducis fultis, bractea rotundata centro peltata subsessili, ovario impresso oblongo, stigmate carnosopuberulo, bacca subglobosa basi impressa apice oblique submucronulata. ☉ In umbris temperatis inter Sondorillo et flumen Guancabamba (Humb. et B. l. c., Bonpl. I in h. Francov. et Humb. I in h. Berol.). — Kunth synops. v. 1. p. 114, A. Dietr. sp. v. 1. p. 165, Miq. syst. p. 552. *Piper polybotryon* Spreng. syst. v. 1. p. 117, Roem. et Sch. syst. v. 1. p. 324. Caulis teres, glaber. Limbi 0,09 longi, 0,065 lati. Petioli 0,04-0,05 longi. (v. s.)

90. *P. TRISTACHYA* (Kunth! in nov. gen. l. p. 61. t. 6), foliis longiuscule petiolatis basi ad  $\frac{1}{4}$  longitud. peltatis subrotundato-ovatis apice breviter acuminatis, acumine acuto, basi rotundatis subsinuoso-cordulatis utrinque glabris membranaceis pellucido-punctulatis subpellucidis 11-13-nerviis et reticulato-nervulosis, centrali nervo ad apicem ducto, lateralibus nervis utrinque 2 ad apicem usque et 2 ad medium ductis, nervulo juxta marginem currente, petiolo glabro, amentis apice pedunculi communis ternis breviter pedunculatis quam folia triplo brevioribus filiformibus densifloris, pedunculo singulo quam petiolum multum brevior, bractea rotundata glandulosa centro peltata, ovario impresso oblongo, stigmate minuto, bacca subglobosa, apice oblique breviter mucronulata. ☉ In regni Novo-granatensis montosis prope Pandi alt. 570 (Kunth l. c.). — Kunth synops. v. 1. p. 114, Bonpl. I in h. Francov., A. Dietr. syst. v. 1. p. 159, Miq. syst. p. 553. (v. s.)

91. *P. LANCEOLATO-PELTATA* (Cas. DC. in Seem. journ. 1866. p. 136), foliis longe petiolatis paulo supra basin peltatis ovato-lanceolatis apice parum protractis acutis ciliatis supra glabris subtus pilosis membranaceo-pellucidis septem-

- nerviis et reticulato-nervulosis, amentis solitariis filiformibus longe pedunculatis pedunculo rhachique ciliolatis, bractea rotundata centro peltata, ovario semimmerso, stigmate minuto, bacca semimmersa ovato-acuta. (1) In Venezuela prope coloniam Tovar (Fendler n. 1149! in h. DC.), regione tempestata prov. Caraccasanæ (Moritz n. 1979! in h. Francov.), Costa-Rica (Hoffmann n. 414! in h. Berol.). — *Pep. ciliata* Mor.! in h. Francov. Herba subacaulis stolonifera. Limbi 0,06 longi. Petioli 0,04 longi. Pedunculi 0,04 longi. Amenta circ. 0,1 longa. (v. s.)
92. *P. TARAPOTANA* (Cas. DC. in Seem. journ. 1866. p. 137), foliis longissime petiolatis basi peltatis ovato-cordatis apice acuminatis basi cordatis utrinque glabris membranaceo-pellucidis 10-nerviis sub lente tenuiter reticulato-nervulosis, petiolo glabro submembranaceo, amentis axillaribus terminalibusque 3-5-approximatis singulis basi squamula plus minusve foliacea fultis, bractea rotundata centro peltata, ovario emerso, stigmate discoideo-papilloso. (1) In Peruvia orientali prope Tarapoto (Spruce n. 4570! in h. DC.). Planta stolonifera ex foliorum axillis scapos elongatos margine membranaceos amentiferos mittente. Limbi circ. 0,12 longi. Petioli circ. 0,14 longi. (v. s.)
93. *P. CORPULATA* (Cas. DC. in Seem. journ. 1866. p. 137), foliis petiolatis basi subpeltatis ovato-rotundis vel ovato-acutis basi cordulatis utrinque glabris membranaceis subpellucidis subtiliter 11-nerviis, petiolo glabro, amento terminali cylindrico densifloro, bractea rotundata centro peltata subsessili, ovario emerso impresso apice acuto, stigmate minuto, bacca ovato-acuta. (1) In Isthmo Panama (Fendler n. 265! in h. Kew.). Caulis glaber. Limbi 0,07 longi, 0,065 lati. Petioli 0,01 longi. Amentum florens 0,15 longum, 0,003 crassum. (v. s.)
94. *P. SUBPALTATA* (Cas. DC. in Seem. journ. 1866. p. 136), foliis longissime petiolatis rotundo-cordatis basi subpeltatis apice acutiusculis membranaceis pellucidis utrinque glabris 5-nerviis et crebre reticulato-nervulosis nervulo juxta marginem currente, petiolo glabro, amentis axillaribus longe pedunculatis subaxillifloris, pedunculo glabro, bractea rotundata centro peltata, ovario emerso, stigmate minuto. (1) In Pichincha (Jameson n. 62! in h. Kew.). Herba glabra repens? scandens? Caulis filiformis. Limbi 0,02 longi, 0,02-0,03 lati. Petioli 0,04-0,08 longi. Pedunculi 0,03-0,035 longi. (v. s.)
95. *P. DEPENDENS* (Ruiz et Pav. ! flor. Per. 1. p. 33. t. 48. f. a), foliis longe petiolatis basi ima subpeltatis ovato-cordatis apice acuminatis subinæquilateralibus glabris pellucido-membranaceis 9-13-nerviis reticulato-nervulosis nervuloque juxta marginem currente, petiolo glabro, amentis longe pedunculatis terminalibus densifloris, bractea obovata centro peltata, ovario emerso, stigmate orbiculari exiguo. (1) In Peruvia locis umbrosis ad Pozuzo (R. et P. l. c., Pav. ! in h. Boiss. et Ruiz ! in h. Willd. n. 707!), prov. Pasto in Nôva Granada alt. 1200 m. (Triana n. 64! in h. suo). — *Miq. ! syst. p. 83. Piper dependens* h. Willd. ! l. c., Vahl enum. v. 1. p. 338. *Pep. Opiziana* A. Dietr. sp. v. 1. p. 162, *Miq. in Linn. v. 20. p. 121, et syst. p. 85* (vide *Miq. in Linn. l. c.*). *Pep. cordifolia* Opiz in Reliq. Hænk. fasc. 3. p. 144. Herba pedatis glabra. Limbi 0,15-0,16 longi. Petioli 0,1 longi. Amenta matura pedunculis longiora, apice scapi aphylli squamis fulta. (v. s.)
96. *P. PSEUDO-DEPENDENS* (Cas. DC. in Seem. journ. 1866. p. 137), foliis longissime petiolatis basi ima subpeltatis rotundato-ovatis apice subattenuatis acutiusculis basi profunde cordatis utrinque glabris tenuissime membranaceis pellucidis epunculatatis plerumque 11-nerviis reticulato-nervulosis nervuloque juxta marginem currente, petiolo glabro membranaceo, amentis axillaribus terminalibusque solitariis subaxillifloris, bractea rotundata centro peltata, ovario impresso, stigmate minuto, bacca globosa apice subacuta basi immersa. (1) In Venezuela prope La Victoria alt. 2100 m. (Fendler n. 181! in h. DC.). Herba tenera. Limbi 0,11 in diam. Petioli 0,09 longi. Pedunculi vix 0,02 longi. Amenta florentia 0,09 longa, 0,002 crassa. (v. s.)
97. *P. BELANGERI*, foliis petiolatis lanceolato-ellipticis apice acuminatis, acumine obtusiusculo, basi acutis utrinque crebre nigropunctulatis supra glabris subtus adpresse piloso-pubescentibus pallidioribus ciliatis membranaceis

- subpellucidis 7-nerviis, nervis exterioribus subtilissimis, nervulo obscuro juxta marginem currente, petiolo piloso-pubescente, amentis terminalibus axillaribusque, pedunculo piloso-pubescente petiolum subæquante, amento folium æquante vel parum superante, bractea rotundata centro peltata subsessili, ovario emerso. ☉ In ins. Martinica (Belanger n. 1371 in h. Boiss.). Herba basi procumbens. Caulis piloso-pubescent. Limbi 0,046 longi, 0,026 lati. Petioli 0,01 longi. (v. s.)
98. P. MIQUELIANA (Cas. DC. in Seem. journ. 1866. p. 138), foliis sessilibus subsessilibusve ovatis apice attenuatis obtusiusculis basi cordatis supra glabris subtus præsertim ad nervos pubescentibus membranaceo-rigidis opacis 7-nerviis, nervis subtus prominulis centrali nervo ad apicem usque lateralibus supra medium ductis, amentis axillaribus terminalibusque solitariis breviter pedunculatis filiformibus subdensifloris, bractea rotundata centro peltata, ovario immerso, stigmate minuto carnosulo, bacca immersa ovato-acuta apice brevissime mucronulata. ♀ In Andibus Quitensibus alt. 6000 ped. (Jameson n. 737 in h. Kew.). Caulis glaber. Limbi 0,02-0,025 longi, media parte 0,02 lati. Pedunculi 0,005 longi. Folia inferiora cordato-ovata, apice obtusa. (v. s.)
99. P. LONGE-PEDUNCULATA (Opiz in Reliq. Hænk. fasc. 3. p. 164), foliis longe petiolatis cordatis obtusiusculis subundulatis glabris 7-nerviis, petiolo glabro folium æquante, amentis oppositifoliis terminalibusve solitariis filiformibus quam pedunculi brevioribus. ♀ In montanis Huanocensibus. — A. Dietr. sp. v. 1. p. 161, Miq. syst. p. 84. Caulis herbaceus glaber carnosus. (Ex Opiz l. c.)
100. P. RHOMBOIDEA (Hook. et Arn. in Beech. voy. bot. p. 70), foliis breviter petiolatis rotundatis vel rhombeo-rotundatis apice breviter acuminatis acutiusculis glabris membranaceis pellucidis remote reticulato-nervulosis 7-nerviis, centrali nervo ad apicem ducto lateralibus utrinque 2 supra medium et exterioribus sub medium ductis. ☉ In insulis Coralicis, Nohau (Beechey in h. Kew.). — Miq. syst. p. 103 et in Hook. Lond. journ. bot. v. 4. p. 418. Piper acuminatum Forster prodr. n. 23<sup>9</sup>. Caulis herbaceus, glaber, teres, basi radicans, 0,002-0,003 crassus. Limbi 0,06 longi, 0,045 lati. Petioli 0,005 longi. (v. s.)
101. P. HIRTA, foliis longiuscule petiolatis rhombeo-ellipticis apice acutis vel obtusiusculis obtusive basi subcuneato-acutis utrinque pubescentibus ciliatis membranaceo-subpellucidis 5-7-nerviis et reticulato-nervulosis, petiolo hirsuto, amentis axillaribus terminalibusque pubescentibus, bractea rotundata ciliata centro peltata, ovario semi-immerso, stigmate discoideo. ☉ In Cuba (Wright n. 512 in h. DC.). Pep. hirtella Miq. (fide Griseb. pl. Wright. Cub. p. 175), Griseb. Cat. pl. Cub. p. 64, non Miq. in Hook. Lond. journ. bot. Herba carnosa hirsuta. Limbi 0,05 longi. Petioli 0,03 longi. (v. s.)
102. P. SAN-CARLOSIANA (Cas. DC. in Seem. journ. bot. 1866. p. 138), foliis longiuscule petiolatis ovato-rhombeis apice acutiusculis obtusiusculisve basi acutis utrinque puberulis et parce nigro-punctulatis tenuiter membranaceis pellucidis reticulato-nervulosis 5-7-nerviis, nervis 3 centralibus fere usque ad apicem ductis cæteris medio evanidis subtilibus, nervulo juxta marginem currente, petiolo glabro, amentis terminalibus elongatis folia multoties superantibus filiformibus subaxillifloris, bractea rotundata centro peltata, ovario subimpresso fusco-punctato, stigmate minuto. ☉ In Venezuelæ valle San-Carlos (Fendler n. 1151 in h. DC.). Herba glabra. Caulis erectus simplex membranaceus subpellucidus. Limbi 0,045 longi. Petioli 0,02 longi, media usque ad 0,2 longa. (v. s.)
103. P. JAMESONIANA, foliis longe petiolatis rotundato-cordatis reniformibusve basi cordatis glabris membranaceo-pellucidis 5-7-nerviis reticulato-nervulosis, amentis axillaribus longe pedunculatis subdensifloris, pedunculo membranaceo glabro, bractea rotundata centro peltata subsessili, ovario obtuso, stigmate minuto. ☉ In Pichincha (Jameson n. 641 in h. Kew.). Acrocarpidium Jamesonianum Miq. in Hook. Lond. journ. bot. v. 4. p. 412. Herba repens glabra. Caulis filiformis. Limbi 0,02 longi, 0,03 lati. Petioli 0,06 longi. Pedunculi 0,06 longi. (v. s.)
- ♂, *microphylla*, foliis multo minoribus, bacca mucronulata (in specimine

sessili). In Andibus Ecuadoriensibus (Spruce n. 6857! in h. Kew. et DC.).  
Acrocarp. Jamesonianum Miq. 1 mss. in h. Kew. Limbi 0,008 longi. (v. s.)

104. P. ENDLICHERI (Miq. 1 in Hook. Lond. journ. bot. v. 4. p. 418), foliis petiolatis e basi cuneata obovatis vel obovato-ellipticis vel ellipticis apice acutiusculis glabris membranaceis pellucidis sparse nervulosis 7-5-nerviis, nervo centrali nervisque lateralibus supra medium ductis exterioribus infra medium ductis vel obsoletis nervulo juxta marginem currente, amentis terminalibus axillaribusque, pedunculo glabro subpellucido folia multoties superante, bractea rotundata centro peltata, ovario semi-immerso apice subacuto, stigmate discoideo minuto, bacca ovato-acuta basi immersa. ☉ In insulis Norfolk (Cunningh. ! in h. Kew. et Endlich. ! in h. Berol. Mac Gillivray n. 971 ! in h. Kew.), Nova Zelandia (in eod. h. !). — Miq. syst. p. 102 in Hook. Lond. journ. bot. v. 4. p. 417. Piper simplex Endl. ! in Prodr. Norf. p. 37, Bauer ill. Norf. t. 12 α (teste Endl. l. c.). Pep. Urvilliana (A. Rich. in d'Urv. voy. Astrol. p. 356), Hook. f. flor. Nouv. Zeal. v. 1. p. 228. Herba repens. Caulis glaber, subpellucidus, 0,002-0,003 crassus. Limbi 0,026 longi, 0,02 lati. Petioli 0,003-0,004 longi. Pedunculi 0,01 longi. (v. s.)
105. P. RUPERTIANA, foliis breviter petiolatis ellipticis vel elliptico-lanceolatis vel obovato-ellipticis apice breviter subattenuatis obtusiusculis obtusive basi subacutis acutisve utrinque glabris nigro-punctatis membranaceis subpellucidis remote nervulosis 7-nerviis centrali nervo ad apicem usque lateralibus utrinque 2 alte ductis exterioribus subtilioribus sæpe obsoletis, petiolo glabro, amentis axillaribus solitariis densifloris filiformibus, pedunculo glabro petiolum subæquante, bractea rotundata centro peltata, ovario impresso oblongo, stigmate minuto. ☉ In ins. Dominicæ super Princip. Rupert caput (in h. Kew.), et Jardin ! in h. Lenorm.). Pep. acuminata Miq. 1 in Hook. Lond. journ. bot. v. 4. p. 416. Caulis fere tetragonus glaber. Limbi 0,05 longi, 0,03 lati. Petioli et pedunculi 0,007 longi. Amenta florentia 0,06-0,07 longa. (v. s.)
106. P. CUBANA, foliis breviter petiolatis elliptico-lanceolatis apice obtusiusculis basi acutis utrinque glabris membranaceis subpellucidis enervulosis 5-nerviis, centrali nervo ad apicem ducto lateralibus 2 paullo validioribus supra medium ductis exterioribus subtilioribus, petiolo glabro, amentis axillaribus solitariis densifloris filiformibus, pedunculo petiolum subæquante, bractea rotundata centro peltata. bacca ovato-acuta mucronulata basi subimmersa. ☉ In Cuba (Wright n. 504 ! in h. Kew.), Martinica (Sieber n. 61 in h. Berol. cum Pep. nigro-punctata), S. Crux (Rich. ! in h. Francov.), Ind. occ. (West ! in h. DC.). Caulis subquadrangulus glaber. Limbi 0,065 longi, 0,035 lati. Petioli pedunculique 0,005-0,007 longi. (v. s.)
107. P. DEFOLIATA (Cas. DC. in Seem. journ. bot. 1866. p. 138), foliis petiolatis subrhombeo-elliptico-lanceolatis, superioribus subrotundato-rhombis utrinque hirsuto-pubescentibus rigidulis subopacis fere enervulosis 5-nerviis centrali nervo ad apicem usque lateralibus 2 supra medium ductis exterioribus subtilioribus, petiolo hirsuto, amentis axillaribus terminalibusve pedunculatis maturis folia fere quadruplo superantibus filiformibus subdensifloris, bractea rotundata glandulosa centro peltata breviter pedicellata, ovario emerso subovato glanduloso, stigmate carnosulo pilosulo, bacca emersa ovata globosa apice mucronulata. 2 In Andibus Bogotensibus alt. 2650 ped. (Triana n. 51 ! in h. DC.), Ecuador (Fraser ! in h. DC.). Caulis basi aphyllus, hirsutus, subquadrangulus, fere 0,002 crassus. Limbi 0,025 longi, 0,015 lati. Petioli 0,006 longi. (v. s.)
108. P. DASYSTACHYA (Miq. 1 in Hook. Lond. journ. bot. v. 4. p. 416), foliis breviter petiolatis superioribus majoribus elliptico-lanceolatis, acumine obtusiusculo, mediis ellipticis basi et apice subattenuatis, infimis minoribus ellipticis apice obtusiusculis basi obtusis omnibus utrinque dense appresse subtilissime pubescentibus subtilissime ciliolatis membranaceis subpellucidis subtiliter fusco-punctulatis reticulato-nervulosis, nervulo juxta marginem currente, 5-nerviis, centrali nervo paullo validiore ad apicem usque ducto lateralibus subtilibus quorum 2 supra medium ducti exteriores subtiliores, petiolo adpresse subtilissime pubescente, amentis terminalibus filiformibus

densifloris, pedunculo petiolum æquante, bractea rotundata pilosula ciliolata centro peltata brevissime pedicellata, ovario emerso obtuso, stigmate carinatosulo penicillato. ☉ In Peruvia prope Chachapoyos (Mathews n. 3229! in h. Kew.). Caulis herbaceus, erectus, subtilissime pubescens. Limbi 0,08 longi, 0,035 lati. Petioli et pedunculi 0,005-0,007 longi. Amenta florentia vix 0,002 crassa. (v. s.)

109. *P. BERTEROANA* (Miq. ! syst. p. 114), foliis sessilibus cuneato-spathulatis apice paullo protractis obtusiusculis supra glabris subtus hirtellis membranaceis vel subcoriaceis subpellucidis reticulato-nervulosis et 5-nerviis, centrali nervo validiore ad apicem usque ducto, lateralibus tenuioribus, nervulo tenui juxta marginem subciliolatum currente, amentis in axillis 4-6 pedunculo communi cymosæ confertis squamis subtus longe hirtis fultis, pedunculo hirsuto, bractea rotundata centro peltata, ovario subimmerso, stigmate minuto. ☉ In ins. Juan Fernandez (Bertero n. 1492! in h. DC.). — Miq. ill. p. 17. t. 11. Herba erecta, 0,3-0,4 alta. Caulis glaber, vix 0,01 crassus. Limbi majores 0,11-0,12, minores 0,04-0,05 longi. (v. s.)

110. *P. BAUERIANA* (Miq. ! syst. p. 120), foliis petiolatis alternis vel oppositis lanceolatis vel rhombeo-ellipticis apice acutiusculis obtusisve basi cuneatis vel e basi cuneata obovatis supra glabris subtus præsertim ad nervos pilosulis apicem versus ciliolatis membranaceis subopacis reticulato-nervulosis et 5-nerviis, centrali nervo ad apicem ducto lateralibus 2 alte ductis exterioribus tenuissimis, petiolo pilosulo, amentis filiformibus laxifloris, bractea rotundata centro peltata sessili, ovario impresso, bacca globosa impressa. ☉ In ins. Norfolk (Bauer ! in h. Endl., Endl. ! in h. Berol., Cunningham ! in h. Kew.). Piper adscendens Endl. ! prodr. Norf. p. 36. Herba inter muscos crescens 0,1 alta. Caulis basi radicans, pilosulus vel glaber. Limbi 0,03 longi, 0,015 lati. Petioli 0,01 longi. Nomen *adscendens* characteribus contrarium videtur. (v. s.)

♂, *Brisbantana*, foliis supra subtusque pubescentibus vel pilosulis subpellucidis, nervulo juxta marginem currente. ☉ Ad Brisbane flum. (Ferd. Müller ! in h. Kew.). Caulis dense pubescens. Petioli pubescentes. Limbi majores 0,04 longi, 0,027 lati. Petioli 0,005 longi. Pedunculi 0,01 longi. (v. s.)

111. *P. PURPURELLA*, foliis alternis vel apice caulium oppositis suboppositisve petiolatis subrhombico-lanceolatis utrinque acutis aut apice obtusiusculis vel apice caulium sterilium e basi cuneata subobovatis omnibus utrinque adpresse pubescentibus vivis subpurpurellis siccis membranaceis subpellucidis parce reticulato-nervulosis et 5-nerviis, centrali nervo ad apicem usque ducto lateralibus 2 supra medium 2 ad medium usque ductis, nervulo juxta marginem currente, petiolo dense adpresse pubescente, amentis axillaribus solitariis vel terminalibus sparsis vel apice ramulorum approximatis densifloris filiformibus florentibus folia æquantibus, bractea rotundata centro peltata subsessili, ovario impresso obovato, stigmate minuto. ♀ In Venezuela ad flumen Tuy infra coloniam Tovar alt. 5000 p. (Fendler n. 1161 ! in h. DC.). Suffrutescens basi radicans, parce ramulosus. Caulis ramulique dense adpresse puberuli. Limbi 0,06-0,075 longi, 0,025-0,035 lati. Petioli 0,015 longi. Amenta 0,001 crassa. (v. s.)

112. *P. OTTONIANA* (Miq. ! syst. p. 100), foliis alternis vel apice ramulorum oppositis petiolatis ellipticis utrinque subattenuatis acutiusculis utrinque glabris apicem versus ciliolatis membranaceis subpellucidis reticulato-nervulosis et 5-nerviis, centrali nervo ad apicem usque lateralibus ad medium ductis nervulo juxta marginem currente, petiolo glabro, amentis terminalibus densifloris filiformibus folia fere duplo superantibus filiformibus, pedunculo glabro petiolum duplo superante, bractea rotundata centro peltata brevissime pedicellata, ovario semiimmerso ovato, stigmate penicillato, bacca ovata semimmersa apice oblique mucronulata. ♀ In horto Berol. culta. Fructulus glaber, ramosus. Limbi 0,025 longi, 0,015 lati. Petioli 0,004 longi. (v. s.)

113. *P. PETIOLARIS* (Cas. DC. in Seem. journ. 1866. p. 138), foliis inferioribus longe superioribus modice petiolatis inferioribus e basi cuneata obovatis superioribus subovato-rhombicis apice acutiusculis obtusiusculisve et sæpe basi

- cordatis cordulatisve utrinque glabris membranaceis reticulato-nervulosis 5-nerviis, centrali nervo ad apicem usque lateralibus alte supra medium ductis exterioribus subtilioribus, petiolo glabro, amentis axillaribus terminalibusve solitariis filiformibus folia multoties superantibus subaxillaris tenuibus pedunculatis, pedunculo quam petiolos brevior, bractea rotundata centro peltata subsessili, ovario subovato post anthesin impresso, bacca ovata-acutiuscula. ☉ In Cuba (Wright n. 2261! in h. DC.), Costa-Ricæ sylvis (Hoffmann. n. 861 et 823! in h. Berol.). Herba 0,25 propemod. alta, basi radicans, glabra. Limbi 0,025 longi, 0,017 lati. Petioli 0,025 longi. (v. s.)
114. *P. ANGULARIS*, foliis petiolatis elliptico-lanceolatis apice breviter acuminate obtusiusculis basi subattenuatis infimis e basi cuneata obovatis vel rotundatis utrinque glabris membranaceis subopacis non reticulato-nervulosis 5-nerviis, centrali nervo ad apicem usque lateralibus subtilibus supra medium ductis exterioribus subtilissimis, petiolo glabro, amentis axillaribus solitariis filiformibus densifloris, pedunculo glabro petiolum multum superante, bractea rotundata centro peltata, ovario impresso, stigmatibus minuto. ☉ In Michoaca (Jameson! in h. Kew.). Caulis erectus, 0,2 altus, basi radicans, glaber, tetragenus. Limbi majores 0,045 longi, 0,025 lati. Pedunculi 0,025 longi. (v. s.)
115. *P. VENEZUELIANA* (Cas. DC. in Seem. journ. 1866 p. 139), foliis breviter petiolatis elliptico-lanceolatis apice acutiusculis mucronulatis basi acutis utrinque glabris vel apicem versus ciliolatis membranaceis subpellucidis parce pellucido-punctulatis utrinque crebre nigro-punctulatis nervis nervisque ægerrime discernendis 3-nerviis nervulorum juxta marginem apice currente, petiolo membranaceo nigro-punctulato ciliolato, amento terminali densifloro breviter pedunculato folia æquante, bractea rotundata centro peltata subsessili, ovario emerso, stigmatibus minuto paullo supra medium ovarii inserto, bacca emersa ovata nigra hirtella. ☉ In Venezuela prope coloniam Tovar alt. 6500 p. (Pendler n. 2618! in h. DC.). Herba procumbens. Caulis complanatus, 0,004 crassus, glaber, nigro-punctulatus. Limbi 0,035-0,04 longi, 0,02-0,025 lati. Petioli 0,008 longi. (v. s.)
116. *P. HIRCINA*, foliis petiolatis forma variantibus aliis elliptico-lanceolatis basi acutis apice imo obtusiusculis submarginulatis aliis rhombico-oblongis aliis rotundatis omnibus glabris apice ciliolatis submembranaceis pellucidis non pellucido-punctulatis parce reticulato-nervulosis 5-nerviis, centrali nervo subtus subprominulo ad apicem usque lateralibus supra medium ductis exterioribus subtilioribus, nervulo juxta marginem currente, petiolo glabro, amento terminali solitario glabro, bractea rotundata centro peltata, ovario immerso, stigmatibus carnosulo. ☉ In ins. Bourbon sylvis altis (Bory de St Vincent! in h. DC.). Limbi 0,04 longi. Petioli 0,008 longi. Pedunculi 0,015 longi. (v. s.)
117. *P. MOULMEINIANA* (Cas. DC. in Seem. journ. 1866. p. 140), foliis breviter petiolatis lanceolatis apice obtusiusculis basi acutis membranaceis pellucidis glabris reticulato-nervulosis et 5-nerviis, centrali nervo ad apicem usque lateralibus 2 alte ductis exterioribus tenuioribus brevioribus, nervulo tenui juxta marginem currente, amentis apice ramulorum subpaniculatum approximatis breviter pedunculatis glabris, bractea rotundata centro peltata, ovario impresso, stigmatibus minuto. ☉ In Moulmein (Parish n. 118! in h. Kew.). Herba repens vel scandens? glabra. Caulis subsulcatus, glaber, quadrangulus. Limbi 0,055 longi, 0,025 lati. Petioli 0,005 longi. Pedunculi 0,007-0,008 longi. (v. s.)
118. *P. NEMOROSA*, foliis petiolatis ovato-ellipticis subinæquilateralibus apice acuminate, acumine acuto obtusiusculove, basi acutiusculis æqualiter in petiolum decurrentibus utrinque glabris membranaceis subpellucidis pellucido-punctulatis sparse nigro-punctulatis atque reticulato-nervulosis 5-nerviis, nervulo juxta marginem currente, amentis axillaribus solitariis folia fere duplo superantibus, bractea rotundata centro peltata subsessili, bacca semiimmersa ovato-acuta. — In Jamaica, Antigua, Dominica (Griseb. flor. W. Ind. p. 165), Barbadoes (Maycock flor. Barb. p. 40), Guyana gallica (Aublét hist. pl. Guy. v. p. 1. p. 21). In hort. Berol. culta. Herb. Willd. n. 1211

- Saururus alius humilium Plum. Am. p. 540, 71. Piper acuminatum L. sp. p. 42, Aublet l. c., Pers. syn. p. 35. P. nemorosum Vahl enum. v. 1. p. 341. P. acuminata A. Dietr. sp. v. 1. p. 148 (exclus. syn. R. et P.), Miq.† syst. p. 95 (exclus. syn. R. et P.), Griseb. l. c. (excl. syn. P. Hamiltoniana Miq. et obliqua R. et P.). Caulis erectus, glaber, tetragonus. Limbi 0,07 longi, 0,035 lati. Petioli 0,012 longi. (v. s.)
119. P. HIRTELLA (Miq.† in Hook. Lond. journ. bot. v. 4. p. 414), foliis petiolatis ovatis ellipticisve acuminatis vel acutis, acumine obtusiusculo, basi obtusis vel rotundatis utrinque adpresse puberulis subtusque glanduloso-punctulatis 3-5-nerviis, petiolo adpresse hirtello, amentis axillaribus solitariis pedunculatis, subdensifloris quam folium plerumque brevioribus, pedunculo petiolum parum superante. ☉ In Dominica (Imray n. 244! in h. Hook.). — Griseb. flor. w. Ind. p. 165. Planta pedalis, inferne crebre radicans decumbens. Limbi 0,05 longi, 0,028-0,026 lati. Petioli 0,002-0,008 longi. (v. s.)
120. P. RUSCIFOLIA (Kunth! in H. et B. nov. gen. v. 1. p. 63), foliis breviter petiolatis elliptico-ovatis apice acuminatis, acumine acuto, basi rotundatis vel acutiusculis utrinque glabris coriaceis 5-nerviis, petiolo glabro, amento oppositifolio? folium æquante vel terminali solitario. ☿ In locis calidissimis, in ripa fluminis Manzanares, prope Cumana (Kunth l. c. et Humb. in h. Willd. n. 718!), Brasilia (Weddell! in h. DC.). — Kunth syn. v. 1. p. 116, A. Dietr. sp. v. 1. p. 155, Miq.† syst. p. 93. Piper ruscifolium Roem. et Sch. syst. v. 1. p. 326. Caulis glaber, sublignosus. Limbi 0,035 longi, 0,015-0,016 lati. Petioli 0,003 longi. (v. s.)
121. P. GUADALOUPENSIS (Cas. DC. in Seem. journ. 1866. p. 139), foliis breviter petiolatis inferioribus cuneato-obovatis superioribus obovato-ellipticis basi obtusiusculis subattenuatis basi subcuneato-acutis utrinque glabris rigidulis subpellucidis pellucido-punctulatis sparse reticulato-nervulosis et 5-nerviis vel raro 7-nerviis, centrali nervo ad apicem usque ducto, lateralibus alte ductis exterioribus sæpe obsoletis, nervulo juxta marginem currente, petiolo glabro, amentis terminalibus filiformibus densifloris folia superantibus, rhachi puberula, bractea rotundata centro peltata subsessili ovario impresso ovato, stigmate minuto, bacca globosa immersa apice submucronulata. ☿ In ins. Guadeloupe (Jardin n. 340! in h. Lenorm. et DC.), Cuba (Wright n. 504! in h. DC.), Ecuador (Fraser! in h. DC.), Sta Crux (Springard! in h. DC.). Pep. acuminata Griseb.† Cat. pl. Cub. p. 63. P. dendrophila Griseb.† Cat. pl. Cub. p. 64, quoad n. 1888. Glabra. Caules subquadranguli. Limbi 0,03 longi, 0,025 lati. Petioli 0,003 longi. (v. s.)
- β, pubescens, foliis supra glabris subtus parce pilosis rigidulis subopacis. In Cuba orientali prope villam Monte Verde dictam (Wright n. 511!, 1688! in h. DC.). Caulis apicem versus alterne uno latere pilosulo. Limbi 0,035-0,05 longi, 0,015-0,02 lati. (v. s.)
122. P. WALLICHII (Miq.† syst. p. 82), foliis longe petiolatis ovato-rotundatis æquilateris basi cordatis truncatisve, glabris junioribus ad imam basin pilis microscopicis citissime delitescens instructis 5-nerviis et reticulato-nervulosis, amento terminali solitario breviter pedunculato crasso subulxifloro, bractea rotundata centro peltata, stigmate puberulo, bacca globosa atra. ☉ In India Orientali (Wall. n. 6665! in h. Wall.). Piper basellæfolium Wall. list n. 6665. Planta succulentissima, membranacea, sicca pellucida, tota sordide fuscescens. Caulis glaber. Limbi 0,025-0,035 longi, 0,023-0,033 lati. Petioli 0,015-0,02 longi. (v. s.)
123. P. RUFESCENS, foliis longe petiolatis late ellipticis apice obtusis basi cordatis membranaceis pellucidis pellucido-punctatis reticulato-nervulosis et 5-nerviis, nervis subtus prominulis, nervulo interrupto juxta marginem currente, amentis axillaribus? vel oppositifoliis?, bractea rotundata centro peltata, ovario emerso fusco-ruguloso, stigmate minuto. ☉ In Peruvia (Pav. n. 277! in h. Boiss.). Herba decumbens. Limbi 0,03-0,035 longi. Petioli 0,02-0,05 longi. (v. s.)
124. P. METAPALCOENSIS, foliis petiolatis obovatis vel obovato-spathulatis rarisime subrotundis utrinque hirtis subpellucidis fusco-punctulatis 3-5-nerviis,



nervulo juxta marginem currente, petiolo hirtio, bractea rotundata centro peltata subsessili, ovario immerso, bacca ovata apice oblique breviter submucronulata. ☉ In regione calida Mexici inter Metapalco et Cuapa (Schiede! in h. Berol.), Guyana (Leprieur! in h. Francov.), Equador (Fraser! in h. DC.), Cuba (Wright n. 2262! in h. DC.). *Acrocarpidium mexicanum* Miq.! syst. p. 53. *Pep. Miqueliana* Griseb. cat. pl. Cub. p. 63 (non C. DC.). *Herbula repens dichotome ramosa*. Caulis ramique longe setuloso-hirti. Limbi 0,005-0,01 longi, 0,004 lati. Petioli 0,003 longi. (v. s.)

125. *P. PURPURASCENS*, foliis caulinis longe petiolatis rhombeis apice obtusiusculis basi cuneatis vel elliptico-rhombeis vel obovatis supra glabris subtus pubescentibus membranaceis subpellucidis reticulato-nervulosis et 3-5-nerviis, centrali nervo ad apicem usque lateralibus nervis alte ductis omnibus subtilibus, petiolo hirtio, amentis terminalibus, bractea rotundata centro peltata subsessili, ovario immerso, stigmatibus minuto, bacca ovata apice acuta, amentis filiformibus densifloris folia triplo superantibus. ¼ In ins. Sandwich (Nutt.! in h. Brit. Mus. sub nom. *Pep. purpurascens*). Ramuli hirtio-pubescentes. Limbi 0,04 longi, 0,02 lati. Petioli caulini usque ad 0,03 longi. (v. s.)

126. *P. GLABELLA* (A. Dietr. sp. v. 1. p. 156), foliis petiolatis ellipticis apice breviter acuminatis, acumine acutiusculo, basi acutiusculis utrinque glabris crebre nigro-punctulatis ciliolatis rigidulis opacis 5-nerviis, centrali nervo validiore ad apicem usque ducto, petiolo superne ciliolato, amentis ramulos terminantibus densifloris solitariis, pedunculo glabro petiolum æquante, bractea subrotundata centro peltata subsessili nigro-punctulata, ovario immerso apice subacutato, bacca ovato-subglobosa basi immersa. ☉ In Jamaicae montosis lapidosi (Swartz!) in h. DC.). — A. Dietr. syst. p. 156. *Piper glabellum* Sw.! prodr. p. 16, flor. Ind. Occ. v. 1. p. 68, Vahl enum. v. 1. p. 345, Willd. sp. v. 1. p. 165. *P. scandens* Sw. prodr. p. 16. *Herba e basi radicans*. Ramuli uno latere biserialiter ciliolati. Limbi 0,02 longi, 0,01 lati. Petioli 0,006 longi. (v. s.)

127. *P. TRINERVIS* (Ruiz et Pav. p. 32. t. 50. fig. b), foliis alternis vel apice ramulorum oppositis petiolatis plerumque elliptico-lanceolatis apice acuminatis, acumine acuto, basi acutis inferioribus basi rotundatis omnibus supra ad nervos pubescentibus ciliolatis coriaceo-pellucidis utrinque crebre nigro punctatis reticulato-nervulosis 3-5-nerviis, nervis lateralibus tenuissimis, petiolo pubescente, amentis axillaribus terminalibusve densifloris folia duplo triplove superantibus, bractea rotundata centro peltata, ovario impresso, stigmatibus discoideo minuto. ☉ In Peruvia (Pav.! in h. Boiss.), Brasilia (Steven! in h. DC.), prov. Rio-Janeiro rupibus, humidis sylvarum (Mart.! in h. monac.), prope Mandioccam (Mart.! ibid.), Venezuela ad Maya alt. 3000 (Fendler n. 1162! in h. DC.), Nova Hispania (Pav.! in h. Boiss.), h. Willd. n. 724! hort. Berol. cult. sub nom. *Pep. leuconeura*. — A. Dietr. sp. v. 1. p. 149. Miq.! syst. p. 94, ill. p. 13. t. 8. fig. a. *Piper trinervum* Vahl enum. 1. p. 342. *Pep. caulibarbis* var. *pilosior* Miq.! in Mart. flor. Bras. fasc. 11. p. 12 et in Linn. v. 20. p. 123 et in h. monac. *Pep. punctulata* Regel! in Bull. Acad. Mosc. 1858. p. 542 et h. Pe-trop. *Herba basi decumbens radicans*. Caules 0,2 alti, inferne aphylli, nigro-punctulati. Ramuli pubescentes. Limbi 0,03-0,035 longi. Petioli 0,01-0,015 longi. Pedunculi 0,01 longi. (v. s.)

β, *brachyphylla*, foliis elliptico-ovatis vel rotundato-ovatis inferioribus ovatis omnibus apice breviter acuminatis, acumine obtuso acutiusculove, utrinque glabris apicem versus ciliolatis utrinque crebre nigro-punctulatis obsolete nervulosis, petiolo glabro ciliatove nigro-punctulato, pedunculo petiolum paullo superante glabro nigro-punctulato, rhachi nigro-punctulata, amentis folia multoties superantibus. ☉ In Brasilia ad Petropolim (Wawra et Maly n. 437! fide Wawra *Ergeb.* p. 132. t. 82. fig. 2), Cuba (Wright n. 506! in h. DC.), saxosis montosis ins. Antioea (Wulfschlegel n. 537 b! et 539! in h. monac.), hort. Berol. cult. (h. Berol.! et h. Willd. n. 722!). — *Pep. brachyphyllum* A. Dietr. sp. v. 1. p. 157, Miq.! syst. p. 99. *Piper brachyphyllum* Spreng. syst. v. 1. p. 115. *Herba semipedalis, succulenta, basi procumbens, in cortice arborum parasitica*. Limbi 0,04 longi, 0,02 lati. Petioli 0,005 longi. (v. s.)

- γ. minor*. Ramuli glabri. Limbi 0,016-0,02 longi, 0,006 lati. ☉ In Brasilia ad Minas Geraes inter muscos et ad arborum truncos (Weddell n. 781 in h. DC. et Lhotsky! ibid.), (v. s.)
128. *P. FALLIDA* (A. Dietr. sp. γ. 1. p. 153), foliis petiolatis obovato-rhombicis vel elliptico-rhombicis apice obtusiusculis basi cuneatis subcoriaceis utrinque glabris 3-5-nerviis nervis 3 centralibus ad apicem usque ductis exterioribus subtilioribus sæpe abortivis, petiolo glabro, amentis axillaribus filiformibus densifloris, pedunculo glabro petiolum paullo superante, bractea rotunda centro peltata stigmatibus minuto, bacca semi-immersa ovata. ☉ In Otahiti (Forster! in h. Brit. Mus., Sidney et Park. icon. t. 8. in Brit. Mus. ubi *P. polymorphum* dicta), Viti (Seem. n. 565! in h. Brit. Mus.), Sandwich (Hook. et Arn. in Beach, voy. p. 96). — Miq. syst. p. 103, Hook. et Arn. l. c. *P. pallidum* Forster! prodr. n. 24, Willd. sp. v. 1, p. 165. Caulis glaber. Limbi 0,06 longi, 0,02 lati. Petioli 0,005 longi. Amenta 0,05 longa. Pedunculi 0,015 longi. (v. s.)
129. *P. ALATA* (Ruiz et Pav.! Flor. Peruv. v. 1. p. 31. t. 48. fig. b), foliis breviter petiolatis oblongo-ellipticis apice acuminatis basi acutiusculis in petiolum decurrentibus membranaceis pellucido-punctulatis utrinque glabris apice ciliolatis reticulato-nervulosis et 3-5-nerviis, centrali nervo validiore, nervis 2 centralibus interdum paullo supra basin solutis, nervulo tenui juxta marginem currente, petiolo glabro, amentis axillaribus terminalibusve solitariis globosis breviter pedunculatis densifloris maturis folia multoties superantibus, bractea subrotunda centro peltata, ovario immerso postea semi-immerso, stigmatibus punctiformi. ☉ In Peruvia nemoribus ad Pozuzo (R. et P. l. c. et Pav. in h. Boiss., Ruiz! in h. Berol.), Andibus (Spruce n. 4576! in h. DC.), Peruvia subandina (Poeppig! in h. Berol.), Nova Granata (Holton n. 232! in h. Boiss.), Venezuela (Fendler n. 1150! 1160! 2403! in h. DC.), Mexico (F. Müller n. 246! in h. Kew., Andreux n. 97! in h. DC.), Cuba (Wright n. 593! et 500! in h. DC.), Santo Domingo (Ham. prodr. p. 2). — A. Dietr. sp. v. 1. p. 149. Miq.! syst. p. 87. *Pep. dendrophila* Griseb. cat. pl. Cub. p. 64 exclus. n. 1688. *P. alatum* Vahl enum. v. 1. p. 342. Planta basi radicans, 0,25 alta. Caulis glaber, anguste tetrapterus. Limbi 0,06-0,02 longi. Petioli 0,005 longi. Pedunculi vix 0,005 longi. Amenta post anthesin 0,11-0,12 longa. (v. s.)
- β, *pterocaulis*, foliis breviter petiolatis latioribus brevioribusque. In Trinidade (Fischer! in h. DC. et h. Petrop.), Brasiliæ monte Cercovado et prov. Rio Janeiro (Mart.! in h. monac.), Santo Domingo (Poiteau! in h. Deless.), *Pep. pterocaulis* Miq.! syst. p. 86. *P. Velloziana* Miq.! in Linn. v. 20 p. 122, et in Mart. flor. Bras. fasc. 11. p. 11.
- γ, *angustifolia*, foliis paullo angustioribus magis acuminatis nervis subtilioribus. In Cuba (Wright n. 505! in h. DC.).
130. *P. FOLIOSA* (Kunth! in Nov. gen. v. 1. p. 63), foliis petiolatis plerumque subbinatis vel subalterne confertis ovatis ellipticisve apice obtusiusculis basi rotundatis utrinque glabris membranaceis subpellucidis obsolete nervulosis 3-nerviis vel obsolete 5-nerviis nervuleque tenui apice juxta marginem currente, petiolo glabro, amentis axillaribus terminalibusve densifloris folia superantibus filiformibus, bractea rotunda centro peltata brevissime pedicellata, ovario emerso. ☉ In vexis frigidis montium Quindiu inter El Moral et La Palmilla alt. 1150 hexap. (Kunth l. c. et Humb. in h. Willd. n. 723!), Peruvia subandina (Poeppig! in h. Berol.). — Kunth synops. v. 1. p. 116, A. Dietr. sp. v. 1. p. 155, Miq.! syst. p. 111. *P. foliosum* Roem. et Sch. syst. v. 1. p. 326. Herba succulenta, erecta, vix spithamea, viridis, basi simplex, sursum conferte ramulosa. Limbi 0,01 longi, 0,007 lati. Petioli 0,002 longi. (v. s.)
131. *P. ACUMINATISSIMA* (Miq.! in Hook. Lond. journ. bot. v. 4. p. 416), foliis petiolatis ovato-ellipticis summis elliptico-lanceolatis æquilatis apice longe acuminatis, acumine angusto acuto juniore ciliolato, utrinque glabris membranaceis atro-viridibus subtus pallidis 3-5-nerviis, nervis 3 centralibus subtus prominentibus exterioribus tenuibus submarginatis, amentis axillaribus solitariis densifloris folia superantibus. ☉ In prov. Minas Geraes (Gardner n. 5187! in h. Kew.). — Miq.! in Mart. Flor. Bras. fasc. 11. p. 12. Caulis pedalis, parce ramosus, succulentus, angulatus (an fere alatus), glaber. Limbi 0,045 longi,

0,026 lati, superiores 0,08-0,1 longi, 0,03-0,02 lati. Petioli 0,003-0,005 longi. Amenta 0,06-0,1 longa. (v. s.)

132. *P. BILINEATA* (Miq.) Syst. p. 106 et Fl. Ind. bat. 1. pars 2. p. 433), foliis petiolatis retundato-ovatis apice breviter acuminatis, acuminē obtusiusculo, basi rotundatis utrinque glabris membranaceis pellucidis laxè reticulato-nervulosis et 3-5-nerviis, centrali nervo ad apicem usque lateralibus 2 alte supra medium exterioribus subtilioribus ad medium usque ductis nervulo tenui apice juxta marginem currente, petiolo glabro, amentis terminalibus folia æquantibus filiformibus subdensifloris, bractea subrotundata centro peltata subsessili, ovario emerso vel subimpresso ovato, stigmatè minuto. 2 In Java monte Salak locis humidis super arborum truncos prostratos (Bl. in h. Berol.), Manilla (Miq. in nov. act. v. 19. p. 485). *P. bilineatum* Bl. in Verh. der Bat. Genoots. v. 11. p. 231, enum. pl. Jav. fasc. 1. p. 75. *Micropiper bilineatum* Miq. in comp. phyt. p. 59. t. 8. fig. B. Caulis succulentus, glaber. Limbi 0,03 longi, 0,025 lati. Petioli 0,008 longi. (v. s.)
133. *P. KUHLIANA* (Miq. syst. p. 104), foliis petiolatis ellipticis ovato-vel rotundato-ellipticis apice subattenuato-obtusis rotundatisve basi rotundatis glabris junioribus apicem versus subtilissime ciliolatis membranaceis subpellucidis reticulato-nervulosis et 3-5-nerviis, centrali nervo ad apicem usque ducto, lateralibus 2 supra medium ductis exterioribus subtilioribus sepe obsolete nervuloque juxta marginem apice currente, petiolo glabro, pedunculo glabro petiolum superante, amentis axillaribus filiformibus folia quintuplo superantibus subaxillifloris, bractea rotundata centro peltata subsessili, ovario semi-immerso ovato, stigmatè discoideo puberulo, bacca ovato-globosa semi-immersa glabra nigra. 2 In ins. Java (Schmidt in h. Berol.). Planta succulenta, ramosa, decumbens, 0,25-0,3 alta. Caulis ramique glabri. Limbi 0,01-0,015 longi, 0,01 lati. Petioli 0,005 longi. (v. s.)
134. *P. SERPENS* (Cas. DC. in Seem. journ. bot. 1866. p. 136), foliis brevissime petiolatis ovatis vel inferioribus ovato-rotundis obtusis utrinque hirsutis rigidulis opacis, petiolo hirsuto, amentis terminalibus solitariis pedunculatis folia multoties superantibus hirtellis, bractea rotundata centro peltata subsessili ovario immerso subobovato, stigmatè carnosulo. ☉ In Nova Grapata prov. Barbacoas ad viam Tuquerras alt. 6000 m. (Triana n. 581 in h. DC.). Herba repens e nodis radicans. Caulis filiformis, subnigrosus, vix 0,001 crassus, hirtellus. Limbi 0,003-0,005 longi, 0,003-0,004 lati. Petioli 0,001 longi. (v. s.)
135. *P. CANDIDA* (Miq.) syst. p. 105), foliis plerumque alternis petiolatis ovalibus vel obovatis apice obtusis basi cuneatis supra glabris subtus puberulis ciliolatis rigidulis vel membranaceis opacis subopacisve utrinque fuscescentibus subtus nigro-punctulatis 2-5-nerviis, nervis haud conspicuis, nervulo juxta marginem currente, petiolo adpresse puberulo, amentis axillaribus terminalibusque folia duplo superantibus subdensifloris, pedunculo petiolum fere duplo superante, bractea rotundata centro peltata subsessili, bacca impressa subglobosa. 2 In ins. Java (Bl. in h. DC. ex h. Lugd. bat.), monte Salak ad ripas cataractæ fluvii Tjapus super arborum truncos valuosos et saxa basaltica (Bl. Zolling. n. 1502 in h. Francov.), Borneo austr. (Miq. in Ann. bot. Mus. Ludg. bat. v. 1. fasc. 5 p. 130), Sumatra (Miq. l. c. p. 138). — *Miq. flor. Ind. bat. v. 1. pars 2. p. 433*, Linn. v. 21. p. 460. *P. candidum* Bl. in Verh. der Bat. Genoots. v. 11. p. 227. fig. 32, enum. pl. Jav. fasc. 1. p. 74, enum. hort. Bogor. p. 33. *Micropiper candidum* Miq. comp. phyt. p. 58. t. 8. fig. D. *Spathæa*, suberecta, basi radicans. Rami superne pubescentes. (v. s.)
136. *P. LEVIFOLIA* (Miq.) syst. p. 107), foliis breviter petiolatis inferioribus cuneato-obovatis superioribus lanceolatis vel ovato-oblongis apice obtusiusculis subtilissime subserrulatis glabris subciliolatis pellucido-punctatis 5-nerviis, nervis lateralibus ante medium evanescentibus, nervulo juxta marginem currente, amentis axillaribus geminatis vel terminalibus 4-approximatis breviter pedunculatis subaxillifloris, bractea subrotundata centro peltata subsessili, ovario post anthesin paullo immerso, stigmatè minuto puberulo. 2 In ins. Java monte Salak super arborum truncos et ad ripas fluvii Tjapus (Bl. in h. DC. ex h. Lugd. Bat.), ins. Java (Zolling. n. 1789 in h. DC., 1960 fide Miq. in Zoll. cat. p. 83), sylvis montis Merapi ins. Java alt. 3-4000 ped. (Miq. pl.

Jungh. fasc. 1. p. 14). — Miq. flor. Ind. Bat. v. 1. pars 2. p. 434, in Linn. v. 21. p. 480. *P. lævifolium* Bl. in Verh. Bat. Genoots v. 9. p. 200, v. 11. p. 229, et Enum. pl. Jav. fasc. 1. p. 74. *Micropiper lævifolium* Miq. comm. phyt. p. 56. t. 9. fig. c. Glaberrima. Limbi 0,03-0,05 longi, 0,02-0,03 lati. Petioli 0,005 longi. (v. s.)

β, *suboppositifolia*, foliis sæpe oppositis. In ins. Java monte Pangerango, prope Hariang (Miq. in Ann. bot. Mus. Lugd. Bat. v. 1. fasc. 5. p. 135).

γ, *parvifolia*, foliis omnibus duplo minoribus. In ins. Java prope Manara (Miq. l. c.).

137. *P. CONVEXA* (Miq. syst. p. 107), foliis petiolatis rotundatis vel ovalibus utrinque obtusis inferioribus obovatis basi cuneatis superioribus oblongo-rhomboides subacuminatis apice subtilissime subserulatis supra glabris subtus pubescentibus membranaceo-rigidulis parce venulosis et 3-nerviis, centrāli nervo ad apicem usque lateralibus nervis supra medium ductis, petiolo ciliolato, amentis apice caulis approximatis filiformibus folia duplo superantibus densifloris, pedunculo petiolum paullo superante glabro, bractea rotundata centro peltata subsessili, ovario ovato postea immerso, stigmate minuto, bacca exigua globosa. ♂ In ins. Java monte Salak super arborum truncos et ad ripas fluvii Tjapus (Bl. in Verh. der Bat. Genoots v. 11. p. 236 et in h. Lugd. Bat. fide Miq. l. c.), ins. Java ad rupes montis Batu Lanke alt. 5000 (Zolling. l. n. 1502? in h. Francov.). — Miq. flor. Ind. Bat. v. 1. pars 2. p. 433. *P. convexum* Bl. l. c., enum. pl. Jav. fasc. 1. p. 76 (except. var.). *Micropiper convexum* Miq. comm. phyt. p. 60. t. 8. fig. e. Caulis basi repens, pubescens. Limbi 0,012 longi, 0,006 lati. Petioli 0,003 longi. (v. s.)

138. *P. WIGHTIANA* (Miq. in Hook. Lond. journ. bot. v. 5. p. 548), foliis alternis vel superioribus oppositis petiolatis inferioribus rotundato-obovatis apice brevissime acutatis basi rotundatis superioribus obovato-ellipticis ellipticisve apice brevissime acutatis basi acutis glabris pellucido-punctatis apice ciliolatis 3-nerviis, centrāli nervo validiore lateralibus subtilibus, amentis axillaribus terminalibusque folia duplo superantibus subaxillifloris, ovario rhachi impresso ovato apice acutato. ☉ In Malabar (Wight ic. t. 1924). — Miq. in Wight ic. l. c. Herba succulenta, erecta, basi radicans. Caulis pubescens. Limbi inferiores 0,022 longi, 0,016 lati. Petioli 0,006 longi. (v. s.)

139. *P. TRINERVULA*, foliis petiolatis ovatis apice breviter attenuatis acutis vel inferioribus ellipticis apice obtusis infimis elliptico-rotundatis utrinque pilosopubescentibus supra demum glabris membranaceo-firmis opacis 3-nerviis, centrāli nervo ad apicem usque ducto, lateralibus nervis subtilibus, petiolo pubescente, amentis axillaribus terminalibusque solitariis vel apice ramulorum approximatis subdensifloris filiformibus folia paullo superantibus, pedunculo petiolum superante pilosulo, bractea elliptico-rotundata centro peltata, ovario emerso apice obtuso, stigmate minuto. ♀ In Venezuela prope coloniam Tovar (Fendler n. 1167! in h. DC. et Moritz! in h. Francov.), Caracas (Gallmer n. 332! in h. Berol.). Pep. purpurascens Moritz! mss. in h. Francov. Caulis pubescens, basi procumbens, radicans. Limbi vix 0,03 longi, 0,02 lati. Petioli 0,007-0,01 longi. (v. s.)

140. *P. MORITZII*, foliis petiolatis subrhombeis apice acutiusculis basi acutis supra glabris subtus ad nervos pubescentibus rigidis subopacis 3-nerviis, petiolo pubescente, amentis axillaribus terminalibusque florentibus quam folia triplo longioribus densifloris, pedunculo petiolum æquante. ♀ In Venezuela (Moritz n. 1865! in h. Brit. Mus.). Ramuli pubescentes. Limbi 0,02 longi, 0,015 lati. Petioli 0,008 longi. (v. s.)

141. *P. NUMMULARIFOLIA* (Kunth! in H. et B. nov. gen. v. 1. p. 66), foliis petiolatis ovato-orbicularibus vel orbicularibus utrinque tenuiter setulosis pellucido-membranaceis sæpe subtus nigro-punctulatis enerviis vel 3-nerviis, amentis terminalibus brevissime pedunculatis subdensifloris, bractea rotundata centro peltata, ovario impresso obtuso, stigmate minuto. ④ In Brasilia super montes Orgaos (Lhotzky! in h. DC., Sello! in h. Berol.), Ilheos (Wawra et Maly n. 385 fide Wawra Ergeb. bot. p. 133), prope Saigado prov. Minarum (h. monac.!), prope San Gabriel (Spruce n. 2399! in h. DC.), Suri-

nam (Hostman n. 117! in h. Berol., Weigelt! in h. DC.), Guyana gallica (Aublet hist. pl. Guy. v. 1. p. 21), sylvis maximis prope Mariepaston (Kegel n. 1357 fide Miq. in Linn. v. 22. p. 76), Venezuela ad Valencia et Campanero alt. 6000 p. (Fendler n. 2404! in h. DC.), fissuris rupium prope cænobium Caripense alt. 420 hexap. (Humb. in nov. gen. v. 1. p. 65 sub nomine *Pep. rotundifolia*), ad flumen Magdalenæ juxta San Bartolomeo (Humb. in h. Willd. n. 733! et 731! spec. super.), San Carlos (Spruce n. 3086! in h. Francov.), Nova Granata (Holton n. 229! in h. Boiss.), ins. Trinitatis (Sieber n. 16! in h. Berol.), Sta Lucia (Schreber in Linn. v. 20. p. 117), Guadaloupa (Bertero! in h. DC., Balbis! in h. Berol.), Jamaica (Sagra! in h. Kew., Griseb. flor. w. Ind. p. 164), Cuba (Griseb. l. c., Wright n. 4181, 5201, 521! in h. Kew., Sagra! Sant. s! in h. DC.), Panama (Seemann n. 898 fide Miq. in bot. Herald. p. 198), Mexico (Schiede! in h. Berol.). — Kunth syn. v. 1. p. 118, A. Dietr. sp. v. 1. p. 160, Miq.! in Mart. flor. Bras. fasc. 11. p. 18; Griseb. flor. w. Ind. p. 164, Cat. pl. Cub. p. 63. *Piper* longum minimum herbaceum scandens Sloane hist. p. 136 et in Coll. orig. icon. t. 90. *Saururus repens* Plum. Am. ed. Par. p. 52. t. 69. *P. rotundifolium* L.! sp. p. 42 et herb., Aublet l. c., Lam. ill. p. 82, Sw. prodr. p. 16, obs. p. 20, flor. Ind. occ. v. 1. p. 72, Vahl enum. v. 1. p. 348, Willd. sp. v. 1. p. 166, Spreng. syst. p. 118. *Pep. rotundifolia* Kunth in nov. gen. v. 1. p. 65, Griseb.! flor. w. Ind. p. 164. *Acrocarpidium nummularifolium* Miq.! syst. p. 52. *Acrocarpidium rotundifolium* Miq. syst. p. 62 (excl. spec. Willd.) et in Hook. Lond. journ. bot. p. 411 et in Linn. v. 20. p. 117. *Herbula repens*. Caulis filiformis, valde debilis, crispulus. Limbi 0,006-0,01 longi. Petioli 0,001-0,003 longi. (v. s.)

β, *pubescens*, caule dense piloso-pubescente. ④ In umbrosis montibus supra saxa et arborum truncos alt. 5000 ped. in Andibus Quitensibus (Jameson n. 244! et 430! in h. DC.), prope Xalapa (Miq. in Linn. v. 18. p. 710). (v. s.)

γ, *obcordata* (Miq.! in Linn. v. 20. p. 118), foliis plerumque obovatis inferioribus obcordatis vel apice emarginatis. ④ In Brasilia (h. Monac.). (v. s.)

142. *P. PARASITICA*, foliis petiolatis rotundatis vel rotundato-ovatis utrinque puberulis subcoriaceis subopacis non reticulato-nervulosis 3-nerviis nervo centrali validiore, nervulo obscuro juxta marginem currente, petiolo puberulo, amentis axillaribus solitariis brevibus, pedunculo puberulo, bractea rotundata centro peltata, ovario immerso, stigmatibus discoideo-puberulo. ☉ In Pichinchæ prov. alt. 10 000 ped. ad arborum truncos (Jameson n. 721! in h. Boiss.). Caulis erectus, puberulus. Limbi 0,013 longi. Petioli 0,008 longi. Pedunculi 0,01 longi. (v. s.)

143. *P. CORDIFOLIA* (A. Dietr. sp. v. 1. p. 154), foliis breviter petiolatis e basi attenuata obovatis apice emarginatis vel obcordatis supra puberulis membranaceis pellucidis remote nervulosis tenuiter 3-nerviis, amentis terminalibus breviter pedunculatis, bractea rotundata centro peltata, ovario emerso, stigmatibus minuto. ☉ In Dominica (Imray! in h. Kew.), Jamaica (Sw. in h. Willd. n. 730! Wilson! in h. Kew., Bertero! in h. Berol., Griseb. flor. w. Ind. p. 164). *P. cordifolium* Sw. prodr. p. 16, flor. Ind. occ. v. 1. p. 71, Vahl enum. v. 1. p. 345, Willd.! sp. v. 1. p. 165. *Acrocarpidium cordifolium* Miq.! syst. p. 56, in Hook. Lond. journ. bot. v. 4. p. 411, in Linn. v. 20. p. 118. *Herba repens*. Caulis glaber. Limbi 0,011 longi. Petioli 0,007 longi. Pedunculi 0,005 longi. (v. s.)

144. *P. ESTRELLENSIS*, foliis petiolatis elliptico-lanceolatis subrhombeis apice obtusis basi rotundatis vel (in eodem specimine) acutis utrinque glabris vel apicem versus subtilissime pilosulis membranaceis subpellucidis pellucido-punctulatis 3-nerviis obsolete nervulosis nervulo interrupto juxta marginem currente, petiolo glabro juniore subtilissime puberulo, amentis terminalibus longe pedunculatis folia duplo triplo superantibus densifloris, pedunculo petiolum multoties superante subtilissime puberulo, bractea rotundata centro peltata subsessili glanduloso-punctulata, ovario oblongo acuto, stigmatibus punctiformi. ☉ In Brasiliæ variis locis ad Mantiquira (Sello! in h. Berol.), monte Corcovado et ad Mandioccam (Mart. fide Miq. in Linn. v. 20. p. 122),

Serra d'Estrella Parasalsa (Pohl n. 4702! in h. Vindeb.), hort. Petrop. cult. (Fisch. ! in h. Dö.). *Pep. myrtifolia* A. Dietr. sp. v. 1. p. 147?, Miq. ! in Lih. l. c. t. 1. fig. 2, syst. p. 92 et in Mart. flor. Bras. fasc. 11. p. 11. t. 2. fig. 7. Planta longa repens. Caulis ramique glabri subterragani. Limbi 0,013 longi, 0,007 lati. Petioli 0,066 longi. (v. s.)

145. *P. MANNI* (Hook. fil. ! ined. in h. Kew.), foliis petiolatis ellipticis basi acutis apice obtusis vel inferioribus obovatis obovato-rhombicisve glabris ciliatis membranaceis pellucidis reticulato-nervulosis subtiliter 3-nerviis nervulo juxta marginem apice currente; petiolo glabro, amentis terminalibus solitariis filiformibus, pedunculo glabro, bractea rotundata centro peltata, ovario impresso, stigmate punctiformi. ☉ In Fernando-Po (Mann n. 2016! in h. Kew.). Herbulis glabra gracilis erecta. Caulis filiformis; basi radicans, 0,1 alta. Limbi 0,01 longi, 0,007 lati. Petioli 0,007 longi. (v. s.)

♂; *Fernandopana*, foliis superioribus elliptico-orbicularibus. ☉ In Fernando-Po (Mann n. 291! in h. Kew.). (v. s.)

146. *P. HAMILTONIANA* (Miq. ! in Hook. Lond. Journ. bot. v. 4. p. 417), foliis brevissime petiolatis ellipticis apice subattenuatis obtusiusculis basi acutis in petiolum subdecurrentibus membranaceis subopacis 3-nerviis; centrali nervo ad apicem usque ducto, nervis lateralibus 2 fere ad apicem usque ductis, petiolo glabro, amentis terminalibus filiformibus densifloris breviter pedunculatis, bractea rotundata centro peltata, ovario subacuminato, stigmate minuto. ♂ ? In ins. Jamaica (Fadyen ! et Purdie n. 108! in h. Kew.). Ramuli glabri. Limbi 0,035 longi, 0,017 lati. Petioli 0,003 longi. Pedunculi 0,005-0,007 longi. Amenta florentia 0,06-0,07 longa. (v. s.)

147. *P. OAHUENSIS*, foliis petiolatis oblongo-ellipticis apice acutis basi cuneatis rigidulis glabris subopacis reticulato-nervulosis 3-nerviis?, petiolo glabro, amentis axillaribus densifloris filiformibus pedunculatis. ♂ In ins. Oahu (Beechey ! in h. Kew.). — *Pep. pallida* A. Dietr. ? sp. v. 1. p. 153. Ramuli glabri. Limbi 0,025 longi, 0,007-0,008 lati. Petioli 0,008 longi. Pedunculi 0,008 longi. (v. s.)

148. *P. MONTICOLA* (Hook. fil. ! mss. in h. K.), foliis longiusculis petiolatis rhombico-ellipticis apice obtusiusculis basi acutis glabris membranaceis pellucidis irregulariter reticulato-nervulosis 3-nerviis; centrali nervo validiore ad apicem usque lateralibus ad medium ductis nervulo juxta marginem currente, amentis axillaribus terminalibusve solitariis, pedunculo glabro, bractea rotundata centro peltata, ovario impresso, stigmate minuto. ☉ In Camerobon montibus alt. 8000 ped. (Mann n. 1303! in h. Kew.). Herba erecta. Caulis glaber, basi radicans, 0,3 altus. Limbi maj. 0,045 longi, 0,025 lati. Petioli 0,015 longi. Pedunculi 0,015 longi. (v. s.)

149. *P. CORCOVADENSIS* (Gardner ! in Hook. Lond. Journ. bot. 1842. p. 187), foliis petiolatis ovato-ellipticis obtusis carnosio-coriaceis 3-nerviis, amentis terminalibus solitariis. ☉ In rupibus umbratis prope aquæductum montis Corcovado (Gardner n. 1192 in h. Kew.), Ilheos ad arborum corticem (Wawra et Maly n. 220 fide Wawra bot. Ergeb. bot. p. 131). — Miq. in Mart. flor. Bras. fasc. 11. p. 12. Caulis procumbens radicans quadrangulus glaber. Limbi 0,006 longi, 0,003 lati. Petioli vix 0,001 longi. (v. s.)

150. *P. ERNSTIANA*, foliis petiolatis inferioribus obovatis ellipticisve superioribus subovato-lanceolatis apice acuminatis, acumine acuto, basi acutis acutiusculisve utrinque glabris nigro-punctatis membranaceis subopacis 3-nerviis; centrali nervo ad apicem usque lateralibus supra medium ductis, petiolo pilosulo vel glabro, amento terminali maturo folia duplo superante filiformi laxifloro, pedunculo petiolum æquante, bractea rotundata centro peltata, bractea subimpressa subglobosa apice suboblique subacutata glabra nigra. ☉ In Venezuela prope coloniam Tovar (Fendler n. 1163! in h. DC.). Caulis basi radicans, hirsuto-pubescentis. Limbi 0,075 longi, 0,025 lati. Petioli 0,007 longi. Pedunculi 0,01 longi. Amenta 0,22 longa. (v. s.)

151. *P. MARTIANA* (Miq. ! syst. p. 189), foliis petiolatis obovatis, basin versus cuneatim angustatis apice rotundatis aliquando emarginatis glabris rigidis

- 1-3-nerviis, centrali nervo ad apicem usque ducto subtus prominulo, lateralibus haud conspicuis, amentis terminalibus brevibus densifloris folia duplo triplove superantibus, pedunculo folia fere duplo superante, bractea rotundata centro peltata, ovario semiimmerso apice oblique rostellato, stigmate punctiformi. ☉ In Brasilia (Sello! in h. Berol.), in arboribus viciis prope Mandioccam (Mart. in h. monac.), in Nava Granata prope La Paila (Horton! in h. Kew. foliis membranaceis diversa). — Miq. in Linn. v. 20. p. 128 et Mart. flor. Bras. p. 23. t. 1. fig. 2. Herba glabra, basi radicans, inter muscos vel in arboribus crescens. Limbi circ. 0,02 longi. Petioli 0,008-0,009 longi. (v. s.)
152. *P. BORBONENSIS* (Miq. syst. p. 109), foliis petiolatis superioribus paulo minoribus subrotundis vel rotundatis apice paulo protracto-obtusis inferioribus saepe paulo majoribus rhombico-oblongis vel rhombico-rotundatis apice obtusiusculis omnibus coriaceis subpellucidis præter apicem ciliatum glabris glandulosis 3-nerviis, centrali nervo crasso subtus prominente ad apicem usque ducto, lateralibus nervis subtilioribus haud conspicuis, nervulis quam exilissimis, petiolo glabro lævi, amento terminali brevissimo laxifloro, pedunculo petiolatus superante, bractea rotundata centro peltata, ovario subimmerso. ☿ In ins. Bourbon (h. Mus. Par. n. 474 et 478. fide Miq. l. c.). Erecta glabra lævis. Limbi 0,03 longi, 0,02-0,02½ lati. Petioli 0,005 longi. (Ex Miq. l. c.)
153. *P. SUBROTUNDIFOLIA* (Cas. DC. in Seem. journ. bot. 1866. p. 136), foliis alternis vel apice oculis oppositis breviter petiolatis subreniformi-rotundatis inferioribus e basi subcordata ovatis omnibus utrinque glabris membranaceis tenuibus pellucidis reticulato-nervulosis 3-nerviis, centrali nervo ad apicem usque lateralibus alte ductis, nervulo apice juxta marginem currente, petiolo glabro membranaceo, amentis axillaribus terminalibusque filiformibus subdensifloris folia multoties superantibus, pedunculo glabro petiolum æquante, bractea rotundata centro peltata subsessili, ovario impresso subovato, stigmate carnosulo-puberulo. ☉ In Cuba (Wright n. 2263! in h. DC.). *P. serpens* Griseb. Cat. pl. Cub. p. 63. Herba basi radicans, 0,01 alta, glabra. Limbi 0,011 longi, 0,013 lati. Petioli 0,005 longi. (v. s.)
154. *P. RIEDELIANA* (Regel! in Gartenfl. 1859. p. 229. t. 265. f. 2. 6. 7. et in Cat. hort. Petrop. 1858. p. 35), foliis breviter petiolatis ovato-acuminatis, acumine acutiusculo, basi rotundatis cordulatisve glabris apicem versus ciliatis nigro-punctulatis 3-nerviis, centrali nervo validiore ad apicem usque et lateralibus supra medium ductis, amentis terminalibus axillaribusque folia superantibus subdensifloris, ovario effuso oblique rostellato nigropunctulato. ☉ In Brasilia (Riedel et Langsdorff! in h. Petrop.), in hort. Petrop. culta. Herba repens radicans. Limbi 0,03 longi, fere 0,02 lati. Petioli 0,005 longi. (v. s.)
155. *P. COLUMBIANA* (Miq. in Hook. Lond. journ. bot. v. 4. p. 429), foliis sessilibus lineari-oblongis utrinque attenuatis apice obtusiusculis glabris rigidulis uninerviis, centrali nervo ad apicem usque ducto subtus prominulo, amentis axillaribus terminalibusque apice ramulorum congestis filiformibus subdensifloris, pedunculo glabro, bractea rotundata centro peltata, ovario basi immerso. ☉ In Colombia (1) (Hartweg n. 1397! in h. Kew.). Caulis glaber 0,003 crassus. Limbi 0,025 longi. Peduncul. 0,015 longi. (v. s.)
156. *P. DEMISSA*, foliis petiolatis ovato-lanceolatis apice obtusiusculis plerumque emarginulatis glabris apice ciliolatis pellucido-membranaceis remote nervulosis uninerviis vel obsolete 3-nerviis nervuloque juxta marginem currente, petiolo glabro subtiliter puberulo, amentis terminalibus solitariis densifloris, bractea rotundata centro peltata, ovario emerso apice attenuato, stigmate minuto. ☉ Ad basin Andium prope Punta Playa (Jameson n. 743! 744! in h. Boiss.), Venezuela (Moritz n. 1940! in h. brit. Mus.). *P. Jamesoniana* C. DC.

(1) Ex nomine celeb. detectoris Americæ rectius est *Columbia* quam more anglico *Columbia*. (Alph. DC.)

- spathulatis, centrali nervo nervos utrinque 4-5 mittente,  $\frac{2}{4}$  In Jamaica (Mac Farlen! in h. Kew.). Caulis angulatus glaber. Limbi 0,07-0,08 longi, 0,02-0,025 lati. Pedunculi 0,02 longi. (v. s.)
168. *P. AMPLEXIFOLIA* (A. Dietr. sp. v. 1. p. 145), foliis sessilibus oblongis lanceolatisque, apiculo brevissimo, basi attenuatis nervoso-venosis glaberrimis carinosis, amentis terminalibus longissimis, pedunculo brevi crasso.  $\frac{2}{4}$  In Jamaica. *P. amplexifolium* Link enum. v. 1. p. 38. *P. talinifolia* Link Jahrb. v. 3. p. 64 (vide R. et Sch. v. 3. p. 243). Caulis brevis, ramis brevibus approximatis. Folia 1-4 poll. longa,  $1\frac{1}{2}$  poll. lata. Amenta 4-5 poll. longa. Pedunculi 6 lin. longi. — Verisimillime eadem ac præcedens.)
169. *P. ACUTIFOLIA* (Cas. DC. in Seem. Journ. 1866. p. 142), foliis petiolatis oblongo-lanceolatis apice acutis basi in petiolum decurrentibus utrinque glabris margine ciliolatis membranaceis subpellucidis reticulato-nervulosis, centrali nervo ad  $\frac{2}{3}$  longitud. usque nervos alternos subadscendentes mittente nervuloque apice juxta marginem corrente, petiolo glabro, amentis apice caulis binatis (uno squama fulto altero terminali) filiformibus densifloris folia æquantibus, bractea rotundata centro peltata subsessili, ovario subimpresso apice triangulariter attenuato, stigmate carnosulo antice supra medium inserto, bacca cylindrica oblique mucronulata. ☉ In Peruvia orientali prope Tarapoto (Spruce n. 4094! in h. DC.). Herba circ. 0,06 alta, glabra. Limbi 0,045 longi, 0,006 lati. Petiolaris pars 0,007 longa. Amenta vix 0,001 crassa. (v. s.)
170. *P. MYRIOCARPA* (Miq. ! syst. p. 185), foliis longiuscule petiolatis elliptico-lanceolatis summisque lanceolatis basi acutis (vel obtusis? Miq. syst. l. c.) tenuiter coriaceis pellucido-punctatis utrinque glabris, centrali nervo ad apicem usque ducto nervos subtiles adscendentes utrinque 3-4 mittente, petiolo ciliato, amento terminali folia subtriplo superante densifloro, pedunculo glabro petiolum subæquante, bacca patente ovato-cylindrica apice acutata. ☉ In Brasilia prov. Minas-Geraes (Clausen! in h. Deless.), Guyana gallica prope Harouang (Sagot n. 538! in h. Kew.). Herba scandens? e nodis radicans. Caulis lignosus. Limbi 0,066 longi, 0,026 lati. Petioli 0,015 longi. Pedunculi 0,01 longi. (v. s.)
171. *P. PALLESCENS* (Miq. in Hook. Lond. Journ. bot. v. 8. p. 460), foliis alternis vel verticillato-confertis petiolatis elliptico-ovatis acuminatis basi leviter connatis cordatis margine denticulato-undatis 3-4-costatis supra saturate viridibus marginibusque puberulis subtus pallidis glandulosis inque costis partissimè pilosulis, amentis elongatis plerumque terminalibus geminatisque densifloris, bractea rotundata pedicellato-peltata glandulosa, ovario immerso vertice plano, stigmate puberulo antice medio ovarii inserto. ☉ Ex Guatemala in horto Societ. Hort. Londin. culta. Rami griseo-fuscescenti-pallidi. Petioli 2-1 poll. longi. Folia 4-3 poll. longa. (Ex Miq. l. c.)
172. *P. LANCIFOLIA* (Hook. icon. plant. t. 332), foliis brevissime petiolatis oblongo-lanceolatis apice cuspidatis basi in petiolum cucullatim decurrentibus utrinque glabris membranaceis subpellucidis margine hyalino-attenuatis reticulato-nervulosis nervoque centrali subus subprominulo ad  $\frac{2}{3}$  longitud. usque nervos suboppositos adscendentes utrinque 5-6 mittente, nervulo juxta marginem apice corrente, amentis apice ramulorum aphylorum plerumque binatis (uno terminali altero axillari squama minuta fulto) quam folia minoribus filiformibus densifloris, bractea rotundata centro peltata subsessili, ovario impresso oblongo apice acuminato antice medio stigmatifero. ☉ In Costa-Ricæ sylvis ad Barba-Vulcan (Hoffmann n. 61! in h. Berol.), prope Xalapa (Hook. l. c.). Herba e nodis radicans, scandens? Caulis glaber. Limbi 0,13 longi, 0,045 lati. Petioli 0,005 longi. (v. s.)
173. *P. MONTANA*, foliis petiolatis elliptico-oblongis utrinque acutis glabris membranaceo-subcoriaceis subpellucidis reticulato-nervulosis nervoque centrali ramuloso, petiolo glabro, amentis terminalibus solitariis, pedunculo glabro petiolum superante, bractea rotundata centro peltata, ovario semi-immerso brevissime rostellato, stigmate minuto antico. ☉ In Peruviae subandinae sylvis montosis (Poeppig n. 1519! in h. Boiss.). Caulis simplex, glaber, 0,1-0,15 altus, radicans, decumbens? Limbi vix 0,05 longi. Petioli 0,008 longi. Pedunculi vix 0,02 longi. (v. s.)



174. *P. SPECTABILIS* (Miq. l in Hook. Lond. journ. bot. v. 4. p. 417), foliis alternis suboppositivis longiusculis petiolatis ovatis apice breviter acuminatis, acuminē acuto obtusovē, supra pilosulis subtus adpresse pubescentibus membrænacels opacis, centrali nervo ad apicem usque ducto subtus vix prominulo ad  $\frac{1}{2}$  longitud. usque nervos alternos adscendentes utrinque 6-8 mittente, petiolo adpresse puberulo, amentis in axillis foliorum squamiformium plerumque ternatis (unde inflorescentia pseudo-paniculata) filiformibus subdensifloris, pedunculo adpresse puberulo, bractea rotundata centro peltata, ovario e basi cuneata apice triangulari-acuto, stigmatē minuto antico.  $\frac{1}{4}$  In Peruvia (Mathews n. 1635! in h. Kew.). Caulis subligustus, nodulosus, supernè adpresse pubescens. Limbi 0,14 longi, 0,065 lati. Petioli 0,635 longi. Pedunculi 0,01 longi. Amenta 0,1-0,12 longa. (v. s.)

175. *P. MAGNOLLEFOLIA*, foliis petiolatis obovatis vel obovato-spathulatis vel obovato-subrotundis apice breviter subattenuatis obtusiusculis saepe levissime emarginulatis basi in petiolum contractis utrinque glabris rigidis subopacis pellucido-punctulatis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos adscendentes suboppositos utrinque 3-4 mittente nervulose juxta marginem currente, amentis terminalibus densifloris folia parum superantibus, pedunculo glabro petiolum æquante, bractea rotundata centro peltata pedicellata, ovario immerso apice suboblique rostrato infra rostrum antice stigmatifero. ☉ In Paramaribo (Kegel n. 806 fide Miq. in Linn. v. 22. p. 76), Surinam (Hostmann n. 2601 in h. Berol.), Cayenna (Aubl. fl. Guy. p. 22), Caracas (Gallmer n. 338! in h. Berol.), Costa-Rica (Hoffmann n. 799! in h. Berol.), in arboribus vetustis ad Xalapa (Miq. in Linn. v. 18. p. 713), ad Zacapan alt. 2-3000 ped. (Galeotti n. 6003 fide Mart. et Gal. in Bull. Acad. Brux. v. 10. p. 18), in Trinitate (Sieber! in h. Deless.), Martinica (Sieber n. 255! et 355!, 349! in h. DC., in h. Berol., Belanger n. 327! in h. Francov.), Sto Domingo (Poiteau! in h. Deless.), Antigua (Poiteau! ibid.), St-Vincent (Miq. in Hook. Lond. Journ. bot. v. 4), Jamaica (Sloane! in h. Brit. Mus.), Cuba (Wright n. 500! et 509! in h. DC.), Guadalupe (Perrottet! in h. Deless.), Barbados (Maycock! flor. Barb. p. 40 quoad synonym. Jacquin). P. longum humilis fructu e summitate caulis prodeunte Sloane! hist. p. 136 et coll. icon. orig. in Brit. Mus. t. 86 cum planta! Saururus humilis Plum. Am. ed. Par. p. 53. Piper obtusifolium Aublet l. c. P. magnoliaefolium Jacq. coll. 3. p. 210, ic. rar. 2. t. 213, Vahl en. 1. p. 338, Loud. encycl. p. 29. n. 529. Pep. magnoliaefolia A. Dietr. sp. v. 1. p. 153, Miq.! syst. p. 97 et in Kruick v. 1. p. 221. Pep. subrotunda A. Dietr. sp. v. 1. p. 154. Pep. obtusifolia A. Dietr. sp. v. 1. p. 154 pro parte, Miq.! syst. p. 194 pro parte, in Linn. v. 20: p. 128, fide Mart. flor. Bras. fasc. 11. p. 24, Griseb. flor. W. Ind. p. 166, cat. pl. cub. p. 64. Suffruticulus pedalis, succulentus, basi radicans, erectus, pauci-ramosus, glaber. Limbi 0,06 longi, 0,035 lati. Petioli 0,015 longi. (v. s.)

β, *emarginulata*, foliis obovato-subrotundis apice emarginulatis coriaceis. In Venezuela prope coloniam Tovar alt. 6500-7000 ped. (Fendler n. 1152! in h. DC.), Nova Granata prov. Popayan (Triana n. 62! in h. DC.), Cuba orientali (Wright n. 1152! in h. DC.). (v. s.)

γ, *parvifolia*, foliis minoribus obovato-spathulatis apice emarginulatis. In Cuba (Ossa! in h. DC.). (v. s.)

176. *P. GALEOTTII* (Miq. syst. p. 190), foliis longe petiolatis obovatis apice lato-rotundatis basi in petiolum contractis utrinque glabris coriaceis pellucido-punctulatis supra viridibus subtus glaucescentibus, centrali nervo subtus prominulo ad basin et altius nervos numerosos adscendentes parallelè subtus prominulos mittente, amentis terminalibus cylindricis folia æquantibus, bractea rotundata margine erosula fusco-glandulosa centro peltata pedicellata, ovario immerso antice infra apicem stigmatifero, bacca profunde immersa obovata vel oblonga apice sterili acuminato nunc flexuoso aucta. In Cordilleris prope Vera Cruz alt. 3000 ped. supra rupes et arborum truncos vetustos (Galeotti n. 6003 in h. Deless.). Herba succulenta, erecta, glaberrima. Limbi 0,08-0,08 longi, 0,045-0,055 lati. Petioli 0,03 longi. (Ex Miq. l. c.)

177. *P. PYRIFOLIA* (Humb. et Bonpl. l. pl. æq. v. 1. p. 61), foliis longiusculis petiolatis late subrhombéo-ellipticis apice acutis acutiusculis basi cuneatis

utrinque glabris coriaceis opacis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos subtiles suboppositos adscendentes utrinque 6-8 mittente, petiolo glabro, amentis terminalibus densifloris longiuscule pedunculatis folia æquantibus, bractea rotundata glanduloso-punctata centro peltata breviter pedicellata, ovario semi-immerso apice suboblique longe rostrato, stigmate carnosulo-puberulo infra rostrum inserto.  $\zeta$  In fissuris rupium et ad arbores Novæ Andalousiæ prope cœnobium Caripense (Humb. et Bonpl. l. c., Humb. ! in h. Willd. n. 709 !), Venezuela (Moritz n. 1783 ! in h. Brit. Mus.). Piper murinum Roem. et Sch. syst. v. 1. p. 325. Pep. pyrifolia Kunth synops. v. 1. p. 115; A. Dietr. sp. v. 1. p. 152, Miq. ! syst. p. 191. Suffructulus glaber, carnosus. Limbi 0,1 longi, 0,065 lati. Petioli 0,02 longi. (v. s.)

178. P. EMARGINATA (Ruiz et Pav. ! flor. Peruv. v. 1. p. 30. t. 49. fig. a), foliis longiuscule petiolatis oblongo-spathulatis apice emarginatis basi cuneatim in petiolum angustatis utrinque glabris membranaceis subpellucidis subtiliter creberrime pellucido-punctulatis remote nervulosis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos suboppositos plerumque 5 adscendentes quorum inferiores magis approximati mittente nervuloque juxta marginem apice currente, petiolo glabro, amentis terminalibus densifloris longiuscule pedunculatis folia subæquantibus, pedunculo glabro petiolum subæquante, bractea rotundata glabra centro peltata breviter pedicellata, ovario immerso apice suboblique longe rostrato, stigmate infra rostrum antice inserto.  $\odot$  In Peruvia (Ruiz ! in h. Berol.). — A. Dietr. sp. v. 1. p. 146; Miq. ! syst. p. 192. Piper emarginatum Vahl enum. v. 1. p. 339. Planta basi repens? succulenta, glabra. Limbi 0,14 longi, 0,055 lati. Petioli 0,025 longi. (v. s.)

179. P. PSEUDO-VARIEGATA, foliis longe petiolatis elliptico-lanceolatis utrinque acutis supra glabris subtus piloso-pubescentibus rigido-coriaceis opacis remote pellucido-punctulatis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos alternos adscendentes utrinque 3 quorum utrinque 2 parum supra basin soluti mittente, petiolo hirtello, amento terminali folio scarioso fulto filiformi densifloro folium superante, bractea rotundata centro peltata breviter pedicellata, ovario immerso apice oblique longe rostrato, stigmate carnosulo infra rostrum inserto.  $\zeta$  In Novæ Granatæ prov. Pasto. (Triana n. 63 ! in h. Triana). Ramuli sublignosi hirtelli. Limbi 0,14 longi, 0,06 lati. Petioli 0,06 longi. (v. s.)

180. P. DISTACHYA (A. Dietr. sp. v. 1. p. 156), foliis petiolatis vel longiuscule petiolatis inferioribus ovatis apice acuminatis, acumine acuto, basi rotundatis cordulatisve superioribus oblongo-ellipticis apice acuminatis, acumine acuto, basi attenuatis subacutisve utrinque glabris junioribus ciliolatis membranaceis subpellucidis pellucido-glandulosis fere enervulosis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos adscendentes alternos suboppositosve mittente nervuloque juxta marginem currente, petiolo glabro, amentis terminalibus axillaribusque plerumque apice ramuli binatis densifloris filiformibus folia duplo superantibus, bractea rotundata centro peltata subsessili, baccis subremotis patentibus cylindricis apice scutatum auctis, scutello subrotundo sursum acutato, stigmate medio scutelli inserto.  $\odot$  In Brasilia (Sello ! in h. Berol. ubi prave Pep. rupestris dicta, Spruce ! in h. Kew.), ex arboribus vetustis dependens ad Orellanakreek (fide Miq. in Linn. v. 18. p. 226), supra arbores ad Panama (Seemann n. 897 fide Miq. in bot. Herald. p. 198), ins. Cayenna (Aublet pl. Guy. v. 1. p. 22), Guyana gallica (Poiteau ! in h. Deless.), Sto Domingo (Poiteau ! ibid.). — Miq. ! syst. p. 184, ill. p. 24. t. 23, Griseb. flor. W. Ind. p. 166. Saururus hederaceus minor Plum. Am. ed. Par. p. 51. t. 67. Piper distachyon L. sp. p. 42, Aublet l. c., Lam. ill. p. 81, Vahl enum. v. 1. p. 344, Willd. sp. v. 1. p. 166, Spreng. syst. v. 1. p. 118. ? Pep. rupestris Miq. ! syst. p. 183 pro parte. Herba scandens vel repens, e nodis radicans. Caulis glaber. Limbi 0,06 longi, 0,04 lati. Petioli 0,02 longi. (v. s.)

181. P. MARISIANA, foliis petiolatis lanceolatis vel subovato-lanceolatis apice acuminatis, acumine acuto, basi acutis utrinque glabris ciliolatis membranaceis pellucidis pellucido-punctulatis parce nervulosis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos adscendentes utrinque 3 quorum inferiores utrinque 2 paullo supra basin soluti mittente nervuloque apice juxta marginem currente, petiolo superne ciliolato, amentis terminalibus vel apice caulis squamis

fulvis et tum binatis densifloris folia superantibus, bractea rotundata centro peltata subsessili, ovario impresso oblique ovato apice scutatum aucto, scuto sursum acutato, stigmata medio scuti inserto, bacca ovato-oblonga vertice oblique breviter mucronata.  $\zeta$  In sylvis ad arbores prope Maribí ditionis Japura prov. Rio Negro (Mart. ! in h. monac.). *Pep. myriocarpa* Miq. ! in Linn. v. 20. p. 128 et in Mart. flor. Bras. fasc. 11. p. 22. t. 1. fig. 5, non Syst. Scandens. Caulis lignosus glaber. Limbi 0,065 longi, 0,025 lati. Petioli 0,006 longi. (v. s.)

182. *P. MACROPODA* (Miq. ! in Linn. v. 20. p. 118), foliis longe petiolatis obovatis vel oblongo-obovatis apice late rotundatis vel obtusis basi attenuatis vel cuneatis utrinque glabris junioribus glandulosis crasse coriaceis pellucidopunctulatis centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos alternos oppositosve adscendentes utrinque 3 sursumque nervulos mittente, amentis terminalibus basi pedunculi bibracteatis vel apice pedunculi communis (id est ramuli) terminatis pedunculatis densifloris folia subæquantibus, pedunculo singulo basi squama fulto, bractea rotundata centro peltata subsessili, ovario immerso elliptico-cylindrico, baccae fuscae nitidae apice filiformi pallido flexuoso.  $\odot$  In Brasilia (Pohl n. 4790 ! in h. Vindob.). — Miq. ! in Mart. Flor. Bras. fasc. 11. p. 24. Caulis ramosus, repens, radicans, glaber. Limbi 0,06-0,065 longi, 0,025-0,03-0,035 lati. Petioli 0,03-0,01 longi. (v. s.)

183. *P. OBTUSIFOLIA*, foliis breviter petiolatis obovato-spathulatis apice rotundatis leviter emarginulatis vel inferioribus apice acutiuscule protractis omnibus basi in petiolum brevem decurrentibus utrinque glabris rigidis subopacis pellucido-punctulatis parce nervulosis et nervo centrali ad  $\frac{1}{2}$  longitud. usque nervos utrinque 5 approximatos mittente, amentis terminalibus solitariis vel apice ramulorum subpaniculatim confertis breviter pedunculatis folia superantibus densifloris filiformibus, bractea rotundata glandulosa centro peltata breviter pedicellata, ovario immerso glanduloso apice longe rostrato, stigmata antice infra rostrum inserto.  $\odot$  In Venezuela inter Maracai et Choroui (Fendler n. 2401 ! in h. DC.), Trinitate (Sieber n. 365 ! in h. DC.), Sainte-Croix (h. DC.), Cuba (Wright n. 1689 ! in h. DC.), in Hort. Berol. cult. (h. Willd. n. 715 !). — A. Dietr. sp. p. 154 pro parte. *Piper obtusifolium* L. ! sp. p. 42 et h. P. *cuneifolium* Jacq. coll. v. 4. p. 127, ic. rar. v. 2. t. 214. Vahl enum. v. 1. p. 340. *Pep. obtusifolia* var. *grandifolia* Miq. ! syst. p. 196. *Pep. cuneifolia* Miq. syst. p. 194, *Pep. Swartziana* Griseb. flor. W. Ind. p. 166. Limbi 0,12 longi (cum parte petiolari), 0,05 lati. Petioli 0,005 longi. (v. s.)

$\beta$ , *clusiæfolia*, foliis subtus cauleque rubescentibus minus carnosius. In caldariis culta. *Piper clusiæfolium* Jacq. coll. v. 3. p. 219, icon. rar. v. 2. p. 2. t. 212. *Piper obtusifolium* Lam. ill. p. 81, Vahl enum. v. 1. p. 338 (exclus. synonym. Pluk. et L.). *Piper marginatum* Haw. suppl. pl. succ. p. 3. *Pep. obtusifolia* Miq. syst. p. 194 pro parte var.  $\alpha$ , Griseb. flor. W. Ind. p. 166.

184. *P. SUCCULENTA*, foliis breviter petiolatis suboblongo-lanceolatis apice obtusiusculis basin versus cuneatis in petiolum decurrentibus margine hyalino-subextenuatis utrinque glabris apicem versus subtilissime ciliolatis membranaceis subpellucidis pellucido-punctulatis reticulato-nervulosis et centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos suboppositos adscendentes utrinque 4 mittente, nervuloque juxta marginem currente, petiolo glabro, amentis axillaribus terminalibusque apice ramulorum approximatis adspectu binatis filiformibus densifloris folia subduplo superantibus, bractea rotundata centro vel (in monstis) paullo supra basin peltata brevissime pedicellata (vel in monstis) non peltata oblonga margine subtilissime ciliolata, ovario semi-immerso apice subrostrato, stigmata subantico, bacca emersa ovata apice suboblique flavicanti-rostrata.  $\odot$  In Venezuela prope coloniam Tovar (Fendler n. 1157 ! et 1158 ! in h. DC.). Herba glabra ramulosa erecta. Ramuli subquadranguli, anguste alati. Limbi 0,09 longi, 0,027 lati. Petioli 0,015 longi. (v. s.)

185. *P. ALPINA* (A. Dietr. sp. v. 1. p. 157), foliis longe petiolatis obovatis vel elliptico-obovatis apice subattenuato-acutiusculis basi subcuneato-acutis in petiolum subdecurrentibus utrinque glabris junioribus apicem versus basique subtilissime puberulis rigidis opacis reticulato-nervulosis et centrali nervo ad

- $\frac{1}{4}$  longitud. usque nervos utrinque 3 mittente, petiolo glabro, am-  
 ribus filiformibus densifloris longe pedunculatis folia æquantibus  
 superantibus, pedunculo glabro petiolum superante, bractea rotunda  
 glandulosa centro peltata subsessili, ovario immerso apice oblique  
 glandulari-rostrato, stigmate punctiformi puberulo antice infra rostru-  
 m. In Jamaicae summis alpinis (Swartz in h. Willd. n. 716!), Cul-  
 n. 507 fide Griseb. cat. pl. Cub. p. 64). Piper alpinum Sw. prod-  
 Ind. occ. v. 1, p. 62, Vahl enum. v. 1, p. 340, Willd. sp. v. 1, p. 11  
 syst. v. 1, p. 115. Pep. alpina Miq. syst. p. 188, Griseb. l. c. et fl.  
 p. 167. Suffructiculus carnosus, sublignescens, suberectus, basi radi-  
 lis subangulosus, glaber. Limbi 0,025 longi, 0,019 lati. Petioli  
 (v. s.)
186. *P. NEMATOSTACHYA* (Link Jahrb. v. 1. 3. p. 6, fide Rœm. et Sch. |  
 p. 248), foliis petiolatis elliptico-lanceolatis apice acuminatis, acutis  
 basi obtusis vel rotundatis supra pilis minutis stellatis adpressis  
 longis aparsis hirtellis junioribus apicem versus ciliolatis rigidulis  
 cœcis non reticulato-nervulosis, centrali nervo ad  $\frac{1}{4}$  longitud. usque  
 oppositos utrinque 2 mittente, petiolo glabro, amentis binatis breviter  
 ciliolatis folia fere triplo superantibus filiformibus densifloris, bractea  
 glandulosa centro peltata subsessili, ovario semi-immerso apice  
 scuto glanduloso punctulato oblongo-elliptico, stigmate medio sc  
 bacca basi subimmersa cylindrica patente apice rostellata.  $\frac{1}{2}$  In  
 arborum truncos (Hoffmansegg in h. Willd. n. 720!), prov. Par  
 h. monac.). — A. Dietr. sp. v. 1, p. 156, Miq. syst. p. 183. (Pip-  
 stachyon Rœm. et Sch. l. c. Pep. rupestris Miq. in Linn. v. 20.  
 Mart. flor. Bras. fasc. 11, p. 22 pro parte. Caulis, basi lignosus  
 ramosus, glabriusculus, sursum hirtellus. Limbi 0,05-0,07 longi  
 lati. Petioli 0,002-0,008 longi. (v. s.)
187. *P. ELONGATA* (Kunth in H. et B. nov. gen. v. 1. p. 6), foliis  
 latius suboblongo-ellipticis apice anguste et subfalcatim acuminatis  
 acutis, basi rotundatis subtiliter ciliolatis basi et apice subtiliter  
 subcoriaceis subopacis pellucido-punctulatis septuplinerviis, ca-  
 paullo supra basin nervos suboppositos utrinque 3 quorum super-  
 apicem usque ducti mittente, nervulo juxta marginem cur-  
 superne ciliolato, amentis terminalibus longiuscule pedunculatis  
 densifloris folia triplo superantibus, bractea rotundata centro  
 sessili glanduloso-punctata, ovario oblongo-elliptico glanduloso  
 apice scutato, stigmate minuto medio scuti inserto, bacca o-  
 drica basi subimmersa apice rostellata.  $\odot$  In convalle Carip-  
 hexap. ad saxa (Kunth l. c. et Humb. in h. Willd. n. 717!), l.  
 n. 5884! in h. Berol.). — Kunth synops. v. 1, p. 115, Miq. syst.  
 tensum Rœm. et Sch. syst. v. 1, p. 325. Caulis erectus, circ. 0,8  
 lentus, glaber. Limbi 0,065 longi, 0,03 lati. Petioli 0,006 longi.
188. *P. PRODUCTA* (Griseb. flor. W. Ind. p. 166), foliis longe pe-  
 acuminatis apice imo acutis mucronulatisque basi rotundatis in-  
 latis utrinque glabris apicem versus ciliolatis membranaceis vel i-  
 dulis pellucido-punctulatis pellucidis subpellucidisve parce  
 centrali nervo ad  $\frac{1}{4}$  longitud. usque nervum utrinque unum adsc-  
 terne mittente nervisque lateralibus utrinque 3-4 fere e basi  
 dentibus subtilibus, petiolo glabro vel apicem versus ciliolato, an-  
 pedunculatis quam folia duplo brevioribus densifloris filiformibus  
 undata centro peltata glandulosa, ovario semi-immerso oblongo  
 tato, stigmate medio scuti inserto, bacca patente cylindrica  
 trita.  $\odot$  In Cuba (Wright n. 501! 508! 1419! 1420! in h. fl.  
 (Mac Fad. fide Griseb. l. c.). — Griseb. in Mem. am. Ac. v. 8, p.  
 Cub. p. 64. Pep. distachya Griseb. Cat. pl. Cub. p. 64 exclus. i-  
 scandens, e nodis radicans, glabra. Limbi 0,08 longi, 0,043 lati.  
 longi. (v. s.)
- 2° Folia multiplinerviæ.
189. *P. ELEGANS*, foliis longe petiolatis elliptico-lanceolatis utrinque

submembranaceis subpellucidis novemnerviis, centrali nervo validiore & longitud. usque nervos alternos adscendentes utrinque 4 quorum inferiores e basi soluti mittente, petiolo glabro, amentis terminalibus vel axillaribus binatis densifloris quam folia brevioribus vel ea subaequantibus, bractea undata centro peltata, ovario immerso apice rostrato, stigmate minuto c. o. ☉ In Peruvia orientalis montibus prope Tarapoto secus flum. Mayo rance n. 48711 in h. DC.). Herba glabra. Limbi 0,07-0,09 longi. Petioli 1-0,07 longi. Pedunculij 0,01 longi. (v. s.)

P. SEPTUPLINERVA (Cas. DC. in Seem. Journ. 1866. p. 142), foliis longe ciliatis oblongo-ovatis apice breviter acuminatis, acumine acuto, basi rotundatis paullo supra basin peltatis utrinque pubescentibus dein supra glabris subcoriaceis opacis septuplinerviis, centrali nervo validiore ad & longitud. usque nervos subuliores adscendentes utrinque 3 mittente, petiolo pubescentibus, amentis apice caulis plerumque binatis (uno terminali altero squamato fulto) pedunculatis filiformibus densifloris folia superantibus, pedunculo quam petiolus fere quadruplo brevior, bractea rotundata centro peltata sessili, ovario post anthesin immerso subovato apice rostrato, stigmate antheris infra rostrum inserto, bacca subovata apice rostrata. ☉ In Cuba (Wright 2360 in h. DC.). Pep. monstrosifolia Griseb. Cat. pl. Cub. p. 64. Fruticosa. Caulis hirtello-pubescent. Limbi 0,16 longi, 0,08 lati. Petioli 0,095 longi. Bacca matura 0,004 crassa. (v. s.)

P. AUSTRALE (Kunth in M. et B. nov. gen. v. 1. p. 62), foliis petiolatis ovato-oblongis apice acuminatis basi rotundatis obtusius utrinque glabris compositis opacis sub lente verruculis obsitis septuplinerviis, centrali nervo nervos adscendentes aegre discernendos paullo supra basin utrinque 2 mittente, laterali nervo utrinque uno e basi soluto, petiolo glabro, amentis axillaribus terminalibusque folia superantibus densifloris, bractea rotundata centro peltata subsessili, ovario oblique ovato scutato antice et apice aucto, stigmate medio scuti inserto, bacca cylindrica basi immersula apice oblique mucronata. ☉ In Brasilia ad Petropolim, Corcovado ad arborum corticem (Wawra et My p. 437, 534, fide Wawra bot. Ergeb. p. 185. t. 83. fig. 1), prov. Para ad arborum fructuosos (Spruce n. 18491 in h. Kew.), Surinam (Seaman n. 6081 in Kew.), prope Quarta et Tourtonne (Kegel n. 1037 fide Miq. in Linn. v. 22. 76), in rupibus regionis temperatae prope Cumana (Humb. in h. Willd. 711'), umbrosis Misantlae, Xalapae (Schiede in h. Schlecht. fide Miq. in An. v. 18. p. 713). — Kunth synopsis. v. 1. p. 115, A. Dietr. sp. v. 1. p. 451, q. l. syst. p. 182, et pro parte in Mart. flor. Bras. fasc. 11. p. 22). Caulis molis, glaber. Limbi 0,04-0,08 longi, 0,03 lati. (v. s.)

P. cordifolia (Wawra l. c. t. 83. fig. 2), foliis inferioribus basi cordatis. In Schomburgkii outa.

P. MACULOSA (A. Dietr. sp. v. 1. p. 163), foliis longe petiolatis late ovato-ipticis apice brevissime acuminatis basi rotundatis subcordatisve subpeltatis utrinque utrinque glabris centrali nervo supra basin nervos adscendentes subpositos utrinque 3 quorum inferiores 2 fere e basi soluti mittente, petiolo abro rubescenti-maculato, amentis terminalibus densifloris folia fere duplo perantibus, bractea rotundata centro peltata, ovario emerso ovato oblique pecto, stigmate antice inserto exiguo. ☉ In Peruvia (Pav. in h. Boiss.), jenna Aubl. hist. pl. (Guy. p. 22.), Stio Domingo (Descourties flor. Ant. 417). Saururus hederaceus Plum. Am. p. 50. t. 66. Piper maculosum L. p. 42, Aublet l. c., Lam. ill. p. 81, Vahl enum. v. 1. p. 344, Willd. v. 1. p. 166, Spreng. syst. v. 1. p. 118. Piper variegatum Pers. syn. 35. Pep. variegata R. et Pav. flor. Peruv. v. 1. p. 33. t. 52. fig. 4, Miq. st. p. 198. Hook. exot. flor. t. 92, Miq. comm. phyt. p. 64. Poirrier maculé secourtils l. c. v. 6. p. 153. Suffructulus e nodis radicans. Caulis glaber, sce vel violaceo maculatus. Limbi 0,12-0,15 longi, 0,1 lati. Petioli 0,15 ngi. (v. s.)

P. WRIGHTIANA, foliis longiuscule petiolatis late ovato-lanceolatis apice acuminatis basi cordatis rotundatisve glabris tenuissimis ciliolatis membranaceis subcoriaceis septupli-novenerviis laterali nervo utrinque uno supra sin soluto ceteris e basi solutis tenuioribus omnibus subus subpromi-

nulis, petiolo glabro, amentis axillaribus pedunculatis densifloris, pedunculo squama fulto, bractea rotundata centro peltata, ovario emerso ovato-rostellato, stigmatibus minuto antico. ☉ In Cuba orientali (Wright n. 498! in h. DC.). Scandens? e nodis radicans. Limbi circ. 0,12 longi, 0,104 lati. Petioli 0,1-0,12 longi. (v. s.)

194. P. STENOCARPA (Regel! in Gartenfl. 1859. p. 291. t. 271), foliis petiolatis ovato-ellipticis apice acuminatis, acumine acuto, basi ima acutis rigidis subopacis utrinque pubescentibus ciliolatisque demum glabris quintupli-novenerviis, centrali nervo ad  $\frac{1}{2}$  longitud. usque nervos alternos subadscendentes utrinque 1-2 mittente, lateralibus nervis utrinque 2-3 e basi ortis supra medium ductis quorum infimi subtiliores, petiolo pubescente, amentis axillaribus terminalibusque folia multoties superantibus densifloris bractea rotundata centro peltata breviter pedicellata, bacca cylindrica basi immersula apice oblique flavide rostellata, stigmatibus antice infra rostellum inserto. ☉ In saxosis humidis umbrosis prope Rio-Janeiro (Riedel! in h. Petrop.) Herba scandens. Caulis junior pilis brevibus pubescens, dein glaber. Limbi 0,05 longi, 0,027 lati. Petioli 0,01 longi. (v. s.)

195. P. PSEUDO-AMPLEXICAULIS, foliis petiolatis cuneato-ovatis basi amplexicaulibus glabris coriaceo-obscuris noveninerviis, centrali nervo validiore subprominulo ad  $\frac{1}{2}$  longitud. usque nervos utrinque 4 oppositos quorum inferiores utrinque 2 e basi soluti mittente, petiolo glabro, amentis axillaribus terminalibusque densifloris sordide ochraceis, bractea rotundata centro peltata fusco-punctulata, ovario subimmerso ovato apice longe rostrato, stigmatibus minuto antice infra rostrum inserto. ☉ In Nova Hispania (Pav. in h. Boiss.), Cuba prope Havanam (Ossa! in h. DC.). Herba 0,2 circ. alta, basi subligulosa, erecta. Limbi 0,035-0,06 longi. Petioli 0,012 longi. (v. s.)

196. P. INCANA (A. Dietr. sp. v. 1. p. 166), foliis petiolatis orbicularibus basi subcordatis vel ovatis crassis carnosius coriaceis opacis utrinque albo-tomentosis supra nitidis noveninerviis, centrali nervo validiore nervos utrinque 4 suboppositos quorum inferiores utrinque 3 e basi cæteri sub medio soluti mittente, petiolo albo-tomentoso, amentis terminalibus elongatis folia triplo superantibus densifloris, pedunculo albo-tomentoso, bractea rotundata centro peltata, ovario subimmerso apice scutato triangulari aucto, stigmatibus minuto pilosulo medio scuti inserto. ☿ In Brasilia (Riedel! in h. hort. Petrop.), rupibus graniticis aridis ad Sebastianopolim (Mart. in h. monac.), in caldariis culta (h. DC.). — Miq.! syst. p. 197, comm. phyt. p. 63. t. 4. fig. A, Linn. v. 20. p. 129, in Mart. flor. Bras. fasc. 11. p. 24. t. 1. fig. 9, Hook. exot. flora t. 66, Schnizl. icon. t. 81. fig. 11-17 (analysis). Piper incanum Spreng. syst. v. 1, p. 117, Loud. encycl. p. 29. n. 545, Haworth Suppl. pl. succ. p. 2, Link et Otto icon. p. 17. t. 7. Suffruticulus carnosus, albo-tomentosus. Limbi 0,05 longi. Petioli 0,035 longi. Amenta 0,1-0,15 longa. (v. v.)

### 3° Folia multinervia.

197. P. MAJOR, foliis longe petiolatis ovato-rotundis apice breviter attenuatis basi rotundatis vel leviter cordatis ciliolatis junioribus utrinque piloso-pubescentibus coriaceo-opacis 7-10-nerviis nervis subprominulis, amentis axillaribus apice ramuli communis geminatis, bractea rotundata flavo-punctata centro peltata, ovario subimmerso apice longe rostellato, stigmatibus carnosulopapilloso infra rostellum antice inserto. ☉ In Brasilia (Sello n. 1204! et 591! in h. Berol.), Rio de Janeiro (Gaudich. n. 88! in h. DC., Weddell n. 298! et 575! in h. DC.), monte Corcovado (Schott! in h. Vindob.) et in Nova Hispania (Pav. in h. Boiss.). Acrocarpidium majus Miq.! syst. p. 60, in Linn. v. 20. p. 118. Pep. hederacea Miq.! in Mart. flor. Bras. fasc. 11. p. 20. Herba repens vel scandens? e nodis radicans. Caulis adpresse piloso-pubescent. Limbi 0,04 longi. Petioli circ. 0,04 longi. Ramuli amenta gerentes 0,05 longi. (v. s.)

β, *hispida*, baccis hispidis. Piper monostachyon Vell. flor. flum. t. 63, non Vahl.

198. P. PONTHEUI (Miq. syst. p. 186), foliis longe petiolatis  $\frac{1}{2}$  longitud. supra

basin peltatis ovato-rotundatis vel ovatis apice acuminatis basi rotundatis supra glabris subtus subtilissime puberulis dein glabratiss subcoriaceis reticulato-nervulosis 7-9-nerviis, centrali nervo validiore subtus subprominulo, lateralibus nervis subtilioribus ægre discernendis, amentis axillaribus longe pedunculatis, bractea rotundata centro peltata, ovario subimmerso apice scutato, stigmatibus minuto antice medio scuti inserto. In India occid. (Ponthieu in h. Deless. ex Miq.), in Venezuela (Moritz n. 342! in h. Brit. Mus.), Guadaloupa (Rich. in h. Francov.), Cuba (Wright n. 497! in h. Kew. et DC.). *Pep. hernandiæfolia* Griseb. Cat. pl. Cub. p. 64. Herba decumbens e nodis radicans. Limbi 0,06-0,08 longi. Petioli 0,05-0,08 longi. Pedunculi 0,02-0,03 longi. (v. s.)

199. *P. STELECHOPHILA*, foliis petiolatis ovato-acuminatis apice oblique acuminatis basi rotundatis vel deltoideis cordatis omnibus basi ima subpeltatis utrinque basi puberula excepta glabris margine extenuato ciliolatis firmule membranaceis subpellucidis pellucidis punctulatis 7-nerviis nervulose obscure juxta marginem currente, petiolo superne ciliolato, amentis terminalibus solitariis, pedunculo puberulo quam petiolus brevior, bractea rotundata centro peltata subsessili, ovario impresso apice peltatim rhomboidali, stigmatibus medio rhombi inserto. ☉ In Andibus Quitensibus declivitate occid. montis Pichinchæ alt. 8000 ad arborum truncos (Jameson n. 400! in h. DC.). Herba scandens. Ramuli uno latere ciliati. Limbi 0,04 longi, 0,022 lati. Petioli 0,011 longi. (v. s.)

200. *P. EMARGINULATA*, foliis petiolatis cuneato-obovatis vel cuneato-lanceolatis apice emarginulatis subcoriaceis opacis glabris 7-nerviis, nervis subtus subprominulis, petiolo brevi glabro, amentis axillaribus plerumque geminatis pedunculatis, bractea rotundata fusco-punctulata centro peltata, ovario subimmerso ovato-attenuato. stigmatibus antico orbiculari carnosulo minuto. ☉ In Peruvia (Pav. in h. Boiss.), Casapi (Mathews n. 1687! in h. Kew.), sylvis Andium occident. (Jameson n. 837! in h. Boiss.). *Pep. obtusifolia* Miq. in Hook. Lond. journ. bot. v. 4 quo ad spec. Mathews. Herba glabra erecta. Caulis glaber tetragonus. Limbi 0,03-0,05 longi. Petioli 0,005 longi. Ramuli amenta gerentes circ. 0,02 longi. (v. s.)

201. *P. CUNEATA* (Miq. in Hook. Lond. journ. bot. v. 4. p. 429), foliis longiuscule petiolatis cuneato-vel-obovato-spathulatis apice obtusis raro retusis basi cuneatis in petiolum attenuatis utrinque glabris vel apicem versus pilosulis carnosis 7-nerviis, centrali nervo validiore lateralibus ægre discernendis, petiolo glabro, amentis terminalibus axillaribusve solitariis longe pedunculatis densifloris, pedunculo plerumque squama fulto, ovario oblique acuminato, stigmatibus antico, bacca acuminatissima. ☉ In ins. St-Vincent (Guilding in h. Hook. fide Miq. l. c.). Herba erecta, glabra, basi radicans. Limbi 0,04-0,07 longi, 0,015-0,025 lati. Petioli 0,02 longi. Pedunculi 0,04-0,05 longi. (Ex Miq. l. c.).

202. *P. UROCARPA* (Fisch. et Mey. in ind. sem. horti Petrop. 1837), foliis petiolatis ovato-cordatis apice acutiusculis utrinque puberulis subtus citius glabris 5-7-nerviis, amentis ramulos axillares terminalesque monophyllos terminantibus, bacca mucrone longo setaceo apiculata, stigmatibus antice infra rostrum inserto. ☉ In Brasilia (Fisch. in h. DC. et Petrop.). — Regel in Gartenfl. 1859. p. 22. t. 265. *Acrocarpidium urocarpum* Miq. in syst. p. 60. Caulis herbaceus, procumbens, radicans, glabriusculus. Limbi 0,015 longi latique. Petioli 0,007 longi. (v. s.)

203. *P. CUBENSIS* (Cas. DC. in Seem. journ. 1866, p. 142), foliis longiuscule petiolatis deltoideo-cordatis glabris margine subciliolatis membranaceis pellucidis 5-7-nerviis non reticulato-nervulosis, petiolo glabro, amentis apice ramuli axillaris petiolo brevioris plerumque geminatis quam folia brevioribus (uno terminali altero laterali squama fulto) breviter pedunculatis, bractea rotundata centro peltata, ovario immerso apice oblique rostrato, stigmatibus globoso-puberulo antico. ☉ In Cuba (Wright n. 499! in h. DC.). *Pep. distachya* Griseb. Cat. pl. Cub. p. 64, quoad n. 499. Herba scandens. Limbi 0,05 longi. Petioli 0,03 longi. (v. s.)

204. *P. REPTANS* (Cas. DC. in Seem. journ. 1866, p. 143), foliis longiuscule petiolatis e basi cordata ovato-rotundis supra hirsutis subtus ad nervos pubescentibus rigidulis subopacis pellucido-punctulatis 5-nerviis parce nervulosis, nervis subtilibus, petiolo hirsuto, amentis axillaribus terminalibusve solitariis longiuscule pedunculatis maturis folia æquantibus, pedunculo hirsuto petiolum æquante, bractea rotundata centro peltata breviter pedicellata, ovario subimpresso oblongo apice subito in rostrum brevem subulato pubescente, stigmatē anticō. ☉ In Novæ Granatæ prov. Barbacoas ad viam Tuqueras alt. 600 m. (Triana n. 581 in h. DC.). Herba repens, hirsuta, e nodis radicans. Caulis filiformis. Limbi 0,012 longi, 0,015 lati. Petioli 0,013 longi. (v. s.)

205. *P. SCANDENS* (Ruiz et Pav. flor. Per. v. 1. p. 32. t. 51. fig. b), foliis longiuscule petiolatis ovato-rotundatis vel subreniformibus apice obtusiusculis supra adpressis subtus ad nervos puberulis membranaceo-subcoriaceis 5-nerviis et reticulato-nervulosis, nervis subtus conspicuis 3 centralibus validioribus, petiolo puberulo limbum superante, amentis solitariis ramulos oppositifolios vel axillares unisquamosos terminantibus pedunculatis, bractea ovato-æuta centro peltata subsessili, ovario oblongo apice rostrato, stigmatē minuto antico, bacca ovato-cylindrica mucronata. ☉ In Peruvia (Pav. in h. Brit. Mus. et Berol.), nemoribus Chinchao, Pozuzo et Macora (Pav. in flor. l. c.). Brasiliâ boreali prope San Gabriel da Cachoeira ad Rio Negro (Spruce n. 2221 in h. DC.), Margaritiva S. Angra dos Reys (Pohl n. 47921 5041 in h. Vindob.), sylvis aboriginibus ad Ilheos (Mart. in h. monac.), Guyana gallica (Leprieux in h. Deless.), Cayenna (Martin in h. Brit. Mus.), Surinam (Miq. syst. p. 57). Nova Andalusia (Humb. et B. nov. gen. v. 1. p. 65 et Humb. in h. Willd. n. 728!), St. Thomas (Friederichsthal in h. Lenorm.), St. Vincent ubi copiose in rupestribus et ad arbores ad orientem montis Morne Souffrier alt. 2000 ped. (Thoms. in Linn. Trans. v. 5. p. 203. t. 21. fig. 2), Sto-Domingo (Poiteau in h. Willd. n. 727!). Piper scandens Vahl enum. v. 1. p. 346. Pep. repens Kunth! l. c. et synops. v. 1. p. 118. P. Myosuros h. Willd. P. Myosurus A. Dietr. sp. v. 1. p. 159. P. repens A. Dietr. l. c. P. Guildingiana A. Dietr. sp. v. 1. p. 162. P. reniformis Miq. in Mart. flor. Bras. fasc. 11. p. 19. Acrocarpidium repens Miq. l. syst. p. 57, illustr. p. 10. t. 3. fig. b in Linn. v. 20. p. 118 et in Nederl. Kruidk. v. 1. p. 220. Acroc. pulicare Miq. syst. p. 57. Acroc. Guildingianum Miq. in Hook. Lond. journ. bot. v. 4. p. 412. P. pulicaris Opiz in Reliq. Hænk. fasc. 3. p. 162. t. 30. fig. 1. Piper bracteatum Thoms. l. c. Pep. repens Griseb. flor. w. Ind. p. 165. Herba repens vel scandens. Caulis filiformis, puberulus. Limbi 0,015 longi. Petioli 0,015-0,025 longi. Amenta 0,02 longa. (v. s.)

β, *fertilior*, foliis reniformibus vel rotundatis apice subattenuatis supra pubescentibus ciliolatis subcoriaceis opacis 3-nerviis, petiolo pubescente. amentis axillaribus. ☉ In Peruvia orientali prope Tarapoto (Spruce n. 3953 in h. DC.), Ecuador (Fraser in h. DC.), Guyana gallica (Sagot n. 539 in h. Kew, et DC.), Portorico (Blaumer n. 204 in h. DC.), Cuba (Wright n. 517 in h. DC.), Pep. repens Griseb. l. Cat. pl. Cub. p. 64. (v. s.)

206. *P. CASARETTI* (Cas. DC. in Seem. journ. bot. 1866. p. 141), foliis longiuscule petiolatis subrotundis apice rotundatis basi leviter subattenuatis rotundisve supra adpressis puberulis subtus glabris membranaceis vel rigidulo-membranaceis subopacis obsolete pellucido-punctulatis reticulato-nervulosis 5-nerviis, centrali nervo ad apicem usque lateralibus supra medium exterioribus ad medium usque ductis, nervulo juxta marginem currente, petiolo adpresso puberulo, amentis oppositifoliis folia æquantibus filiformibus densifloris, pedunculo petiolum æquante adpresso puberulo, bractea rotundata glandulosa centro peltata subsessili, ovario semimmerso oblongo acuminato, stigmatē puberulo supra medium inserto. ☉ In Brasiliâ prope Rio Janeiro (Casaretto n. 1041 in h. DC.). Herba procumbens e nodis radicans. Ramuli adpressi puberuli, filiformes. Limbi 0,025 longi latique. Petioli 0,015 longi. (v. s.)

207. *P. PSEUDO-PELTOIDEA*, foliis longe petiolatis subrotundato-ovatis apice acuminatis, acumine acuto, basi rotundatis cordulatisve supra glabris subtus



piloso-pubescentibus rigidulo-membranaceis subpellucidis pellucido-punctulatis enervulosis 5-nerviis, nervis subtus albicantibus, centrali nervo ad apicem usque lateralibus duobus fere ad apicem exterioribus supra medium ductis, nervulo juxta marginem currente, petiolo pubescente, amento terminali densiflora folia æquante, pedunculo pubescente petiolum fere æquante, bractea rotundata centro peltata subsessili, ovario impresso apice mucronulato subscutellato. ☉ In Venezuela (Fendler n. 1150  $\beta$ ! in h. DC.). Caulis scandens, e nodis radicans, piloso-pubescent. Limbi 0,03-0,05 longi, 0,035-0,04 lati. Petioli 0,03-0,07 longi. (v. s.)

208. *P. PELTOIDEA* (Kunth! in H. et B. nov. gen. v. 1. p. 61. t. 7), foliis longe petiolatis basi ima subpeltatis ovato-acutis vel subrotundo-ovatis apice subprotracto-acutis membranaceis subpellucidis supra glabris subtus puberulis margine ciliolatis 5-nerviis nervuloque juxta marginem extenuatum currente, petiolo pubescente, amentis terminalibus. bractea rotundata centro peltata, ovario impresso apice oblique rostrato, stigmate discoideo infra rostrum inserto. ☉ In Venezuela prope coloniam Tovar (Fendler n. 1150! in h. DC.), ad saxa et arborum truncos prope Bordonos et Cumana (Kunth! l. c. et Humb. in h. Willd. n. 708!). — Kunth synopsis. v. 1. p. 114, A. Dietr. sp. v. 1. p. 150, Miq.! syst. p. 187. *Piper peltioideum* Roem. et Sch. syst. v. 1. p. 324, Spreng. syst. v. 1. p. 118. Herba repens, scandens, e nodis radicans. Caulis puberulus. Limbi 0,06 longi. Petioli 0,03-0,07 longi. (v. s.)

209. *P. CARDIOPHYLLA* (Cas. DC. in Seem. Journ. 1866, p. 141), foliis petiolatis e basi cuneata obovatis apice rotundatis apice imo emarginatis utrinque glabris apicem versus ciliolatis membranaceo-rigidulis subopacis parce nervulosis 5-nerviis, centrali nervo ad apicem usque lateralibus 2 supra medium ductis exterioribus subtilioribus, nervulo juxta marginem apice currente, petiolo glabro, amento terminali densiflora filiformi folia æquante longiuscule pedunculato, pedunculo glabro petiolum fere duplo superante, bractea rotundata centro peltata subsessili, ovario impresso ovato apice suboblique complanato rostrato, stigmate minuto subattico. ☉ In valle Del Conca Novæ Granatæ (Triana! in h. suo). Herba suberecta, basi decumbens radicans. Caulis 0,1 altus, glaber. Limbi 0,02-0,035 longi, 0,012-0,02 lati. Petioli 0,007 longi. (v. s.)

210. *P. SYLVATICA*, foliis petiolatis subrhombeo-lanceolatis interdum ellipticis utrinque acutis raro obovatis utrinque glabris nigro-punctatis margine brevissime et tenuissime ciliolatis membranaceo-subcoriaceis subpellucidis reticulato-nervulosis et 5-nerviis nervulo juxta marginem apice currente, petiolo puberulo, amentis solitariis axillaribus vel terminalibus folia triplo superantibus, bractea rotundata centro peltata, ovario subimmerso apice breviter rostellato, stigmate minuto apice oblique inserto. ☉ In Peruvix Andibus inferioribus prope Cuchero (Præpfig! in h. Boiss.). Caulis erectus, pubescens. Limbi 0,06 longi. Petioli 0,015 longi. Amenta ante anthesin 0,16-0,2 longa. (v. s.)

211. *P. SUMIDORIANA*, foliis longiuscule petiolatis subovato-lanceolatis apice acuminatis, acumine acuto, basi attenuato-acutis utrinque glabris membranaceis subpellucidis 5-nerviis, centrali nervo validiore ad apicem usque ducto sursum nervulos utrinque 2 mittente, lateralibus utrinque 2 alte ductis subtilioribus, nervulo juxta marginem apice currente, petiolo glabro, amentis terminalibus filiformibus densifloris folia duplo superantibus, pedunculo quam petiolus brevior, bractea rotundata centro peltata breviter pedicellata, ovario impresso apice scutello sursum acuto aucto, stigmate medio scutelli inserto. ☉ In Brasilia ad Sumidorio (Langsdorff! in h. Petrop.). Herba glabra e nodis radicans. Limbi 0,07 longi, 0,02 lati. Petioli 0,025 longi. (v. s.)

212. *P. PROCUMBENS* (Cas. DC. in Seem. Journ. 1866, p. 141), foliis longissime petiolatis rotundis vel subovato-rotundis utrinque glabris subcoriaceis 5-nerviis, centrali nervo ad apicem usque ducto, lateralibus 2 fere ad apicem ductis exterioribus tenuioribus, amento terminali, pedunculo petiolum subæquante, bractea rotundata centro peltata breviter pedicellata, ovario apice acuto breviter aucto, stigmate medio scutelli inserto punctiformi, bacca ovato-attenuata mucronulata. ☉ In Peruvia orientali prope Tarapoto (Spruce

n. 4279! in h. Kew.). Herba procumbens. Limbi 0,065 longi, 0,07 lati. Petioli 0,08 longi. Pedunculi 0,045 longi. Amenta 0,05 longa, 0,003 crassa. (v. s.)

β, *brachystachya*, amentis minoribus, bacca ovato-cylindrica apice rostrata Prope Tarapoto (Spruce n. 4279 a! in h. Kew. (v. s.)

213. P. *FLEXICAULIS* (Wawra bot. Ergeb. p. 134. t. 19. fig. A), foliis petiolatis ovato-ellipticis obtusiuscule acuminatis basi breviter acutatis glabris vel junioribus apice puberulis supra saturate viridibus subtus pallidioribus opacis 3-5-nerviis, centrali nervo ad apicem usque ducto subtus prominente, lateralibus nervis subtilioribus, amentis axillaribus sæpe geminatis ternisve, pedunculo petiolum æquante, bractea rotundata excentrice peltata, ovario impresso acutiuscule producto, stigmatibus antice. ☉ In horto Schœnbrunn culta. Succulentia, ramosa. Caules pedales, glabri. Limbi 0,04 longi, 0,021 lati. Petioli 0,005 longi.

β, *microphylla*, foliis quam in præcedente plus quam triplo brevioribus. In h. Schœnbrunn culta. (Ex Wawra l. c.)

214. P. *IONOPHYLLA* (Griseb. ! pl. Wright. pars 1. p. 175), foliis longe petiolatis late deltoideis apice acutis basi obtusis vel subcordatis supra adpresse pubescentibus subtus pilosis fusco-punctulatis vel glabris membranaceis vel subpellucidis 3-nerviis, nervis parum conspicuis, nervulo juxta marginem currente, petiolo pubescente, amentis axillaribus, bractea rotundata centro peltata, ovario immerso apice oblique subtiliter rostellato, stigmatibus pilosis infra rostellum inserto. ☉ In Venezuela inter Maracai et Choroni alt. 6500 ped. (Fendler n. 2405! in h. DC.), ins. Cuba (Wright n. 516! in h. DC.), St.-Thomas (h. DC.). Pep. ionophylla Griseb. ! cat. pl. Cub. p. 64. Herba repens. Limbi 0,035 longi. Petioli 0,02 longi. Pedunculi 0,025 longi. (v. s.)

215. P. *SYLVESTRIS* (Cas. DC. in Seem. journ. 1866, p. 142), foliis longe petiolatis ellipticis vel ovato-attenuatis utrinque acutis et pubescentibus rigidulis subopacis non reticulato-nervulosis 3-nerviis, centrali nervo ad apicem usque ducto lateralibus interdum subtilissimis, amentis axillaribus solitariis densifloris longissime pedunculatis, pedunculo puberulo, bractea rotundata centro peltata, ovario immerso, stigmatibus carnosulo-puberulo subantice, bacca cylindrica apice rostrata. ☉ Ad basin Cordillerarum (Jameson! in h. Kew.). Herba repens. Caulis filiformis pubescens. Limbi 0,01 longi, 0,007 lati. Petioli 0,01 longi. Pedunculi 0,025 longi. Amenta 0,007 longa. (v. s.)

216. P. *CHORONIANA* (Cas. DC. in Seem. journ. 1866, p. 141), foliis longe petiolatis ad  $\frac{1}{2}$  longitud. a basi peltatis ovato-rotundatis vel omnino rotundis apice breviter protractis coriaceis margine ciliatis, centrali nervo ad apicem usque ducto subtus conspicuo, cæteris nervis fere inconspicuis, petiolo glabro, amentis axillaribus densifloris, bractea rotundata centro peltata, ovario semiimmerso peltatis gibboso apice rostellato, stigmatibus minuto antice. ☉ In Venezuela inter Maracai et Choroni (Fendler n. 2402! in h. DC.). Herba procumbens. Limbi 0,1-0,12 longi, 0,09-0,1 lati. Petioli 0,09 longi. (v. s.)

217. P. *HERNANDIÆFOLIA* (A. Dietr. sp. v. 1. p. 157), foliis longe petiolatis ad  $\frac{1}{2}$ - $\frac{1}{4}$  longitud. a basi peltatis ovato vel lato-ovatis apice abrupte acutecue acuminatis basi lato-rotundatis præsertim subtus et ad nervum centralem patule et subretrosum pubescentibus supra glabriusculis coriaceis, centrali nervo subtus prominente, nervulis obsolete, petiolo patule et subretrosum pubescente, amentis axillaribus longe pedunculatis (pedunculo ad  $\frac{1}{2}$  longitud. unibracteato simplici vel bifido) brevibus densifloris, stigmatibus antice, bacca emersa ovata longe rostrata verrucosa. ☉ In ins. St.-Vincent (Goulding in h. Hook. fide Miq. in Hook. Lond. journ. bot. v. 4. p. 428). — Miq. syst. p. 72 et in Hook. l. c. Piper hernandiæfolium Vahl enum. v. 1. p. 344, Spreng. syst. v. 1. p. 118. Radicans, repens. Caulis patule et subretrosum pubescens. Limbi 0,04-0,08 longi, fere 0,05 lati. Petioli 0,04-0,05 longi. (Ex Miq. l. c.)

218. P. *FLAVESCENS*, foliis longe petiolatis ovato-acutis coriaceis subtus pubescentibus margine ciliatis supra glabris conspicue uninerviis, centrali nervo ad apicem usque currente flavescente, amentis terminalibus longe peduncu-

latis, bractea rotundata centro peltata, ovario basi immerso ovato-acuto, stigmate carnosulo antico, bacca rostellata patente. ☉ In Peruvia (Pav. in h. Boiss.). Ramuli pubescentes. Limbi 0,04 longi, 0,02-0,03 longi. Pedunculi 0,065 longi. (v. s.)

219. *P. CILIARIS* (Cas. DC. in Seem. journ. 1866, p. 144), foliis inferioribus longe superioribus modice petiolatis elliptico-rotundis utrinque obtusis utrinque glabris margine subfusce ciliatis coriaceis opacis, petiolo ciliato, amentis terminalibus solitariis longe pedunculatis, pedunculo petiolum superiorem subsuperante pilosulo, amentis maturis folia subæquantibus, bractea rotundata centro peltata breviter pedicellata, ovario impresso antice et apice triangulariter subscutato complanato apice rostrato, stigmate minuto carnosulo medio scutelli inserto. ☉ In Novæ Granatæ prov. Buenaventura alt. 100 m. (Triana n. 59! in h. DC.). Suffruticulus procumbens e nodis radicans. Caulis filiformis, pilosulus. Limbi 0,02-0,03 longi, 0,02-0,023 lati. Petioli inferiores 0,04, super. 0,013 longi. (v. s.)
220. *P. CILIOSA* (Cas. DC. in Seem. journ. 1866, p. 144), foliis longe petiolatis ellipticis vel elliptico-rotundatis utrinque obtusis et glabris margine incanescens ciliatis coriaceis opacis, petiolo glabro, amentis terminalibus solitariis longe pedunculatis maturis folia subsuperantibus filiformibus densifloris, pedunculo glabro petiolum æquante, bractea rotundata centro peltata breviter pedicellata, ovario semimmerso oblongo apice rostrato, stigmate antico supra medium inserto. ☉ In Novæ Granatæ prov. Barbacoas ad viam Tuquerras alt. 800 m. (Triana n. 60! in h. DC.). Planta repens, e nodis radicans. Caulis glaber, tetragonus. Limbi 0,06 longi fere 0,05 lati. Petioli 0,04 longi. (v. s.)

*Species affines, non satis notæ.*

221. *P. EMARGINELLA*, foliis petiolatis orbicularibus apice emarginatis seu quasi obcordatis glabris margine tantum hirtellis saturate viridibus lineam circiter longis latisque, petiolo limbum fere æquante glabro, amentis terminalibus solitariis filiformibus semiuncialibus glabris floribus distinctis diandris. ☉ In Guadaloupa. *Piper emarginellum* Sw. (fide Wickstr. plant. Guad. p. 56). Plantula cæspitosa. Radix filiformis. Caulis repens filiformis. (Ex Wickstr. l. c.)
222. *P. OBLIQUA* (Ruiz et Pav. flor. Peruv. v. 1. p. 32. fig. c), foliis breviter petiolatis oblique oblongo-ovatis apice attenuatis acutis glabris carnosis, centrali nervo ad apicem usque ducto fere tota longitudine nervos patulo-subadscendentes alternos utrinque 5 mittente, amentis axillaribus terminalibusque densifloris folia superantibus. ☉ In nemoribus Cuchero, Pozuzo et Pillao (Ruiz et Pav.). — A. Dietr. sp. v. 1. p. 148 (exclus syn. L.). Miq. syst. p. 96. *Piper acuminatum* Sw. obs. p. 20. ? Vahl enum. v. 1. p. 342 (exclusis synonym. Plum. et L.). Willd. enum. p. 47, Spreng. syst. v. 1. p. 115, Bot. mag. 1822, Pers. synopsis. p. 35. Radix fibrosa. Caulis teres, geniculatus. (Ex R. et P. l. c.)
223. *P. PURPUREA* (Ruiz et Pav. flor. Peruv. v. 1. p. 32. t. 49 fig. b), foliis petiolatis ovatis apice acuminatis acutis supra venis purpureis notatis subtus purpureis, amentis axillaribus terminalibusque filiformibus purpureis folia triplo superantibus pendulis densifloris, pedunculo petiolum superante. ☉ In montibus Chinchao umbrosis et locis saxosis. — A. Dietr. sp. v. 1. p. 151, Miq. syst. p. 85. *Piper purpureum* Vahl enum. v. 1. p. 343. Planta fere pedalis. Caulis flexuosus, summitate divisus, inferne nudus, purpureus, teres, erectus. Rami patentes. Limbi 0,07 longi, 0,058 lati. Petioli 0,005 longi. (Ex R. et P. l. c.)
224. *P. MONOSTACHYA* (Ruiz et Pav. flor. Peruv. v. 1. p. 33), foliis petiolatis obovatis ovatisve enerviis supra viridibus subtus purpureis, lineis quinque obscure purpureis longitudinalibus lateralibus duobus brevioribus. ☉ In montibus Chinchao supra arborum vetustorum truncos. — A. Dietr. sp. p. 1. p. 152, Miq. syst. p. 86. *Piper monostachyon* Vahl enum. v. 1. p. 343. Planta bipalmaris. Caulis teres glaber radicans. (Ex R. et P. l. c.)
225. *P. SALIGNA* (Kunth in H. et B. nov. gen. v. 1. p. 62), foliis subsessilibus approximatis lanceolatis utrinque angustatis obtusis simpliciter venosis car-

nosis glabris, amentis crassitie pennæ passerinæ pedicellatis densifloris quam folia brevioribus. In declivitate occidentali montis Quindiu in ripa fluminis Boquira alt. 920 hexap. — Kunth synops. v. 1. p. 115. Piper salignum Rœm. et Sch. syst. v. 1. p. 325. Pep. saligna A. Dietr. sp. v. 1. p. 144, Miq. syst. p. 94. Caulis erectus, ramosus, pedalis, carnosus. Folia subbipollicaria. Amenta semipollicaria.

226. *P. DIVERSIFOLIA* (Kunth in H. et B. nov. gen. v. 1. p. 64), foliis petiolatis subrotundo-obovatis apice basique obtusis supra glabris subtus adpressè pilosis ciliolatis subpellucidis membranaceis parce nervulosis et quinquenerviis, centrali nervo ad apicem usque ducto, lateralibus nervis alte supra medium ductis, infimis subtilioribus brevioribus, nervulo juxta marginem currente, amento filiformi folium duplo superante densifloro, bractea rotundata centro peltata subsessili, bacca ovata basi immersa apice rostrata. Ad saxa in deversis Andium Pastoensium inter Paramo de Apohte et Sebondy alt. 760 hexap. (Kunth l. c. et Bonpl. in h. Francov.). — Kunth synops. v. 1. p. 117, A. Dietr. sp. v. 1. p. 158, Miq. syst. p. 81. Piper diversifolium Rœm. et Sch. syst. v. 1. p. 333, Spreng. syst. v. 1. p. 117. Limbi 0,025 longi, 0,023 lati. (v. s.)
227. *P. MOLLE* (Kunth in H. et B. nov. gen. v. 1. p. 65), foliis petiolatis ovatis apice acutiusculis incano-pubescentibus subtus sparse mollior pubescentibus carnosus subcoriaceis subpellucidis trinerviis nervis ramulosis nervulis raris, petiolo mollior pubescente. ☉ In regni Novo-Granatensis devexis montis Quindiu alt. 980 hexap. (Kunth l. c. et Humb. in h. Willd. n. 729! et in h. Berol.). — Kunth synops. v. 1. p. 118, A. Dietr. sp. v. 1. p. 155. Piper molle Rœm. et Sch. syst. v. 1. p. 326. Acrocarpidium molle Miq. syst. p. 62. Caulis repens radicans pilis incanis patulis vestitus. Limbi 0,015 longi, 0,011 lati. Petioli 0,003 longi. (v. s.)
228. *P. BIUNCIALIS* (Kunth in H. et B. nov. gen. v. 1. p. 65), foliis petiolatis subrotundato-ovatis obtusis basi rotundatis tenuissime membranaceis glabris, amentis oppositifoliis, pedunculo glabro. In temperatis Andium Novo-Granatensium valle de Quilquase. — A. Dietr. sp. v. 1. p. 158, Miq. syst. p. 82. Piper biunciale Rœm. et Sch. syst. v. 1. p. 322. Spreng. syst. v. 1. p. 117. Caulis bipalmaris, dichotomus, glaber. Folia 4-5 lin. longa. Petioli 2-3 lin. longi. Differt a *Pep. pellucida* præter puritatem foliorum nervo medio tantum venosâ cauleque dichotomo. (Ex Kunth l. c.)
229. *P. PERFORATA* (Opiz in Reliq. Haenk. fasc. 3. p. 162. t. 30. fig. 2), foliis petiolatis lanceolatis carnosis pilosis pellucido-punctatis trinerviis, petiolo piloso, amentis axillaribus terminalibusque pedunculatis filiformibus solitariis geminisve folia multum superantibus. ♀ In Mexico. — Miq. syst. p. 93. Caulis herbaceus, ramosus, pilosus. Limbi 0,03 longi, 0,01 lati. Petioli 0,001 longi. (Ex Opiz l. c.)
- β, *glabrata* (A. Dietr. sp. v. 1. p. 145), foliis glabris. In Mexico.
230. *P. PUMILA* (Opiz in Reliq. Haenk. fasc. 3. p. 163. t. 30. fig. 3), foliis petiolatis ovatis obtusiusculis trinerviis pilosis carnosis, petiolo pubescente, amentis axillaribus terminalibusque filiformibus solitariis pedunculosis foliisque superantibus. ♀ In Peruviae montanis Huanocensibus. — A. Dietr. sp. v. 1. p. 158, Miq. syst. p. 93. Caulis brevissimus erectus pilosus, basi radicans. Limbi 0,02 longi, 0,015 lati. Petioli 0,006 longi. (Ex Opiz l. c.)
231. *P. FURCATA* (Opiz in Reliq. Haenk. fasc. 3. p. 164), foliis petiolatis ovatis vel rotundato-ovatis obtusis glauris ciliatis 3-7-nerviis, amentis oppositifoliis pedunculatis conjungatis distachyis, pedunculo glabro. ♀ In Peruviae montanis. — A. Dietr. sp. v. 1. p. 158, Miq. syst. p. 84. Caulis herbaceus, carnosus, repens, glaber. (Ex Opiz l. c.)
232. *P. ROXBURGHIANA* (A. Dietr. sp. v. 1. p. 144), foliis sessilibus oblique lanceolatis glaberrimis, amentis oppositifoliis breviter pedunculatis. In Archipelago Malayano. P. Roxburghianum Rœm. et Sch. syst. mant. 1. p. 242. Piper angustifolium Roxb. Fl. Ind. v. 1. p. 163 (non Ruiz et Pav.). Pep. angustifolia Miq. syst. p. 109. Caulis scandens, repens, glaber. (Ex Roxb. l. c.)
233. *P. MYRTIFOLIA* (A. Dietr. sp. v. 1. p. 147), foliis rhombeo-oblongis basi

- acutiusculus succulentis enervulosis subtus trinerviis, petiolo brevi, pedunculis longitudo petiolorum axillaribus solitariis terminalibus geminis, amentis filiformibus acutis. In ins. Sta Cruz. Piper myrtifolium Vahl enum. v. 1. p. 341. *Pep. myrtifolia* Miq. syst. p. 92. Caulis pedalis, simplicissimus, purpurascens, superne foliosus, flexuosus. Folia pollicaria. Amenta bi-tripollicaria. (Ex Vahl l. c.)
234. *P. MACROSTACHYA* (A. Dietr. sp. v. 1. p. 149), foliis petiolatis remotis ovato-oblongis acuminatis obscure multinerviis, amentis axillaribus gracilibus, pedunculo quam petiolus brevior, bacca oblonga sessili vix magnitudine seminis Mili. In Cajenna (Rich. Æde Vahl). — *Miq. syst.* p. 91. Piper macrostachyon Vahl enum. v. 1. p. 341. Caulis erectus crassitie pennæ oblongum, glaber uti tota planta. Folia subtripollicaria. Petioli semipollicares. Amenta pedalia. (Ex Vahl. l. c.)
235. *P. CONCINNA* (A. Dietr. sp. v. 1. p. 164), foliis patulo-pendulis cordatis obtusis petiolis decurrentibus lucidis quinquenerviis internodio his laminae petioli brevioribus, ramulis gracilibus flexuosis.  $\S$  In Jamaica. — *Miq. syst.* p. 82. Piper concinnum Haw. succ. rev. p. 198. Nuda, lævis. (Ex Haw. l. c.)
236. *P. ? PÆPPIGH* (Miq. syst. p. 84), foliis petiolatis rotundato-ovatis ovatisve apice breviter acuminatis acutis basi rotundatis vel subattenuato-rotundatis utrinque glabris ciliatis, ciliis setulosis, membranaceo-rigidulis regulariter nervulosis, nervulis parallelis transversis, novemnerviis, nervis centralibus a paulo supra basin solutis, amentis in specim. deficientibus, petiolo late amplexicauli hirtello.  $\odot$  In Peruvia subandina (Pæppig in h. Berol.). Caulis glaber, a nodis cristas longasque radices fibrilloso-ramosas emittens. Limbi 0,09 longi, 0,065 lati. Petioli 0,01 longi. Specieservatione Piperas referens. (v. s.)
237. *P. KRAWANGENSIS* (Miq. syst. p. 108. et Fl. Ind. bat. v. 1. p. 2. p. 434), foliis longius petiolatis superioribus ovato-oblongis obtusioribus trinerviis, amentis folio brevioribus. In cacumine montis Burangrang et prope montem Ignivomum Tankwan-prah, in provincia Krawang Java ad arborum truncos vetustiss. Piper lævifolium Bl. in Verh. der Bat. Genoots. v. 9. p. 200 et 11. p. 229, var.  $\gamma$ . latifolium *Miq. comm. phyt.* p. 57.
238. *P. NERVOSO-VENOSA* (Miq. syst. p. 104), foliis alternis oppositisve petiolatis inferioribus minoribus suborbiculatis superioribus majoribus ovatis ovalibusve omnibus utrinque glabris trinerviis reticulatis subtus albicantibus, amentis axillaribus terminalibusque pedunculatis solitariis geminisve vel ternis, pedunculo petiolum fere æquante, bractea reniformi persistente, ovario immerso ovoideo, stigmate tenuissime penicillato. In regionibus altioribus montium Salak et Gédé insulæ Java super truncos putridos et inter saxa basaltica. — *Miq. Flor. Ind. Bat. v. 1. pars 2. p. 432.* Piper umbrosum Bl. in Verh. der Bat. Genoots. p. 201. *P. nervoso-venosum* ejusdem l. c. v. 9. p. 225, enum. pl. Jav. fasc. 1. p. 74. Micropiper nervoso-venosum *Miq. comm. phyt.* p. 57. t. 8. fig. d. Limbi inferiores circ. 0,015 longi, supremi 0,04 longi, 0,02 lati. Petioli 0,01 longi. Pedunculi 0,03-0,04 longi. (Ex *Miq. l. c.*)
239. *P. TITHYMALOIDES* (A. Dietr. sp. v. 1. p. 152), foliis subovatis acutiusculis basi acutis crassis coriaceis exsiccatis supra novemnerviis, nervis suboppositis, petiolo subtus carinato, pedunculis solitariis terminalibus geminis, spathis duobus lanceolatis caducis, bractea ovata.  $\mathcal{Z}$  In Santo Domingo. — *Miq. syst.* p. 192. Piper tithymaloides Vahl enum. v. 1. p. 340. Caulis inferne radicans, crassitie pennæ aserina. Folia bi-tripollicaria. Petioli pollicares. Pedunculi tripollicares. Amenta crassitie filii emporetici tenuioris. (Ex Vahl l. c.)
240. *P. MARGARITIFERA* (Hook. icon. plant. v. 1. p. 91), foliis petiolatis spatulato-cuneiformibus apice breviter acuminatis acutis subcoriaceis supra ad nervos parce subtus confertius hirtellis tri-quinquennerviis, nervis 3 centralibus magis conspicuis subtus prominulis, nervulo juxta marginem currente, petiolo flavo-verruculoso, amentis in axillis cymose confertis singulo squama parva fulto pedunculo densifloro, pedunculis hirtellis, bractea rotundata centro peltata sessili, cæteris nondum efformatis.  $\S$  In ins. Juan Fernandez (Bertero n. 14931 in h. DC.). — *Miq. syst.* p. 112. ill. p. 16. t. 10. E radice

fasciculata caulis lignosus, rectus, 0,25 altus, teres, circ. 0,003 crassus, tuberculis crassiusculis instructus. Limbi 0,05-0,07 longi, 0,02-0,025 lati. Petioli 0,005 longi. (v. s.)

241. *P. PORTOBELLENSIS* (Beurling in flor. Portobell. p. 147), foliis brevissime petiolatis e basi subobtusa oblongo-acuminatis obscure quintuplinerviis, amentis oppositifoliis subulato-filiformibus pedunculatis pendulis. ☉ In umbris ins. Portobello. Herbacea, caulescens, subsimplex, ad nodos radicans, glabra. Folia 2-2½ poll. longa, 1-½ poll. lata. Amenta circ. 4 poll. longa.
242. *P. PSEUDO-RHOMBEA*, foliis longe petiolatis elliptico-lanceolatis basi acuminatis margine integerrimis vel subserrulatis glabris membranaceo-pellucidis reticulato-nervulosis trinerviis nervuloque juxta marginem currente, amentis terminalibus pedunculatis solitariis glabris maturitate sublaefloris, bractea rotundata centro peltata, bacca globosa lævi. ☉ In ins. Ceylan prov. centrali (Thwaites n. 2172 fide Thwaites enum. p. 292). Pep. Courtallensis var. β. Thwaites l. c. Herba repens scandens. Limbi vix 0,05 longi. Petioli circ. 0,01 longi. Pedunculi 0,015 longi. (Ex Thwaites l. c.)

#### B. Folia opposita.

##### 1° Folia penninervia.

243. *P. RECURVATA* (Miq. syst. p. 141), foliis breviter petiolatis oppositis vel supremis quaternis inferioribus minoribus subrotundis superioribus ellipticis apice obtusiusculis integerrimis vel apicem versus obsolete crenulatis supra glabriusculis subtus imprimis ad nervum hirsutiusculis utrinque imprimis subtus nigro-punctulatis, centrali nervo nervulos paucos parum ramulosos mittente, petiolo hirsuto, amentis terminalibus, pedunculo hirsuto petiolum superante, bacca ovata. ☉ In sylvis elevatioribus montis Gédé in ins. Java ad truncos arborum (Bl. in h. Lugd. Bat. fide Miq. l. c.), sylvis montis Patusha (Miq. plant. Reinw. in Linn. v. 21. p. 480), truncis arborum in sylvis montis Javæ alt. 5000 ped. (Miq. l. c.). Philippinis (Cuming n. 1920 in h. Hook. — Miq. in flor. Ind. Bat. v. 1. pars 2. p. 435 et in Hook. Lond. journ. bot. v. 4. p. 423 et ll. cc. *P. recurvatum* Bl. in Verh. der Bat. Gen. v. 11. p. 233. f. 38. Suffruticulus humilis, repens, e nodis parum tumidis radicans, pilis mollibus sparsis vestitus. Limbi 0,02 longi, 0,01 lati. Petioli 0,005 longi. (Ex Miq. l. c.)

β, *minor*, foliis e basi cuneata rhombeo-ellipticis apice obtusiusculis majoribus ultrapollaribus. In Borneo australi (Miq. Ann. bot. Mus. Lugd. Bat. v. 1. fasc. 5), Java monte Pangerango (Miq. l. c.).

244. *P. DECAISNEI* (Miq. syst. p. 183), foliis longiuscule petiolatis rotundatis vel ovato-rotundatis rameis minoribus omnibus carnosius coriaceis glandulis obscuris punctatis glabris tenuissime uninerviis nervuloso-costulatis vel subtrinerviis, amentis axillaribus oppositis vel internodiis summis valde contractis subcymose confertis longe pedunculatis filiformibus, bacca semimmersa ovato-globosa subverruculosa. ☉ In ins. Bourbon (h. Mus. Par. n. 705, fide Miq. l. c.). Herba glabra, decumbens. Caulis teres, lævis. Limbi 0,005-0,01 longi. Petioli 0,003-0,004 longi. (Ex Miq. l. c.)
245. *P. ELLIPTICA* (A. Dietr. sp. v. 1. p. 167), foliis breviter petiolatis ellipticis vel obovato-subrotundis rotundatisve utrinque glabris atque nigro-punctulatis membranaceis subpellucidis, centrali nervo ad apicem usque ducto sparse alterne ramuloso, nervulo juxta marginem currente, petiolo glabro, amentis terminalibus axillaribusve longiuscule pedunculatis folia duplo triplove superantibus filiformibus subdensifloris, pedunculo glabro nigro-punctulato, bractea rotundata centro peltata subsessili, ovario subimpresso vel impresso rhachique nigro-punctulatis, stigmate penicellato apice inserto. ☉ In ins. Maurice (Sieber n. 377! et Bouton! in h. DC. Heyne in h. Willd. n. 750!), Bourbon (Boivin! in h. Boiss. et h. Mus. Par. n. 636! in h. Le-norm.). — Link Jahrb. 1. 3. p. 64, fide R. et Sch. i. c. Miq. l. syst. p. 127. *P. ellipticum* Lam. ill. p. 83. Vahl enum. v. 1. p. 350, Spreng. syst. v. 1. p. 119. *P. mauritianum* R. et Sch. Mant. v. 1. p. 245. Planta digitalis vel subpalmaris,

basi decumbens, e nodis radicans. Caulis glaber, 0,003 circ. crassus. Limbi 0,01-0,02 longi, 0,007 lati. Petioli 0,003 longi. Pedunculi 0,006 longi. (v. s.)

### 2° Folia multiplinervia.

246. *P. MARGINATA* (Miq.† syst. p. 129), foliis raro ternis vel apice ramorum quaternis obovatis vel rotundatis succulentis pellucido-membranaceis utrinque nigro-punctulatis glabrisque apice subciliolatis quintuplinerviis, centrali nervo ad apicem usque ducto fere e basi nervos utrinque 2 alte ductos quorum infimi sæpe obsoleti mittente vel trinerviis, nervulo obscuro juxta marginem corrente, petiolo glabro, amentis axillaribus terminalibusque longiuscule pedunculatis, pedunculo petiolum superante, bractea rotundata fusco-punctata centro peltata subsessili, ovario subimmerso rhachique fusco-punctulatis, stigmatibus minuto puberulo apice imo inserto. ☉ In ins. Bourbon (h. Mus. Par. n. 686! et Sieber n. 165! in h. DC. Sieber. n. 167! in h. Berol.) Pep. Perrottetiana Sieber in h. Berol. et h. Deless. Herba repens. Limbi 0,025 longi. Petioli 0,006-0,008 longi. (v. s.)

247. *P. PICHINCHÆ* (Cas. DC. in Seem. journ. of bot. 1866. p. 144), foliis breviter petiolatis glabris orbiculato-reniformibus apice obtusis basi in petiolum subdecurrentibus reticulato-nervulosis quintuplinerviis. ☉ In vallibus Pichinchæ (Jameson n. 747! in h. Kew.). Herba repens, e nodis radicans, glabra. Caulis teres. Limbi 0,005 longi. Petioli 0,0015 longi. (v. s.)

248. *P. UNDENINERVIA*. foliis breviter petiolatis subelliptico-lanceolatis apice longiuscule acuminatis, acumine acuto, basi attenuato-acutis utrinque glabris supra nigro-punctulatis subopacis firmulis undeninerviis nervis exterioribus subtilibus, petiolo glabro, amentis axillaribus terminalibusque florentibus folia æquantibus densifloris, pedunculo glabro petiolum paullo superante, bractea ovata centro peltata pedicellata, ovario impresso, stigmate subantico. ☉ In Peruvia (Pav.† in h. Boiss.). Caulis glaber teres pedalis. Limbi 0,1 longi, 0,05 lati. Petioli 0,006 longi. (v. s.)

### 3° Folia multinervia.

249. *P. SEPTEMNERVIS* (Ruiz et Pav. flor. Peruv. p. 31. t. 47. fig. c.), foliis oppositis rarius ternis breviter petiolatis lanceolatis apice acuminatis, acumine acuto, basin versus attenuato-acutis coriaceis septemnerviis, amentis terminalibus axillaribusque folia superantibus, bractea rotundata centro peltata brevissime pedicellata, ovario ovato, stigmate carnosulo apice imo inserto, bacca basi immersa vertice mucronata. ☉ In nemoribus ad Pozuzo supra saxa (R. et P. l. c.), Jamaica (Griseb. flor. w. Ind. p. 168, Purdie! in h. Kew.) et Fisch.† in h. Petrop. sine patria. — Griseb. l. c. Piper septemnerve Vahl enum. v. l. p. 349. Planta pedalis. Radix fasciculata, fibrosa. Caules plurimi erecti radicales, nodis purpurascenscentibus. Limbi 0,08 longi, 0,028 lati. Petioli 0,005 longi. (Ex R. et P. l. c.)

250. *P. SEEMANNIANA* (Miq. in Seem. bot. of Herald, p. 198. t. 31), foliis breviter petiolatis inferioribus elliptico-oblongis superioribus lanceolato-oblongis apice acuminatis basi acutis crassius coriaceis subtus pallidis, 7-5-nerviis, amentis axillaribus oppositis modice pedunculatis densifloris, ovario immerso nondum efformato, stigmate antico? ☉ In Volcano Chiriqui, Veraguas supra arbores (Seem. n. 896). Herba circ. pedalis. Ramuli subelongati. Petioli 2 lin. longi. Folia 2½-3 raro 4 poll. longa. Pedunculi 2-4 lin. longi.

251. *P. CILIATA* (Kunth! in H. et B. nov. gen. v. 1. p. 68), foliis oppositis summisve ternis petiolatis elliptico-oblongis utrinque acutis supra pubescentibus subtus ad nervos pubescentibus margine ciliolatis membranaceis subpellucidis pellucido-punctulatis 5-nerviis, centrali nervo ad apicem usque ducto, lateralibus 2 supra medium evanidis, exterioribus subtilibus nervulose juxta marginem corrente, petiolo ciliolato, amentis axillaribus solitariis, ovario emerso, stigmate discoideo-papilloso apice inserto. ☉ In Peruvia (Pav.† in h. Boiss.): convalli juxta cœnobium Caripense ad arborum truncos alt. 430 hexap. in Nova Andalousia (Kunth l. c. et Humb.† in h. Willd. n. 744!),

- regno Quitensi (Jameson n. 59! in h. Kew.), Antiochia et Trinitate (Griseb. l. c.) — Miq.! syst. p. 119, Griseb. pl. W. Ind. p. 168. Piper ciliatum R. et Sch. syst. v. 1. p. 333. *Pep. Quitensis* Miq.! in Hook. Lond. Journ. bot. v. 4. p. 421. Caulis hirtellus, 0,002 crassus. Limbi 0,03-0,05 longi. Petioli vix 0,02 longi. Pedunculi vix 0,02 longi. (v. s.)
252. *P. ARABICA* (Miq.! syst. p. 121), foliis petiolatis elliptico-rhomboides vel oblongo-rhomboides apice obtusiusculis basi cuneatis adpresse pubescentibus margine ciliatis fusco-punctatis subpellucidis reticulato-nervulosis et 5-nerviis nervulo juxta marginem currente, petiolo pubescente, amentis axillaribus terminalibusque folia paullo superantibus post anthesin subulaxifloris, pedunculo puberulo, bractea rotundata centro peltata brevissime pedicellata, ovario emerso, stigmatibus exiguo puberulo. ☉ In promontorio Bonæ Spei prope Natal (Guenzius! in h. Boiss. et Deless., Drège! in h. Petrop.). — Miq. ill. p. 18. t. 12. *Pep. Caffra* E. Mey.! pl. Dreg. Herba erecta circ. 0,3 alta. Caulis dense pubescens. Limbi 0,03-0,035 longi, 0,025 lati. Petioli 0,007-0,01 longi. (v. s.)
- β, *floribunda* (Miq.! syst. p. 122), foliis anguste rhombeo-ellipticis. Circa Tananarivo in Madagascar (Goudot! in h. Deless.). (v. s.)
253. *P. ROTUNDATA* (Kunth! in H. et B. nov. gen. v. 1. p. 67. t. 12), foliis petiolatis suborbiculato-reniformibus vel elliptico-suborbiculatis apice obtusis vel breviter protractis acutiusculis basi truncato-retusis membranaceis supra subtusque ad nervos hirtis margine ciliatis imprimis supra nigro-punctatis 5-nerviis, centrali nervo ad apicem usque ducto, exterioribus inconspicuis, nervulo juxta marginem currente, petiolo hirtio, amentis axillaribus terminalibusque densifloris interdum squamis fultis, pedunculo hirtio, bractea rotundata centro peltata subsessili, ovario emerso, stigmatibus minuto subantico, bacca ovato-acuta. ☉ In excelsis regni Novo-Granatensis inter Pansitara et vallem Yacanatu alt. 980 hexap. (Kunth l. c. et Humb. in h. Willd. n. 736!), Andibus quitensibus (Jameson n. 759! 59! in h. Kew.). — Kunth synopsis. v. 1. p. 119, Miq.! syst. p. 118. Piper rotundatum Rcm. et Sch. syst. v. 1. p. 333. *P. villosum* Rich! mss. in h. Francov. Caules caespitiosi procumbentes hirti. Limbi 0,015-0,03 longi. Petioli 0,015 longi. Pedunculi 0,015 longi. (v. s.)
254. *P. DINDYGULENSIS* (Miq.! syst. p. 122), foliis petiolatis inæqualibus summis ternis ellipticis obovato-rhombicis obovatisve vel radicalibus rotundatis utrinque puberulis membranaceo-subcoriaceis pellucidis reticulato-nervulosis et 5-nerviis, amentis axillaribus terminalibusque folia paullo superantibus filiformibus, pedunculo glabro petiolum æquante, bractea subrotundata centro peltata, ovario emerso, stigmatibus papilloso-puberulo apice inserto. ☉ In rupibus provinciæ Dindygul alt. 2500 ped. (Wall. n. 6663! in h. DC.). — Miq. ill. p. 19. t. 13, Wight icon. t. 1921. Piper ovalifolium Heyn! in Wall. h. Planta succulenta, erecta, 0,25 alta. Caulis ramulique puberuli. Limbi 0,045 longi. Petioli 0,008 longi. (v. s.)
255. *P. MAYPURENSIS* (Kunth! in H. et B. nov. gen. v. 1. p. 66), foliis longe petiolatis obovato-ellipticis apice obtusis subretusis basi æquali cuneatis carnosio-succulentis utrinque glabris coriaceis opacis 5-nerviis, nervis subtus prominulis, centrali nervo ad apicem usque ducto, lateralibus nervis 2 fere ad apicem usque exterioribusque alte supra medium ductis, petiolo glabro, amentis filiformibus subdensifloris, bractea rotundata centro peltata subsessili, ovario rhachi impresso, stigmatibus discoideo-puberulo apice imo inserto. ☿ In sylvis ad flumen Orinocco juxta El Raudal de Maypure (Humb. in h. Willd. n. 734!). — Kunth synopsis. v. 1. p. 118, Miq. syst. p. 135, ill. p. 20. t. 16. Piper maypurens Rcm. et Sch. syst. v. 1. p. 332. Suffructulus 1-2 pedalis (Miq. l. c.). Limbi 0,07 longi, 0,085-0,105 lati. Petioli 0,07 longi. (v. s.)
256. *P. PETIOLATA* (Hook. fil. in Trans. linn. soc. v. 20. p. 181), foliis parvis longe petiolatis late elliptico-ovatis vel rotundatis utrinque obtusis planis carnosius obscure 5-nerviis, amentis 2-4 terminalibus elongatis. ☉ In James Irs. Arch. Gallapagos. Glaberrima. Folia vix  $\frac{1}{2}$  unc. longa, remota. Petioli æquilong. Amentum 4 unc. longum.
257. *P. MARIANNENSIS*, foliis petiolatis elliptico-ovatis utrinque obtusis obtusius-



culisve utrinque glabris vel apicem versus margine ciliolatis membranaceis subpellucidis pellucido-punctatis 5-nerviis, centrali nervo ad apicem usque ducto, lateralibus ad medium usque ductis nervulo juxta marginem apice currente, petiolo glabro, amento terminali filiformi densifloro folia æquante, pedunculo glabro petiolum æquante, bractea rotundata centro peltata subsessili, ovario impresso ovato, stigmate oblique inserto. ☉ In insulis Mariannis (Gaudich. l in h. Berol.) *Pep. membranacea* Miq. syst. p. 120? (non Hook. et Arn.). Herba glabra, erecta, circ. 0,1 alta. Caulis 0,002 crassus, margine anguste extenuatus. Limbi 0,02-0,03 longi; 0,013-0,02 lati. Petioli 0,003 longi. (v. s.)

258. *P. MARMORATA* (Hook. fil. in Bot. mag. t. 5568), foliis petiolatis confertis ovatis apice acutis basi profunde cordatis carnosio-coriaceis utrinque glabris supra albo-marmoratis subtus pallidioribus 5-nerviis (in icone 7-nerviis), amentis apice caulis confertis squamis fultis folia æquantibus pedunculatis subdensifloris, bractea rotundata centro peltata breviter pedicellata, ovario glanduloso, stigmate decido oblique inserto. ☉ In Brasilia meridionali a Weir collect. et in horte Societ. hortic. Lond. culta. Herba robusta glaberrima. Caulis teres ramosus. Limbi 0,12 longi, 0,07 lati. (v. s.)

259. *P. CHILENSIS*, foliis interdum alternis summis oppositis petiolatis ellipticis utrinque obtusis apice aliquando brevissime protractis pellucido-membranaceis epunctatis supra glabris subtus ad nervos puberulis apicem versus ciliatis reticulato-nervulosis et 3-5-nerviis, nervulo juxta marginem currente, petiolo puberulo, amentis axillaribus solitariis pedunculatis, bractea rotundata centro peltata pedicellata, ovario post anthesin subimmerso glabro rostellato, stigmate globoso minuto antice inserto. ☉ In Chili (Lechler n. 3020! in h. Boiss.). Herba ad arborum truncos ascendens. Caulis puberulus. Limbi 0,025 longi. Petioli 0,008 longi. Pedunculi circ. 0,008 longi. (v. s.)

260. *P. PERROTETIANA* (Miq. syst. p. 146), foliis oppositis vel ternis modice petiolatis obovatis basi attenuatis vel inferioribus minoribus subrotundo-obovatis utrinque glabris nigro-punctatis membranaceo-rigidis, petiolo glabro reticulato-nervulosis et trinerviis vel subquinenerviis, centrali nervo validiore, cæteris nervis subtilioribus, amentis axillaribus terminalibusque longiuscule pedunculatis folia æquantibus filiformibus. pedunculis folia paullo superantibus, bractea rotundata centro peltata, ovario basi immerso, stigmate carnosulo apice fere imo inserto. ☉ In ins. Mauritio (Sieber n. 165! in h. Kew. et Berol.). Herba repens. Caulis glaber. Limbi 0,02 longi. Petioli 0,005 longi. (v. s.)

β, *macrophylla*, foliis majoribus basi cuneatis. Limbi 0,03 longi. Petioli 0,006 longi. (v. s.)

261. *P. PORTULACOIDES* (A. Dietr. sp. v. 1. p. 172), foliis superioribus plerumque ternis cæteris oppositis obovato-oblongis vel subspathulatis basi acutis glabris fusco-punctatis membranaceo-pellucidis 3-subquinenerviis nervuloque tenui plus minusve interrupto juxta marginem currente, amentis axillaribus longe pedunculatis, bractea rotundata centro peltata brevissime pedicellata, ovario emerso, stigmate punctiformi puberulo apice imo inserto. ☉ In insulis Bourbon (Sieb. n. 167! in h. DC.), Mauritio (Sieb. n. 164! in h. DC. et Berol.), in India (Ventenat! in h. Deless.). — Miq. syst. p. 130, Wight's ic. t. 1922. *Pep. candolleana* Miq. syst. p. 146. *Piper portulacoides* Lam. l. ill. p. 82, Vahl enum. v. 1. p. 350. *Piper trifolium* h. Willd. n. 737! (spec. Lam.). Herba repens e nodis caulinis radicans. Limbi 0,015-0,04 longi. Petioli 0,005-0,012 longi. (v. s.)

262. *P. LANGSDORFFII* (Miq. syst. p. 116), foliis petiolatis superioribus alternis cæteris oppositis obovato-vel elliptico-rhomboides apice parum attenuatis obtusiusculis basi fere cuneatis utrinque pubescentibus subcarnosio ciliolatis reticulo-nervulosis et trinerviis, lateralibus nervis subtilioribus, amentis terminalibus breviter pedunculatis, bractea oblonga basi truncata medio peltata, ovario emerso-impresso, stigmate penicillato apice oblique inserto. ☉ Verisimiliter e Brasilia in caldariis Berolin. et Kew. culta. — Miq. l in Linn. v. 20. p. 124 et in Mart. flor. Bras. fasc. 11. p. 13. *Micropiper Langsdorffii* Miq.

comm. phyt. p. 52. t. 8. fig. F. Suffruticulus vix pedalis. Ramuli pallide rubelli, breviter pubescentes. Limbi 0,045 longi. Petioli 0,015. Pedunculi vix 0,01 longi. Amenta 0,1-0,12 longa. Verisimiliter forma *Pep. blandæ*. (v. s.)

β, *increscens*, caulis dense hirtopubescentibus, foliis superioribus rhombeo-obovatis attenuato-obtusis 3-5-nerviis utrinque præsertim subtus ad nervos pilosis siccis nigro-punctatis. Ad Eugenio de Vargas (Pohl n. 5040! in h. Vindob.). *Pep. increscens* Miq.! in Linn. v. 20. p. 124. Limbi superiores 6-9 lin. longi. Petioli  $\frac{1}{2}$  - 1  $\frac{1}{2}$  lin. longi. (v. s.)

263. *P. INSULARUM* (Miq.! in Hook. Lond. journ. bot. v. 4. p. 422), foliis petiolatis ovato-attenuatis apice acutis basi rotundatis vel subattenuatis rigidulis subopacis junioribus supra pubescentibus subtus præsertim ad nervos pubescentibus non reticulato-nervulosis trinerviis, centrali nervo ad apicem usque ducto, lateralibus ad medium usque ductis, petiolo pubescente, amentis axillaribus terminalibusque filiformibus densifloris, rhachi pilosula, bractea rotundata centro peltata, ovario emerso, stigmate minuto apice subantice inserto. ☉ In ins. Oahu (Diell n. 53! in h. Kew.). Herba basi radicans. Caulis pubescens, ad nodos tumidulus. Limbi 0,015 longi, 0,008 lati. Petioli 0,002 longi. Pedunculi 0,005 longi. (v. s.)

β, *glabrata*, foliis paullo majoribus glabris, amentis crassioribus. In ins. Sandwich (Diell in h. Kew.). Caulis glaber. Limbi 0,03 longi, 0,012 lati. Petioli 0,005-0,008 longi. Pedunculi 0,005 longi. (v. s.)

264. *P. CATHARINÆ* (Miq.! syst. p. 127), foliis brevissime petiolatis e basi cuneata obovatis apice rotundatis utrinque subtiliter puberulis rigidulo-membraneis subpellucidis pellucido-punctulatis fere enervulosis trinerviis, nervulo apice marginem currente, petiolo subtiliter puberulo, amentis terminalibus filiformibus folia duplo superantibus, pedunculo petiolum multoties superante puberulo, rhachi subtiliter glanduloso-puberula, bractea rotundata centro peltata subsessili, ovario profunde immerso ovato apice mucronato, stigmate apice imo inserto, bacca patula profunde immersa oblongo-ovata apice mucronata. ☉ In Santa-Catharina (Gaudich. n. 282! in h. DC. et Deless.). Herba basi decumbens radicans, dein erecta, spithamea, ramulosa. Ramuli puberuli. Limbi 0,01-0,015 longi, 0,007-0,01 lati. Petioli 0,002 longi. Amenta matura 0,001 crassa. (v. s.)

265. *P. CYCLOPHYLLA*, foliis brevissime petiolatis orbicularibus piloso-pubescentibus ciliatisque membranaceis trinerviis vel enerviis, amentis adspectu axillaribus ramulos axillares squamas 2 gerentes terminantibus, bractea rotundata centro peltata, amentis folia multoties superantibus, ovario immerso, stigmate carnosulo apice imo inserto. ☉ In Peruvia orientali prope Tarapoto et Obidos (Spruce n. 517! in h. Francov.), Brasiliæ prov. Para (Spruce n. 4987! in h. DC.), Panama (Seemann n. 607! in h. Kew.). (v. s.)

266. *P. CIRCINATA* (Link Jahrb. v. 1. 3. p. 64 fide R. et Sch. Mant. v. 1. p. 245), foliis brevissime petiolatis orbicularibus carnosius utrinque sparse puberulis subopacis rigidulis subtiliter trinerviis, amento terminali squamis 2 fulto, bractea rotundata centro peltata, ovario emerso, stigmate minuto apice imo inserto. ☉ In Brasilia (Pohl! in h. Vindob., Spruce! in h. DC.), Paramaribo (Pohl! in h. Vindob.), Sto-Domingo (Poiteau! in h. Deless. et DC., Desfont. in h. Willd. n. 741! et 740!), India occidentali (Swartz obs. p. 20). — Miq.! in Linn. v. 20. p. 127 et in Mart. flor. Bras. fasc. 11. p. 21. *Pep. rotundifolia* Griseb.! Cat. pl. Cub. p. 63. Piper longum minimum herbaceum scandens Sloane! hist. p. 187 et icon. orig. n. 90 cum planta. Piper circinatum h. Willd.! n. 740, R. et Sch. l. c. *P. circulare* h. Willd.! n. 741. *P. rotundifolium* Sw. l. c. (non L.). *Acrocarpidium rotundifolium* Miq.! syst. p. 62. Herba filiformis longissime repens. Caulis sparse patentim puberulus, pilis mollibus sensim deciduis. Limbi 0,006 in diam. (v. s.)

267. *P. MONINI*, foliis petiolatis ellipticis utrinque subattenuatis subtus ad nervos pubescentibus subcoriaceis opacis trinerviis, amentis axillaribus longe pedunculatis densifloris, pedunculo pubescente, rhachi glabra, bractea rotundata centro peltata subsessili, ovario emerso, stigmate exiguo puberulo apice inserto. ☉ In ins. Bourbon (Monin! 1833. in h. DC.). Herba erecta pubescens

circ. 0,2 alta. Limbi 0,035 longi. Petioli circ. 0,01 longi. Pedunculi circ. 0,025 longi. (v. s.)

268. *P. MANDIOCCANA* (Miq. ! in Linn. v. 20. p. 125), foliis summis ternis cæteris oppositis petiolatis ellipticis vel rhombéo-ellipticis vel sublancoolato-ellipticis apice obtusis basi acutis aut obtusiusculis utrinque glabris membranaceo-pellucidis fusce punctatis indistincte trinerviis, centrali nervo ad apicem usque ducto, lateralibus subtilioribus ægre discernendis supra medium evanidis, nervulo obscuro juxta marginem currente, petiolo glabro amentis terminalibus densifloris folia superantibus. ☉ In arboribus ad Mandioccam (Mart. ! in h. monac.) — Miq. ! in Mart. flor. Bras. fasc. 11. p. 17. Herba glabra. Limbi 0,02 longi, 0,008 lati. Petioli 0,003 longi. (v. s.)
269. *P. SERPYLLIFOLIA* (A. Dietr. sp. v. 1. p. 174), foliis oppositis vel infimis aliquando alternis petiolatis subrotundis vel ellipticis pellucidis membranaceis tenuissimis utrinque glabris apicem versus plerumque ciliolatis fusce glandulosis obsolete trinerviis, nervis ramulosis nervuloque juxta marginem currente, amentis terminalibus pedunculatis folia parum superantibus sub-laxifloris, bacca subimmersa ovata. ☉ In ins. Mauritio (Neraud ! in h. Deless.) — Miq. ! syst. p. 131 (exclus. synonym. Lam.) *Piper serpyllifolium* Vahl en. v. 1. p. 351 ? quo ad specim. Commerson. Herba succulenta, tota pellucida. Caulis filiformis, nigro-punctulatus. Limbi 0,004-0,008 in diam. Petioli 0,001-0,005 longi. (v. s.)
270. *P. FERNANDEZIANA* (Miq. ! syst. p. 139), foliis oppositis raro ternis petiolatis superiorgibus ovato-oblongis apice acutiusculis infimis caulinis subrotundis truncato-retusis omnibus glabris apicem versus ciliatis utrinque nigropunctulatis membranaceis subcoriaceis enervulosis trinerviis, centrali nervo ad apicem usque ducto, lateralibus nervis supra medium evanidis, petiolo margine hirtello, amentis axillaribus sparsis vel oppositis ternis sub-laxifloris, bractea rotundata centro peltata, ovario impresso obtuso, stigmate minuto, bacca mucronulata. ☉ In sylvis umbris frigidis ins. Juan Fernandez (Bertero ! n. 1491 ! in h. DC., h. Deless.), in Chili ad lacum Ranco (Lechler n. 3020 ! in h. DC.), prope Valdivia in Chili (Philippi n. 87 ! in h. DC. et 775 ! in h. Francov.), Pichincha (Jameson ! in h. Kew.), Wahoo (Nuttall ! in h. Brit. Mus.). — Miq. ! in Hook. Lond. journ. bot. v. 4. p. 423, in Wight icon. t. 1923. Herba erecta ramulosa. Caulis sublignosus, pubescens et ad nodos hirsutus. Limbi 0,02-0,025 longi. Petioli 0,005-0,015 longi. Pedunculi 0,01 longi. Amenta post anthesin 0,03-0,04 longa. (v. s.)
271. *P. RUBRINODIS* (Kunth et Bouché ! in ind. semin. hort. Berol. 1848. p. 12), foliis oppositis apice ramorum 3-6-verticillatis confertis inæqualibus petiolatis oblongis apice obtusis emarginulatis basi cuneatis crasso-carnosiss membranaceis subopacis trinerviis, nervis supra nitidulis subtus pallidioribus, centrali nervo subtus prominulo, amentis terminalibus ternis longissime pedunculatis densifloris folia triplo superantibus, pedunculo petiolum multoties superante, bractea ovata obtusa infra apicem peltata, ovario oblongo vertice ampliore obtuso, stigmate punctiformi antico. ☉ In Guatemala (Warszewicz ! in h. hort. Berol.). Erecta, carnosissima, glabra. Rami rubro-lineati. Limbi 0,02-0,08 longi, 0,01-0,02 lati. Petioli 0,005 longi. Pedunculi 0,085 longi. (v. s.)
272. *P. MUSCOSA* (Link Jahrb. v. 1. 3. p. 64, fide R. et Sch. Mant. v. 1. p. 245), foliis brevissime petiolatis oblongo-ellipticis apice obtusis basi acutis utrinque glabris coriaceis opacis trinerviis, nervis subtus carinato-prominulis, centrali nervo ad apicem usque ducto, lateralibus alte supra medium ductis, petiolo glabro, amentis axillaribus filiformibus densifloris folia æquantibus longe pedunculatis, pedunculo folium multum superante glabro, bractea rotundata centro peltata sessilibus, ovario semimmerso, stigmate carnosulo-papilloso puberulo apice imo inserto. ☉ In Brasilia (Hoffmansegg in h. Willd. n. 7391), in arboribus prope Taguados urb. Para et S. Ludov. prov. Paraens. et Maragn. (Mart. ! in h. monac.). — Miq. syst. p. 181, in Linn. v. 20. p. 127 in Mart. flor. Bras. fasc. 11. p. 21. t. 1. fig. 8. *P. quadrangularis* Griseb. flor. w. Ind. p. 168. *Piper muscosum* Rem. et Sch. Mant. l. c. Coriaceo-succulenta glaberrima nitidula cartilagineo-coriacea. Caulis longe

repens ad nodos radicans. acute quadrangulus, 0,001 crassus. Limbi 0,025-0,035 longi, 0,01-0,019 lati. Petioli 0,0015 longi. (v. s.)

273. *P. ANGULATA* (Kunth! in H. et B. nov. gen. v. 1. p. 86), foliis breviter petiolatis obovatis vel elliptico-obovatis apice obtusis basi cuneatis utrinque glabris coriaceis opacis trinerviis, nervis subtus carinato-prominulis, centrali ad apicem usque ducto, lateralibus supra medium ductis, petiolo glabro, amentis longe pedunculatis filiformibus densifloris folia triplo superantibus pedunculo squamulis fulto sulcato subtiliter sparsim hirtello, rhachi foveolata sulcata, bractea rotundata centro peltata subsessili glanduloso-punctata, ovario immerso oblongo, stigmate capitellato carnosio-puberulo apice imo inserto. ☉ In locis umbriosis prov. Cumanensis ad ripam fluminis Jagua (Kunth l. c. et Humb. in h. Willd. n. 738! et in h. Berol.), in arboribus inter muscos prope Maribi prope Rio Negro (Mart. in h. monac.), Surinam (Hostmann n. 784! in h. Boiss. Miq. in Nederl. Kruidk. v. 1. p. 211). — Kunth synops. v. 1. p. 119, Miq. l. syst. p. 180, in Linn. v. 20. p. 127, in Mart. flor. Bras. fasc. 11. p. 21, in Hook. Lond. journ. bot. v. 4. p. 427. *P. angulatum* R. et Sch. Mant. v. 1. p. 332. Herba repens, rigide coriacea, subcartilaginea, nitidula, glaberrima. Caulis acute quadrangulus, repens, e nodis radicans. Limbi 0,025 longi, 0,016 lati. Petioli 0,003 longi. Caulis 0,001 crassus. (v. s.)

274. *P. VENTENATII* (Miq. l. syst. p. 125), foliis petiolatis obovato-oblongis carnosissimis pellucido-punctatis glabris subtus albido-pallidis et nigro-punctatis subtrinerviis, petiolo rarissime puberulo, amentis axillaribus terminalibusque folia triplo superantibus longiuscule pedunculatis, pedunculo folia triplo superante, bractea oblonga centro peltata, ovario oblongo, stigmate crassiusculo puberulo, bacca semiimmersa. In ins. Java (Ventenat! in h. Deless.). ins. Manilla ad Rio Passy (Miq. in nov. act. nat. cur. v. 19. p. 486). Herba carnosae coriaceae. Caulis repens ad nodos radicans. Limbi majores 0,02 longi, 0,007-0,015 lati. Petioli 0,005-0,007 longi. (v. s.)

β, *pubescens* (Miq. flor. Ind. Batav. v. 1. pars 2. p. 434), caule minute puberulo, foliis apice ciliolatis opacis 0,011 longis, 0,006 latis. In ins. Manilla (h. Berol!). (v. s.)

275. *P. BERNIERIANA* (Miq. l. syst. p. 124), foliis petiolatis ovato-spathulatis apice rotundatis basi cuneatis utrinque glabris rigidulis trinerviis, centrali nervo validiore, amentis terminalibus axillaribusque folia fere duplo superantibus, pedunculo petiolum duplo superante. ☉ In Madagascar latitud. Sta-Maria (Bernier in h. Deless. l.). Herba 0,1 propemod. alta, basi aphylla repens radicans. Caulis glaber. Limbi 0,02 longi, 0,01 lati. Petioli 0,008 longi. (v. s.)

276. *P. QUADRANGULARIS* (A. Dietr. sp. v. 1. p. 169), foliis breviter petiolatis ovalibus obtusiusculis basi rotundatis glabris opacis rigidis trinerviis, amentis axillaribus densifloris breviter pedunculatis. ☉ In Trinitate. — Miq. syst. p. 135, Griseb. flor. w. Ind. p. 168. Piper quadrangulare Thoms. in Trans. Soc. Lin. v. 9. p. 202. t. 21. fig. b. Caulis repens spithameus, simplex vel parce ramosus, glaber. tetragonus. Folia 12" longa, 8" lata. Petioli 1" longi 6"-8" longa. (Ex Griseb. l. c.)

277. *P. RETUSA* (A. Dietr. sp. v. 1. p. 155), foliis alternis vel apice ramulorum oppositis inferioribus majoribus e basi cuneata obovato-ellipticis apice obtusis apice imo saepe emarginatis superioribus minoribus rotundato-obovatis membranaceis subpellucidis utrinque glabris reticulato-nervulosis uninerviis vel obsolete trinerviis triplinerviisve, nervis lateralibus ad medium usque ductis saepe obsoletis, petiolo glabro, amentis terminalibus longiuscule pedunculatis filiformibus subaxillaribus folia triploquadraplo superantibus, bractea rotundata centro peltata subsessili, ovario impresso ovato, stigmate minuto puberulo apice imo inserto, bacca impressa subglobosa nigra. ☉ In Promontorio Bonæ Spei ad Kerstenbloch (Bergius! in h. Berol.). — Miq. l. syst. p. 132. Piper retusum L. fil. supp. p. 91, Thunb. flor. cap. p. 443, Vahl enum. v. 1. p. 346, Willd. sp. v. 1. p. 165. Herba repens. Rami adscendentes erecti, glabri, spithamei, circ. 0,002 crassi. Limbi 0,02 longi, 0,01 lati. Petioli 0,005 longi. (v. s.)

β, *alternifolia*, foliis omnibus alternis subrotundis apice emarginatis.

⊙ In Promontorio Bonæ Spel (Ecklon et Zeyher! in h. Berol.). Limbi 0,005 lati. (v. s.)

γ, *ciliolata*, foliis apice ciliolatis interdum emarginulatis. In monte Diaboli (Fisch. ! in h. Petrop.). (v. s.)

278. P. ANTILLARUM, foliis petiolatis inferioribus alternis superioribus oppositis omnibus subrhombico-ellipticis apice acutis obtusiusculisve basi cuneatis supra dense pubescentibus subtus dense villosulis ciliatis rigidis opacis pellucido-punctulatis subtus pallidioribus uninerviis, petiolo dense villosulo, amento terminali densifloro folium duplo superante, pedunculo villosulo petiolum superante, rhachii hirtella, bractea rotundata glandulosa margine ciliata centro peltata breviter pedicellata, ovario impresso glanduloso, stigmatibus apice oblique inserto. ⊙ In Cuba orientali (Wright n. 5111 in h. Boiss.). Herba erecta. Caulis villosulus. Limbi 0,027 longi, 0,014 lati. Petioli 0,01 longi. (v. s.)

279. P. REINWARDTIANA (Miq. syst. p. 141), foliis petiolatis oppositis summis ternis inferioribus minoribus subrotundis superioribus majoribus obovato vel ovato-rotundatis omnibus pubescentibus uninerviis, nervo parum conspicuo nervisque lateralibus subtilissimis vix cernendis, petiolo densius pubescente, amentis terminalibus laxifloris, pedunculo petiolum fere æquante, bractea peltata immersa ovata glabra. ⊙ In Borneo (Miq. in Ann. Mus. Lugd. Bat. v. 1. fasc. 5. p. 135), cacumine montis Tjerimai ins. Java ad trunco arborum (Bl. in h. Lugd. Bat. fide Miq. l. c.), supra Kaboendoengang (Miq. in Ann. Mus. Lugd. l. c.). — Miq. l. c. et in flor. Ind. bat. v. 1. pars 2. p. 434. Piper subrotundum Bl. in Verh. der Bat. Genoots. v. 11. p. 235. fig. 39. Suffructiculus carnosus, dimidium pedem altus, pubescens, inferne radicans, subprostratus, purpurascens, pennam corvinam crassus. Limbi superiores 0,015 longi, 0,003 lati. Petioli 0,005 longi. (Ex Miq. l. c.)

280. P. CILIOLATA (Miq. in Hook. Lond. Journ. bot. v. 6. p. 459), foliis oppositis quaternisve modice petiolatis obovato vel elliptico-orbicularibus obtusis emarginatis vel brevissime acutis subtus pallidis supra saturate viridibus nitidis margineque dense ciliatis uninerviis. ⊙ In horto Kewensi culta ab horto ducis Bedford communic. cum commentatione e Mexico australiore? Carhosa succulenta præter margines foliorum glabra. Petioli aliquot lineas longi. Folia in eodem verticillo si plus quam duo adsunt diversiformia  $\frac{1}{2}$ –1 pollicaria.

*Species affines non satis notæ.*

281. P. PUBESCENS (Ruiz et Pav. flor. peruv. v. 1. p. 31), foliis oppositis petiolatis obovatis subtus pubescentibus trinerviis, amentis ternis quorum medium longissimum. In umbrosis montium Huassahuassi supra saxa. — Miq. syst. p. 135. Piper saxicolum Vahl enum. v. 1. p. 349. Piper pubescens Pers. syn. p. 34. Planta bipedalis. Caulis erectus ramosus teres pubescens. (Ex R. et P. l. c.)

282. P. HAWORTHIANA (A. Dietr. sp. v. 1. p. 171), foliis oppositis rarissime ternis rhombeis coriaceo-carnosis minutissime pubescentibus ciliatisque trinerviis, centrali nervo ramoso. ✕ In India? — Miq. syst. p. 180. Caulis erectus. (Ex Dietr. l. c.)

283. P. HUMILIS (A. Dietr. sp. v. 1. p. 168), foliis oppositis oblongis acutis utrinque villosis subtus pallidioribus, petiolo quam folium quadruplo brevior, amentis terminalibus solitariis vel geminis filiformibus. In ins. S. Crucis (West fide Vahl). — Miq. syst. p. 134. Piper humile Vahl enum. v. 1. p. 349. Caulis basi repens, demum erectus, spithameus, tetragonus, villosus, superne parum divisus. Folia semipollicaria. Amenta bipollicaria. Pedunculi semipollicares. (Ex Vahl l. c.)

284. P. ALSINOIDES (A. Dietr. sp. v. 1. p. 168), foliis oppositis petiolatis ovato-oblongis subtus quinquenerviis, amentis axillaribus brevibus. In Carolina meridionali (Fraser fide Lam.) sed in Floris amer. ignota. — Miq. syst. p. 134. Piper alsinoides Lam. ill. v. 1. p. 82, Spreng. syst. v. 1. p. 120. (Ex Lam. l. c.)

285. P. VABLI (A. Dietr. sp. v. 1. p. 169), foliis petiolatis oblongis obtusiusculis

- β, *pilosior*, foliis plerumque ternis utrinque hirtis-pilosis cōmactis opacis, amentis terminalibus solitariis densifloris folia triplo superantibus, ovario semi-immerso ovato, stigmate cartosulo apice imo inserto. ☉ In *Brasilia* (Selle! in h. Berol.); in arboribus ad Salgado prope flumen S. Paulisci et ad Guidowald (Wart. I in h. monac.), prope Petropoli ad arborum corticem (Wawra et Maly n. 469b, fide Wawra bot. Ergebn. p. 133). Herba coriacea, radicans. Caules juniores pilosi, subtragoni, plicato-sulcati. Limbi 0,006 longi. Petioli vix 0,001 longi. (v. s.)
298. P. *COCKIANA*, foliis ternis quaternisve petiolatis subrotundis superioribus elliptico-lanceolatis omnibus membranaceo-coriaceis opacis subtus pubescentibus trinerviis, petiolo pubescente, amentis axillaribus, pedunculo pilosulo, bractea rotundata centro peltata, ovario subimmerso, stigmate papilloso-carnosulo. ☉ In ins. Sandwich (Gaudich. I in h. DC.). Nuncupata in honorem cel. detectoris insularum. Caulis herbaceus, basi radicans, pubescens, 0,001 crassus. Limbi superiores 0,015 longi. Petioli 0,005-0,01 longi. (v. s.)
299. P. *JAVANICA* (Miq. in plant. Jungh. fasc. 1. p. 15), foliis plerumque ternis brevissime petiolatis obovato-cuneatis vel e basi acutiuscula ellipticis subtus pubescentibus pellucido-punctulatis, amentis apice ramulorum ternis quinise pedunculatis laxifloris, bractea oblonga vel rotundata centro peltata breviter pedicellata glabra fusco-punctata, ovario emerso fusco-punctato, stigmate piloso-papilloso, bacca ovata. In ins. Java ad arbores (Zoll. n. 2016 I in h. DC.), monte Ungaran alt. 3-4000 ped. (Miq. l. c.). — Miq. I in Zoll. cat. p. 83, flor. Ind. Bat. v. 1. pars 2. p. 426. Planta  $\frac{1}{2}$ -1 ped. alta, basi nuda radicans, superne ramosa. Limbi circ. 0,008 longi. Petioli circ. 0,005 longi. (v. s.)
300. P. *QUATERNATA* (Miq. in Seem. bot. of Herald p. 197), foliis quaternis brevissime petiolatis obovato-vel-elliptico-rotundatis utrinque obtusis nitidis supra glabris subtus dense pilosis supra impresso-punctatis trinerviis. Ad Volcano Chiriqui (Seemann n. 895). Pep. reflexæ A. Dietr. proxima. Ramosa succulenta, basi radicans. Rami tetragoni, piliferi. Folia 2-2 $\frac{1}{2}$  raro 3 lin. longa. Petioli  $\frac{1}{2}$  lin. longi.
301. P. *EBULIS* (Miq. in Lign. v. 18. p. 711), foliis 3-5-verticillatis longiuscule petiolatis obovato-rotundatis plerumque retusis coriaceis haud pellucido-punctatis supra præsertim apicem versus tenere puberulis subtus pallidis glandulose punctatis trinerviis, petiolo tenere puberulo, amentis terminalibus solitariis densifloris glabris, bractea subrotundata carnosa centro peltata pedicellata, ovario apice stigmatifero. ☉ In regione frigida Mexici ad Perote (Schiede in h. Schlecht. fide Miq. l. c.). Pep. reflexæ A. Dietr. similis. Planta aromatica, Coriandri saporem aliquatenus æmulans, a mexicanis cruda coctaque comeditur. Limbi 0,007-0,012 plerumque 0,008-0,01 longi. Petioli 0,002-0,008 longi. Pedunculi 0,015 longi.
302. P. *OVALIFOLIA* (Hook. I flor. exot. t. 165), foliis oppositis ternisve brevissime petiolatis ovalibus apice retusis membranaceo-rigidis. ciliolatis junioribus puberulis trinerviis, centrali nervo ad apicem usque ducto, lateralibus tenuibus, petiolo hirtello, amentis axillaribus terminalibusque longe pedunculatis, pedunculo puberulo, bractea rotundata centro peltata subsessili, ovario semi-immerso apice acuminato, stigmate orbiculari, bacca globosa rostellata lævi. ☉ In ins. St Vincent (h. Kew.), Guadalupe (Krauss I in h. DC.), Dominica (Imray n. 334 I in h. Kew.), Trinifate (Sieber n. 299 I in h. DC.). — Miq. syst. p. 154 et in Hook. Lond. Journ. bot. v. 4. p. 424. Herba repens. Caulis hirtellus circ. 0,001 crassus. Limbi 0,015 longi, 0,01 lati. Petioli 0,06 longi. Pedunculi 0,035 longi. (v. s.)
303. P. *NUMMULARIOIDES* (Griseb. I in Syst. Bemerck. üb. Phil. p. 44), foliis petiolatis inferioribus alternis superioribus 3-5-verticillatis omnibus elliptico-obovatis apice obtusis basi subattenuatis tenuissime puberulis membranaceo-rigidius inconspicue trinerviis non reticulato-nervulosis, nervulo juxta marginem corrente, amentis terminalibus axillaribusque subdensifloris longiuscule pedunculatis, bractea rotundata centro peltata, ovario impresso apice imo stigmatifero. ☉ In Cordilleris Chilensibus ad arborum truncos (Lechler n. 838 I in h. DC. et Philippi in h. Kew.). Herba e nodis radicans, glabra. Limbi 0,02 longi. Petioli 0,005 longi. (v. s.)

304. *P. FILIFORMIS* (Ruiz et Pav.) flor. peruv. v. 1. p. 31. t. 47. fig. b), foliis quaternis cuneato-obovatis ciliolatis utrinque pilosiusculis non reticulato-nervulosis trinerviis, petiolo minutissime velutino-puberulo, rhachi setulosa, bractea ovato-attenuata centro peltata, ovario immerso, stigmatibus carnosulo minuto. ☉ In nemoribus Chinchao supra saxa (Pav.) l. c. et in h. Boiss., Lechler n. 2409! in h. Kew.). — A. Dietr. sp. v. 1. p. 181. *Piper brachystachyon* Vahl enum. v. 1. p. 354. *P. filiforme* Pers. syn. p. 34. Herbula. Limbi 0,007 longi, 0,005 lati. (v. s.)

305. *P. REFLEXA* (A. Dietr. sp. v. 1. p. 180), foliis ternis quaternisve (raro ad dichotomias senis) petiolatis pellucido-punctatis supra glabris subtus pilosis coriaceis obsolete trinerviis nervulosis vel enervulosis nervulosis saepe apice juxta marginem currente, petiolo hirtello, amentis terminalibus pedunculatis folia 2<sup>o</sup>-5<sup>o</sup> superantibus, pedunculo tenuiter hirtello; *rhachis* hirtella foveolata, bractea rotundata centro peltata subsessili glabra carnosula, ovario profunde immerso ovato-acuto, stigmatibus capitellato puberulo; bractea ovato-acuta subpatente. ☉ In utroque orbe, nempe: in Promontorio Bonae spei ex gr. ad Vitenhage valle Krokakama (Zeyher! in h. Franch.) in h. DC. (Hügel n. 4324! in h. Endl.), Ceylon prov. centrali alt. 6000 ped. (Thwaites n. 2462 fide Thwaites enum. p. 292), Javae monte Pangerango (Miq. in Ann. bot. Mus. Lugd. Bat. v. 1. fasc. 5. p. 135. Bl.) in h. DC. ex h. Lugd. Bat.), Novae Hollandiae regione fluminis Hastings et sinus maris Moreton (F. Müller fide Miq. in Pip. nov. Novae Hollandiae in Meded. der Koning. Akad. Van Veenensch, Afdeel. Natur. 2<sup>e</sup> Reeks Deel 2. p. 7), Tahiti (D'Urville! in h. Berol.), Sandwich (Macrae! in h. Brit. Mus. et h. DC. Mann n. 242! in h. DC.); in America ad Caracas (Gallmer n. 331! et 335! in h. Berol.), Costa-Rica (Hostmann n. 796! in h. DC.), Brasilia ad Petropolim (Wawra et Maly n. 469; fide Wawra Ergeb. bot. p. 133), Trinitate (Sieber n. 290! in h. Endlich.), Jamaica (Linden n. 1794 fide Griseb. Cat. pl. Cub.). — Miq. syst. p. 169, in Mart. flor. Bras. fasc. 11. p. 17, in Hook. Lond. journ. bot. v. 4. p. 426, Wight icon. t. 1933. fig. 1, flor. Ind. Bat. v. 1. pars 2. p. 436; Griseb. flor. W. Ind. p. 168, Cat. pl. Cub. *Piper reflexum* L. fl. suppl. p. 91, Lam. ill. p. 83, Thunb. flor. cap. p. 443, Vahl enum. v. 1. p. 353. *P. tetraphyllum* Forster prod. n. 25, Willd. sp. v. 1. p. 168, Sw. obs. p. 23. *P. saxatile* Wall! herb. P. pusillum Bl. in Verh. der Bat. Genoot. v. 11. p. 382. fig. 37. *Pep. tetraphylla* Hook. et Arn. in Beech. bot. p. 95. *Herba succulenta repens*. Caulis sulcatus, praesertim ad nodos pilosulus. Folia magnitudine et consistentia secundum varietates sequentes variabilia. (v. s.)

α. *Capensis*, foliis plerumque longius petiolatis, petioli longitudine variabiles), utrinque glabris subcoriaceis subpellucidis trinerviis, nervulo juxta marginem apice currente, pedunculo glabrato folia duplo superante. In Promontorio Bonae Spei (Zeyher n. 3873!), prope Natal (Gouinzius!). Petioli plerumque 0,002 longi. *Pép. reflexa* forma *capensis* Miq. syst. p. 169. (v. s.)

β. *aemula*, foliis ut in praecedente, pedunculo hirtello folia aequante. In ins. Norfolk (Bauer! in h. Endl. et in h. Berol. Cunningham! in h. Kew.). *Piper aemulum* Endl. prod. flor. Norf. p. 36, Bauer ill. Norf. ined. t. 127, fide Endl. l. c. *Pép. reflexa*, forma *Norfolkensis* Miq. syst. p. 172. (v. s.)

γ. *membranacea*, foliis ovatis submembranaceis subpellucidis trinerviis, pedunculo fere glabro folia duplo superante, petiolo vix 0,001 longo. In ins. Bourbon (h. DC. ex h. Mus. Par.) et Mauritio (h. Kew.). (v. s.)

δ. *nervulosa*, foliis ellipticis apice subattenuatis basi obtusis membranaceis brevissime petiolatis trinerviis laxo nervulosis, pedunculo folia duplo triplo superante brevissime hirtello. In ins. Oahu (Beechey! et Barclay! in h. Kew.). *Pép. reflexa*, forma *Forsteriana*, Miq. syst. p. 173. (v. s.)

ε. *pusilla*, foliis longius petiolatis submembranaceis glabris trinerviis, nervis lateralibus conspicuis. In ins. Java (Zollinger n. 1985!). *Piper pusillum* Bl. in Verh. der Bat. Genoot. v. 11. p. 232. *Micropiper pusillum* Miq. comm. phyt. p. 62. t. 5. fig. b. *Pép. reflexa* forma *Javana* Miq. syst. p. 172. (v. s.)

paullo tenuioribus alte supra medium ductis, petiolo glabro, amentis terminalibus longiuscule pedunculatis filiformibus folia multoties superantibus subsensifloris, bractea oblongo-elliptica utrinque obtusa vel apice acutiuscula centro peltata subsessili, ovario post anthesin semiimmerso ovato apice contracto, stigmate capitellato carnosulo-penicillato. 5 In Peruvia (Spruce n. 4285! in h. DC.), Sta Catharina Brasiliæ (Miq. in Hook. Lond. Journ. bot. v. 4. p. 424), Venezuela ad Caracas alt. 30-700 hexag. (Kunth l. c. Gállmer n. 328! in h. Berol. Moritz n. 298! in h. Brit. Mus.), in hort. Berol. cult. (h. Willd. n. 746!). — Miq. syst. p. 150, in Mart. flor. Bras. fasc. 11. p. 15, Piper pereskiaefolium Jacq. Coll. v. 4. p. 126, icon. rar. v. 2. p. 2. f. 219, Vahl enum. v. 1. p. 352, Willd. sp. v. 1. p. 167. P. rubricaulis Link enum. alt. v. 1. p. 39, R. et Sch. Mant. v. 1. p. 246. P. stellatum Vell. flor. flum. t. 64 (non Jacq.). Pep. rubricaulis A. Dietr. syst. p. 182, Miq. syst. p. 149, in Mart. flor. Bras. fasc. 11. p. 14. Caulis suberectus, basi lignescens, parce dichotome ramosus. Rami coriaceo-succulenti, teretiusculi, profunde sulcati, subangulosi. Limbi 0,03-0,055 longi, 0,015-0,035 lati. Petioli 0,004 longi. Amenta florentia 0,12 longa. (v. s.)

314. *P. FONTINALIS* (Kunth! in H. et B. nov. gen. v. 1. p. 71), foliis inferioribus quaternis senise superioribus oppositis alternisve omnibus sessilibus linearilanceolatis apice acutiusculis basi lata affixis margine subrevolutis utrinque glabris carnosius coriaceis subopacis indistincte trinerviis, amentis terminalibus pedunculatis filiformibus densifloris folia multoties superantibus, bractea rotundata centro peltata subsessili glandulosa, ovario obovato post anthesin impresso, stigmate minuto? ☉ In convallibus fontibus madentibus juxta ripam fluminis Amazonum inter pagum Tomependac et Jaen de Bracamoros (Kunth l. c. Humb. in h. Willd. n. 758!). — Miq. syst. p. 162, A. Dietr. sp. v. 1. p. 176. P. fontinale Rœm. et Sch. syst. v. 1. p. 335. Pep. reflexa Kunth! l. c. p. 70 (Humb. in h. Berol.), synopsis v. 1. p. 122. Herba fontinalis Callitriches speciem majorem fere referens, pallide viridis, succulenta, fragilis, 0,2 propem. alta. Caulis simplex, teretiusculus, glaber. Limbi 0,01 longi, 0,002 lati. (v. s.)

315. *P. RAINEURA* (Miq. syst. p. 175), foliis quaternis et pluribus plerumque septenis et octonis breviter petiolatis rhombeo-vel lanceolato-ellipticis utrinque attenuatis apice obtusiusculis basi acutis utrinque glabris subtilissime ciliolatis coriaceis opacis glanduloso-punctulatis trinerviis, nervis subtus carinatis, centrali nervo ad apicem usque lateralibus paullo supra medium ductis, petiolo glabro, amentis longiuscule pedunculatis filiformibus glabris folia quintuplo superantibus densifloris, pedunculo glabro folia subtriplo superante, bractea ovato-rotundata centro peltata subsessili, ovario profunde immerso, stigmate carnosulo. ☉ In Brasilia (Sello! in h. Berol., Pohl in h. Vindob.), ad Santa-Gallo (Peckott! in h. Mart.). — Miq. in Linn. v. 20, p. 126, in Mart. flor. Bras. fasc. 11. p. 18. t. 2. fig. b. Herba coriaceo-carnosa, cartilaginea, supra muscoso-decumbens et radicans. Rami glabri, subtetragoni, 0,04-0,05 longi, 0,004 crassi. Limbi 0,006-0,007 longi, 0,003 lati. Petioli 0,001 longi. (v. s.)

316. *R. QUADRIFOLIA* (Kunth! in H. et B. nov. gen. v. 1. p. 69), foliis quaternis e basi cuneata obovatis apice obtusis emarginatis brevissime petiolatis coriaceis glabris trinerviis vel subtriplinerviis, centrali nervo ad apicem usque ducto utrinque nervos 1-2 alternos mittente, amentis terminalibus solitariis longiuscule pedunculatis filiformibus densifloris folia triplo quadruplo superantibus, rhachi glabra, bractea rotundata centro peltata subsessili, ovario per anthesin immerso ovato-oblongo, stigmate carnosulo-puberulo, bacca immersa ovata apice mucronata. ☉ In Peruvia (Pav. in h. Boiss.), Andibus quitenibus (Spruce n. 6118! in h. DC.), Venezuela prope Tovar (Fendler n. 1179! in h. DC.), inter Vera-Cruz et Orizaba (h. Kew.), Nova-Andalousia (Humb. in h. Willd. n. 754! et in h. Berol.), Guyana gallica (Aubl. hist. v. 1. p. 21), ad rupes humidæ Mirador alt. 8000 ped. (Martens et Gal. in bull. Acad. Brux. v. 10. n. 2. p. 19), Jamaica (Mac Fadyen! in h. Kew.). Piper foliis quaternis Plum. pl. Am. ed. Burm. p. 238. t. 242. fig. 3. Piper quadrifolium L. sp. p. 43, Aublet l. c. Willd. dict. v. 26. p. 20. fig. 4, Lam. ill. p. 82, Vahl enum. v. 1.



- p. 355, Willd. sp. v. 1. p. 168, Sw. obs. p. 22. *Piper affine* Martens et Gal. (fide Miq. syst. p. 552) non C. DC. supra. *Piper inciaum* Rich. l. in h. Francov. Pep. quadrifolia Kunth synops. v. 1. p. 121, Miq. l. syst. p. 159, Griseb. flor. W. Ind. p. 168, Hook. exot. flora t. 22. *Peper. alpina* Martens et Gal. l. c. (non Dietr.). Caulis basi radicans, erectus, glaber. Limbi 0,006-0,008 longi. Petioli 0,0015 longi. (v. s.)
317. *P. REGELII*, foliis ternis quinisque breviter petiolatis et basi cuneata obovatis utrinque glabris rigidulis subopacis trinerviis, nervis haud conspicuis, petiolo glabro, amentis terminalibus folia fere quadruplo superantibus filiformibus subdensifloris, bractea rotundata centro peltata subsessili, ovario immerso ovato apice mucronulato, apice mucronuli stigmatifero. ♀ In Brasilia (Riedel et Langsdorff! in h. Petrop.). Ramosus. Rami sublignosi, tetragoni, glabri. Ramuli tetragoni, glabri. Limbi 0,006 longi, 0,005 lati. Petioli vix 0,001 longi. (v. s.)
318. *P. OBVERSA* (A. Dietr. sp. v. 1. p. 173), foliis ternis breviter petiolatis elliptico-obovatis apice obtusis vel levissime emarginulatis basi acutiusculis utrinque glabris rigidulis subpellucidis indistincte trinerviis, petiolo glabro, amentis terminalibus folium multoties superantibus filiformibus densifloris, pedunculo folium æquante glabro, bractea rotundata centro peltata subsessili, ovario subimmerso, stigmate penicillato. ☉ In Cuba orientali (Wright n. 6151 in h. Deless. et DC.), ins. Monserrat (Vahl l. c. et Griseb. l. c.). — Griseb. l. in mem. Am. Acad. v. 8. p. 176, flor. W. Ind. p. 168. *P. obversum* Vahl eolog. p. 5, enum. v. 1. p. 354. — Secundum Vahl accedit ad figuram *P. trifolii* Plum. Herba repens e nodis radicans. Caulis glaber 0,002 crassus. Limbi 0,01 longi, 0,008 lati. Petioli 0,002 longi. (v. s.)
319. *P. TOMENTOSA* (A. Dietr. sp. v. 1. p. 172), foliis 3-4 verticillatis plerumque quaternis subsessilibus ovatis vel ovato-ellipticis obtusiusculis utrinque sericeo-hirtis obsolete uninerviis, amentis axillaribus terminalibusque sæpe apice ramulorum cymose confertis, bractea glabra rotundata centro peltata pedicellata, ovario immerso ovato, stigmate puberulo, bacca immersa. ☉ In Java supra arborum truncos. — Miq. syst. p. 142. *P. tomentosum* Vahl enum. v. 1. p. 351. Herba succulenta, tota hirta. Limbi inferiores 0,025-0,03 longi, 0,015-0,01 lati. Petioli inferiores 0,001-0,003 longi. Limbi superiores 0,01-0,015-0,02 longi. (Ex Miq. l. c.)
- β, *carnea* (Miq. l. syst. p. 143), foliis oppositis raro ternis petiolatis ovato-ellipticis vel subrhombeis apice obtusis obtusiusculisve basi subattenuato-obtusiusculis utrinque et petiolo denso et molli tomento vestitis subcoriaceis opacis, centrali nervo vix conspicuo, amentis axillaribus terminalibusque folia duplo superantibus densifloris, pedunculo vestito petiolum æquante. ♀ In Java (Zollinger n. 1986! in h. DC., Francov. et Deless.). *Piper tomentosum* var. *carnosum* Bl. in Verh. der Bat. Genoots. v. 11. p. 237. fig. 41. *Micropiper tomentosum* Miq. comm. phyt. p. 63. t. 9. fig. b. *Pep. tomentosa* var. β Miq. ill. p. 21. t. 18, flor. Ind. Bat. v. 1. pars 2. p. 435. *Carnosa*, molliter hirsuta. Limbi majores 0,04 longi, 0,012 lati. Petioli 0,006 longi. (v. s.)
- γ, *pusilla* (Miq. l. Zoll. Verz. p. 83), forma duplo minor singulis partibus. In Java super arbores (Zoll. n. 2243, 2593, 1906, fide Zoll. l. c., n. 2593! in h. Francov.). (v. s.)
320. *P. THRICHOCARPA* (Miq. l. syst. p. 176), foliis ternis quaternisque breviter petiolatis lanceolato-ellipticis apice obtusis basin versus angustatis utrinque puberulis obsolete uninerviis, petiolo puberulo, flora subimmerso, bacca ovato-acuminata hirtella. ☉ Prope Bahiam (Blanchet n. 1444! in h. Deless.). — Miq. in Mart. flor. Bras. fasc. 11. p. 18. Herba spithamea et altior, dichotome ramosa, decumbens et radicans. Caulis et ramuli puberuli. Limbi 0,01 longi, 0,003-0,004 lati. Petioli 0,001 vix 0,002 longi. (v. s. et pro parte ex Miq. l. c.)
321. *P. AGAPATENSIS*, foliis ternis quinisque brevissimis petiolatis elliptico-oblongis vel oblongo-ovatis subcoriaceis puberulis obscure uninerviis non reticulato-nervulosis pellucido-punctatis, amentis terminalibus solitariis, bractea rotundata centro peltata, ovario immerso, stigmate carnosulo exiguo, bacca

- globo-attenuata. 3 In Nova Hispania (Pav. in h. Boiss.) et Peruvia ad Agapata in rupibus (Lechler n. 1934 in h. DC.). *Pep. heterophylla* Griseb. in Lechl. exsicc. per. Caulis lignosus, erectus, ramulosus, puberulus, 0,002 crassus. Limbi superiores majores 0,015 longi. (v. s.)
322. *P. FLAGELLIFORMIS* (Miq. in Hook. Lond. Journ. bot. v. 4. p. 423), foliis oppositis ternisve subsessilibus oblongo-obovatis apice obtusis basi attenuatis subtus tenerrime subpuberulis membranaceis tenuiter uninerviis tenuissime nervulosis, amentis axillaribus terminalibusque filiformibus subulaxifloris pedunculatis, pedunculo glabro, bractea rotundata centro peltata, ovario subimmerso, stigmate orbiculari. ☉ In Archipelagi Gallapagos ins. James (Darw. in h. Kew.). Herba erecta. Caulis glaber, filiformis, vix 0,001 crassus. Limbi 0,018 longi, 0,007 lati. Pedunculi vix 0,01 longi. Amenta circ. 0,05 longa. (v. s.)
323. *P. TENERRIMA* (Schlecht. in Linn. v. 6, p. 353), foliis binis-aenis brevissime petiolatis oblongo-spathulatis apice emarginulatis glabris apice ciliolatis opacis rigidis pellucido-punctulatis penninerviis, centrali nervo subtus prominulo nervos utrinque 4 alte ductos mittente, amentis terminalibus solitariis longius pedunculatis glabris subdensifloris filiformibus folia fere quadruplo superantibus, bractea rotundata centro peltata, ovario immerso oblongo-attenuato, stigmate capitellato carnosulo, bacca semi-immersa subglobosa apice mucronata. ☉ In Mexico inter Vera-Cruz et Orizaba (Meiss. n. 1328 in h. Kew.), in regione temperata Mexici ad Costa Grande de Jalacingo ad arborum truncos (Schiele in h. Berol.). — Miq. syst. p. 168, ill. p. 23. t. 21. *Pep. schiedeana* Schlecht. in Linn. v. 7. p. 138. Herbula caespitosa gregaria. Limbi 0,005-0,006 longi. Petioli vix 0,001 longi. (v. s.)
324. *P. LEPTOPHYLLA* (Miq. in Linn. v. 18. p. 712), foliis ternis quaternisve petiolatis patulis vel reflexis spathulato-vel oblongo-linearibus obtusis vel raro acutiusculis pallidis coriaceis haud crassis nec pallido-punctatis supra praesertim versus marginem petiolumque obsolete puberulis sensim glabratibus subtus pallidis et glanduloso-punctatis obscure uninerviis et enervulosis, amentis terminalibus breviter pedunculatis filiformibus densifloris, ovario immerso, bacca subglobosa punctata, stigmate terminali. ☉ In regione temperata Mexici (Schiele in h. Schlecht. fide Miq. l. c.). Succulenta. Caules inferne glabri, an tetragoni? nascentes pilis brevissimis densis patulis vestiti. Limbi 0,005-0,02 longi, 0,003-0,004 lati. Petioli 0,001-0,003 longi. Pedunculi 0,006 longi.
325. *P. HARTWEGIANA* (Miq. in Hook. Lond. Journ. bot. v. 4. p. 425), foliis quaternis quinisque brevissime petiolatis lato-vel subovato-ellipticis apice obtusis basi acutiusculis apice raro emarginatis vel mucronulatis crasse carnosius supra praesertim basin versus puberulis sensim glabratibus subtus glabriusculis et glanduloso-punctatis prope basin uninerviis, petiolo dense hirtello-puberulo, amentis terminalibus solitariis geminatisve pedunculatis densifloris, ovario oblongo acuminatim attenuato, bacca ovata acuta punctata. ☉ In Colombia (Hartweg n. 1401 in h. Kew.). Caulis spithameus, inferne e nodis radicans. Ramuli hirtello-incani. Limbi 0,008-0,015 longi, 0,006-0,01 lati. Petioli 0,001-0,002 longi. (v. s.)
326. *P. OBCORDATA* (Presl Epimel. p. 223), foliis quaternis breviter petiolatis obcordatis carnosulis planis anguste acutatis basi acutis cartilagineo-marginatis glaberrimis in emarginatura ciliolatis uninerviis enervulosis, pedunculo terminali compresso folium duplo superante, amento cylindrico obtuse apiculato pedunculum aequante, bractea minuta rotundata centro peltata subsessili ciliolata, ovario glabro, stigmate orbiculari colorato subsessili terminali, bacca ovato-subglobosa obliqua. ☉ In Cordilleris Peruviae. Glaberrima, ad nodos radicans, semidigitalis, repens. Folia semipollicem longa, 3 lin. lata. Petioli lineam longi.
327. *P. PORTULACAEFOLIA* (Kunth in H. et B. nov. gen. v. 1. p. 70), foliis quaternis brevissime petiolatis oblongo-obovatis vel spathulatis apice emarginulatis basi obtusiusculis utrinque glabris pellucido-punctulatis membranaceo-rigidulis uninerviis, nervulis nullis uno tamen apice juxta marginem currente, petiolo glabro, amentis terminalibus filiformibus folia quintuplo

- superantibus subdensifloris, pedunculo glabro petiolum superante, bractea rotundata glandulosa centro peltata subsessili, ovario obovato semi-immerso stigmatate carnosulo-puberulo apice fere imo vel subantice inserto. ☉ In convallibus umbrosis humidis rupium Popayanensium juxta villam La Erre et pagum La Cruz alt. 1140 hexap. (Humb. et B. l. c. et Humb. in h. Willd. n. 755 l.), Bogota alt. 2670 ped. (Triana, in h. DC.), prope Puerto Montt (Philippi in h. DC.). — Kunth synops. v. 1. p. 121, Miq. l. syst. p. 157. Piper portulacæfolium Roem. et Sch. syst. v. 1. p. 331. *Pep. australis* Philippi in Linn. v. 30. p. 200. Planta succulenta, 0,2 alta, radicans, decumbens, an demum erecta? glabra. Caulis coriaceus, subtetragonus. Limbi 0,013-0,025 longi, 0,006-0,008 lati. Petioli 0,001-0,002 longi. (v. s.)
328. *P. CEYLANICA* (Miq. in Hook. Lond. journ. bot. v. 5. p. 550), foliis oppositis ternis quaternisve breviter petiolatis ellipticis vel obovatis obtusis vel apice rotundatis basi plerumque acutis fere omnino glabris obsolete uninerviis et parce nervulosis centrali nervo infra apicem in anastomoses dissoluto, amentis axillaribus terminalibusque solitariis vel geminis densifloris, pedunculo subglabro, bractea rotundata centro peltata pedicellata, ovario breviter immerso, stigmatate penicillato. ☉ In ins. Ceylan (Walker n. 1904 fide Miq. l. c.). Herba digito parum altior, basi inter muscos radicans nuda, sursum caulibus ramisque pubescentes cito glabrescentes. Limbi 0,008-0,015 longi, 0,008 lati. Petioli 0,001-0,002 longi.
329. *P. DEPPEANA* (Schlecht. et Cham. in Linn. v. 5. p. 75), foliis ternis quaternisve petiolatis rotundato-obovatis basi acutiusculis membranaceis opacis glabris pellucido-punctulatis tenuiter reticulato-nervulosis uninerviis vel trinerviis, centrali nervo ad apicem usque ducto, nervis lateralibus 2 e basi ortis sæpe adjunctis, nervulo juxta marginem currente, amentis terminalibus longiuscule pedunculatis filiformibus densifloris folia triplo-quadruplo superantibus, pedunculo rhachique hirtello-puberulis, pedunculo folia triplo superante, bractea rotundata centro peltata, ovario immerso, bacca basi immersa ovato-mucronata. ☉ In Brasilia inter muscos (Sello in h. Berol.), Mexico (Deppe in h. Berol.). *Pep. Deppeana* Miq. l. syst. p. 160, in Hook. Lond. journ. bot. v. 4. p. 426, in Mart. flor. Bras. fasc. 11. p. 16. *Pep. sellowiana* Miq. l. syst. p. 161, in Hook. Lond. journ. bot. v. 4. p. 426. Herba basi decumbens radicans, erecta 0,02-0,06 alta. Ramuli subangulosi, sursum puberuli. Limbi 0,004-0,007 longi, 0,004 lati. Petioli 0,005 longi. (v. s.)
330. *P. LOXENSIS* (Kunth in H. et B. nov. gen. v. 1. p. 70), foliis quaternis breviter petiolatis obovato-ellipticis apice obtusis retusisve basi acutiusculis utrinque glabris rigidulis enerviis vel subuninerviis, centrali nervo obscure discernendo, petiolo glabro, amentis terminalibus densifloris filiformibus folia multoties superantibus, pedunculo glabro folia æquante, bractea rotundata glandulosa centro peltata subsessili, ovario immerso ovato apice acuminato, stigmatate capitellato puberulo. ☉ In locis umbrosis prope Loxa alt. 1059 hexap. (H. et B. l. c. et Humb. in h. Willd. n. 756 l.), Nova Hispania (Pav. in h. Boiss.), Mantiquirea in Brasilia (Sello in h. Berol.). — Kunth synops. v. 1. p. 122, Miq. l. syst. p. 158 et in Mart. flor. Bras. fasc. 11. p. 16. Herba digitalis vel spithamea, glabra, basi decumbens radicans, simplex vel dichotome ramosa. Caulis coriaceus, plicato-rugulosus, subquadrangulus. Limbi 0,006 longi, 0,005 lati. Petioli 0,001 longi. (v. s.)
- ++ Ovarium apice oblique stigmatiferum.
331. *P. MACRAENA* (Cas. DC. in Seem. journ. bot. 1866, p. 145), foliis ternis longe petiolatis elliptico-lanceolatis apice acutiusculis basi acutis supra glabris subtus ad nervos tenuissime puberulis membranaceo-rigidulis subpellucidis penninerviis, centrali nervo ad apicem usque ducto subtus prominulo ad  $\frac{1}{2}$  longitud. usque nervos utrinque 3-4 subadscendentes quorum supremi ad apicem usque fere ducti infimi subtiliores mittente, petiolo glabro, amentis filiformibus densifloris folia æquantibus, pedunculo glabro quam petiolum multum breviori, bractea rotundata centro peltata subsessili, ovario impresso obovato, stigmatate minuto. ☉ In Archipelagi Sandwich ins. Owhyhee

- (Macrae! in h. Brit. Mus.). Ramuli glabri vel ochraceo-puberuli. Limbi 0,07-0,09 longi, 0,03-0,06 lati. Petioli 0,04 longi. Pedunculi 0,01 longi. (v. s.)
332. P. DOMINICANA (Cas. DC. in Seem. journ. 1866, p. 146), foliis ternis vel apice ramulorum oppositis breviter petiolatis subtriangulo-ellipticis apice attenuato-obtusiusculis utrinque glabris coriaceis opacis septemnerviis, centrali nervo ad apicem usque et lateralibus 2 fere ad apicem usque ductis, petiolo glabro, amentis terminalibus solitariis filiformibus densifloris, pedunculo glabro folia æquante, bractea rotundata centro peltata subsessili, ovario emerso apice acutato. 5 In ins. Sancti-Domingi (Rich. in h. Francov. et Deless.). Ramuli coriacei glabri. Limbi 0,045 longi, 0,03 lati. Petioli 0,006 longi. (v. s.)
333. P. VITIANA, foliis inferioribus longe petiolatis alternis oppositisve superioribus minoribus ternis illis e basi cuneata lanceolatis his ellipticis utrinque attenuatis omnibus utrinque pubescentibus membranaceis obscuris reticulato-nervulosis quinquenerviis, petiolo pubescente, amentis axillaribus terminalibusque filiformibus densifloris folia paulo superantibus maturitate subaxillaribus pedunculatis pubescentibus, bractea rotundata centro peltata brevissime pedicellata, ovario emerso, stigmate carnosulo. ☉ In ins. Viti (Seeman p. 565! in h. DC. et in h. Kew. cum foliis omnibus alternis). Pep. pallida Seem. Fl. Viti. 259. Limbi 0,04 longi. Petioli circ. 0,01 longi. Pedunculi 0,015 longi. (v. s.)
334. P. CASAPIANA (Cas. DC. in Seem. journ. 1866, p. 146), foliis ternis petiolatis ovato-acuminatis utrinque piloso-pubescentibus membranaceis pelliculis reticulato-nervulosis quinquenerviis, centrali nervo ad apicem usque ducto, lateralibus 2 supra medium ductis externis subtilioribus, petiolo piloso-pubescente, amentis axillaribus verticillatis filiformibus subaxillaribus pedunculatis interdum apice ramuli axillaris binatis, bacca semi-immersa ovata apice brevissime mucronulata. ☉ In Peruvia prope Casapi (Mathews n. 1689! in h. Kew.). Herba. Caulis teres, piloso-pubescent. Folia superiora minora, squamiformia, unde umbella amentorum terminalia. Limbi majores 0,06 longi, vix 0,03 lati. Petioli 0,008 longi. Pedunculi 0,015 longi. Amenta vix 0,04 longa, vix 0,001 crassa. (v. s.)
335. P. MOERENHOUTI, foliis oppositis ternisve puberulis ciliolatisque inferioribus cuneato-obovatis apice parum protractis obtusis superioribus e basi cuneata lanceolatis apice obtusiusculis omnibus quinquenerviis, amentis axillaribus terminalibusque elongatis subaxillaribus pedunculatis glabris, pedunculo puberulo, bractea rotundata glabra fusco-punctulata centro peltata, ovario ovato, stigmate papilloso. ☉ In ins. Tahiti (Moerenhout! in h. DC.). Ramuli petiolique puberuli. Limbi superiores 0,03 longi. Petioli circ. 0,006 longi. Pedunculi vix 0,02 longi. Amenta circ. 0,13 longa. (v. s.)  
 β, petiolata, petioli 0,015 longi. In ins. Tahiti (h. Kew. l.). (v. s.)
336. P. BLANDA (Kunth! in H. et B. nov. gen. v. 1. p. 67), foliis oppositis ternis quaternisve breviter petiolatis elliptico-rhombicis vel rhombico-ellipticis utrinque acutis inferioribus elliptico-rotundis rotundisve omnibus utrinque pubescentibus ciliatisque fusco-punctulatis membranaceis subpellucidis laxè reticulato-nervulosis quinquenerviis, petiolo puberulo, amentis axillaribus terminalibusque sæpe apice ramulorum verticillatim confertis filiformibus folia multoties superantibus, bractea ovata carnosula centro peltata, ovario emerso, stigmate puberulo, bacca ovato-acuta lævi. ☉ In Boliviae prov. Larecaja vicinis Sorata prope Ladrilloni ad rivulum in umbrosis alt. reg. temp. 2650 m. (Mandon n. 1120! in h. DC.). Pichincha (Jameson n. 680!, 681! in h. Boiss.). Tarapoto (Mathews n. 837! in h. Vindob.). Caracas (Gallmer n. 334! in h. Berol.), ad arborum truncos inter Caracas et Rio Guayre alt. 460 hexap. (Humb. l. c. et in h. Berol.). Caracas (h. Willd. n. 7471), Venezuela (Fendler n. 1171! in h. DC.), Colombia (Hartweg n. 1395! in h. Kew.), in alta planitie Latunga prope Cotopaxi (h. monac. l.), locis humidis ad Curucuti alt. 2-3000 ped. (Wagner n. 126 in Linn. v. 26. p. 632), Nova Hispania (Pay. in h. Boiss.). — Kunth synops. v. 1. p. 120, Hook. exot. flor. t. 21, Miq. l. syst. p. 115. Piper blandum Jacq. icon. rar. v. 2. t. 218, coll. v. 3. p. 210, Lam.

ill. p. 82, Vahl enum. v. 1. p. 352, Willd. ! sp. y. 1. p. 164. *Pep. Quitensis* Benth. plant. Hartw. p. 253. Herba succulenta. Caulis erecti, teretes, viridi-rubentes, pilosi. Limbi 0,03-0,055 longi. Petioli 0,005-0,01 longi. (v. s.)

β, *reticulata*, foliis membranaceo-pellucidis crebre reticulato-nervulosis, elliptico-acutis apice brevissime attenuatis sæpe obtusis. — Frope Caracas (Vargas ! in h. DC.), in Mexico ad Orizaba (Sallé et Botteri ! in h. DC.), Carolina meridionali (Michaux ! in h. Francov. ex h. Rich.).

γ, *dissimilis*, foliis inferioribus elliptico-rotundis obtusis superioribus oblongo-ellipticis apice acutiusculis basi acutis membranaceis 3-5-nerviis obsolete nervulosis. In Venezuela (Fendler n. 11701, 11721, 11731 in h. DC.), in asperis et umbrosis juxta Valle de Juanambu et El Tablon alt. 760 hexap. regni Novo-Granatensis (H. et B. nov. gen. v. 1. p. 67. t. 13, Humb. ! in h. Berol. et in h. Willd. n. 7431). *Pep. dissimilis* Kunth ! l. c. et synops. v. 1. p. 119, Miq. ! syst. p. 117. *Piper dissimile* Roem. et Sch. syst. v. 1. p. 333. (v. s.)

337. *P. BOTTERII* (Cas. DC. in Seem. journ. 1866. p. 146), foliis ternis quaternisve petiolatis ovato-acuminatis apice obtusiusculis basi rotundatis vel elliptico-lanceolatis subrhombeisve apice obtusiusculis basi acutis utrinque præsertim ad nervos pilosulo-pubescentibus membranaceis reticulato-nervulosis subpellucidis pellucido-punctulatis quinquenerviis, centrali nervo ad apicem usque ducto lateralibus 2 alte supra medium ductis externis subtilioribus brevioribus nervulose juxta marginem apice currente, petiolo pubescente, amentis axillaribus terminalibusque filiformibus folia duplo superantibus densifloris, pedunculo pilosulo-pubescente, bractea rotundata centro peltata subsessili, ovario semi-immerso ovato, stigmatem papillosulo pilosulo. ☉ In Mexico (Botteri ! et Sallé ! in h. DC.). Caulis simplex ? 0,002 crassus, pubescens. Limbi 0,04 longi, 0,025 lati. Petioli 0,015 longi. (v. s.)

338. *P. LANCEOLATA* (Cas. DC. in Seem. Journ. 1866, p. 145), foliis quaternis-senis breviter petiolatis lanceolatis basi in petiolum subcurrentibus supra ad nervos pubescentibus dein nervis exceptis glabris subtus glabris rigidulo-membranaceis pellucido-punctulatis enervulosis quinquenerviis, nervis 3 centralibus subtus subcarinato-prominulis, nervo centrali ad apicem usque ducto lateralibus alte supra medium ductis externis subtilioribus nervulo pellucido juxta marginem currente, petiolo supra pubescente, amentis axillaribus breviter pedunculatis filiformibus subdensifloris folia duplo superantibus, pedunculo glabro petiolum æquante, bractea rotundata centro peltata subsessili, ovario postea semi-immerso apice oblique acutato, stigmatem puberulo. ☿ In Andibus Quitensibus alt. 9000 ped. (Spruce n. 6110 ! in h. DC., Jameson n. 343 ! in h. DC.). Ramuli teretiusculi, glabri, ad nodos hirtelli, 0,005 crassi. Limbi 0,035 longi, 0,012 lati. Petioli 0,003 longi. Amenta vix 0,001 crassa. (v. s.)

339. *P. HYPOLEUCA* (Miq. ! syst. p. 136), foliis superioribus ternis quaternisve longiuscule petiolatis obovato-ellipticis apice acutiusculis acuminatisve basi acutis vel obovato-rotundatis infimis oppositis minoribus omnibus supra atro-viridibus glabris subtus ad nervos parce hirtellis albicantibus quinquenerviis, petiolo sursum hirtello. ☿ In ins. Sandwich (Gaudich. ! in h. Deless.). Fruticulus 0,3 altus, basi e nodis radicans, fuscus, subligneus, pennam corvinam crassus. Limbi 0,02-0,045 longi, 0,015-0,02 lati. Petioli 0,01-0,015 longi. (v. s.)

340. *P. MEMBRANACEA* (Hook. et Arn. ! in Beech. bot. p. 96, non Miq.), foliis oppositis ternisve breviter petiolatis elliptico-lanceolatis utrinque acutis apice acuminatis glabris tenuiter membranaceis reticulato-nervulosis, quinquenerviis, centrali nervo ad apicem usque ducto lateralibus 2 alte externis ad medium ductis nervulo juxta marginem apice currente, amentis axillaribus pedunculatis filiformibus densifloris, bractea rotundata centro peltata, ovario emerso, stigmatem minuto. ☉ In ins. Oahu (Beechey ! in h. Kew.). ins. Mariannis (Gaudich. n. 216 ! in h. Deless.). — Miq. ! syst. p. 120. Herba ? ramosa. Ramuli glabri teretes. Limbi vix 0,04 longi, 0,015 lati. Petioli 0,005-0,007 longi. Pedunculi 0,01 longi. (v. s.)

341. *P. RHEXIFOLIA* (Moritz mss. in h. Brit. Mus.), foliis quaternis longiuscule petiolatis elliptico-lanceolatis apice acuminatis, acumine acuto, basi acutis utrinque glabris membranaceis subpellucidis quinquenerviis, petiolo glabro, amentis axillaribus terminalibusque folia æquantibus?, ovario oblique stigmatifero. ♀ In sylvis umbrosis ad saxa muscosa rarior prope coloniam Tovar (Moritz n. 1868! in h. Brit. Mus.). Suffructulus. Caulis glaber. Limbi 0,06 longi, 0,03 lati. Petioli 0,02 longi. (v. s.)
342. *P. GAUDICHAUDII* (Miq.! syst. p. 137), foliis ternis quaternisve breviter petiolatis elliptico-lanceolatis utrinque acutis vel inferioribus rotundatis obovatisve omnibus utrinque glabris rigido-coriaceis reticulato-nervulosis quinquenerviis, centrali nervo ad apicem usque ducto, lateralibus fere ad apicem externis ad medium usque ductis nervulo juxta marginem currente, petiolo glabro, amentis axillaribus terminalibusque pedunculatis filiformibus densifloris, pedunculo glabro, bractea rotundata centro peltata, bacca semiimmersa ovoidea. ♀ In ins. Oahu (Seemann n. 2258! in h. Kew. et DC., Meyen! in h. Berol., Gaudich! in h. DC.). Caulis erectus, basi radicans, glaber, dichotome ramosus. Limbi 0,04 longi, 0,022 lati. Petioli 0,008 longi. Pedunculi 0,008 longi. (v. s.)
343. *P. PACHYPHYLLA* (Miq.! syst. p. 137), foliis oppositis quaternisve petiolatis obovatis vel obovato-orbicularibus supra hirsutis subtus puberulis ciliolatis vel ætate glabratibus rigidis opacis trinerviis vel quinquenerviis plerumque trinerviis, centrali nervo ad apicem usque ducto, lateralibus alte externis sub medium ductis, amentis axillaribus longe pedunculatis filiformibus densifloris glabris, bacca basi immersa ovato-acuta. ♀ In ins. Oahu (Beechey!, Seemann n. 2259! in h. Brit. Mus. et DC., Chamisso! in h. Berol.). Caulis glaber, 0,2 altus, basi radicans, dichotome ramosus. Ramuli juniores pubescentes. Limbi 0,035 longi, 0,023 lati. Petioli 0,01-0,015 longi. Pedunculi 0,025 longi. Amenta matura 0,025 longa. (v. s.)
344. *P. PAVONIANA*, foliis oppositis quaternisve petiolatis elliptico-lanceolatis glabris supra ad nervos pilosis apicem versus ciliolatis pellucido-subcoriaceis pellucido-punctatis reticulato-nervulosis trinerviis vel quinquenerviis, nervis 3 centralibus magis conspicuis subtus subprominulis nervulo juxta marginem currente, petiolo glabro, amentis axillaribus filiformibus glabris breviter pedunculatis, bractea rotundata centro peltata, ovario per anthesin emerso glabro, stigmatibus exiguos, bacca globosa hirtella. ☉ In Peruvia (Pav.! in h. Boiss.). Caulis glaber, ad nodos puberulus. Limbi vix 0,06 longi. Petioli 0,015 longi. Pedunculi 0,008-0,01 longi. (v. s.)
345. *P. SANDWICENSIS* (Miq.! syst. p. 126), foliis oppositis ternisve longiuscule petiolatis obovato-spathulatis utrinque pilosulis ciliolatis membranaceis subopacis pellucido-punctulatis reticulato-nervulosis trinerviis, centrali nervo ad apicem usque ducto lateralibus alte ductis nervulo interrupto juxta marginem apice currente, petiolo piloso-hirtello, amentis axillaribus terminalibusque longe pedunculatis folia triplo quadruplove superantibus filiformibus subdensifloris, bractea rotundata centro peltata sessili, ovario semi-immerso, stigmatibus minuto, bacca impressa ovato-globosa. ☉ In ins. Oahu (Meyen! in h. Berol.). — Miq. ill. p. 19. p. 14. Herba succulenta 0,1 alta, caules e radice fibrosa agens plures simplices vel parce ramosos subadpresse hirtellos. Limbi 0,02-0,025 longi, 0,015 circ. lati. Petioli 0,009 circ. longi. Pedunculi 0,02 longi. (v. s.)
346. *P. FENDLERIANA*, foliis oppositis ternisve oblongo-ellipticis apice acutiusculis basi attenuatis vel e basi attenuata oblongo-obovatis ovatisve vel minoribus orbicularibus tenuiter pubescentibus membranaceis pellucidis subtiliter trinerviis, petiolo glabro, amentis axillaribus terminalibusque subdensifloris pedunculatis, bractea rotundata centro peltata subsessili, ovario impresso, stigmatibus minuto. ☉ In Venezuela prope Tovar alt. 6000 ped. (Fendler n. 1174!, 1175! in h. DC.). Herba erecta. Caulis basi radicans, teres, glaber. Ramuli pilis reflexis dense puberuli. Limbi 0,035 longi. Petioli 0,02 longi. Pedunculi 0,012 longi. (v. s.)
347. *P. HIRSUTA* (Miq.! syst. p. 143), foliis ternis quaternisve vel apice ramu-

lorum oppositis petiolatis ellipticis apice attenuato-obtusiusculis basi obtusis vel inferioribus obovatis vel spatulato-ellipticis utrinque adpresse villosis coriaceis trinerviis, amentis axillaribus terminalibusque breviter pedunculatis filiformibus subdensifloris folia multoties superantibus, bractea rotundata centro peltata subsessili, ovario emerso attenuato, stigmatе minuto. ☉ In Brasilia (Sello! in h. Berol.). — Miq. ! in Mart. flor. Bras. fasc. 11. p. 14. Herba erecta, 0,3 circ. alta. Caulis hirsutus. Limbi 0,03 longi. Petioli 0,006-0,007 longi. Pedunculi 0,01 longi. (v. s.)

348. *P. POLYSTACHYA* (Miq. ! syst. p. 116), foliis ernis petiolatis e basi cuneata obovato-subrhombeis rhombeisve apice breviter subattenuatis obtusiusculis utrinque sparse pilosis ciliolatis membranaceis reticulato-nervulosis et triner-viis, centrali nervo ad apicem usque ducto lateralibus 2 alte supra medium ductis nervulo juxta marginem apice currente, petiolo piloso-pubescente, amentis axillaribus terminalibusque folia duplo superantibus densifloris, bractea rotundata centro peltata brevissime pedicellata, ovario oblongo semiimmerso, stigmatе penicillato. ☉ In Antillis (Rich. ! in h. Francov. et DC. Bonpl. ! in h. Berol., h. Willd. n. 748 !), Barbadoes (Maycock plant. Barb. p. 40 quoad synonym. Jacq.), in Brasilia ad Petropolim super arborum truncos (Wawra et Maly n. 427, fide Wawra Ergeb. bot. p. 131); in horto Florentino culta. — Miq. in Mart. flor. Bras. fasc. 11. p. 14. t. 14. fig. 7, Hook. exot. flor. t. 23. Piper obtusifolium Jacq. icon. rar. v. 1. p. 2. t. 9, Lam. ill. p. 82. *P. polystachyon* Vahl enum. v. 1. p. 354, Willd. ! sp. v. 1. p. 167, Loud. encycl. p. 29. n. 537, Ait. hort. Kew. v. 1. p. 49. *Micropiper Blumei* Miq. comm. phyt. p. 53. t. 8. fig. g. Caulis herbaceus pubescens. Limbi majores 0,03-0,04 longi, 0,02 lati. Petioli usque ad 0,005 longi. (v. s.)
349. *P. PEDUNCULATA* (Cas. DC. in Seem. journ. 1866. p. 146), foliis plerumque ternis breviter petiolatis subrhombico-ellipticis ellipticisve apice obtusis basi acutiusculis supra subtusque pubescentibus rigidulo-membranaceis subopacis nigro-punctulatis trinerviis, centrali nervo ad apicem usque lateralibus supra medium ductis, nervulo apice juxta marginem currente, petiolo pubescente, amentis axillaribus terminalibusque sparsis vel apice caulis confertis folia duplo triplove superantibus longe pedunculatis, pedunculo petiolum multoties folium duplo superante hirtello-pubescente, bractea rotundata centro peltata subsessili, ovario impresso ovato, bacca globosa. ♂ ? In ins. Bourbon (Rich. ! in h. Francov.). Inter detritus foliorum et muscos in sylvis cres-cens. Limbi 0,03 longi, 0,016 lati. Petioli 0,007 longi. (v. s.)
350. *P. CAMPTOTHRICHA* (Miq. ! syst. p. 144), foliis quaternis subsessilibus infe-rioribus obovato-rotundatis superioribus obovato-rhombis oppositis omnibus utrinque dense pubescentibus subtus glanduloso-punctulatis obscure triner-viis inferioribus coriaceis opacis superioribus membranaceis, amentis axilla-ribus terminalibusque filiformibus subdensifloris folia per anthesin multoties superantibus, bractea rotundata centro peltata subsessili, ovario semiimmerso, stigmatе carnosulo-puberulo. ☉ In Mexico (Schiede ! in h. Berol., Buchinger n. 545 ! in h. Lenorm. et DC.), in arboribus ad Hacienda de la Laguna (Miq. in Linn. v. 18. p. 711). — Miq. ill. p. 21. t. 19. Planta succulenta, erecta, 0,25 alta. Radix fascicularis, crassiuscule fibrosa. Caulis teretiusculus, sul-catus, pilis recurvatis longis nitidis inferne aureo-fuscis dense villosus. Limbi 0,013 longi latique. (v. s.)
351. *P. RUBELLA* (Hook. ! flor. exot. t. 58), foliis 2-5-verticillatis plerumque quaternis petiolatis ellipticis vel elliptico-lanceolatis subcoriaceis brevi-pilosis trinerviis, centrali nervo subtus prominente lateralibus tenuissimis, amentis axillaribus, pedunculo puberulo dilute rubello, bractea rotundata centro peltata, ovario emerso, stigmatе paullo antico minuto penicillato. ☉ In Mexico inter Vera-Cruz et Orizaba (Meissner ! in h. Kew.). — Miq. syst. p. 164, Piper rubellum Haw. Herbula caespitosa-gregaria, pusilla. Caulis carnosus, ramosissimi, rubelli, pilosi. Limbi 0,015 longi. Petioli 0,001-0,002 longi. (v. s.)
352. *P. KLOTZSCHIANA* (Miq. syst. p. 145), foliis quaternis-senis breviter petio-latis inferioribus obovato-spathulatis apice rotundatis truncatisve superioribus lanceolato-spathulatis apice obtusis acutiusculisve basi cuneatis omnibus

utrinque incano-hirsutis rigidis trinerviis, nervis subtus prominulis, petiolo hirsuto, amentis axillaribus apice ramulorum umbellatim confertis brevissime pedunculatis densifloris folia superantibus, pedunculo communi petiolum sequante hirsuto, bractea rotundata centro peltata albo-marginata, cæteris nondum efformatis. ☉ In Peruvia subandina ad Maynas (Poeppig! in h. Berol.). Caulis 0,35 altus, hirsutus, erectus, basi simplex, apice ramulosus. Limbi 0,025 longi; 0,01 lati. Petioli 0,005 longi. (v. s.)

353. *P. MATHEWSIANA* (Miq. in Hook. Lond. journ. bot. v. 4. p. 423), foliis ternis quinis septenisve brevissime petiolatis rhombeo-spathulatis apice attenuato-obtusiusculis basi cuneatis utrinque pubescentibus ciliolatis membranaceis pellucidis reticulato-nervulosis et trinerviis, amentis axillaribus terminalibusque subdensifloris breviter pedunculatis, bractea rotundata centro peltata subsessili, ovario impresso apice breviter attenuato, stigmatè minuto. ☉ In Peruvia ad Zacapota (Mathews n. 1688! in h. Kew.). Suffrutescens. Caulis erectus, teres, pubescens, verticillatim ramulosus. Limbi 0,03 longi. Petioli 0,003-0,005 longi. Pedunculi 0,005-0,007 longi. (v. s.)

354. *P. VARIIFOLIA*, foliis plerumque quinis brevissime petiolatis, e basi acuta oblongo-obovatis apice rotundatis vel e basi cuneata obovato-ellipticis, apice attenuato-obtusis acutiusculisve utrinque dense subincano-tomentosis rigidulis opacis trinerviis, nervis subtus prominulis, amentis axillaribus verticillatis per anthesin folia dimidio superantibus, pedunculo petiolum duplo superante, bractea rotundata centro peltata subsessili, ovario emerso acutato. ☉ In Peruvia (Poeppig! in h. Petrop.). Herba erecta; circ. 0,4 alta, ramosa. Caulis dense tomentosus. Limbi majores circ. 0,02 longi fere 0,01 lati. Petioli 0,001 longi. (v. s.)

355. *P. PALLENS* (Kunth! in H. et B. nov. gen. v. 1. p. 69), foliis ternis breviter petiolatis ellipticis utrinque obtusis atque subtilissime puberulis membranaceis subpellucidis enervulosis, petiolo glabro, amentis axillaribus terminalibusque folia circit. triplo superantibus filiformibus densifloris; bractea rotundata fusco-punctata centro peltata, ovario subimmerso oblongo, stigmatè minuto. ☉ In scopulosis Andium Popayanensium locis altis opacatis inter Pisone et convallem Cucunuco alt. 1200 hexap. (H. et B. l. c. et Bonpl. in h. Francov.). Piper pallens Roem. et Sch. syst. v. 1. p. 334. Herba. Caulis repens glaber. Limbi 0,016 longi, 0,01 lati. Petioli 0,005 longi. (v. s.)

356. *P. TRIPFOLIA* (A. Dietr. sp. v. 1. p. 173), foliis ternis petiolatis ellipticis vel subrotundato-ellipticis utrinque obtusis apice imo emarginulatis, supra glabris subtus ad nervos pilosis ciliatis rigidulo-membranaceis pellucido-punctulatis enervulosis trinerviis, nervis subtus prominulis, centrali ad apicem usque ducto, lateralibus nervis 2 supra medium ductis, amentis terminalibus longiuscule pedunculatis folia triplo-quadruplo superantibus longiuscule pedunculatis filiformibus densifloris, pedunculo petiolum multum superante, bractea rotundata glanduloso-punctata centro peltata subsessili, ovario semiimmerso oblongo apice acutato, stigmatè discoideo-puberulo, bacca subimmersa subglobosa apice oblique mucronata. In ins. Guadaloupa (Rich. in h. Francov., Duchassaing! in h. DC., Bertero! in h. DC.), Martinica (h. Berol.). Cajenna (Aubl. pl. Guy. v. 1. p. 22). — Miq. syst. p. 148, Griseb. flor. w. Ind. p. 168. Saururus hederaceus triphyllus Plum. Am. ed. Par. p. 52. t. 68. Piper trifolium L. sp. p. 43, Hill dict. v. 26. t. 20. fig. 3, Aublet l. c., Lam. ill. p. 83, Vahl enum. v. 1. p. 351, Willd. sp. v. 1. p. 167. Herba ad truncos arborum reptando-scandens. Caulis tetragonus, piloso-pubescentis. Limbi 0,011 longi, 0,012 lati. Petioli 0,003 longi. (v. s.)

357. *P. PULCHELLA* (A. Dietr. sp. v. 1. p. 177), foliis ternis-quinis breviter petiolatis oblongo-obovatis subspathulatisve apice obtusis obtusiusculisve basi subcuneatis crasse carnosius subtus concavis supra subtusque minutissime puberulis subtus ætate rubescentibus rigidulis opacis trinerviis, petiolo minute puberulo, amentis terminalibus axillaribusque sæpe verticillatis pedunculatis subdensifloris filiformibus folia duplo triplo superantibus, pedunculo petiolum multum superante minute puberulo, bractea rotundata centro peltata, ovario emerso ovato, stigmatè minuto puberulo. ☉ In Jamaica (Griseb. flor. w. Ind. p. 167); Cuba (Wright n. 2265 fide Griseb. Cat. pl. Cub. p. 64).



- in caldariis culta (h. DC.), Sandwich (Beech. bot., p. 96); h. Willd. n. 753! fol. 1. spec. infer. et fol. 3 exceptis, 754! — Miq. syst. p. 166; Griseb. flor. W. Ind. p. 167. Saururus n. 8. Brown Jam. p. 204. Piper pulchellum Vahl Enum. v. 1. p. 353. P. verticillatum L. f. sp. ed. p. 43 (h. Lin.); Hill dict. v. 26. p. 20. f. 20. p. 5, Lath. ill. p. 83, Willd. sp. v. 1. p. 168; Alt. hort. Kew. v. 1. p. 50. Pep. verticillata Miq. syst. p. 166; Griseb. l. c. Herba palmaris, succulenta, pallide viridis, verticillatim ramosa, pube minute brevissima pilisque majoribus patentibus intermixtis vestita. Limbi 0,015 longi, 0,006-0,008 lati. Petioli 0,001-0,002 longi. (v. s.)
358. P. LINNÆI (Cas. DC. in Seem. jourd. 1866. p. 145), foliis plerumque quinis petiolatis linearibus apice obtusis supra pilosulis membranaceo-pellucidis trinerviis, amentis terminalibus subdensifloris filiformibus longiusculis pedunculatis folia duplo superantibus, bractea rotundata carnosa centro peltata, ovario subimmerso, stigmate minuto, bacca semi-immersa apice oblique mucronulata flavicante. ☉ In sylvis Andium Quinsium (Jameson n. 89!), Venezuela prope coloniam Tovar (Fendler n. 1176! in h. DC., Moritz n. 803! in h. Brit. Mus.), Caracas (Gallmer n. 333! in h. Berol.). Herba repens. Caulis filiformis. Limbi 0,01 longi, 0,002 lati. Petioli 0,003-0,006 longi. Pedunculi 0,005-0,007 longi. (v. s.)
359. P. DISCOLOR, foliis ternis petiolatis ellipticis utrinque subacutis glabris subtus ad nervos subtiliter velutino-puberulis membranaceis rigidulis subopacis laxe reticulato-nervulosis trinerviis, centrali nervo ad apicem usque ducto subtus subprominulo, lateralibus nervis supra medium ductis subtilibus, amentis filiformibus tenuibus subaxillifloris. ☉ In ins. Jamaica (Mac-Fadyen f. in h. Kew.). Herba. Caulis basi radicans, 0,05-0,06 altus, teres, tenuis, glaber. Limbi 0,015 longi, 0,007-0,008 lati. Petioli 0,004 longi. Pedunculi 0,004 longi. (v. s.)
360. P. ELLIPTICIFOLIA, foliis ternis petiolatis ellipticis apice breviter attenuatis acutiusculis basi acutiusculis glabris ciliatis basin versus pilosulis membranaceis trinerviis nervis tenuibus, petiolo hirtello, amentis axillaribus terminalibusque pedunculatis, pedunculo hirtello, bractea rotundata membranacea centro peltata subsessili, ovario emerso attenuato, stigmate minuto. ☉ In Venezuela ad coloniam Tovar (Fendler n. 1171! in h. DC.). Herba erecta, basi radicans. Caulis hirtellus. Limbi 0,025 longi, 0,015 lati. Petioli circ. 0,01 longi. Pedunculi 0,008 longi. (v. s.)
361. P. COURTALLIENSIS (Miq. in Hook. Lond. journ. bot. v. 5. p. 549), foliis plerumque ternis brevissime petiolatis rhombeo-ellipticis apice acutiusculis basi cuneatis vel oblongo-obovatis apice obtusis basi acutis glabris rigidis subopacis trinerviis, centrali nervo ad apicem usque lateralibus ad medium usque ductis, amentis axillaribus terminalibusque brevibus pedunculatis densifloris, bractea rotundata centro peltata, ovario impresso apice obtuso, stigmate punctiformi. ☉ In ins. Ceylon (Thwaites n. 831! in h. Kew. et 2173! in h. DC.). — Miq. in Wight icon. t. 1923. f. 2, Thwaites enum. p. 292. Herba repens. Caulis glaber, sulcatus, quadrangulus. Limbi 0,025 longi, 0,007-0,008 lati. Petioli 0,003-0,008 longi. Pedunculi 0,01-0,012 longi. (v. s.)
362. P. GALIODES (Kunth! in H. et B. nov. gen. v. 1. p. 71), foliis quaternis-quinis sessilibus superioribus oblongo-spathulatis vel oblongis apice obtusis inferioribus obovatis membranaceis pellucido-punctatis subpellucidis parce nervulosis vel enervulosis inconspicue trinerviis nervulose juxta marginem currente, amentis axillaribus terminalibusque folia multoties superantibus subaxillifloris breviter pedunculatis, plerumque apice ramulorum verticillatis pedunculo glabro, bractea rotundata centro peltata subsessili, ovario semi-immerso, stigmate glanduloso carnosulo-papilloso, bacca ovata apice subrotellata. ☉ In Brasilia (Sellow in h. Berol.), Andibus quitenisibus alt. 10000 ped. (Jameson n. 444! in h. DC. et 814! in h. Kew.), regno Novo-Granatensi juxta catractam Tequendana (H. et B. l. c., Humb. in h. Willd. n. 762 et in h. Berol.); Nova-Granata (Linden n. 231! et 233! in h. Kew.), prov. Pasto alt. 2500 ped. (Piana n. 56! in h. suo), Venezuela (Fendler n. 1171! 1178! 1180! in h. DC.; Moritz n. 1782! in h. Brit. Mus., Gallmer n. 830! in h. Be-

rol.), Jamaica (Griseb. flor. W. Ind. p. 167.). — Humb. pl. æq. v. 2. p. 58, Kunth synopsis. v. 1. p. 123, Miq. l. syst. p. 156, illust. p. 22. t. 20, in Mart. flor. Bras. fasc. 11. p. 15. t. 2. fig. 2, Griseb. flor. W. Ind. p. 167. P. galioides Rœm. et Sch. syst. v. 1. p. 335. Herba erecta. Caulis teretiusculus, subtilissime puberulus, sulcatus, subangulosus, 0,004-0,005 crassus. Limbi 0,015-0,02 longi. (v. s.)

α, *longifolia*, foliis lineari-oblongis apice obtusis basi subattenuatis subsessilibus, limbis 0,02-0,03 longis. ☉ In Boliviae prov. Ladrilloni ad rivulum in umbrosis reg. temp. alt. 2650 ped. (Mandon n. 1120 ! pro parte in h. DC.), Ecuador (Fraser! in h. DC.), Nova-Granata (Holton n. 231! in h. DC.), Costa-Rica prope San Jose (Hoffmann n. 187! in h. Berol.).

β, *nigro-punctulata*, foliis e basi cuneata obovatis glabris nigro-punctulatis fere enervulosis, limbis 0,008 longis. In regione æquatoriali (Jame-son! in h. Kew.), Brasilia prope Petropolim ubi ad arborum truncos (Wawra et Maly n. 441! fide Wawra Ergeb. bot. p. 131).

γ, *Menkeana*, foliis rigidis crebre pellucido-punctulatis, limbis 0,1 longis. In prov. Minas Geraes (Mart.! in h. monac.). Pep. Menkeana Miq. l. in Linn. v. 20. p. 124 et in Mart. flor. Bras. fasc. 11. p. 15.

363. P. INÆQUALIFOLIA (Ruiz et Pav. l. flor. per. v. 1. p. 30. t. 46. fig. a), foliis quaternis senisve petiolatis inæqualibus superioribus longioribus oblongo-spathulatis apice obtusis basi in petiolum decurrentibus inferioribus obovatis omnibus glabris rigidulo-membranaceis pellucido-punctulatis subopacis aromaticis trinerviis, centrali nervo ad apicem usque ducto, lateralibus ad medium usque ductis ægre discernendis, petiolo glabro, amentis axillaribus terminalibusque verticillatis subdensifloris folia multoties superantibus, bractea rotundata centro peltata subsessili, ovario semi-immerso obovato, stigmatibus carnosulo. ☉ In Peruvia Limæ, Huanuci, Cantæ locis saxosis et supra arbores (R. et P. l. c., Pav. l. in h. Boiss.), Caracas (Humb. in h. Willd. n. 759! et 760! et Gallmer! in h. Berol.), in hort. Madrit. cult. (h. DC.) — Kunth! in H. B. nov. gen. v. 1. p. 71, synopsis. v. 1. p. 122, Miq. l. syst. p. 148. Piper inæqualifolium Vahl enum. v. 1. p. 355. P. aromaticum Willd. l. enum. hort. Berol. supp. p. 4. Suffruticulus herbaceus, aromaticus, pedalis, e basi radicante gregarie erectus. Caulis glaber vel velutino-puberulus. Limbi 0,035-0,05 longi, 0,008-0,015 lati. Petioli 0,005 longi. (v. s.)

364. P. OLIVACEA (Cas. DC. in Seem. journ. 1866, p. 146), foliis plerumque quinis breviter petiolatis subpathulato-oblongis apice obtusis obtusiusculisve basi subcuneatis utrinque glabris coriaceis subopacis subpellucido-punctulatis subtus olivaceis rugulosis trinerviis, centrali nervo ad apicem usque ducto, lateralibus alte supra medium ductis, petiolo glabro, amentis axillaribus terminalibusque folia multoties superantibus filiformibus densifloris, pedunculo glabro petiolum multoties superante, bractea rotundata pellucido-punctata centro peltata subsessili, ovario emerso ovato apice oblique acutiusculo. ♀ In Costa-Rica (Hoffmann n. 810! in h. Berol.). Suffrutex erectus, basi radicans. Rami ramulique dense pubescentes, nodosi, cylindrici. Limbi 0,017 longi, 0,006 lati. Petioli 0,004 longi. Amenta florentia 0,15 longa, 0,001 crassa. (v. s.)

365. P. BOIVINII (Cas. DC. in Seem. journ. 1866, p. 146), foliis ternis quaternisve breviter petiolatis elliptico-obovatis apice obtusis basi cuneatis utrinque glabris rigidis opacis trinerviis, petiolo glabro, amentis axillaribus folia multum superantibus subdensifloris, pedunculo glabro petiolum superante, bractea rotundata centro peltata subsessili, ovario impresso apice obtuso, stigmatibus carnosulo, bacca glabra ovata acuta. ☉ In insul. Moohely (Boivin! in h. Brit. Mus.). Suffruticulus procumbens, glaber. Limbi 0,02 longi, 0,011 lati. Petioli 0,01 longi. Pedunculi 0,01 longi. (v. s.)

366. P. KUNTHIANA, foliis quaternis breviter petiolatis elliptico-rotundatis coriaceis pellucido-punctatis puberulis vel glabris uninerviis, nervo subtus prominulo, amentis axillaribus terminalibusque breviter pedunculatis, pedunculo puberulo, bractea rotundata carnosula pellucido-punctata centro peltata, ovario profunde immerso, stigmatibus carnosulo penicillato, bacca basi immersa

- ovata oblique rostellata. ☉ In Andibus Quitensibus alt. 10000 ped. super arborum truncos (Jameson n. 668! in h. Boiss.). Caulis herbaceo-coriaceus, puberulus, e nodis radicans, dichotomus. Limbi 0,011-0,012 longi. Petioli 0,003 longi. Pedunculi 0,01 longi. (v. s.)
367. *P. MICROPHYLLA* (Kunth! in H. et B. nov. gen. p. 69. t. 15), foliis quaternis quinque brevissime petiolatis oblongo-lanceolatis apice acutiusculis basi in petiolum attenuatis utrinque minutissime puberulis apicem versus minutissime ciliolatis rigidulis subopacis glanduloso-punctulatis enervulosis uninerviis, petiolo ciliolato, amentis terminalibus filiformibus densifloris pedunculatis folia duplo superantibus, bractea rotundata glandulosa centro peltata subsessili, ovario ovato, stigmate carnosulo post anthesin semi-immerso. ☉ In lapidosis frigidis Andium Quinduensium juxta El Bogueon del Parano alt. 1650 hexap. (H. et B. l. c., Humb. in h. Willd. n. 749! et in h. Berol.). — Kunth synops. v. 1. p. 121, Miq! syst. p. 167. *Piper microphyllum* Rœm. et Sch. syst. v. 1. p. 334. Herba perennans, succulento-coriacea, basi aphylla, repens, radicans vel terræ immersa, vix spithamea. Caulis ramique sub quadranguli, plicato-sulcati, juniores minutissime puberuli. Limbi 0,004-0,007 longi, 0,0015-0,002 lati. Petioli vix 0,001 longi. (v. s.)
368. *P. APHYLLA*, foliis plerumque quaternis brevissime petiolatis oblongo-ellipticis apice obtusis basi subattenuatis minutis glabris ciliolatis membranaceis pellucido-punctulatis obsolete nervulosis uninerviis nervuloque juxta marginem corrente, petiolo puberulo, amentis terminalibus subdensifloris, pedunculo subtiliter hirtello petiolum superante, bractea rotundata centro peltata, ovario subimmerso, stigmate minuto ☉ In sylvis Andium Quinduensium (Jameson n. 690! in h. Boiss.). Herba basi aphylla, repens, radicans, spithamea. Ramuli juniores subtiliter puberuli. Limbi 0,004-0,01 longi, 0,0015-0,004 lati. Petioli 0,001 longi. Pedunculi 0,003-0,005 longi. (v. s.)
369. *P. GALLAPAGENSIS* (Miq! in Hook. Lond. journ. bot. v. 4. p. 426), foliis ternis quaternisve vel oppositis subsessilibus ovato-oblongis apice obtusis utrinque glabris membranaceis uninerviis, amentis axillaribus terminalibusque brevibus densifloris pedunculatis, bractea rotundata centro peltata, ovario subimmerso. ☉ In Archipelagi Gallapagos insulis Chatham et James (Wood et Darwin! in h. K. w.). Herba. Caulis velutino-puberulus, striatus. Limbi 0,005 longi. Pedunculi 0,004 longi. Amenta vix 0,02 longa. (v. s.)
370. *P. ORBICULARIS*, foliis oppositis ternisve minutis subsessilibus orbicularibus utrinque pubescentibus pellucido-membranaceis enervulis, amentis terminalibus axillaribusque densifloris glabris, bractea rotundata centro peltata, ovario immerso, stigmate carnosulo-puberulo, bacca basi immersa ovata brevi. ☉ In Brasilia ad Rio-Janeiro (Gaudich. in 1096! in h. DC.). Herbula repens. Caulis filiformis tenuissimus glaber. Limbi 0,003 in diam. Petioli vix 0,001 longi. (v. s.)

*Species affines non satis notæ.*

371. *P. RHOMBEA* (Ruiz et Pav. flor. Peruv. v. 1. p. 31. t. 46. fig. c.), foliis quaternis brevissime petiolatis oblongo-rhombeis subtus convexis supra planis purpurascenscentibus eveniis, amentis terminalibus purpureis cylindricis folia triplo superantibus densifloris. In sylvis Pillao et Chacahuassi supra arbores et saxa. *Piper rhombeum* Vahl enum. v. 1. p. 353. Caules nonnulli gemini furcati, teretes, ad nodos nitidi, aliquando procumbentes radicales. (Ex R. et P. l. c.)
372. *P. CONGESTA* (Kunth! in H. et B. nov. gen. v. 1. p. 68. t. 14), foliis quino-septenis subsessilibus caule simplici dense approximatis ovato-acutis basi plerumque contractis utrinque glabris rigidulo-membranaceis subopacis pellucido-punctulatis enerviis, amentis apice caulis dense congestis filiformibus densifloris folia paulo superantibus brevissime pedunculatis. In declivitate Andium Peruvianorum inter flumen Chulucanas et pagum Guancabamba alt. 950 hexap. (Kunth l. c. et Bonpl. in h. Francov.). — Kunth synops. v. 1. p. 120, Miq. syst. p. 166. *Piper congestum* Rœm. et Sch. syst. v.

- p. 334. Caulis bipedalis, erectus, ramosus, carnosus, crassitie pennæ cygneæ. Limbi 0,025 longi, fere 0,02 lati. (v. s.)
373. *P. ANGUSTATA* (Kunth) in H. et B. nov. gen. v. 1. p. 68), foliis ternis quaternis? longiuscule pedunculatis rhombeo-lanceolatis apice acuminatis acutiusculis basi angustatis acutis utrinque glabris rigidulo-membranaceis subopacis pellucidis punctulatis 4-5-nerviis, nervulis raris nervulo tamen juxta marginem apice currente sæpe interrupto, nervis externis ante medium evanidis, petiolo glabro. In umbrosis prov. Cumanæ prope Cumanacoa et San Ferdinando alt. 100 hexap. (Humb. l. c. et in h. Willd. n. 745!). Caulis e nodis fibrillas radicales crassiusculas mittens. Caulis coriaceo-succulentus, subangulosus. Limbi 0,06 longi, 0,02 lati. Petioli fere 0,02 longi. (v. s.)
374. *P. PEEPULOIDES* (Kunth in H. et B. nov. gen. v. 1. p. 69), foliis quaternis petiolatis obovato-subrotundis obtusis glabris junioribus pubescentibus carnosius eveniis, amentis axillaribus solitariis. In radicibus montis Cotopaxi locis frigidis subapricis scopulosis juxta pagum Mubalo alt. 1590 hexap. — Kunth synops. v. 1. p. 121, Miq. syst. p. 157. *P. peepuloides* Roem. et Sch. syst. v. 1. p. 335. (non Roxb.). Caulis spithameus, procumbens nec repens, ramosus, tetragonus, glaber. Ramuli tenuissime pubescentes. Folia 4-5 lin. longa. Petioli lineam longi. (Ex Kunth l. c.)
375. *P. ROLLINGO* (Link, Jahrb. 1, 3. p. 65, fide Roem. et Sch. Mant. v. 1. p. 247), foliis quaternis lanceolatis obtusis basi attenuatis apiculatis ciliatis nervoso-venosis, amentis terminalibus fasciculatis longissimis. — Miq. syst. p. 154. *Piper Rollingo* Roem. et Sch. l. c. Folia 9-10 lin. longa. Amenta 3-4 poll. longa. (Ex Roem. et Sch. l. c.)
376. *P. Plicata* (Opiz in Reliq. Hænk. fasc. 3. p. 163), foliis ternis brevissime petiolatis late lanceolatis margine reflexis coriaceis glabris supra excavato-punctatis subtus subpulverulentis, pedunculis terminalibus solitariis brevissimis, amentis pedunculos multo superantibus erectis filiformibus, bractea peltata.  $\frac{1}{2}$  In Acapulco. — Miq. syst. p. 153. *P. Pereskiaefolia* valde affinis; an eadem? Caulis glaber, triangularis, angulis acutis. Limbi 0,04 longi, fere 0,02 lati. Petioli 0,002 longi.
377. *P. SUAVEOLENS* (Hamilt. prodr. p. 2), foliis subternis ovato-oblongis glaberrimis, amentis solitariis terminalibus. In Cuba. — Miq. syst. p. 168. *Piper suaveolens* Desf. Cat. hort. Par. v. 3. p. 414. Verisimiliter eadem ac *P. inæqualifolia* R. et P. Caulis perennis, carnosus-subtripetalis. Rami subsimplicis. Folia fricata odorem gratum Cariophylli aromatici spirant. (Ex Ham. l. c.)
378. *P. SWARTZIANA* (Miq. syst. p. 155), foliis inferioribus oppositis decussatis superioribus quaternis omnibus petiolatis lineari-oblongis obtusis integris subtus pallide rubro-maculatis pallidioribus utrinque glabris, amentis terminalibus pedunculatis, pedunculo folia æquante glabro, bractea subrotundata, bacca oblonga minuta. Inter muscos ad radices arborum in montibus altis Jamaicae. *Piper filiforme* Sw. (non Pers.) prodr. p. 16, icon. t. 5, flor. Ind. occ. v. 1. p. 73, Vahl enum. v. 1. p. 351, Willd. sp. v. 1. p. 169. *Pep. filiformis* A. Dietr. sp. v. 1. p. 177 (non Ruiz et Pav.), Griseb. flor. W. Ind. p. 167. Herbacea. Caulis filiformis, longissime repens, tetragonus, glaber, striat-maculatus. (Ex Sw. l. c.)
379. *P. ORBICULATA* (A. Dietr. sp. v. 1. p. 180), foliis subquaternis minimis sessilibus orbiculatis hirsutis. In Carolina (ubi nunc ignota). — Miq. syst. p. 169. *Piper orbiculatum* Poir. encycl. v. 5. p. 476. Caulis hispidus radicans. (Ex Dietr. l. c.)
380. *P. TRICARINATA* (A. Dietr. sp. v. 1. p. 181), foliis subquinis elliptico-ovatis concavo-canaliculatis, acumine obtusato, subtus præcipue inferne nervoso-tricarinatis. — Miq. syst. p. 152. *Piper tricarinatum* Haw. suppl. pl. succ. p. 102. Caulis subpedalis ramosus. Rami teretes striati. Petioli sublineares. (Ex Dietr. l. c. et Haw.)
381. *P. JUNGHUIANIANA* (Miq. pl. Jungh. fasc. 1. p. 14), foliis ternis breviter petiolatis ellipticis vel sublanceolato-ellipticis basi obtusiusculis carnosius glanduloso-punctatis utrinque rariter puberis glabrescentibus uninerviis vel nervulo

utrinque adjuncto subtrinerviis, amentis breviter pedunculatis apice ramulorum ternis vel (foliis depauperatis) paniculato-confertis densifloris, ovario semi-immerso, bacca semi-immersa ovoidea punctata. In Javæ regione Djampang. — Miq. flor. Ind. bat. v. 1. pars 2. p. 435. Stirps  $\frac{1}{2}$ - $\frac{1}{4}$  ped. alta, robusta. Petioli lineam longi. Folia  $\frac{3}{4}$ -1 $\frac{1}{4}$  poll. longa, 2 $\frac{1}{2}$ -4 lin. lata. Amenta  $\frac{1}{4}$  vulgo 2 $\frac{1}{2}$  poll. longa, 1 lin. crassa.

382. *P. RAMULOSA* (Anders. enum. Gallap. p. 57), foliis oppositis vel quaternis quinisque breviter petiolatis ovato-oblongis apice rotundatis obtusis basi in petiolum subangustatis utrinque brevissime puberulis, inflorescentia ignota. — Ad truncos arborum in cacumine summi montis insulæ Charles archipelagi Gallapagos. Caulis vix digitalis, inferne crassitie pennæ passerinæ, mox divisus, obscure viridis, succosus, teretiusculus, sursum fuscior, tetragonus et puberulus. Folia 2 lin. longa lineam lata. Petioli semilineam longi.

*Peperomiæ non satis notæ et incertæ sedis.*

383. *P. CRYSTALLINA* (Ruiz et Pav. flor. Peruv. v. 1. p. 30), foliis oblongis inde oblongo-ovatis subtus punctato-excavatis sessilibus integerrimis, amentis oppositifoliis filiformibus densifloris laxifloris. In collibus Limæ locis petrosis. — A. Dietr. sp. v. 1. p. 147, Miq. syst. p. 90. *Piper crystallinum* Vahl enum. v. 1. p. 341. Caulis dichotomus, tetragonus, hyalinus, striis rubris maculatus, nodis ciliatus. Tota planta gaudet odore fragrantissimo et suavissimo aniso valde simili. (Ex Ru. et P. l. c.)
384. *P. HEMIONITIDIFOLIA* (Ham. prodr. pl. Ind. occ. p. 2), foliis ovato-oblongis subspathulatis crassiusculis obscure venosis scapo subramoso, amentis alternis. In Sto Domingo. *Pep. hemionitidifolia* Miq. syst. p. 75. Acaulis, glabra. (Ex Ham. l. c.)
385. *P. LUNARIA* (Ham. prodr. pl. Ind. occ. p. 2), foliis longissime petiolatis lato-reniformibus submarginatis, amentis axillaribus subcapillaribus laxifloris. In Sto Domingo. *Pep. lunaris* A. Dietr. sp. v. 1. p. 167, Miq. syst. p. 49. Verisimiliter *Verhuelia cordifolia*. Glabra, repens, filiformis. (Ex Ham. l. c.)
386. *P. SPATHULATA* (Ham. prodr. pl. Ind. occ. p. 2), foliis longe petiolatis in petiolum attenuatis obovatis submarginatis, amentis terminalibus filiformibus. In Jamaica. — A. Dietr. sp. v. 1. p. 153, Miq. syst. p. 102. Verisimiliter *Pep. cordifolia*. Glabra, filiformis, repens. (Ex Ham. l. c.)
387. *P. HÆNKEANA* (Opiz in Reliq. Hænk. fasc. 3. p. 162), foliis terminalibus numerosis longe petiolatis cordatis inæqualiter repandis supra margineque glabris subtus pubescentibus basi floriferis multinerviis, petiolo pilosiusculo folium æquante, amentis aggregatis folio brevioribus capillaribus laxis, floribus distinctis, baccis glabris. In montanis Peruviae. — Miq. syst. p. 76. *Phyllobryon Hænkeanum* Presl Epim. bot. p. 222. Radix horizontalis, dentato-tuberosa, fibrosissima. Caulis erectus, pubescens. (Ex Opiz l. c.)
388. *P. SIMPLEX* (Ham. prodr. flor. Ind. occ. p. 2), foliis ovato-oblongis subacutis trinerviis, amento terminali solitario folium triplo superante. In Jamaica. — A. Dietr. sp. v. 1. p. 152, Miq. syst. p. 101. Glabra, caule erecto simplici foliis caducis inferne cicatrizzato. (Ex Ham. l. c.)
389. *P. SEMIRUENSIS*, foliis alternis inferioribus lato-suborbicularibus superiori-  
bus ovalibus vel ovatis obtusis lævibus margine apicem versus ciliolatis quin-  
quenerviis, amentis axillaribus terminalibusque solitariis petiolo duplo longio-  
ribus. In graminosis montis Semiru prope Widadarin alt. 6000 pæd. *Pep.*  
*sp. nova* Zoll. in Flora 1847. p. 470 (Zoll. n. 2324). Caulis infra radicans.

*Species exclusæ.*

*P. BLATTARUM* A. Dietr. = *Piper blattarum*.

*P. GRANDIFOLIA* A. Dietr. = *Piper subpeltatum* Willd.

- P. KNOBLECHERIANA Schott = Verhuelia Knoblechteriana C. DC.  
 P. LANATA A. Dietr. = Piper lanatum Roxb.  
 P. PEPULOIDES A. Dietr. = Chavica pepuloides Miq.  
 P. PELTATA A. Dietr. = Piper peltatum L.  
 P. PRUINOSA Kunth = Piper peltatum L.  
 P. RENIFORMIS A. Dietr. = Ve huellia elegans Miq.  
 P. RUBELLA Hook. et Arn. = Piper purpurascens Desf.  
 P. SARMENTOSA A. Dietr. = Piper sarmentosum Roxb.  
 P. SCUTATA A. Dietr. = Piper peltatum L.  
 P. SPECIOSA Kunth = Piper peltatum L.  
 P. SIDÆFOLIA A. Dietr. = Piper subpeltatum Willd. var. β.  
 P. UMBELLATA A. Dietr. = Piper umbellatum L.  
 P. VARIEGATA R. et P. = Piper maculosum Hill.?

### Index generis Peperomix.

(Numeri ad numerum speciei respondent.)

Peperomia Abyssinica Miq. 63, *acuminata* Dietr. 118, *acuminata* Griseb. 121, *acuminata* Miq. 105, *acuminata* Ruiz et Pav. 68, *acuminatissima* Miq. 131, *arutifolia* C. DC. 169, *ardescens* C. DC. 70, *Agapateris* C. DC. 321, *alata* Ruiz et Pav. 129, *alismæfolia* Presl 29, *alpina* Dietr. 185, *alpina* Griseb. 293, *alsinoides* Dietr. 284, *amplexicaulis* Dietr. 167, *amplexifolia* Dietr. 168, *Andinacea* C. DC. 55, *angularis* C. DC. 114, *angulata* Kunth 273, *angustata* Kunth 373, *angustifolia* Miq. 232, *Antillarum* C. DC. 278, *aphylla* C. DC. 368, *Arabica* Miq. 252, *arifolia* Miq. 34, *arifolia* var. Miq. 33, *asarifolia* Dietr. 26, *australis* Phil. 327.

Pep. Bangroana C. DC. 59, *basellæfolia* Kunth 72, *Baueriana* Miq. 110, *Belangeri* C. DC. 97, *Berlandieri* Miq. 307, ζ. *Bernieriana* Miq. 275, *Berteroana* Miq. 109, *bilineata* Miq. 132, *biuncialis* Kunth 228, *blanda* Kunth 336, *Boivinii* C. DC. 365, *Boliviensis* C. DC. 312, *Borbonensis* Miq. 152, *Botterii* C. DC. 337, *brachyphylla* Dietr. 127, *brevipes* C. DC. 20.

Pep. *Caffra* E. Mey. 252, *Caledonica* C. DC. 58, *camptotrigha* Miq. 350, *candida* Miq. 135, *Candolliana* Miq. 231, *cardiophylla* C. DC. 209, *Carlosiana* C. DC. 81, *Casapiana* C. DC. 334, *Casaretti* C. DC. 206, *Catharinæ* Miq. 264, *caulibasis* Miq. 87, *caulibarbis* Miq. var. 127, *Ceylanica* Miq. 328, *Chachapoyasensis* C. DC. 295, *Chilensis* C. DC. 259, *Choromiana* C. DC. 216, *Chrysotrigha* Miq. 26, *ciliaris* C. DC. 219, *ciliata* Kunth 251, *ciliata* h. Franquev. 91, *ciliolata* Miq. 20, *ciliosa* C. DC. 220, *circinata* Link 266, *claytonioides* Kunth 36, *colorata* Kunth 88, *Columbiana* Miq. 155, *concaua* Ruiz et Pav. 293, *concinna* Dietr. 235, *congesta* Kunth 372, *conjugata* Kunth 41, *convexa* Miq. 137, *Cookiana* C. DC. 298, *Corcoadensis* Gardn. 149, *cordifolia* Dietr. 143, *cordifolia* Opiz 95, *cordulata* C. DC. 93, *Cotyledon* Benth. 39, *Coulteri* C. DC. 161, *Coutallensis* Miq. 361, *Coutallensis* var. Thw. 242, *crystallina* Ruiz et Pav. 383, *Cubana* C. DC. 106, *Cubensis* C. DC. 203, *cuneata* Miq. 201, *cuneifolia* Miq. 183, *cyclophylla* C. DC. 265.

Pep. *dasystachya* Miq. 108, *Decaisnei* Miq. 244, *decurrens* C. DC. 165, *defoliata* C. DC. 107, *demissa* C. DC. 156, *dendrophila* Griseb. 121, 129, *dendrophila* Schlecht. 85, *dependens* Ruiz et Pav. 95, *Deppeana* Schlecht. 329, *dichotoma* Reg. 310, *diffusa* C. DC. 14, *Dindygulensis* Miq. 254, *discolor* C. DC. 359,

*dissimilis* Kunth 336 var., *d'stachya* Griseb. 188, 203, *distachya* Dietr. 180, *diversifolia* Kunth 226, *dolabriformis* Kunth 159, *Dominicana* C. DC. 332.

*Pep. edulis* Miq. 301, *elegans* C. DC. 189, *elliptica* Dietr. 245, *ellipticifolia* C. DC. 360, *elliptico-rhombea* C. DC. 78, *elongata* Kunth 187, *elongata* Miq. 76, *emarginata* Ruiz et Pav. 178, *emarginella* C. DC. 221, *emarginulata* C. DC. 200, *Endlicheri* Miq. 104, *Erasmia* C. DC. 22, *Ernstiana* C. DC. 150, *Estrellensis* C. DC. 144, *exigua* Griseb. 28, *exigua* Miq. 52, *exilis* Griseb. 21.

*Pep. Fendleriana* C. DC. 346, *Fernandeziana* Miq. 270, *Fernandopoana* C. DC. 19, *filiformis* Dietr. 378, *filiformis* Ruiz et Pav. 304, *fimbriata* Miq. 197, *flagelliformis* Miq. 322, *flavescens* C. DC. 218, *flexicaulis* Wawra 213, *foliosa* Kunth 130, *foliiflora* Ruiz et Pav. 16, *fontinalis* Dietr. 314, *fragrans* C. DC. 77, *Fraseri* C. DC. 24, *furcata* Opiz 231.

*Pep. Galeottiana* Hook. 11, *Galeottii* Miq. 176, *galioides* Kunth 362, *Gallapagensis* Miq. 369, *Gardneriana* Miq. 45, *Gaudichaudii* Miq. 342, *gemella* Miq. 44, *glabella* Dietr. 126, *glabella* Griseb. 86, *glabra* C. DC. 162, *Guadaloupensis* C. DC. 121, *Guayaquilensis* C. DC. 37, *Guildingiana* Dietr. 205.

*Pep. Hænkeana* Opiz 387, *Hamiltoniana* Miq. 146, *Hartwegiana* Miq. 325, *Haworthiana* Dietr. 282, *Haycockii* C. DC. 57, *hederacea* Miq. 197, *hemionitidifolia* Ham. 384, *hernandiæfolia* Griseb. 198, *hernandiæfolia* Miq. 217, *heterophylla* Griseb. in Lechl. exs. 322, *heterophylla* Miq. 157, *heterostachya* Miq. 308, *Heyneana* Miq. 309, *Heyneana* Thw. 289, *Hilariana* Miq. 54, *hircina* C. DC. 116, *hirsuta* Miq. 347, *hirta* C. DC. 101, *hirtella* Griseb. 101, *hirtella* Miq. 119, *hispidula* Dietr. 18, *Hoffmannii* C. DC. 15, *humilis* Dietr. 283, *hydrocotyloides* Miq. 31, *hymenophylla* Miq. 52, *hypoleuca* Miq. 339.

*Pep. inæqualifolia* Ruiz et Pav. 363, *incana* Dietr. 196, *increscens* Miq. 262, *insularum* Miq. 263, *ionophylla* Griseb. 214.

*Pep. Jamesoniana* C. DC. 103, *Jamesoniæ* C. DC. olim 156, *Japurensis* C. DC. 76, *Javanica* Miq. 299, *Junghuhniana* Miq. 381.

*Pep. Klotzschiana* Miq. 352, *Kravangensis* Miq. 237, *Kuhliana* Miq. 133, *Kunthiana* Miq. 366.

*Pep. lævifolia* Miq. 136, *lævis* C. DC. 79, *lanceolata* C. DC. 338, *lanceolato-peltata* C. DC. 91, *lanceifolia* Hook. 172, *Lang-dorffii* Miq. 262, *Larecajana* C. DC. 69, *latifolia* Miq. 287, *laxiflora* Kunth 53, *leptophylla* Miq. 324, *leptostachya* Hook. et Arn. 284, *lignescens* C. DC. 75, *Lindeniana* Miq. 311, *linearis* C. DC. 358, *longepedunculata* Opiz 99, *longifolia* C. DC. 67, *Loxensis* Kunth 330, *lunaria* Ham. 385, *Lyalli* C. DC. 73.

*Pep. Microæna* C. DC. 331, *macropoda* Miq. 182, *macrorhiza* Kunth 1, *macrostachya* Dietr. 234, *macrothyrsa* Miq. 158, *maculosa* Dietr. 192, *Madagascariensis* C. DC. 286, *magnoliæfolia* Dietr. 175, *major* C. DC. 197, *Mandiocana* Miq. 268, *Mandonii* C. DC. 12, *Mannii* Hook. f. 145, *margaritifera* Hook. 240, *marginata* Miq. 246, *Mariannensis* C. DC. 257, *Maribiana* C. DC. 181, *marm. rata* Hook. f. 258, *Martiana* Miq. 151, *Mascarena* C. DC. 60, *Mathewiana* Miq. 353, *Mauritiana* Dietr. 245, *Maypurensis* Kunth 255, *melanostigma* Miq. 83, *membranacea* Hook. et Arn. 340, *membranacea* Miq. 257, *Mekeana* Miq. 362 var., *Metapalcoensis* C. DC. 124, *Mexicana* Miq. 7 et ? 1, *microphylla* Kunth 367, *minima* C. DC. 28, *Miqueiæna* C. DC. 98, *Miqueliana* Griseb. 124, *Miradoreaiana* C. DC. 8, *Moerenhoutii* C. DC. 335, *mollis* Kunth 227, *Monini* C. DC. 267, *monostachya* Ruiz et Pav. 224, *monsteraefolia* Griseb. 190, *montana* C. DC. 173, *montica* A. Hook. f. 148, *Moritzii* C. DC. 140, *Moulmeiniana* C. DC. 117, *muscosa* Link 272, *muscophila* C. DC. 10, *mysuroides* Miq. 74, *Myosurus* Dietr. 205, *myriocarpa* Miq. 170, 181, *myrtifolia* Dietr. 233 et ? 144, *Myrtillus* Miq. 294.

*Pep. nana* C. DC. 61, *nematostachya* Link 186, *nemorosa* C. DC. 118, *nervosovenosa* Miq. 238, *nigropunctata* Miq. 86, *navalis* Miq. 160, *nummularifolia* Kunth 141, *nummularioides* Griseb. 303.

*Pep. Oahuensis* C. DC. 147, *obcordata* Presl 326, *obliqua* Ruiz et Pav. 222, *obtusifolia* C. DC. 183, *obtusifolia* Dietr. 175, *obtusifolia* Miq. 183 var., *obtusifolia* Miq. pro parte 200, *obversa* Griseb. in Dietr. 318, *olivacea* C. DC. 364, *Opi-*

*ziana* Dietr. 95, orbicularis C. DC. 370, orbiculata Dietr. 379, *Ottomania* Miq. 112, ovalifolia Hook. 302, ovato-peltata C. DC. 5.

*Pep. pachyphylla* Miq. 343, *pallens* Kunth 355, *pallescens* Miq. 171, *palida* Dietr. 128 et ? 147, *paniculata* Reg. 25, *parasitica* C. DC. 142, *Parkeriana* Miq. 163, *parnassiaefolia* Miq. 49, *parvifolia* C. DC. 2, *patula* C. DC. 84, *Pavoniana* C. DC. 344, *pedunculata* C. DC. 349, *peepuloides* Kunth 374, *pellucida* Kunth 48, *peltoides* Kunth 208, *pereskiaefolia* Kunth 313, *perforata* Opiz 229, *Pernambucensis* Miq. 66, *Perrottetiana* Miq. 260, *Perrottetiana* Sieb. 246, *petiolaris* C. DC. 113, *petiolata* Hook. f. 256, *phyllantha* Opiz 16, *Pichinchæ* C. DC. 241, *pilosa* Ruiz et Pav. 65, *piperea* C. DC. 164, *plicata* Opiz 376, *Poeppigii* Miq. 236, *polybotrya* Kunth 89, *polystachya* Dietr. 348, *Ponthieu* Miq. 198, *Portobellensis* Beurl. 241, *portulacæfolia* Kunth 327, *portulacoides* Dietr. 261, *procumbens* C. DC. 212, *producta* Griseb. 188, *pseudoamplexicaulis* C. DC. 195, *pseudodependens* C. DC. 86, *pseudofurcata* C. DC. 30, *pseudopeltoides* C. DC. 207, *pseudopereskiaefolia* C. DC. 291, *pseudorhombea* C. DC. 2.2, *pseudorhynchophoron* C. DC. 168, *pseudovariegata* C. DC. 179, *pterocaulis* Miq. 129, *pubescens* Ruiz et Pav. 281, *pulchella* Dietr. 357, *pulcaris* Opiz 205, *pumila* Opiz 230, *punctulata* Reg. 127, *purpurascens* C. DC. 125, *purpurascens* Moritz 139, *purpurea* Ruiz et Pav. 223, *purpurella* C. DC. 111, *pyrifolia* Humb. et B. 177.

*Pep. quadrangularis* Dietr. 276, *quadrangularis* Griseb. 272, *quadrifolia* Kunth 316, *quaternata* Miq. 300, *quintuplinervia* C. DC. 247, *Quitensis* Miq. 251, *Quitensis* Benth. 336.

*Pep. ramulosa* Anders. 382, *recurvata* Miq. 243, *reflexa* Dietr. 305, *reflexa* Kunth 314, *Regelii* C. DC. 317, *Reinwardtiana* Miq. 279, *repiformis* Miq. 205, *repens* Kunth 205, *reptans* C. DC. 204, *retusa* Dietr. 277, *rhexiaefolia* Moritz 341, *rhombea* Ruiz et Pav. 371, *rhomboidea* Hook. et Arn. 100, *Riedeliana* Reg. 153, *Rollingo* Link 375, *rotundata* Kunth 253, *rotundifolia* Griseb. 256, *rotundifolia* Kunth 141, *Roxburghiana* Dietr. 232, *rubella* Hook. 351, *rubioides* Kunth 306, *rubricaulis* Dietr. 313, *rubrinodis* Kunth et Bouché 271, *rufescens* C. DC. 123, *Rupertiana* C. DC. 105, *rupestris* Kunth 191, *rupestris* Miq. 186 et 180, *ruscifolia* Kunth 120.

*Pep. saligna* Kunth 225, *San-Carlosiana* C. DC. 102, *Sandersii* C. DC. 33, *Sandwicensis* Miq. 345, *scandens* Ruiz et Pav. 205, *Schiedeana* Schlecht. 323, *Schomburgkii* C. DC. 13, *scutellæfolia* Ruiz et Pav. 4, *secunda* Ruiz et Pav. 33, *Sæmanniana* Miq. 250, *Sellowiana* Miq. 329, *Semiruensis* C. DC. 389, *septemnervis* Ruiz et Pav. 249, *septuplinervia* C. DC. 190, *serpens* C. DC. 134, *serpens* Griseb. 153, *serpentarioides* Miq. 46, *serpyllifolia* Dietr. 269, *simplex* Ham. 388, *spatulata* Ham. 386, *spectabilis* Miq. 174, *Spruceana* Benth. 47, *Sprucei* C. DC. 6, *stelechophila* C. DC. 199, *stellata* Dietr. 307, *stenocarpa* Reg. 194, *stolonifera* Kunth 9, *striata* Ruiz et Pav. 64, *suaveolens* Ham. 377, *subpeltata* C. DC. 94, *subrotunda* Dietr. 175, *subrotundifolia* C. DC. 153, *succulenta* C. DC. 183, *Sumidoriana* C. DC. 211, *Surinamensis* C. DC. 80, *Swartziana* Griseb. 183, *Swartziana* Miq. 378, *sylvatica* C. DC. 210, *sylvestris* C. DC. 215.

*Pep. talinifolia* Kunth 71, *talinifolia* Link 168, *Tarapotana* C. DC. 92, *tenella* Dietr. 17, *tenera* Miq. 18, *tenerrima* Schlecht. 323, *tenuiflora* Opiz 50, *teragona* Ruiz et Pav. 290, *tetraphylla* Hook. et Arn. 305, *thrichocarpa* Miq. 320, *Thyrsitesii* C. DC. 289, *tithymaloides* Dietr. 239, *tomentosa* Dietr. 319, *Tovarianæ* C. DC. 56, *transparens* Miq. 32, *Trianae* C. DC. 27, *tricarinata* Dietr. 380, *trifolia* Dietr. 356, *trinervia* Miq. 316, *trinervis* Ruiz et Pav. 127, *trinervula* C. DC. 139, *trineura* Miq. 315, *tristachya* Kunth 90, *tristachya* Opiz 303, *tuberosa* Opiz 43.

*Pep. umbellata* Miq. 40, *umbilicata* Ruiz et Pav. 3, *undeninervia* C. DC. 218, *urocarpa* Fish. et Mey. 202, *Urvillianæ* Rich. 104.

*Pep. VahlII* Dietr. 285, *valantoides* Miq. 305 var., *variegata* Ruiz et Pav. 192, *varifolia* C. DC. 354, *Velloziana* Miq. 82 et 129 β, *Venezueliana* C. DC. 115, *Ventenatii* Miq. 274, *Verhuellia* C. DC. 23, *verticillata* Miq. 357, *Victoriana* C. DC. 292, *villosa* C. DC. 42, *Vincetiana* Miq. 62, *Vitiana* C. DC. 333, *Vogelii* Miq. 31.



Pep. Wallichii Miq. 122, Weddellii C. DC. 35, Wightiana Miq. 138, Wrightiana C. DC. 193.

*Genera dubia, vix nota.*

X. DUGAGELIA? *Gaudich. in Voy. de Freycinet p. 513 et 514. — Piperis sectio Endl. gen. n. 1820, 6.*

Flores hermaphroditi. Bractea foliacea, basi decurrens. Stigma capitellato-globosum, pilosum. — Planta in h. Juss. ex Gaudich., mihi ign.

XI. SYMBRYON *Griseb. Cat. pl. Cub. p. 64.*

Flores dielines, masc. ignoti. Ovarium unicum, squamis 2 basifixis lateralibus subulatis basi dilatata vaginantibus stipatum. Ovulum basilare. Stigma sessile, nudum, minutissimum. — Frutex podis æqualibus. (Ex Griseb. l. c.)

S. TETRASTACHYUM (Griseb. l. c.). foliis alternis petiolatis ovatis apice abrupte acuminatis utrinque glabris opacis chartaceis impunctatis 5-nerviis, centrali nervo ad apicem usque ducto, sequentibus ante apicem per anastomoses evanidis 2 exterioribus fere marginalibus, omnibus subtus prominulis, nervulis subtransverse parallelis, petiolo glabro tereti, rhachi hirtella, bractea lanceolata apice subulata dorso hirtella, squamis hypogynis 2 ovatis dorso hirtellis, ovario ovato stipitato apice imo stigma minutum gerente. ♂ In ins. Cuba (Wright n. 2268 in h. DC.). Limbi circ. 0,085 longi, 0,042 lati. Petioli 0,018 longi. (v. s.)

---

## Ordo CLXXXVIII (1). CHLORANTHACEÆ (2).

*Loranthearum pars Juss. Ann. Mus. hist. nat. XII. p. 299; Spreng. Syst. veg. I. p. 683. — Caprifoliacearum pars Spreng. Anl. z. Kennln. d. Gew. II. p. 620. — Chloranthæ R. Br. bot. Mag. n. 2190 adnot.; Nees in R. Br. Verm. Schr. I. p. 404; Lindl. Collect. bot. t. 17, Veg. Kingd. p. 519; Kunth syn. pl. IV. p. 200; E. Mey. de Houttuynia p. 51. — Chloranthaceæ Bl. Enum. pl. Jav. t. I. p. 78, Fl. Javæ fasc. 8; Agardh f. Theor. syst. pl. t. XX. p. 240; Lindl. Introd. ed. I. p. 183; Bartling Ord. nat. pl. p. 78 et 85; Endl. gen. pl. p. 204; Endl. Enchirid. bot. p. 148: Clarke on the struct. and affin. of Myric. bull. soc. bot. Fr. 1857. p. 1060; Schnizl. fam. f. 6. t. 80; Mart. Fl. Brasil. f. XI; Miq. Fl. Jav. bat. I. p. 800; Cordemoy in Adans. III. p. 280; Le Maout et Decsne traité gén. de bot. p. 503.*

Flores hermaphroditi, monoici vel dioici, bracteati vel omnino nudi, apetal. — *Hermaphroditi*: Calyx nullus. Stamina 1-3, ad ovarii latus affixa, filamentis connatis, connectivo sæpius ultra loculos producto; antheris et bilocularibus 4-thecis et unilocularibus dithecis, connectivo adnatis, introrsum rima longitudinali dehiscens. Pollen globosum. Ovarium uniloculare, stigmate sessili plano præditum. Ovulum in ovarii cavo unicum, orthotropum ex apice pendulum. — *Masculi*: bracteati vel ebracteati (*Hedyosmum*), monandri, antheram sessilem bilocularem tetrathecam si tentes. — *Feminei*: bracteati. Calyx nullus vel minimus, subtubulosus, gamosepalus, margine tridentatus. Ovarium nudum globosum stigmate sessili plano vel inferum subtrigonum stigmate ligulato subtriobove instructum. Ovulum unicum, orthotropum, ex apice cavi pendulum. Drupa carnosula, unilocularis, putamine perduro. Semen unicum, apice adfixum, testa tenui in fructu. Albumen copiosum, carnosio-oleosum. Embryo in albuminis apice inclusus, minimus, radícula infera, cotyledonibus divaricatis vel vix conspicuis. — *Herbæ* frutices vel arbores in America tropica, in Polynesia, in Asia tropica et subtropica, rarius boreali indigenæ, ex Africa, Europa et Nova-Hollandia ut videtur omnino exules; aromaticum odorem et exsiccatae spargentes; foliis decussatis penninerviis plerumque denticulatis stipulatis, stipulis apice tantum liberis subulatis simplicibus vel pluries divis. basin versus inter se et cum petiolis in vaginas plus minus elongatas sæpius ochreiformes connatis, floribus hermaphroditis et masculis semper, femineis rarius (*Ascarina*) spicatis uni-

(1) Numerus CLXXXVII vacat, propter Saurureas Piperaceis adscriptas.

(2) Auctore Hermanno comite a Solms Laubach.

tribracteatis; femineis plerumque cymosis cymulis floriferis. bractæatis axillaribus vel varie paniculatis.

Ordo naturalis Piperaceis arcta affinitate conjunctus ovarii structura ab iis discrepans, quoad habitus ad Rubiaceas quasdam quodammodo accedens, ab omnibus fere auctoribus hic collocatus, a Cordemoy (Adans. III), Plataneis et Ceratophylleis auctus, genera amplectens tria quam maxime naturalia nempe :

I. Flores hermaphroditi : . . . . . *Chloranthus*

II. Flores diclines.

A. Masculus bracteatus, femineus  
calyce nullo, stigmate plano : . . . . . *Ascarina*

B. Masculus ebracteatus, femineus  
inferus calyce tridentato; stig-  
mate capitato-subtrilobo vel  
lingulato : . . . . . *Hedyosmum*.

I. CHLORANTHUS Sw. in *Philos. transact.* 77. p. 354, *Prodr.* 84; *L'Herit. sert. angl.* 1. p. 35; *Juss. gen. pl.* p. 423; *Schreb. gen. pl.* p. 193. n. 1733; *Willd. Spec. pl.* 1. 503, 248; *Pers. Syn. pl.* 1. p. 148; *Roem. et Schult. Syst. veg.* 3. p. 29, 567; *Spreng. Anleit.* 3. p. 620; *Spreng. Syst. veg.* 3. p. 683. n. 3044; *R. Br. Bot. Mag.* 2190; *Lindley Collect. bot.* t. 171; *Blume Enum. pl. Jaræ fasc.* 1. p. 78, *Flora Jaræ fasc.* VIII; *Endl. gen. pl.* p. 265. n. 1819; *Stieb. et Zucc. in Acad. Vindob. Mem.* 1846 p. 232; *Cordemoy in Adans.* 3. p. 295; *Benth. Fl. Hongk.* p. 334. — *Nigrina Thunb. Nov. gen.* p. 58, *Fl. Jap.* p. 65, *Acta Ups.* VII p. 142; *Lam. Ill. gen.* 1. p. 294; *Poir. in Lam. Encycl. méth. bot.* 4. p. 489. — *Creodus Lour. fl. cochin.* 1. p. 112. — *Peperidia Reichb. Consp. gen.* p. 212. — *Cryphæa Hamilt. in Breust. Edinh. journ. of sc. jan.* 1825. p. 11. — *Ascarina Bl. en. pl. Jar.* 1. p. 79 (non Forst.). — *Sarcandra Gardn. in McClelland's Calc. Journ. of nat. hist.* VII. p. 38. — *Tricercandra A. Gray Acc. of bot. spec. J pan.* p. 318, *On the bot. of Japan* p. 405. — *Saintlegeria Cordemoy in Adans.* 3. p. 300. — *Strophia Noronh. sched. mspt. ex Cordem. Adans.* 3. p. 295.

Flores hermaphroditi, sessiles, bracteati, spicati, monandri vel triandri. Ovarium conicum, stigmate sessili truncato coronatum. Antheræ filamento crasso prælitæ ad ovarii latus affixæ vel pro flore singulæ, vel ternæ filamentis tum in squama carnosa tripartita coalitis; mediana bilocularis 4 theca loculis discretis interdum sterilibus, laterales si adsunt uniloculares 2-thecæ, omnes connectivo valido dorsali instructæ. Drupa carnosa subglobosa, vel subcarnosa poriformis, putamine fragili. Embryo in albumine copioso minimus, cotyledonibus divaricatis. — Frutices et herbæ perennes, aromaticæ, in Asia orientali præcipue tropica et subtropica, rarius boreali indigenæ, ramulis glabris oppositis ad nodos articulatis, foliis decussatis, stipulacris; stipulis lanceolatis tantum ad basim inter se et cum petiolis in ochreas vix distinctas connatis, nodos quasi margine membranaceo circumdantibus; inflorescentiis terminalibus ramosis spicatis.

- § 1. *Fruticosi*. — Frutices vel fruticuli erecti vel subscandentes. Drupa globosa vel subglobosa, carnosa, pallida. — *Chloranthus autor. plurim.* — *Sarcandra* Gardn.

\* *Triandri*. — Antheræ in squamam ad medium tripartitam connatæ, mediana 4-theca loculis discretis, laterales breviores uniloculares 2-thecæ, omnes connectivo crasso rotundato obtuso præditæ.

1. *C. OFFICINALIS* (Bl. en. p. 79, Fl. Jav. v. 3. p. 10. t. 1), foliis maxime variis magnis ovatis elongato-ovatis vel sublanceolatis acuminatis sæpius in apiculum productis, margine serratis vel serrulatis, serraturis acutis plus minus callosis. § Per Asiam tropicam et subtropicam orientalem frequentissimus; in India orientali (Bengalia, Assam, Khasyia mont.), in Tenasserim, ad Pulo Penang, in ins. Andamanis, in ins. Java, Sumatra, in Philippinis ins. (Cuming pl. Phil. n. 521. *C. salicifolius* Presl. — Miq. fl. ind. bat. p. 804, Cord. in Adans. 3. p. 298. *C. elatior* R. Br. bot. Mag. 2190? Link enum. horti Berol. 1. 140. n. 1295? *C. erectus* Sweet hort. sub. Lond. 28? Wall. n. 6881. *C. indicus* Wight ic. t. 1945. *C. sumatranus* Miq. fl. Ind. bat. suppl. p. 343. *C. salicifolius* Presl epim. bot. 231. *C. officinalis* Schnizlein fam. t. 80. fig. 1-6, Cordem. in Adans. 2. 298. *Cryphea erecta* Ham. in Brewst. Edinb. journ. jan. 1825. p. 11. t. 2. — Planta quoad foliorum formam maxime variabilis. Rami strictiusculi vel subscandentes, glabri, ramulis oppositis. Folia mediæ magnitudinis circa 13 cm. longa 5-6 cm. lata, majora 18 cm. longa 6-7 cm. lata, minora quæ sæpissime apud plantam in hortis sub nomine *C. elatioris* culta occurrunt 8-9 cm. longa 4-4,5 lata. Formam singularem foliis parvulis angustioribus 7-8 cm. longis 2-2,5 cm. latis, ex Pulo Penang Wallich retulit, quam tamen haud speciem constituere posse propriam specimen docet Griffithianum in Khasyia lectum foliis salicinis fere 8 cm. longis et 1, 5 latis simul cum normalibus gaudens. Altera foliis fere obovatis vel rotundato-ovatis brevi-acuminatis circa 11 cm. longis et 5 cm. latis insignis in Bengalia et in montibus Sillet inventitur (Wallich 6881, Griffith). Ab insequente specie quam proxima primo visu serraturæ indole differt. (v. v. et s.)
2. *C. INCONSPICUUS* (Sw. in Phil. Transact. 78. p. 359. t. 15), foliis illis præcedentis speciei multo minoribus ovatis vel rotundato-ovatis sæpius subcuneatis vix acuminatis obtusisve remote et obtuse dentato-crenatis, dentibus convexo-rotundatis apice callosis. § In Asia orientali tropica et subtropica, præcedente ut videtur multo rarior: in China meridionali (Millet), ad Macao (Callery), in insul. Java (Zollinger 1845 n. 1949), in Japonia, ubi ex cl. Siebold haud sponte nascens sed e China allatus in hortis tantum colitur. — Gmelin Syst. veg. 1. p. 280; Willd. Sp. pl. 1. p. 688; L'Hérit. Sert. angl. 1. p. 35. t. 2; Sieb. in Nov. Act. Nat. cur. 14. p. 2. pg. 681; Schnizlein ic. fam. t. 80. fig. 8-15 anal. flor.; Miq. Mus. lugd. bat. 3. p. 129; Sieb. et Zucc. mem. Acad. monac. 1846. p. 232; Cordem. in Adans. III. p. 295; Benth. Fl. Hongk. p. 334; Le Maout et Decsne Traité gén. de bot. p. 504 c. icon. *Ch. obtusifolius* Miq. fl. Ind. bat. p. 802? *Creodus odorifer* Lour. fl. cochinch. 1. p. 112? *Nigrine* en épi Lam. Dict. encycl. 4. p. 489. *Nigrina spicifera* Lam. Illustr. n. 1506. t. 71. *Nigrina spicata* Thunberg ab omnibus fere auct. hic refertur notis tamen ut recte observavit cl. Siebold sat gravibus, utut caule herbaceo palmari et ultra, discrepans. A *Chl. officinali* Bl. quam proximo foliis constanter multo minoribus, late ovatis, circa 5 cm. longis et 2, 5-3 cm. latis, maximis 7-8 cm. longis 5 cm. latis facillime distinguenda. Accedit serratura foliorum supra descripta, hic obtuse dorso convexa, in præcedente contra acuta dorso subrecta. Fruticulus humilis ramosus, foliis pallide virentibus, in caldariis horticorum botanicorum Europæ totius quam maxime communis. (v. v. et s.)

\*\* *Monandri*. Anthera bilocularis 4-theca, loculis discretis apicem versus convergentibus, connectivo crasso obtuso rotundato prædita.

3. *C. PERNYANUS*, foliis ex oblongo-lanceolatis acuminatis tenuibus sat longe petiolatis glaberrimis margine grosse serratis, spicis floriferis simplicibus

axillaribus paucifloris (basin versus omnino sterilibus), floribus minimis in bractearum membranacearum axillis fere inclusis, antheris in ovario alte insertis bilocularibus incurvis. § In Chinæ meridionalis prov. Kony-Tschou (Perny 1858). Ch. monostachys R. Br. Bot. Mag. 2190? Suffrutex circiter orgyalis, multicaulis. Caules tenues, quadrangulares, inferne ad nodos bisquamati, apice tantum foliati. Folia 7-9 cm. longa, circiter 3 cm. lata. Spicæ floriferæ tenues, ramis omnino desituta sat numerosæ. Flores minimi vix 1 mm. longi. Fructus ignoti. — Species distinctissima ex habitu, florequè minuto monandro statim recognoscenda. Eadem ut videtur quam Chl. monostachys R. Br. vel eo arcte affinis cujus cum icone et descriptione optime quadrat. Cum vero in specimine unico valde manco Pernyano non contigerit ut flores invenirent triandros, qui tamen floribus monandris in planta Browniana immixti dicuntur, alio nomine designandam esse speciem putabam (v. s. in h. Mus. Paris.)

4. C. BRACHYSTACHYS (Bl. Fl. Jav. fasc. 8. p. 13 et 14. t. 2), foliis subcoriaceis glabris ex oblongo lanceolatis longe acuminatis grosse et argute serratis basin versus integris in petiolum brevem attenuatis, inflorescentia pro more brevi parte ramosa, bracteis floralibus coriaceis e basi naviculari lanceolatis acutis, floribus magnis, antheris crassis cylindraceis rectis, drupis subglobosis carnosulis. § In Asia tropica et subtropica orientali; in India orientali (Bengalia, Khasya mont.), in ins. Ceylon (Leschenault, Thwaites 2207), ad Pulo Penang, peninsulæ Malaccanæ (Wallich n. 6880), in ins. Java (Blume, Zollinger n. 1475), in China meridionali ad Hongkong (Seemann), in Japonia ad Nagasaki (Oldham), in ins. Philippinis (Cuming n. 1392, h. Boiss.), in ins. Formosa (Oldham n. 462). — Miq. fl. Ind. bat. p. 801, Ann. Mus. lugd. bat. 3. p. 129; Benth. Fl. Hongkong p. 334. C. monander R. Br. bot. Mag. 2190 adnot. C. ceylanicus Miq. Fl. Ind. bat. I. p. 802. C. denticulatus Cord. in Adans. 2. p. 296. Ascarina serrata Bl. Enum. pl. Javæ I. p. 79. Sarcandra chloranthoides Gardn. in McClelland's Calc. Journ. of nat. hist. 8, p. 348; Wight & n. 1946; Le Maout et Decsne Traité gén. bot. p. 504 (anal.); Cord. in Adans. 2. p. 301. Rami strictiusculi, glabri, siccitate striati. Ramuli oppositi, internodio basali squamato haud foliato præditi. Folia 8-14 cm. longa, 3-5 cm. lata, serratura arguta maxime variabili, dentibus nunc remotis, nunc approximatis, in apiculum plus minus longum productis. Specimen Hongkongense valde mancum quidem foliorum forma fere ovata et serratura pro more mutica a cæteris discrepat. Inflorescentiæ ramulos terminantes parce ramosæ, fructuum maturitatis tempore circ. 3 cm. longæ, drupas gerentes, subglobosas vel e globoso ovoideas circ. 5 mm. longas et 4-4,5 mm. latas, secundum Blume pallidas, secundum Miquel rubras. A congeneribus jam omnium partium rigiditate coloreque quam exsiccatione induit rufo-fusco, primo aspectu differt. (v. s.)

§ 2. *Herbacei*. — Herbæ perennes, rhizomate repente præditæ. Bractæe floriferæ subtrilobæ. Drupæ pyriformes, subcarnosæ, ut videtur luteo-virides vel fusciculæ.

\* *Brachyuri*. Antheræ subæquilongæ in squamam ad medium tripartitam connatæ, connectivis vix ultra loculos productis crassis rotundatis. — *Chloranthus auct. omn.*

5. C. SERRATUS (Röm. et Schult. Syst. veg. 3. p. 461), caule simplici ultrapedali herbaceo ad nodos bisquamato, foliis 4 approximatis brevipetiolatis magnis elongato-ovatis in acumen plus minus longum desinentibus grosse serratis, spica florifera terminali simplici vel ex basi bipartita, floribus in bractearum subtrilobarum axillis sessilibus vel subpedunculatis, antherarum connectivis crassis muticis rotundatis vix ultra loculos productis, drupa magna pyriformi viridula distincte cicatrisata. ¶ In Japoniæ ins. (Small, Wright, Ringgold and Rodgers, Zollinger n. 165), prope Hakodaki (Wilford 1018), et Jokohama (Oldham n. 801 in h. Boiss.). — Miq. Mus. lugd. bot. 2. p. 201, 3. p. 129; Sieb. et Zucc. Abh. d. Münch. acad. t. 3. p. 232; Cord. in Adans. 3. p. 297.

C. Blumeanus Cord. in Adans. 4. p. 296. *Nigrina serrata* Thunb. Act. nov. ac. Upsal. 7. p. 142. t. 5. f. 1 (icon. mala). Herba perennis, rhizomati horizontali, caule simplici lævi ex paucis internodiis elongatis apice bisquamatis extructo, spica florifera simplici vel e basi di rarissime trichotoma terminato. Folia quatuor binata decussata, internodia bina supra obsidentia. Internodium foliiferum superius abbreviatum, unde rosule fere quadrifoliæ spicam floriferam cingentis habitus. Folia 11-15 cm. longa, 3,5-7 cm. lata, plus minus incise et gross. serrata, apiculo angustato 2-3 cm. longo instructa. Antherarum squama ab illa C. officinalis et inconspicui non nisi antheris lateralibus medianis subæquilongis diversa. Habitus contra fere omnino specierum insequentium præfert. Chl. japonici Sieb. in herbario Musei Paris. specimen asservatum hexaphyllum, tribus internodiis foliiferis præditum. (v. s. in h. DC. Hook. Boiss. Vindob. Monac.)

\*\* *Macrouri*. Antheræ 3 in squamam ad basin usque fere tripartitam coalitæ, connectivis ultra loculos longe productis loricatis præditæ. — *Tricercandra* A. Gray. — *Saintlegeria Cordemoy*.

6. C. FORTUNEI, caule herbaceo ad nodos bisquamato apicem versus foliato, foliis quaternis vel interdum senis decussatis inter se approximatis, spica florifera simplicissima brevi, antheris tribus fertilibus, mediana biloculari tetra-thea, lateralibus unilocularibus dithecis, connectivis planis loricatis subparallelis perangustis prælongis, loculos longitudine multoties superantibus, drupis (ex Cordemoy) crassis obovatis. ♀ In China boreali (R. Fortune). *Tricercandra Fortunei* A. Gray bot. mem. (on the bot. of Japan) p. 405. *Saintlegeria gracilis* Cord. in Adans. 3. p. 300. Caulis semipedalis glaber. Folia secundum cl. Cordem. elliptico-lanceolata, reticulato-venosa, glabra, serrulata. 3-4 cm. longa 1-1,5 cm. lata. Spica florifera terminalis circa 2-3 cm. longa. Flores albi (Fortune). Planta cum flore tempore nondum omnino sit evoluta, foliaque tum vix explicentur ampliori descriptioni haud apta. Specimen fructiferum, ex cl. Cordemoy, drupas obovatas crassas cicatricatas, imul cum foliis adultis præbet neque in herbario Musei Paris. neque alias ut invenirem mihi contigit. (v. s. in herb. Hook., Mus. Paris.)

7. C. JAPONICUS (Sieb. nov. act. nat. Cur. 14. p. 2. p. 681), caule herbaceo. ad nodos bisquamato, foliis 4 decussatis sub spica florifera terminali rosulatis modice petiolatis ovatis glabris acuminatis haud in apiculum productis margine incisi-o-serratis, antheris basi tantum connatis divaricatis, fertilibus binis lateralibus, unilocularibus dithecis, mediana omnino sterili, connectivis et medianæ et lateralium loricatis angustis obtusis, loculis triplo vel subquadraplo longioribus. drupa magna pyriformi vix distincte cicatricata in sicco striata, colore luteo-viridi tincta. ♀ In China boreali, ad Mandschuriam litus latit. 44-45 (Wilford). in ins. Coreana Tsu-sima (Wilford), in Japonia ad Hakodaki (Hodgson). C. mandshuricus Rupr. Dec. pl. Amur. t. 2 (bull. soc. bot. de Fr. 1860. p. 527); Rupr. et Maxim. Prim. fl. Amur p. 463; Ruor. in Maack. *Tricercandra quadrifolia* A. Gray account of bot. spec. Japan. in Perrys Exp. p. 318; bot. mem. (on the bot. of Japan p. 405); Miq. Mus. lugd. bat. 3. p. 130. Herba perennis, rhizomate horizontali, caule simplici lævi spithamæ et internodiis bisquamatis elongatis extructo, spica florifera omnino simplici terminato. Folia quaterna, binata, decussata, internodia terminalia obsidentia. Internodium foliiferum superius tali modo abbreviatum ut folia verticillata videantur esse. Folia 7-8 cm. longa, 3,5-5 cm. lata; spica florifera 4-4,5 cm. longa; drupæ maturæ 4 mm. longæ, apicem versus 2 mm. latæ. Folia ut prioris speciei flore tempore subconvoluta, nondum omnino explicata, e speciminibus fructiferis ergo descripta. (v. s. in h. Hook., Boiss., Mus. Paris.)

\*\*\* Floribus ignotis, si ex affinitate totius plantæ cum C. japonico concludere licet, macrouris.

8. C. OLDHAMI, foliis subsessilibus late ovatis fere subcuneatis acuminatis toto ambitu crenato-dentatis glaberrimis, inflorescentia terminali ramosa elongata, ramis oppositis rectangule patentibus elongatis, drupis pyriformibus in sicco

castaneis, collo tenui rugoso luteo-viridi instructis, floribus ignotis. ♀ In insula Formosa (Oldham n. 466 in h. Boiss.). Nusquam alias reperta. Species etsi præcedenti ut videtur proxima primo intuitu tamen foliorum forma et serratura inflorescentiæ indole ac druparum forma coloreque perbellè distincta. Habitus omnino præcedentis. Folia quaterna, fere rotundato-ovata, 11 cm. longa, 7 cm. lata, apiculo circa 1,5 cm. longo instructa. Inflorescentia 8-14 cm., internodium ejus basale 5-6 cm. longum, cetera breviora. Rami floriferi internodio basali circiter æquilongi. Drupæ maturæ 3 cm. longæ. (v. s.)

*Species haud visæ.*

9. *C. MONOSTACHYS* (R. Br. Bot. Mag. t. 2190), antheris incurvis 3-lobis indivisive spicis solitariis floribus alternis. — E China introd. in hort. Soc. hort. londin.; caulibus jun. simplicissimis, deinde ramosis (Lindl.). — Br. l. c.; Lindley Coll. bot. t. 17. Herba succosa pedalis glaberrima. Caules simplices demum ramosi plures et unus rarior rotundato-tetragoni sæpius deformes. Folia opposita stipitata petiolata nitidissima ovato-oblonga acute serrata, serraturis apiculatis, pinnam concava rugosa demum plana convexa, supra læte viridia, infra pallidiora reticulatim venosa. Petoli sulcati. Stipulæ connectivæ obsoletæ. Spica terminalis pauciflora, solitaria, pedunculata. Flore in dentibus decussatis rhacheos oppositis, bractea margine sphacelata ipsis paulo breviora extus suffulta. Perianthium 0. Stamina extrorsa unilateralia. Filamenta in una massa crassa carnea infra antheras marginata extus longitudinaliter 2-sulcata, varii lateri partim adnata, connata. Antheræ 3 albæ ovatæ innatæ, versus rhachem super stigmate incurvæ, basi connatæ, connectivo maximo carnosò rugoso, loculis introrsis simplicibus 2-valvibus linearibus longitudinaliter dehiscens, iis antherarum lateralium integrarum solitariis, intermediæ apice bifidæ geminis. Ovarium rhachi proximum, uniloculare; ovulo unico a pariete rhachi proxima appenso; stigma subsessile integrum concavum. Pericarpium clausum carnosum turbinatum deciduum extus cicatrice filamentum notatum. Semen subrotundum pendulum; testa ossea. Nucleus femini conformis, membrana propria vestitus et chalaza hilo proxima insignitus. Albumen maximum carnosum. Embryo minimus inversus ab umbilico remotus, radícula brevissima centrifuga, cotyledonibus divaricatis, albumini arctissime adhærens et difficillime extricandus. Planta ut ex descriptione videtur typice monandra, d-in in horto degenera, « ramis sæpius deformibus » et floribus polymeris instructa. Num eadem quam Chl. Pernyanus qui cum icone et descriptione sat bene quadrat vel proxima ejus speciei ignotum. Quoad formam drupæ transitum moderari videtur inter Chl. fruticosos et herbaceos.

10. *C. GRANDIFOLIUS* (Miq. Fl. Ind. bat. p. 802). Ramuli verruculosi. Folia longiuscule ( $\frac{3}{4}$ – $\frac{1}{2}$  poll.) petiolata e basi acuta vel subcuneata obverse oblonga acuminata remotius a basi (inde ab  $\frac{1}{4}$  long.) callosò-mucronato-serrata membranacea 7-8 poll. longa, costulis 8-10 patule erectis. Spicæ brachiatiæ terminales floribus dissitis suboppositis vel alternis, bractea ovata naviculari apice subcallosa. Stamina? — Wight Herb. n. 878 (ex Miquel). Probabiliter forma Chl. officinalis maxime polymorphi, a quo in descriptione omnino nullum discrimen inveniendum.

11. *ASCARINA* Forst. Char. gen. pl. 59; Juss. Gen. pl. p. 482; Willd. Sp. pl. 4. p. 647; Sprengel Gen. pl. 1. p. 11 n. 36, Syst. veg. vol. 1. p. 19. n. 33; R. Br. Bot. nov. 1821 n. 2190 adnot.; Endl. Gen. pl. p. 264. n. 1818, Enchir. bot. p. 148; Cordemoy in Adans. 3. p. 301; Hook in Journ. of Linn. Soc. 1. p. 127, 129; Seemann in Bonpl. 1861 p. 251, Flora Vitiensis t. 74. p. 258.

Flores dioici, spicati, intra bracteam navicularem solitarii, sessiles, bibracteolati; masculi monandri; anthera crassa cylindræa biloculari tetratheca, loculis contiguis parallelis; feminei nudi, ovarium sistentes subglobosum, subapice cicatrice notatum, stigmate sessili truncato præ-

ditum. Drupa nuda subcarnosa putamine fragili. — Frutices vel arbusculæ in insulis Oceanicis indigenæ, aromaticæ, ramulis glabris ad nodos articulatis, foliis decussatis penninerviis obtuse serratis stipulatis, stipulis cum petiolis in vaginas latas breves amplexicaules connatis.

Genus a *Chlorantho* notis levissimis tantum, floribus dioicis scilicet et antheræ loculis parallelis contiguis diversum, nihilominus maxime naturale, et ulterius conservandum, quum nulla transitum et characterum et patriæ moderans forma sit nota.

1. *A. POLYSTACHYS* (Forst. Char. gen. pl. 59), dioica, foliis glabris late ovatis subcuneatisve basin versus integris in petiolum brevem attenuatis apice obtusis sæpius submarginatis crenato-dentatis, spicis masculis foliis subæquilongis, antheris (evacuatis tantum notis) ab axi subrectangule distantibus vel recurvatis ex ovato cylindricis, connectivo ultra thecas in apiculum recurvatum producto præditis. § In ins. Societatis, Taïti (Forster fruct., Vieillard fr. fl. masc., Morrenhout fr. fl. fem.), ad Walpoo (Bidwill fruct., Barclay fl. masc.). Folia late ovata vel subcuneata 7-10 cm. longa supra medium ubi latissima 2,5-4 lata, petiolis 1-1,5 cm. longis instructa, e medio circ. crenato-serrata. Drupæ maturæ parvulæ 1 mm. circ. longæ subglobosæ exsiccatione corrugatæ. Antheræ magnæ, circa 4 mm. longæ, 1 mm. latæ, lutescentes. (v. s. in h. Hook, DC., Mus. Par., Boiss.)
2. *A. LANCEOLATA* (Hook. fil. in Journ. Linn. soc. 1. p. 127 et 129), dioica, foliis glabris ovato-lanceolatis acuminatis, margine ex apice infra medium obtuse serratis, basin versus in petiolum brevem desinentibus integris, spicis floriferis foliis subæquilongis, antheris primum axi appressis cylindraceis, connectivo thecas superante in apiculum rectum muticum producto præditis. § In Oceanis insulis : in ins. Viti (Seemann pl. Vitiens. 564), Samoa (Powell aprili 1862), Kennady, Raoul Island (Gillivray jul. 1854. Voy. of Herald, bot. n. 982). — Seemann in Bonpland. 1861. p. 259, a mission to Viti p. 442 solum nomen, Flora Vitiensis. p. 258. t. 74. Rami teretes in sicco striati, internodiis brevibus. Folia circ. 5,5-7 cm. longa, in medio ubi latissima 1,5-2 cm. lata, distincte et obtuse serrata, acuminata vel brevi-apiculata, petiolo circa 1 cm. longo instructa. Drupæ et antheræ illis prioris speciei paulo minores et tenuiores. Ab *A. polystachya* quam proxima foliis ovatis acuminatis nunquam obtusis subcuneatis, serratura, antheræ forma et connectivi indole sat superque diversa. (v. s.)
3. *A. RUBRICAULIS*, monoica, foliis petiolatis glabris subtus glaucescentibus ex oblongo lanceolatis acuminatis serratis, spicis et masculis et femineis axillari-bus ad basin subfasciculato-ramosis quam folia multo brevioribus, antheris semper ab axi reclinatis patentibus angustis cylindraceis connectivo recto obtuso præditis. § In Novæ Caledoniæ montibus altitud. 400-600 metr. (Panther, Deplanche, Vieillard n. 1212), in Nova Zelandia (Colenso, hb. Hook.). *A. lurida* Hook. fil. (in Journ. Linn. soc. 1. p. 129 solum nomen). Rami teretes subquadranguli exsiccatione corrugati, ut folia colore rufidulo adpersi, unde nomen. Folia coriacea, petiolo 1 cm. longo instructa, 6-9 cm. longa, 1,5-2 cm. lata. Spicæ floriferæ axillares, e basi laxè fasciculatæ, 1-2 cm. longæ. Antheræ 2 mm. longæ, ab axi subrectangule reclinatæ, quoad formam illis *A. lanceolatæ* similes, sed angustiores. — Species distinctissima spicarum indole a prioribus longe abhorrens; antheræ directione et connectivi forma *A. polystachyam* referens, quoad habitum ad *A. lanceolatam* magis accedens. Monoica, sexu vero ut videtur uno semper prævalente. (v. s. in h. Mus. Paris, Hook., DC., Boiss.)

*Species exclusæ.*

- A. SERRATA* Bl. Enum pl. Jav. I. p. 79. = *Chloranthus brachystachys* Bl  
*A. RUBRA* Poir. in Lam. Encycl. 9. p. 475. = *Morella rubra* Lour. fl. cochinch 2 p. 670, planta affinitatis ignotæ.



III. HEDYOSMUM Sw. prodr. fl. Ind. occ. 847, Flora Ind. occ. p. 59; Willd. sp. pl. IV. p. 476. n. 1703; Schreb. gen. pl. n. 1453; Spreng. gen. pl. 2. p. 696 n. 3512; Spreng. syst. veg. 3. p. 865 n. 3129; R. Br. bot. Mag. n. 2190 nov. 1821 adnot.; Vermischte Schr. ed. Nees. 1. p. 404; Endl. gen. pl. p. 264 n. 1817; Enchirid. p. 148; Griseb. Fl. of the West Ind. isl. p. 172; Agardh Theor. syst. pl. p. 240; Dietr. syn. pl. 5. p. 312. n. 4898; Martius fl. bras. fasc. 11; Cordemoy in Adans. 3. p. 302. — Tafalla Ruiz et Pav. Prodr. fl. peruv. t. 29, Syst. veg. pl. peruv. et chil. p. 270.

Flores dioici vel monoici. *Masc.* : spicati, sessiles, monandri, omnino ebracteati. Anthera bilocularis, tetraheca, loculis parallelis, connectivo thecas superante crasso apiculato vel peltato obtuso prædita. *Fem.* : vel in capitula ramosa multiflora conglobati, vel cymosi. Cymæ vel in foliorum axillis singulæ vel in pseudo-racemos plus minus ramosos dispositæ, 1-5-floræ, bracteate, bracteis sæpius totius cymulæ connatis carnosus. Ovarium inferum, nudum, trigonum, perigonio subtubuloso apici tridentato coronatum. Stigma filiforme, ligulatum vel subclavatum, irregulariter trialatum, alis cum perigonii laciniis alternantibus. Drupæ subcarnosæ, putamine perduro, nudæ vel bracteis persistentibus connatis carnosus circumdatæ. Embryo in albumine copioso minutissimus, fere indivisus, cotyledonibus vix perspicuis. — Arbores et arbustulæ Americæ meridionalis necnon Mexici et Indiæ occidentalis, aromaticæ, resinifluæ, ramulis oppositis ad nodos articulatis per spatium plus minus longum cum ramo ex quo oriuntur connatis, foliis decussatis penninerviis, plerumque serratis plus minus petiolatis, stipulaceis, stipulis apice tantum liberis cum petiolis in vaginam permagnam ochreiformam laxam amplexicaulem connatis.

#### SECTIO I. EUHEDYOSMUM.

Cymulæ femineæ in racemos dispositæ. Internodia florifera elongata; flores ergo discreti pedicellati, bractea unica herbacea fulti. Antheræ connectivo in apiculum herbaceum producto præditæ. — Species in Indiæ occidentalis insulis indigenæ.

1. H. NUTANS (Sw. Fl. Ind. occ. 959), monoicum foliis tenuibus lanceolatis sensim in apicem tenuem productis serrulatis, racemis femineis multifloris ramulorum apices obsidentibus, drupis ovoideis exacte trigonis viridi-flavis, spicis masculis axillaribus racemis femineis approximatis longe et tenuipedunculatis primum ovalibus dein elongatis cylindricis, antheris primum coacervatis ovatis connectivo recto herbaceo lanceolato præditis, dein post anthesin remotis subcylindricis linearibus, connectivo tum basi recto apice tantum incurvo. ♂ In Indiæ occidentalis insulis, in Jamaica (Sw. Balb. Wilson), in Cuba (Wright pl. Cub. n. 1466 et 491). — Griseb. Fl. of the brit. West Ind. Isl. p. 173, Pl. Cub. Wright. p. 176; Cord. in Adans. 3. p. 305. Rami graciles ramulosi. Folia tenuiora, sicca valde fragilia, circa 8 cm. longa, basin versus 15-20 mm., ad apicem 3-5 mm. lata, quoad formam illis Salicis vitellinæ admonentia. Spicæ parvulæ, primum ovales, 6-8 mm. longæ, 4-5 mm. latæ, dein post anthesin 10-16 mm. longæ, in pedunculis 4-5 cm. longis subpendulæ. Drupæ circ. 2 mm. longæ 1,5 mm. latæ, carnosulæ. (v. s.)
2. H. GRISEBACHII, dioicum? ramulis validis, foliis subcoriaceis ex ovato lanceolatis acuminatis margine serrulatis, racemis femineis ramuli apicem obsidentibus, drupis illis speciei præcedentis multo majoribus globosis distincte trigonis, spicis masculis longe cylindricis ad ramorum apices singulis vel paucis axillaribus erectis modice pedunculatis, pedunculis validis, antheris subcylindricis connectivo incurvato coronatis. ♂ In Cuba orientali prope villam Monte Verde dictam (C. Wright n. 1415 masc. et 490 fem.). H. arborescens Griseb. Pl. Wrightianæ p. 176 non Sw. Rami parce ramosi strictiusculi. Fo-

lia illis *H. nutantis* multo majora, haud in apiculum longum tenue producta, circ. 10 cm. longa, 3 cm. lata. Spicæ masculæ 2-4 cm. longæ, cylindricæ, 5-6 mm. latæ, pedunculis strictis 5-6 mm. longis erectis insidentes. Antheræ illis *H. nutantis* crassiores æquilongæ. Drupa matura ut videtur purpurea, 3 mm. longa, 2½ lata. (v. s. in h. Hook., DC.)

SECTIO II. TAFALLA. — Genus *Tafalla Ruiz et Pav. Prodr. fl. peruv. t. 29.*

Cymulæ femineæ 1-5-floræ, internodiis abbreviatis. Flores in glomerulum bracteis carnosissimis arcte appressis convolutum coaliti. Antheræ connectivo brevi crasso truncato peltato instructæ.

\* *Macrocarpa*. — Rami floriferi axillares, internodium unicum plus minus elongatum sistentes apice monodiphyllum. Cymulæ fructiferæ crassæ ut et spicæ masculæ in ramulo fructifero singulæ vel binatæ, intra foliorum axillas sessiles.

3. *H. SCABRUM*, dioicum, foliis petiolatis ovatis vel late ovatis basi subrotundatis, apice in apiculum perobtusum productis, toto margine grosse et obtuse serratis, superne glabris, inferne plus minus præsertim secus nervos secundarios remotiusculos pilosulo-scabris, cymulis femineis ut et spicis masculis in ramuli elongati diphylli apice binis axillaribus sessilibus, cymulis fructiferis ovoideis crassis a bracteis connatis subcarnosis arcte involutis drupas gerentibus ovatas obtuse trigonas, spicis masculis ante anthesin crassis ovatis dein longe cylindraceis sæpius curvatis axi crassissimo. 5 In Peruvia (Pavon, Ruiz in herb. brit. Mus., Mathews), ad Chinchao Peruviae (Rivero in Mus. par.), in Novæ Granadæ monte Quindiu (Humb. et Boupl.) *Tafalla scabra* R. et P. Syst. veg. fl. peruv. et chil. p. 270. *H. hirsutum* Kunth in H. et Bonpl. nov. gen. et sp. pl. 7. p. 126. Cord. in Adans. 3. p. 307. *H. latifolium* Cord. Adans. 3. p. 308. Rami subteretes, vaginis glabriusculis. Folia in petiolo 1 cm. longo, 8-12 cm. longa, in medio 4-6 cm. lata. Cymulæ fructiferæ maturitate 1 cm. longæ, 5 mm. latæ; spicæ masculæ post anthesin 10-11 cm. longæ. Arbuscula 4-orygalis (Pavon). 11. *latifolium* Cordem. nil est quam forma *latifolia* minus pilosa. (v. s.)

β, *Pavonii*, foliis multo angustioribus ex oblongo sublanceolatis acuminatis glaberrimis. In Peruvia (Pavon in h. Boissier). Propria forsitan species transitum inter *H. scabrum* et *H. Mandoni* moderans. Specimina multa inter se simillima in herb. Boiss. asservata vidi. Folia 3-4 cm. lata, 8-12 cm. longa, glaberrima, hic illic sublucida. Cymulæ (floriferæ tantum suppetunt) subglobosæ, 6 mm. circ. longæ. Spicis masculis multo minus quam speciei longis, axi vix incrassato præditis gaudere, idæque propriam speciem esse ulteriori examini relinquendum ex fragmentis, quæ adsunt, videtur. (v. s.)

4. *H. MANDONI*, dioicum, foliis oblongo-lanceolatis acuminatis vel in apiculum brevem productis, basin versus integris in petiolum desinentibus, cæterum profunde crenato-dentatis, dentibus ex angusta basi curvulis apice rotundatis obtusis callosis, nervis secundariis numerosis confertis superne glabris inferne parce pilosulo-hirtellis, ramulis floriferis illis prioris speciei similibus internodium unicum elongatum diphyllum foliis lanceolatis crenato-dentatis nervosis mox deciduis cymulis binis axillaribus sessilibus instructum sistens. Cymulis fructiferis crassis subglobosis a bracteis carnosissimis involutis, drupis oblongis trigonis, spicis masculis ignotis. 3 In Bolivia regionibus temperatis ad Cerro de Silusani provincie Larecaja alt. 2600 metr. (Mandon pl. And. boliv. n. 1095). Rami stricti quadranguli internodiis elongatis. Folia circiter 8 cm. longa, 2,5 cm. lata petiolo 5-7 mm., serraturis 1 mm. circ. longis instructa. Nervii secundarii subtus maxime prominuli, 3-5 mm. inter se distantes, pilosulo-hirtelli. Ramulus floriferus 5-6 cm. longus, folia ejus 3-4 cm. longa, 7-9 mm. lata. Cymulæ fructiferæ circ. 1 cm. longæ. *H. scabro* peraffine tamen serraturæ indole sat superque ab eo recedens. (v. s. in h. Hook., DC., Boiss. et Vindob.)

5. *H. PARVIFOLIUM* (Cordem. mon. in Adans. 3. p. 308), dioicum, foliis pro more parvis ovatis vel rotundato-ovatis subsessilibus obtusissimis coriaceis subulcicis glabris sæpius corrugatis margine remote crenato-dentatis, ramulis fructiferis brevibus axillaribus di- vel sæpius monophyllis cymulam sessilem semper fere singulam rarissime binas ferentibus, cymulis fructiferis e globoso-ovoideis crassis bracteis caruosis connatis arcte involventibus instructis, spicis masculis (secundum Cordemoy) brevibus densis. ♀ In Novæ Granatæ montosis, ad Santa Fé de Bogota loco Peña vieja dicto (Goudot 1844), in provincia Ocaña 8000 ped. alt. (Goudot 1846; Schlim n. 1130). Rami valde ramosi, obsolete tetragoni, internodiis brevibus. Folia 5-7 cm. longa, 2 cm. circ. lata, illa ramuli floriferi 1,5 cm. longa, 1 cm. lata, omnia tenacissima coriacea. Cymulæ fructiferæ circa 7 mm. longæ, 5 mm. latæ. Spicæ masculæ in hb. Musei paris. ubi eas vidit Cord. non suppetunt. Species per distincta, foliis parvis coriaceis ramulisque floriferis brevibus unicam cymulam tantum plerumque ferentibus insignis. (v. s. in h. Hook., Mus. par., Boiss.)
6. *H. CUMBALENSE* (Karst. Fl. Columb. 2. f. 4. t. CLXVIII. p. 129), dioicum, foliis oblongis vel obovato-oblongis coriaceis glabris grosse serratis breviter acuminatis brevi-petiolatis, cymulis fructiferis ovoideis carnis in ramis dense foliatis terminalibus, spicis masculis brevibus cylindraceis crassis erectis ramos foliatis terminantibus. ♀ In Colombia ad silvarum margines regionis frigida montis ignivomi Cumbal (Karsten). Arbor 12 pedum. Ab affini *H. parvifolia* Cord. jam spicis et cymulis ramos dense foliatis terminantibus, foliis minoribus angustioribusque sat superque diversum speciem sistens distinctissimam a nullo peregrinatore ulterius observatam. Folia 3-3,5 cm. longa, circiter 1 cm. lata, minora 2 cm. longa, 1 cm. lata; cymulæ fructiferæ 8 mm. circa longæ, 4-5 mm. latæ, bracteis carnis arctissime involutæ. (v. fragment. ab auctore benevole comm.)
- \*\* *Microcarpa*. Rami floriferi feminei paniculato-ramosi, plus minus expansi, cymulis minoribus permultis onusti. Spicæ masculæ vel in foliorum axillis subsolitariæ, vel laxè paniculatæ, vel ad ramulos axillares verticillato-conglobatæ. — Species extricatu difficillimæ præter *H. pseudo-Andromedam* inter se adeo vicinæ ut primo aspectu omnes paucas tantum species constituere diceres, et ex sola diagnosi vix determinandæ, qua de causa synonymia quoque sat confusa.
7. *H. PSEUDO-ANDROMEDA*, dioicum, foliis coriaceis glaberrimis late ovatis in petiolum attenuatis, apice rotundato-submuticis, margine exsiccatione subrecurvo et medio circa ad apicem serrulatis, racemis femineis terminalibus parvis contractis cymulas pro more magnas paucifloras gestantibus, spicis masculis subsolitariis erectis axillaribus brevibus cylindraceis pedunculatis ad ramulorum apices coacervatis. ♀ In Venezuelæ subalpinis prope coloniam Tovar (Moritz n. 1904, Fendler pl. Venezuel. n. 1117 et 1127). Rami quadranguli, pallidi, dense ramulosi. Folia circ. 5 cm. longa, 2 cm. lata, petiolo 1 cm. longo prædita, maxima 7 cm. longa, 3,5 cm. lata, sicca superne sæpius corrugato-rugulosa pallide viridia, inferne præsertim juniora in ferrugineum vergentia vel subferruginea. Cymulæ fructiferæ circa 5 mm. longæ, spicæ masculæ ante anthesin 7 mm. ad 1 cm. longæ, adultiores ignotæ. Planta omnino sui generis cum nulla alia comparanda, foliorum tenacitate illis *H. parvifolii* admonentia. (v. s. in h. Hook., DC., Boiss., Vindob.)
8. *H. BONPLANDIANUM* (Kunth in H. B. n. g. et sp. pl. VII. p. 165. t. 634 et 655), dioicum? foliis firmis subcoriaceis glabris supra distincte punctulatis oblongis vel elongato-ovatis breviter acuminatis, margine ex apice ad imam basin crebre et minute serrulatis, cymulis femineis laxè paniculatis ramorum apice obsidentibus pro more magnis ovoideis uni-bifloris, drupis bracteas superantibus ovatis obsolete trigonis, spicis masculis post anthesin cylindraceis in ramuli supremorum foliorum axillis subsolitariis erectis. ♀ ? In Nova Granata ad montem Quindiu (Humb. et Bonpl.). — Cord. Adans. 3. p. 301. — *H. glabratum* Cord. Adans. 3. p. 303, non Kunth. Folia 9-14 cm. longa, 3-4 cm.

lata. Cymulæ femineæ ovatæ, circ. 5 mm. longæ, 2-3 mm. latæ; bracteæ connatæ drupis breviores (in icone Kunthii nimis breves). Specimina Humboldtiana in herb. Willd., Mus. Paris., et Schlechtendal. asservata, omnia nomine H. glabrati Kunth errore forsân designata, optime cum Kunthii H. Bonplandiani icone et descriptione quadrant. H. glabratum ex auctoris diagnosi et descriptione aliam plantam esse vix dubium videtur. Quum autem H. glabrati H. B. K. specimina originalia cum diagnosi quadrantia nusquam ullo in herbario invenirentur, quam ad speciem referendum sit haud facile extricandum esse censemus. Si forsân duarum vicinarum insequentium specierum synonymum esset, tamen nova nomina statuenda esse pro incerto veteri putamus. (v. s.)

9. *H. CALLOSO-SERRATUM* (Ersted Pl. centro-americo. in Natur. Foren Vidensk. Meddeleser 1856, p. ?), dioicum, foliis firmis subcoriaceis glabris supra subdistincte punctulatis elongato-ovatis breviter acuminatis margine minute calloso-serrulatis, cymulis femineis laxè paniculatis ramorum apices obsidentibus illis præcedentis speciei minoribus unifloris ovoideis, drupis bracteis pro cymula tribus brevioribus, bracteis apice subulato libero callosis, spicis masculis ignotis. ♀ In America centrali ad latera montis Svasu alt. 9000' (Ersted). Rami validi quadranguli glabri. Folia circa 8-13 cm. longæ, in medio ubi latissima 3-4,5 cm. lata, glaberrima, vix lucida. Cymulæ fructiferæ circ. 2 mm. longæ, 2 mm. latæ, bracteis tribus drupam superantibus infra connatis supra liberis subulatis callosis. A proximo H. Bonplandiano differt foliis latioribus margine minus crebre serrulatis, cymulis unifloris, drupis bracteis arcte circumdantibus brevioribus. (v. specimen siccum ab inventore benetole comm.)
10. *H. DOMBEYANUM*, foliis ex ovato-oblongis brevi-petiolatis subcoriaceis levibus supra subscabris distincte punctulatis, margine obtuse calloso-serratis subacuminatis, cymulis pro more magnis laxè paniculatis ovoideis laxius ac longe bracteatis circiter 3-floris, drupis ovatis trigonis, spicis masculis ignotis. ♀ Specimen adest unicum s. nomine Tafallæ glaucæ cum specimenibus H. integri Cord. in eadem charta herb. Musei Par. asservatum a cl. Dombey in Peruvia ad Cuchero lectum. Hed. glaucum Cord. in Adans 3. p. 303 ex parte, non R. et P. Folia 7-10 cm. longæ, 2-3 cm. lata, petiolo 1-1,5 cm. longè instructa. Cymulæ femineæ circ. 6-8 mm. longæ, circ. 3-4 mm. latæ. A præcedente, inflorescentia magis ramosa, cymulis majoribus plerumque 3-floris, bracteis drupis subæquilongis, foliorum serraturæ indole sat superque diversa; cymularum magnitudine quodammodo speciebus macrocarpis admonens, quoad inflorescentiam multiramam omnino ceteris microcarpis similis. (v. s.)
11. *H. GOUDOTIANUM*, foliis firmis subcoriaceis asperulis supra distincte punctulatis oblongis acuminatis brevipetiolatis, toto ambitu crebre et distinctè serrulatis, cymulis femineis ad ramorum apices contracto-plerumque subpaniculatis subglobosis quam præcedentium specierum minoribus 3-5-floris bracteis drupas æquantibus præditis, spicis masculis cylindraceis ad ramorum apices subpaniculatis erectis. ♀ In Nova Granata ad montem Quindiu (Goudot 1844). H. glabratum Cord. in Adans 3. p. 303 ex parte. Rami validi distinctè quadranguli. Folia in petiolo 1 cm. longo 10-15 cm. longæ, 3-5 cm. lata, hirsutæ scabra plus minus adpersa, asperula. Cymulæ femineæ 2-3,5 mm. longæ, circ. 3 mm. latæ, tri-quinquefloræ. Spicæ masculæ cylindricæ post anthesin 3-4 cm. longæ. Habitu H. Bonplandiano admonens, cymularum fabrica et dispositione valde ab eo abhorrens transitumque ad sequentes species affinitate arcte inter se conjunctas aliquando moderans. (v. s. in h. Mus. paris. ex h. Boiss.)
12. *H. ARBORESCENS* (Sw. Fl. Ind. occ. p. 961), dioicum? foliis elongato-ovatis acuminatis brevipetiolatis glabris lucidis tenerrime tantum punctulatis, margine ex apice infra medium remote crenato-denticulatis denticulis obtusis. cymulis floriferis circa 5-floris ad ramorum apices in racemos parce ramosos expansos dispositis, spicis masculis axillaribus subsolitariis subsessilibus erectis ovatis dein elongatis cylindricis ramorum apices obsidentibus. ♀ In montibus ins. Guadaloupe (Perrottet, Funck et Schlim, 89), in insulis Marti-

nica (Sieber n. 342, Belanger n. 591 in h. Boiss.), Porto Rico (Fajardo n. 185 in h. Boiss.), Jamaica (Wiles in hb. Deless.), St-Vincent (Anderson). — Griseb. Fl. of the brit. West Ind. Islands p. 173; Cord. in Adans. 3. p. 304 ex parte. *H. elegans* Cord. in Adans. 3. p. 305. Rami obsolete tetragoni. Folia in petiolo 1 cm. longo, 9-10 cm. longa, in medio 3-3,5 cm. lata, glabra. Lucida fere laurina, subtus sæpius nitore subsericeo prædita, nervis secundariis remotis minime prominulis. Spicæ masculæ post anthesin cylindraceæ 3-3,5 cm. longæ; cymulæ fructiferæ apice divaricatæ subglobosæ circa 3 mm. longæ. (v. s.)

13. *H. INTEGRUM* (Cord. in Adans. 3. p. 302), dioicum, foliis late elongato-ovatis modice petiolatis subcoriaceis supra glaberrimis tenere punctulatis, adultis ad basin plus minus rotundatis margine revolutis, supra plerumque nitentibus sublucidisve, junioribus margine planis subserrulatis ad basin subito in petiolum anguste alatum desinentibus apice in acumen tenue productis sæpius glaucescentibus, cymulis femineis ad ramulorum apices laxè paniculatis tri-quinquefloris, drupis bracteas circumvolventes superantibus parvis exacte trigonis, spicis masculis ad ramorum apices numerosis in ramulis axillaribus vario modo glomerulatis ante anthesin brevibus ovatis dein ? *♂* *H. glaucum* Cord. in Adans. 3 p. 303 ex parte. *H. bolivianum* Cord. l. c. p. 306 ex parte. *H. racemosum* Cord. l. c. p. 304 ex parte. *Tafalla racemosa* R. et P. syst. veg. fl. peruv. et chil. p. 271 ex parte. In Peruvia (Pavon hb. Delessert, Poeppig n. 1023 et 1426); prope Tarapoto Peruviae orientalis (R. Spruce n. 4311), pr. San Juan del Oro (Weddell n. 4677). Rami læves ramosi. Folia in petiolo 1-1,5 cm. longo, 8-11 cm. longa, in medio 3,5-4 cm. lata. Cymulæ bracteis omnino involutæ circiter 4 mm. longæ; spicæ masculæ confertæ ovatæ circ. 1 cm. longæ, evacuatae ignotæ. (v. s.)

14. *H. RACEMOSUM* (G. Don in Edinb. Rev. 3. p. 432), dioicum? foliis brevipetiolatis subcoriaceis glabris hic illic supra sublucidis subdistincte punctulatis margine serrulatis, quoad formam quam maxime variis vel permagnis fere exacte ovatis vel minoribus elongato-ovatis vel ex ovato-lanceolatis sæpius subobliquis acuminatis, nervis secundariis infra prominulis numerosis, basin et apicem versus vel subabruptis vel sensim angustatis, cymulis femineis expanso-paniculatis ramulorum apices obsidentibus 2-3-floris, bracteis circumvolventibus flores superantibus plerumque longius acuminatis apice sæpius recurvato, drupis ovoideis obsolete trigonis, spicis masculis ad ramorum apices expansos paniculatis parvulis primum ovoideis dein cylindraceis. *♂* In Brasilia (Pohl, in hb. reg. monac.), in Peruvia (Pavon in h. Boissier), in Bolivia provinciis Larecaja et Yungas (Weddell). *Tafalla racemosa* R. et P. Syst. veg. fl. peruv. et chil. p. 271. *H. bolivianum* Cord. in Adansonia 3. p. 306, ex parte *H. integrum* Cord. loc. cit. p. 302 ex parte — *Arbor triorgyalis*, ramis obsolete 4-gonis, his et vaginis glabris. Folia in petiolo 4 cm. longo vel 15 cm. longa, 7 cm. lata vel circ. 9 cm. longæ, 2,5-4 cm. lata. Cymulæ femineæ, vel ut spicæ masculæ, circa ramorum apices paniculatæ. A congeneribus præsertim ab *H. integro* cæterum simillimo quocum sæpius commutantur foliorum nervis secundariis multo crebrioribus, bracteisque cymulas circumvolventibus drupas superantibus acumine recto vel incurvo præditis, inflorescentiæ masculæ denique fabrica sat longe abhorret. (v. s. in h. reg. monac., Mus. Paris., Boiss.)

15. *H. SPRUCEI*, dioicum, foliis elongatis acuminatis sæpius subcuneatis basin versus sensim in petiolum modicum attenuatis glabris superne asperulis tenerrime punctulatis margine e basi integra minute denticulatis, cymulis femineis laxè paniculatis ramulorum apices obsidentibus bi-trifloris, drupis maturis bracteas circumdantibus duplo fere longioribus, apice liberis minutis globosis exacte trigonis, spicis masculis ignotis. *♂* In Peruvia orientali (Gay in h. Mus. Par.); prope Tarapoto (R. Spruce n. 4436). *H. arborescens* Cord. in Adans. 3. p. 304 ex parte. Rami tenues, tetragoni, vaginis scaberrimis obsiti. Folia cic. 14 cm. longa, apicem versus ubi latissima 4-5,5 cm. lata, acuminata, vix in apiculum producta, firma, haud coriacea. Cymulæ fructiferæ circ. 3 mm. longæ. Drupæ in speciminibus visis constanter ad angulos tantum carnosæ, subacidulæ, ad latera parenchymate carnosæ fere destitutæ.

pallidæ, bracteis circumdantibus multo longioribus. Species etsi flores masculi desunt ex notis commemoratis ut videtur sat distincta, druparum indole et bractearum brevitate ab omnibus congeneribus recedens. (v. s.)

16. *H. BRASILIENSE* (Mart. Fl. bras. fasc. 11. p. 3), dioicum, foliis brevi-petiolatis magnis oblongis vel ex ovato-oblongis acuminatis subcoriaceis læribus superne subovatis perdistincte punctulatis, margine exclusa basi distincte serratis, serraturis crebris, dorso convexis rotundatis, cymulis femineis in paniculas contractas vel subcontractas ramulorum apices obsidentes dispositis bifloris, drupis bracteis cingentibus subæquilongis ovoideis trigonis, spicis masculis in ramulis axillaribus circa ramorum apices coadunatis dense-verticillato-conglobatis ante anthesin ovatis. ♂ In Brasilia, in prov. Minas Geraes (Clausen n. 383, Martius, Arkermann, Widgren, Casaretto), in prov. St. Paulo Saint-Hilaire n. 769). *H. Bonplandianum* Mart. Syst. mat. med. Brasil. p. 100, non Kunth. *H. racemosum* Cord. in Adans 3. 304 ex parte. *H. acutifolium* Cord. in Adans. 3. p. 306. Omnium congenerum robustissimum. Rami crassi, quadranguli, ochreis maximis subscabris obsiti. Folia subcoriacea, 10-13 cm. circ. longa, in medio 4-5 cm. lata. Serraturæ 4-5 mm. inter se distantes, dorso convexæ. Spicæ masculæ 6-8 mm. longæ, ovatæ, dein? (v. s.)
17. *H. WEDDELLIANUM* (Cord. in Adans. 3. p. 306 ex parte), dioicum, foliis magnis ex ovato-lanceolatis vel sublanceolatis subtenuibus superne glabris subdistincte punctulatis, plerumque longius petiolatis sensim sensimque in petiolum, apice in apiculum plus minus longum attenuatis, margine basin versus integris, cæterum distincte et remote serratis, serraturis dorso rectis, vix convexis, cymulis femineis circa ramorum apices subcontracto-paniculatis 3-5-floris, bracteis drupis ovatis obtuse trigonis vix æquilongis, spicis masculis circa apices ramorum coadunatis in ramulis axillaribus paniculato-ramosis verticillatis ovatis, post anthesin ex ovato-cylindricis. ♂ In Brasilia, ad Rio de Janeiro (Glaziou in herb. Mart.), in montibus Serra dos Orgoes (Casaretto, Gaudichaud n. 1120). *H. arborescens* Cord. in Adans. 3. p. 304 ex parte. Num revera species propria vel potius varietas præcedentis cui peraffine ulterius examinandum. Constanter minus robustum. Rami obsolete quadranguli glabri. Folia longius (0,5-2 cm.) petiolata, circ. 10-12 cm. longæ, in medio 3-3,5 cm. lata, secundum specimen varia, vel 14 cm. longæ et 5 cm. lata, vel 13 cm. longæ et 3 cm. lata apiculata, substantia subtenui insignia, serraturis 6-10 mm. inter se distantibus rectis vix convexis. Spicæ masculæ 5 mm. dein post anthesin 1 cm. longæ. (v. s. in h. DC., Mart., Mus. Paris.)
18. *H. LECHLERI*, foliis pro more parvis lanceolatis brevi-petiolatis glabris supra subdistincte punctulatis margine e medio remote serrulatis, racemis femineis ramorum apices obsidentibus axillaribus contractis pauci-ramosis, cymulis floriferis sat magnis 2-3-floris, drupis magnis ovatis obtuse trigonis, spicis masculis ad apices ramorum subracemosis ovatis dein cylindricis. ♂ In Peruvia ad Talaxara (Lechler n. 2632, 2660, ad El Gavan (Lechler, 2279). Folia circ. 6-7 cm. longæ, 1,5-2 cm. lata, glabra, e vaginis pilosulis orta. Cymulæ fructiferæ ovatæ, magnæ, 4 mm. latæ, 5 mm. longæ, drupis apice liberis instructæ. Foliis parvulis, cymulis præ cæteris magnis, racemi feminei denique indole a præcedentibus diversa species. Descript. ex fragmentis in epistola missis! (v. s. in h. Hook.)

#### SECTIO III. ARTOCARPOIDES.

Flores feminei numerosi in capitulo ovoideo dense congesti, bracteis omnibus inter se connatis persistentibus maturitate carnosus apice calloso tantum liberis circumdati, perigonio breviter tubuloso, apice subtrilobo, stigmatibus sessilibus elongato linguæformi angusto apice incurvo præditi. Flores masculi illis prioris sectionis omnino similes. Species mexicanæ et peruvianæ, quoad habitum femineæ plantæ ad Artocarpeas accedentes, quoad characteres et plantæ masculæ indolem omnino Chloranthaceæ. Tota inflorescentia fructuum maturitate carnosus drupis trigonis numerosis irregulariter facta. — Duæ tantum adhuc innotuerunt species quarum una mexicana altera peruviana. Aliæ forsitan inter Artocarpeas latentes denuo inveniendæ.

19. *H. GLAUCUM*, dioicum, foliis ex elliptico-lanceolatis serrulatis acuminatis brevipetiolatis glabris supra tenui-punctulatis, spicis masculis ad ramorum apices racemosis ovatis dein ex ovato-cylindræis, floribus femineis in capitula carnosa terminalia et axillaria coadunatis, bracteis drupis subæquilongis infra omnibus in massam carnosam coalitis supra liberis ligulatis apice obtusissimo incrassato-callosis, drupis maturis ovatis trigonis lucidis apice liberis. ♀ In Peruvia (Pavon herb. Boissier). Dehinc non amplius repertum. Tafalla glauca R. et P. Syst. veg. fl. peruv. et chil. p. 270, non *H. glaucum* Cord. Rami validi, subquadranguli, ochreis stipulaceis permagnis in margine truncato apicibus multiramosis obsessis. Folia brevissime petiolata 10-15 cm. longa, in media parte circa 3 cm. lata, hic illic subnitidula. Spicæ masculæ, ovatæ, 1,5-2 cm. longæ, dein post anthesin parum elongatæ, subcylindræ, 2-3 cm. longæ. Capitula fructifera carnosa, terminale 3 cm. longum, circa 2 cm. latum, axillaria in pedunculo 1 cm. longo 1,5 cm. longa, 1 cm. lata, bracteis linearibus dentatis suffulta. (v. in h. Boiss.)
20. *H. ARTOCARPUS*, dioicum, foliis ex ovato-elongatis petiolatis glabris tenuissime punctulatis margine serratis serraturis sat profunde incisus obtusis callosis, spicis masculis circa ramorum apices axillaribus vel subracemosis, longe pedunculatis, primum crassis ovoideis, dein maxime elongatis cylindræis axi crassissima præditis, capitulis femineis et terminalibus et axillaribus apicem ramuli obsidentibus arcte glomeratis carnosis, bracteis drupis ovato-3-gonis longioribus infra omnibus inter esse coalitis apicalibus varie incurvis ligulatis obtusissimis callosis. — In mexicani regni montosis provinciæ Jalapa (Hahn, Herb. de la comm. scientif. du Mex. 1866, n. 458). Rami validi, tetragoni, ochreis stipulaceis illis prioris speciei simillimis præditi. Folia in petiolo 1-1,5 cm. longo, 12-20 cm. longa, in media parte 4-6 cm. lata. Spicæ masculæ in pedunculo 2,5-3 cm. longo, 8-13 cm. longæ, 1 cm. circa latæ, axis incrassatæ præditæ. Capitula fructifera ovoidea, terminalia circa 3 cm. longa, 2 cm. lata, collateralia plerumque bina axillaria fere subglobosa 1 cm. longa. Ab *H. glauco* ceterum proximo foliorum forma serratura et magnitudine, spicarum muscularum indole, bracteis denique drupas maturas superantibus, facillime distinguendum. (v. s. in herb. Mus. Par.)

*Species non satis notæ.*

21. *H. MEXICANUM* (Cord. in Adans. 3. p. 307), foliis magnis firmis oblongo-lanceolatis glabris superne tenere punctulatis grosse serratis, spicis masculis subsolitariis? cylindræis, cymulis femineis ignotis. ♀ In prov. mexicana Vera Cruz alt. 3000' (coll. Galeotti 1840). Hujus plantæ nihil præstat nisi fragmenta ad diagnosin specificam constituendam haud apta. Ab omnibus notis tamen certe diversa species. Folia 11-18 cm. longa, 3-6 cm. lata, brevipetiolata. Spicæ masculæ 3-5 cm. longæ. (v. s. in h. Delessert.)
22. *H. ANGUSTIFOLIUM* (Tafalla angustifolia R. et P. Syst. veg. fl. peruv. et chil. p. 272), foliis sublanceolatis serratis acuminatis articulationibus distantibus amentis strobilibusque 2-3-nis. Frut. 3-orgyalis. In Peruvia.
23. *H. LACINIATUM* (Tafalla laciniata R. et P. Syst. veg. fl. peruv. et chil. p. 272), foliis oblongis acuminatis serratis, amentis et strobilibus 1-2-3-nis. Arbor 4-orgyalis. — In Peruvia,

## Ordo CLXXXIX. GARRYACEÆ (1).

Garryaceæ Lindl. bot. reg. 1834. t. 1686, Veg. Kingd. p. 295, c. is.; Endl. gen. p. 288; J. G. Ag. theor. syst. p. 157 (excl. gen. *Simmondsia*, supra p. 22).

Flores dioici. MASC. Perigonium calycinum, quadripartitum, lobis oblongis, apice per anthesin coherentibus. Petala 0. Stamina 4, basi perigonii inserta, ejus lobis alterna (2); filamenta brevía; antheræ oblongæ, erectæ, basi fixæ, biloculares, rimis 2 longitudinaliter dehiscentes. Rudimentum ovarii nullum. FEM. Perigonium ovario adnatum, apice breviter vel obsolete bilobum. Ovarium uniloculare. Ovula 2, apice loculi funiculis brevibus suspensa, sub stylis posita, anatropa, rhaps interiori epitropa (Ag. f. l. c.). Styli 2, lobis perigonii alterni, erga axim antice et postice siti. Fructus baccatus, plerumque hispidus, stylis persistentibus glabris terminatus. Semen sæpius abortu unicum. Albumen copiosum. Embryo miniquis, axillis aut subaxillis; radícula supera, quam cotyledones longiore. — Arbusculæ vel frutices Americæ sept. temperatæ; foliis stricte oppositis, exstipulatis, penninerviis, integris vel subintegris, persistentibus; racemis simplicibus spiciisve axillaribus; floribus masc. aut fem. intra bracteas oppositas basi connatas ternatis solitariisve. Lignum zonis concentricis orbatum dixit Lindley, quod negat cl. Adr. de Jussieu (Taxonomie, in Dict. sc. nat. p. 58). — Primo adspectu, ex inflorescentia, Cupuliferis aliisque Amentiferis proximæ, sed absentia stipularum, præsentia albuminis copiosi, ovarioque stricte uniloculari discrepant; Juglandaceis et Myricaceis quodammodo propiores, tamen evolutione ovulorum, albumine aliisque char. distant. Pessime (ex ovario infero) post Antidesmeas Euphorbiacearum collocavit Endlicher. Cum Cornaceis affinitatem suspicati sunt cl. Brongniart (Enum. genr. 1843, p. xii), Adr. Juss. (l. c.) et Lindley (Veg. Kingd. ed 3. p. 772), obstantibus tamen et defectu petalorum et ovario uniloculari. Hisce neglectis propiores essent Hamamelideis Grubbiaceisque ubi præterea styli duo distincti satis similes adsunt.

GARRYA Lindl. bot. reg. t. 1686 (præunte Dougl. in sched.); Torr. in Explor. Pacif. railr. 4. p. 136. — Garrya et Fadyenia Endl. gen. suppl. 2. p. 30, suppl. 4. p. 38 (excl. plerisq. sp.). — Characteres ordinis (3).

(1) Auctore Alph. de Candolle.

(2) Nec, ut quidam dixerunt, opposita.

(3) Species Benthamianæ cum Torreyanis ex speciminibus authenticis nondum comparatæ,



Sectio I. EUGARRYA. — Genus *Garrya* Endl. gen. suppl. 2. p. 30.

Styli lineares, elongati, erecto-divergentes. — Bractee abbreviatæ, latæ.

A. Flores in axillis bractearum ternati.

1. *G. ELLIPTICA* (Lindl. bot. reg. t. 1686, ex Dougl. msc.), ramis junioribus pubescentibus, foliis breviter petiolatis ovatis vel ovato-oblongis acutis-marginate undulatis supra glabris subtus pilis intertextis pubescentibus, racemis pendentibus folio multo longioribus, bracteis oppositis late ovatis acutis basi connatis externe pubescentibus, floribus pedicellatis. ♂ In Nova California (Dougl. ! Hartw. p. 1953 in h. Boiss. ! ) prope Monterey (Benth. pl. Hartw. p. 266), in Novo Mexico (Torr. in Explor. Pacif. railr. 4. p. 136), et versus ostium Oregon (Hook. fl. bor. am. 2. p. 143). Frutex. Petioli 2 lin. longi; limbi 1-2 poll. longi, 6-12 lin. lati, pilis plerisque fasciculatis. Bractee 3 lin. longæ latæque, pube sericea brevi. Pedicelli fl. masc. bracteam superantes vel æquantes, medio sæpe longiore. Lobi fl. masc. oblongi,  $1\frac{1}{2}$  lin. longi, dorso apice præsertim pilosi, diu ibi adhærentes. Anthera lobis breviores, filamento minimo multo longiores. Ovaria incano-hispidissima. Styli glabri, divergentes, bractee superantes. (v. s.)
2. *G. FREMONTII* (Torr. in Explor. Pacif. railr. 4. p. 136), foliis late ellipticis utrinque acutis vix mucronatis planis glabris supra nitidis integerrimis, racemis masc. ramosis, bracteis ovatis acuminatis supra medium connatis, inferioribus 3-floris, floribus pedicellatis, fem. .... ♂ In Sacramento super. (Fremont). Folia quam in affini *G. Wrightii* latiora, juniora paulo pilosa. Spicæ 2-4 poll. longæ, pendulæ ? Bractee in cupulam bidentatam connatæ, basi et frequenter ubique in axilla 3-floræ.

B. Flores in axillis bractearum solitarii.

3. *G. WRIGHTII* (Torr. in Expl. Pacif. railr. 4. p. 136), foliis elliptico-oblongis utrinque acutis mucronatis crassis planis opacis margine muriculatis, racemis ramosis, bracteis lanceolatis basi connatis interdum foliaceis et vix connatis, floribus masc. solitariis pedicellatis, fem. sessilibus. ♂ In Novo Mexico ad montem San Francisco et Copper mines (Torr. l. c., Wright coll. 1849. n. 634, coll. 1851-52. n. 1789, ex Torr.). Affinis *G. laurifolia*, cujus tamen folia magis obtusa latiora, margine tenuiora nec ibi muricata. Folia  $1\frac{1}{2}$ -2 poll. longæ, 9-12 lin. lata, valide mucronata.
4. *G. LAURIFOLIA* (Benth. plant. Hartw. p. 14. n. 81 et p. 50. n. 384), foliis elliptico-oblongis integerrimis vel minute dentatis subtus pubescentibus, racemis masc. abbreviatis densis, floribus utriusque sexus in quaque bractea solitariis, sessilibus bractea brevioribus. ♂ In Mexici montibus prope Guanajuato, Anganguio et Zimapan. — Lindl. in Bot. mag. 1840. misc. p. 27. *G. Lindleyana* Hartw. ined. fide Benth. *Fadyenia laurifolia* Endl. gen. suppl. 4. p. 38 (nomen sol.). Arbor vel arbuscula. — Ad hanc speciem num. 589 plantarum Societatis Oregon retulerunt botanici qui specimina mecum comm. (anno 1852), num ex autopsia plantæ Hartw. nescio. In specimine meo rami jun. petioli et folia utrinque sub oculo armato pilis sparsis brevibus donata; petioli 3-7 lin. longi; limbi 1-2 poll. longi, 6-12 lin. lati; racemi feminei petiolo subæquales nutantes. Bractee connatæ, 2 lin. longæ latæque, e basi latæ acutæ, puberulæ. Baccæ 4 lin. longæ.
5. *G. OVATA* (Benth. plant. Hartw. p. 14. n. 80), foliis ovatis integerrimis submucronatis subtus dense pubescentibus, racemis masc. abbreviatis, floribus utriusque sexus in quaque bractea solitariis subsessilibus bractee excedentibus. ♂ In Mexici monte Gigante dicto Guanajuato, alt. 9000 ped. (Benth. l. c. p. 31). *Fadyenia ovata* Endl. gen. suppl. 4. p. 38.
6. *G. MACROPHYLLA* (Benth. plant. Hartw. p. 50, sine num.), foliis amplis late ovato-ellipticis junioribus supra araneosis subtus lanatis adultis subbullatis supra glabris nitidis subtus lanatis glabratissive, racemis brevibus in panicula

densa pauciflora dispositis, floribus (utriusque sexus?) in quaque bractea solitariis sessilibus. — In Mexico prope Zimapan. *Fadyenia macrophylla* Endl. gen. suppl. 4. p. 38 (nomen). Folia  $3\frac{1}{2}$  - 4 poll. longa,  $2-2\frac{1}{2}$  poll. lata. Fructus vix grano piperis major. Flores masc. vix obs.

7. *G. LINDEHEIMERI* (Torr. in Expl. Pacif. railr. 4. p. 136), foliis oblongis obovatisve obtusis submucronatis supra glabriusculis subtus pubescentibus margine integris, spicis folio brevioribus, bracteis lanceolatis ovatisve, floribus masc. breviter pedicellatis. — In Texas occid. et Novo Mexico (Lindh., Wright n. 633). Folia incl. petiolo  $2\frac{1}{2}$  - 3 poll. longa. An flores masc. solitarii ? cl. auctor non dixit. Flores fem. ign.
8. *G. OBLONGA* (Benth. plant. Hartw. p. 51. n. 385), foliis oblongo-lanceolatis vel oblongo-ellipticis mucronatis vel rarius submuticis junioribus subtus vel utrinque laxe tomentosis adultis supra glabris nitidis subtus tomentosis glabratissive, racemis abbreviatis, floribus.... ♂ In Mexici collibus prope Regla. *Fadyenia oblonga* Endl. gen. suppl. 4. p. 38 (nomen). Frutex 6-10-ped. Folia circiter  $1\frac{1}{2}$  po'l. longa, 6 lin. lata. Vix cognita.

Sectio II. *FADYENIA*. — Genus *Fadyenia* Endl. gen. suppl. 2. p. 30, non suppl. 4. p. 39. (Non *Fadyenia* Hook. gen. Filic.)

Styli ovato-oblongi, patenti-reflexi, ovario multo breviores. — Limbus calycis obsoletus. Bractæ elongatæ.

9. *G. FADYENII* (Hook. ic. t. 333), foliis elliptico-oblongis apiculatis integris basi acutis glabris, racemis folio brevioribus laxifloris, floribus in axilla bractæ solitariis subsessilibus, bracteis e basi connata oblongis in fl. fem. longioribus. ♀ In Jamaica montibus cæruleis (Mac Fadyen Hartw. n. 1570, ex Benth.) et Cuba orient. (Wright ! pl. Cub. 1856-57. n. 492). *Fadyenia Hookeri* Griseb. fl. w. Ind. p. 286, plant. Wright. p. 189. Rami juniores adpresse pilosi, cito glabrati. Petioli 3-4 lin. longi; limbi  $1\frac{1}{2}$  - 3 poll. longi, 6-9 lin. lati, supra nitidi. Racemi pubescentes, masc. pendentes, fem. erecti. Bractæ floribus longiores, masc. (ex ic.) circiter 3 lin. longæ, patentes, acutiusculæ, glabræ, parum persistentes. Lobi fl. masc. extus pilosi, diu apice connati. Ovarium ovoideo-compressum, subpubescens. Fructus globosus, glaber, 3-4 lin. latus, subbaccatus. (v. s.)

# INDEX NOMINUM ET SYNONYMORUM

## VOLUMINIS XVI, SECT. 1.

*Obs.* Nomina admissa litteris romanis, synonyma italicis traduntur, nempe in utraque classe :

Ordinalia majoribus,  
Tribualia et subtribualia majusculis,  
Generica vulgaribus,  
Sectionalia minoribus.

	Pagina		Pagina
Achudemia Bl.....	163	Cannabis Tourn.....	30
Acrocarpidium Miq.....	392	CANNABISACEÆ Ser.....	28
Androsyce Wedd.....	171	Carpunya C. DC.....	326
Anemia Nutt.....	237	Carpunya Presl.....	240, 326
Anemiopsis Hook. et Arn.'.....	237	Caulobryon Kl. mss.....	240
Apopiper C. DC.....	366	Ceratiola Mich.....	27
Arianthe Miq.... 240, 251, 326,	336	Chamaebainia Bl.....	218
ARTOCARPEÆ.....	32	Chamaebainia Wight.....	218
Antocarpoides Solms.....	484	Chavica Miq.....	388
Ascarina Bl.....	473	CHLORANTHACEÆ Bl.....	472
Ascarina Forst.....	477	CHLORANTHÆÆ Br.....	472
Australina Gaudich.....	235 <sup>39</sup>	Chloranthus Sw.....	473
AUSTRALINEÆ Wedd.....	235 <sup>39</sup>	Coccobryon C. DC.....	336
		Coccobryon Kl. mss.....	240, 336
Bœhmeria Jacq.....	195	Colema Raf.....	26
BŒHMERIÆÆ Gaudich.....	195	Corema D. Don.....	26
Brachystachys C. DC.....	251	Crantzia Sw.....	14
Brocchia Mauri.....	22	Creodus Lour.....	473
BUXACEÆ Klotzsch.....	7	Cryphæa Ham.....	473
BUXEÆ Endl.....	7, 9	Cubeba Miq.....	240
BUXINEÆ Dumort.....	7	Cypholophus Wedd.....	239 <sup>9</sup>
Buxus Tourn.....	13, 17		
		DAPHNIPHYLLACEÆ Mull. Arg.....	1
Caidbeja Forsk.....	235 <sup>34</sup>	Daphniphyllum Bl.....	1
Callianira Miq.....	240, 243	Debregeasia Gaudich.....	235 <sup>23</sup>
CANNABACEÆ Al. Br.....	28	Dendrocnide Miq.....	77
CANNABINACEÆ Lindl.....	28	Dendrocnide Wedd.....	85
CANNABINEÆ Bl.....	28	Didymodoxa E. Mey.....	235 <sup>61</sup>
CANNABINEÆ E. Mey.....	28	Didymogyne Wedd.....	235 <sup>57</sup>

1. Anemopsis in Hook. et Arn. sed rectius, ex Anemia, Anemiopsis ut scripsit Endlicher.

# 490 INDEX NOMINUM ET SYNONYMORUM.

	Pagina		
<i>Discocarpus</i> Liebm.....	77	<i>Helxine</i> Req.....	
<i>Discocarpus</i> Wedd.....	84	<i>Hemistylis</i> Benth.....	
<i>Distemon</i> Wedd.....	235 <sup>62</sup>	<i>Hesperocnide</i> Torr. et Gr.....	
<i>Droguetia</i> Gaudich.....	235 <sup>57</sup>	<i>Houttuynia</i> Thunb.....	
<i>Dubruetia</i> Gaudich.....	104	<b>HUMILINÆ</b> Dumort.....	
<i>Dugagelia</i> Gaudich.....	392, 471	<i>Humulus</i> L.....	
		<i>Hyrtanandra</i> Miq.....	
<i>Elatostema</i> Endl.....	171, 172	<i>Langeveldia</i> Gaudich.....	
<i>Elatostema</i> Forst.....	171	<i>Laportea</i> Gaudich.....	
<b>ELATOSTEMÆ</b> Gaudich.....	104	<i>Laurea</i> Gaudich.....	
<b>EMPETRACEÆ</b> Lindl.....	24	<i>Lecanocnide</i> Bl.....	
<b>EMPETREÆ</b> Nutt.....	24	<i>Lecanthus</i> Wedd.....	
<b>EMPETRIDEÆ</b> S. F. Gray.....	24	<i>Lepidopelma</i> Kl.....	
<b>EMPETRINÆ</b> Dum.....	24	<i>Leptocnide</i> Gaudich.....	
<b>EMPETRIDIA</b> Raf.....	24	<i>Leucocnide</i> Miq.....	
<i>Empetrum</i> Tourn.....	25	<i>Leucococcus</i> Liebm.....	
<i>Enckea</i> Miq. Kunth.....	240, 243, 326	<i>Leucosyke</i> Zoll. et Mor.....	
<i>Enckea</i> C. DC.....	243	<b>LUPULACEÆ</b> Rasp.....	
<i>Endammia</i> Raf.....	26	<i>Lupulus</i> Bauh.....	
<i>Erasmia</i> Miq.....	392		
<b>EUBOEHERIEÆ</b> Wedd.....	195	<i>Macropiper</i> Miq.....	241
<b>EUBUXACEÆ</b> Baill.....	11	<i>Macrostachys</i> C. DC.....	
<b>EUBUXEÆ</b> Plée.....	11	<i>Maoutia</i> Wedd.....	
<i>Eubuxus</i> Baill.....	17	<b>MAOUTIÆ</b> Wedd.....	
<b>EUFORSKOHLEÆ</b> Wedd.....	235 <sup>54</sup>	<i>Margarocarpus</i> Wedd.....	
<i>Eugarrya</i> A. DC.....	487	<i>Mattuschkia</i> Gmel.....	
<i>Euhedysmum</i> Solms.....	479	<i>Memorialis</i> Ham... 235 <sup>2</sup> ,	
<i>Euleuca</i> Raf.....	26	<i>Micropiper</i> Miq.....	
<i>Euleucum</i> Raf.....	26	<i>Mildea</i> Griseb.....	
<i>Expiper</i> C. DC.....	240	<i>Missiessya</i> Gaudich.....	
		<i>Morella</i> Lour.....	
<i>Fadyenia</i> A. DC.....	488	<i>Morocarpus</i> Siebold.....	
<i>Fadyenia</i> Endl.....	488	<i>Muldera</i> Miq.....	
<i>Fleurya</i> Gaudich.....	70	<i>Myriocarpa</i> Benth.....	
<i>Fleurya</i> Endl.....	70		
<i>Forskohlea</i> L.....	235 <sup>54</sup>	<i>Nanocnide</i> Bl.....	
<b>FORSKOHLEÆ</b> Wedd.....	235 <sup>54</sup>	<i>Nematanthera</i> C. CD.....	
<i>Freirea</i> Gaudich.....	235 <sup>51</sup>	<i>Nematanthera</i> Miq.....	
		<i>Neraudia</i> Gaudich.....	
<i>Garrya</i> Lindl.....	486	<i>Nigrina</i> Thunb.....	
<b>GARRYACEÆ</b> .....	486	<i>Nothocnide</i> Bl.....	
<i>Gesnouinia</i> Gaudich.....	235 <sup>49</sup>		
<i>Gesnouiia</i> Endl.....	235 <sup>49</sup>	<i>Oakesia</i> Tuck.....	
<i>Girardinia</i> Gaudich.....	100	<i>Obetia</i> Gaudich.....	
<i>Girardinia</i> Endl.....	100	<i>Oreocnide</i> Miq.....	
<i>Gonostegia</i> Turcz.....	235 <sup>2</sup>	<i>Ottonia</i> Spreng.....	240
<i>Goughia</i> Wight.....	1		
<i>Gymnotheca</i> Decsne.....	237	<i>Pachysandra</i> Mich.....	
<i>Gyandra</i> Wall.....	1	<i>Parietaria</i> Tourn.....	
<i>Gyrotænia</i> Griseb.....	99	<b>PARIETARIEÆ</b> Wedd.....	
		<i>Pellionia</i> Gaudich.....	
<i>Meckeria</i> Kunth.....	240, 331	<i>Peltobryon</i> Miq.....	240
<i>Hedysmum</i> Sw.....	479		

# INDEX NOMINUM ET SYNONYMORUM. 491

Pagina	Pagina
Reichb..... 473	Sclepsion Wedd..... 78
Ruiz et Pav..... 392	Serpentaria Pluk..... 239
RIEÆ Miq..... 391	Serronia Gaudich..... 251
Wedd..... 235 <sup>38</sup>	Simmondsia Nutt..... 22
on Miq..... 392	SIMMONDSIÆ Mûll. Arg..... 22
ndl..... 104, 235 <sup>63</sup>	Soleirolia Gaudich..... 235 <sup>41</sup>
..... 240	Spathium Lour..... 239
E.E. Kunth..... 235	Spharostachys Miq..... 240
E. Miq..... 240	Splitgerbera Miq..... 195
EE DC..... 235	Stachyocnide Bl..... 219
C. DC. (1866)..... 391	Steffensia C. DC..... 251
Wedd..... 235 <sup>10</sup>	Steffensia Kunth..... 240, 251
ermum Zipp. et Miq... 235 <sup>15</sup>	Stropha Nor..... 473
Lour..... 238	Styloceras Adr. Juss..... 9
phe C. DC..... 331	STYLOCEREÆ Baill..... 7, 9
ophe Miq..... 240, 331	STYLOCEREÆ Mûll. Arg..... 9
Gaudich..... 219	Symbryon Griseb..... 471
E.E. Wedd..... 104	
omm..... 191	Tafalla Ruiz et Pav..... 479, 480
pr..... 171	Tafalla Solms..... 480
	Thaumuria Gaudich..... 235 <sup>44</sup>
opsis Miq..... 240	Tildenia Miq..... 392
Gaudich..... 235 <sup>43</sup>	Tillocnide Bl..... 207
Endl..... 235 <sup>43</sup>	Touchardia Gaudich..... 235 <sup>13</sup>
	Tricera Baill..... 14
ria Cordemoy..... 473	Tricera Sw..... 14
a Gardn..... 473	Tricercandra A. Gray..... 473
mys Gaudich..... 235 <sup>41</sup>	Tropis sp. Willd..... 9
ILAMYDEÆ Wedd.... 235 <sup>9</sup>	Tuckermannia Kl..... 26
a Lindl..... 11	
Wedd..... 82	ULMACEÆ..... 32
ACEÆ Lindl..... 237	Urera Gaudich..... 88
E.E. C. DC..... 237	UREREÆ Gaudich..... 39
EE Rich..... 235, 237	Urtica Tourn..... 39, 235 <sup>63</sup>
sis Turcz..... 239	URTICACEÆ Endl..... 32
L..... 238	URTICÆ Juss..... 32, 235
us Wedd..... 98	URTICEÆ Br..... 32
Kuath..... 240, 251, 326	
hos C. DC..... 241	Verhuellia Miq..... 391
hos Griff..... 241	Villebrunea Gaudich..... 235 <sup>20</sup>
skia Endl..... 70	VILLEBRUNEÆ Wedd..... 235 <sup>16</sup>
Gaudich..... 191	
Raf..... 77, 78	Zippelia Bl..... 240, 251

## ERRATA.

Pag. 39, lin. 1. URUREÆ.... *lege* UREREÆ.

Pag. 219, lin. 11. Lencoccus.... *lege* Leucococcus.

Pag. 235<sup>a</sup>, PHENAX STOCKSH Wight = Phenax pentandra.... *lege* = Memorialis pentandra.

PHENAX TERNATA Benn. = Phenax ternata.... *lege* = Memorialis ternata.

PHENAX TOMENTOSA Wight = Phenax hirta.... *lege* = Memorialis hirta.

Pag. 252. PIPER EUCALYPTOPHYLLUM.... *lege* P. EUCALYPTIPHYLLUM.

Pag. 269. PIPER DUMOSUM (ad finem) ex Rudge l. c.... *lege* (v. s.)

Pag. 278. PIPER PSEUDOMOLLICOMA.... *lege* P. PSEUDOMOLLICOMUM.

Pag. 308. PIPER POTOMORPHE.... *lege* P. POTHOMORPHE.

Pag. 361, lin. 7 ante finem. Chavica Sinensis Champ. in Kew journ. of bot. v. 1 p. 106.... *lege* vol. 6 p. 116.

Pag. 362, lin. 22, PIPER INDICUM Miq.... *lege* C. DC.

— lin. 28, ante non Rich. *lege* Miq.! syst. p. 315.

Pag. 379. PIPER TILLEFOLIUM Ham. Species delenda. Est laurea tilizefolia Gaudich. in præ. vol. 6 p. 116.

Pag. 427, lin. 13, P. MAGNOLLEFOLIA.... *adde* Dietr







Stanford University Libraries



3 6105 011 548 364

582  
C219

Genus  
Ficus depurata

Des't

NAME

22 Sept '08

Aug 30 '08

DATE

Candolle,  
Prodromus 872

